Published in final edited form as:

Food Chem. 2018 June 30; 252: 258–264. doi:10.1016/j.foodchem.2018.01.120.

Essential micronutrient and toxic trace element concentrations in gluten containing and gluten-free foods

Tracy Punshon^{1,*} and Brian P. Jackson²

¹Department of Biological Sciences, Class of 1978 Life Sciences Center, 78 College Street, Dartmouth College, Hanover, New Hampshire, 03755 USA

²Department of Earth Sciences, 19 Fayerweather Hill Road, Dartmouth College, Hanover, New Hampshire, 03755 USA

Abstract

For individuals following a gluten-free (GF) diet, rice is commonly the major grain. People following a GF diet have a higher arsenic burden than the general population. We conducted a multielemental market basket study of GF and gluten containing ingredients and prepared foods (Mn, Fe, Ni, Cu, Zn, Cr, Co, Se, Cd, Sb, Pb, total As, As species, total Hg and methylmercury). Foods containing rice were significantly higher in As, Hg and Pb and lower in Se, Fe, Cu and Zn. Wheat-based foods were higher in Cd. Mercury concentrations were low (<3.5 ng/g); speciation was predominantly methylmercury. Arsenic and mercury in rice were correlated. GF foods contained significantly more As and Hg. Eating a wide variety of GF grains may reduce contaminant exposure and increase micronutrient status compared to a rice-based GF diet.

Keywords

Rice; arsenic; mercury; gluten-free; speciation; market basket

1.0 INTRODUCTION

Grains have long been a staple of the human diet, being an excellent source of carbohydrate. The most widely consumed grains – rice, maize and wheat – provide 60% of the world's food energy intake (FAO, 1995). Like other plants, these staples can accumulate non-essential and potentially toxic trace elements from natural or human input to the soil. It is well-established that the concentration of arsenic in rice grain is higher than in the grains of other cereals such as wheat, oats and barley (Cubadda, Jackson, Cottingham, Van Horne, & Kurzius-Spencer, 2017). This is due in equal parts to the practice of rice paddy farming and the physiological requirement of rice for silicon, for which arsenite is a chemical analog.

*Corresponding author.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors declare no conflicts of interest.

Flooding rice paddies promotes the dissolution of iron oxide minerals that otherwise sequester arsenic; the released arsenic is efficiently taken up rice plants via silica transport systems and translocated to rice grain (Punshon, Jackson, Meharg, Warczack, Scheckel, & Guerinot, 2017). Compared to environmental factors, it is the cultivar of rice which has the greatest influence on the amount of arsenic that accumulates in the rice grain (Norton, Pinson, Alexander, McKay, Hansen, Duan, et al., 2012). Grain arsenic concentration varies widely between different cultivars, which has given rise to breeding efforts to develop cultivars with low arsenic accumulation characteristics.

Inorganic arsenic, the most acutely toxic form (Le, Ma, Lu, Cullen, Aposhian, & Zheng, 2000) and consequently the form regulated in food and water, tends to accumulate in the nutrient-rich outer layers of the rice grain (tegumen, pericarp and aleurone layers) known as the bran. Brown rice, therefore, usually contains higher concentrations of inorganic arsenic than white, in which the bran layers have been removed or 'polished' (Meng, Feng, Qiu, Anderson, Wang, & Zhao, 2014). Food products made primarily from rice bran can be particularly high in inorganic arsenic (Sun, Williams, Carey, Zhu, Deacon, Raab, et al., 2008).

The EU has introduced regulations specifically for white rice of 0.2 µg/g for inorganic arsenic and rice products aimed at infants have a lower limit of 0.1 µg/g (EU, 2015). Because of their lower acute toxicity to humans, the organic forms of arsenic are not targeted for regulation. The US FDA has also set an action limit of 0.1 µg/g for inorganic arsenic in infant rice cereals (U. FDA, 2016). The regulatory focus on foods aimed at infants and young children reflects their heightened developmental vulnerability and higher exposure to arsenic (ingestion normalized to body weight) compared to adults (European Food Safety Authority, 2009). Concerns about arsenic exposure from rice-based diets have been raised for Asian populations and for individuals following a gluten-free (GF) diet where processed food products (such as bread and pasta) use rice as a staple grain instead of wheat (Meharg, Norton, Deacon, Williams, Adomako, Price, et al., 2013). A recent statistical analysis of the National Health and Nutritional Examination Study (NHANES) database (2009–2014) of urine and blood concentrations measured in the US population found that urinary arsenic was significantly higher in individuals that self-reported as being on a GF diet (n=74) compared with the total individuals sampled (n=7471) (Bulka, Davis, Karagas, Ahsan, & Argos, 2017).

Other potentially toxic elements have also been found in rice. Cadmium can reach levels of concern (Meharg, et al., 2013) when rice is grown aerobically. In China, mercury contamination of rice grown on contaminated soils can contribute to human mercury exposure (Rothenberg, Windham-Myers, & Creswell, 2014). Arsenic and mercury are among the top three contaminants of concern on the Agency for Toxic Substances and Disease Registry's National Priority list; together with cadmium (at number 7) they are taken up by plants, and are developmental neurotoxins (Grandjean & Herz, 2015).

Food, in general, is the main source of cadmium exposure for non-smokers. In 2012 the European Food Safety Authority reviewed cadmium exposure in the European population and concluded that children and adults at the 95th percentile of exposure could be consuming

cadmium levels in excess of health based guidelines (0.001 mg/kg/day) (European Food Safety Authority, 2009). Subsequently maximum allowable limits for cadmium in certain foods were adjusted down to reduce exposure to the general public. Wheat can be a major source of cadmium to diet and, like arsenic uptake in rice, cadmium uptake in wheat is dependent on the variety of wheat grown (Harris & Taylor, 2004).

There has been a recent focus on mercury uptake by rice and subsequent human exposure; this is primarily an issue with rice grown on mercury polluted soils, and again growing rice anaerobically exacerbates the problem, because sub-oxic conditions promote methylation of mercury to methylmercury, the most toxic form. Hence, when fish consumption is low, rice products can be the major source of methylmercury (MeHg) to the diet (Rothenberg, Yu, Liu, Biasini, Hong, Jiang, et al., 2016). A recent market basket analysis of infant rice products found that rice cereals and rice teething biscuits were on average 61 and 92 times higher in methylmercury respectively than cereals made with wheat or oats (Rothenberg, Jackson, Carly McCalla, Donohue, & Emmons, 2017). Provisional tolerable weekly intake (PTWI) limits for methylmercury exposure from food were established in the EU by the Joint Expert Committee on Food Additives (JECFA) of 1.3 µg mercury/kg body weight (0.18 µg per day) (JECFA, 2007). The US EPA oral reference dose (RfD) for daily intake of methylmercury is 0.1 µg/kg body weight (USEPA, 2001). Analysis of data from NHANES, mentioned above, also found higher blood mercury (inorganic and methylmercury) concentrations in people following a GF diet (Bulka, Davis, Karagas, Ahsan, & Argos, 2017). Increased rice consumption is one possible explanation.

Numerous market basket studies have measured the concentration and speciation of arsenic in whole grain rice, food containing rice, and ingredients processed from rice. This has provided the basis for the food-arsenic regulation mentioned above. Databases of arsenic concentrations and speciation in many rice-containing food products are available (FDA, 2016) and it is clear that some iterations of a GF diet could potentially provide a higher arsenic exposure (Munera-Picazo, Ramirez-Gandolfo, Burlo, & Carbonell-Barrachina, 2014) than is considered safe in terms of increases in life-time cancer risk. Elevated cadmium, lead and nickel concentrations were also found (Orecchio, Amorello, Raso, Barreca, Lino, & Di Gaudio, 2014). However, side-by-side comparisons of the concentration of arsenic and other relevant elements in GF and non-GF products – needed to inform a dietary arsenic exposure risk assessment – have not been extensively carried out. In addition to the 1% of the US population who have celiac disease based on seroprevalence, GF diets are also necessary for those with non-celiac gluten sensitivity and wheat allergies. Despite concerns about the nutritional adequacy of the GF diet (Theethira & Dennis, 2015), it has gained a reputation as being more beneficial for health and weight maintenance than diets containing gluten.

Surveys conducted in 2013 indicated that almost 25% of the US population had adopted a GF diet (DiGiacomo, Tennyson, Green, & Demmer, 2013). In line with this, the GF food retail market more than doubled in value between 2011 and 2016 (Group, 2013), a trend which is expected to continue.

In this study, we measured the concentration of arsenic, mercury, lead, cadmium and other elements including micronutrients manganese, iron, zinc, copper and selenium, in locally-

available rice grains and rice-containing products and compared them to metal concentrations measured in equivalent wheat-based products. Arsenic and mercury speciation was determined in rice grains and rice-containing products, because total concentrations of these elements were high enough in these products to allow reliable speciation analysis and exceed proposed regulatory limits. We examined the elemental content of other readily available GF flours and grains such as amaranth, oat, sorghum, almond and coconut. We show that, consistent with observations of mercury in blood of GF diet followers (Raehsler, Choung, Marietta, & Murray, 2017; Vici, Belli, Biondi, & Polzonetti, 2016), rice-containing products appear to be a source of methylmercury at very low concentrations (up to 3.5 ng/g) in comparison with products made from wheat or other grains.

2.0 METHODS AND MATERIALS

2.1 Sample procurement and preparation

Sixty-seven food products were purchased from local food stores in Hanover and West Lebanon (New Hampshire, USA) and from on-line suppliers from late 2016 through mid 2017. The choice of food products was intended to compare readily available GF cooking ingredients and staple prepared food products and their gluten-containing counterparts. Products were grouped into flours, whole grain rice, and prepared foods. Designation of the products as GF, organic or enriched were made on the basis of information on the packaging. We purchased 19 different rice grains, three rice flours, and 19 non-rice flours (including corn, corn masa, whole wheat, all-purpose wheat, sprouted wheat, spelt, millet, oat, buckwheat, chickpea, coconut, and almond, amaranth) as well as other popular grains (black chia seed and tricolor quinoa). The GF and non-GF prepared food products consisted of gluten free pastas, breads, cakes (Table 1). Most GF products contained rice as one of top 3 listed ingredients.

Whole grains were not rinsed or cooked prior to being prepared for analysis, and were ground in a clean, dry coffee grinder. Flour and powder samples were acid digested without further preparation. Moist prepared food samples were freeze-dried and homogenized prior to analysis. The final results were corrected so they reflected the wet weight concentration. Foods were analyzed for Mn, Fe, Ni, Cu, Zn, Cr, Co, As, Se, Cd, Sb, Hg, and Pb (Table 1). Whole grain rice and rice flour were subject to speciation analysis for arsenic (inorganic As, dimethylarsinic acid (DMA) and monomethylarsonic acid (MMA)) and mercury (methylmercury and total mercury).

2.2 Sample digestion for total metals analysis

Samples were acid digested by closed vessel microwave assisted digestion; 5 ml of 9:1 optima HNO₃:HCl was added to 0.25 g of sample and the digestion temperature was ramped to 220°C over 15 minutes and held for a further 20 minutes. Following digestion, the sample was diluted to 50 ml with deionized water.

For arsenic speciation, 0.25 g of sample was heated to 100°C in 2% HNO₃ following our previous methods (Jackson, 2015). Methyl mercury analysis was performed by species

specific isotope dilution, acid extraction, ethylation, purge and trap concentration, gas chromatographic separation and ICP-MS analysis following our previous methods (Taylor, Jackson, & Chen, 2008).

2.3 Elemental analysis

Samples digested for total elemental analysis were analyzed by collision cell inductively coupled plasma mass spectrometry (ICP-MS) (Agilent, 7900x) operated in helium gas mode for all elements and no gas mode for lead and mercury. For digestions and extractions, one sample per batch (where a batch is denoted as 20 samples) was digested/extracted in duplicate and spiked with analyte and taken through the digestion/extraction process. Additionally, one spiked blank (fortified blank), reagent blank and standard reference material per batch were taken through the digestion/extraction process. For analysis, additional quality control involved duplicate analysis of digested/extracted samples and analyte spikes of digested/extracted samples. For total analysis by ICP-MS, the instrument was calibrated using NIST-traceable multi-element standards prepared on the day of analysis. The calibration was verified using second source NIST-traceable standards after the calibration and repeatedly every 10 samples.

Method detection limits and standard reference material recoveries (%) are given for all elements in Table 1. We also analyzed mercury using the direct mercury analyzer (Milestone, Shelton, CT). This data was used for total mercury and % recovery of the NIST 1568b was $86 (\pm 3) \%$ ($n=4$).

2.4 Arsenic speciation

Arsenic species were analyzed by anion exchange chromatography coupled to reaction cell ICP-MS (Agilent 8800, Santa Clara, CA) as described previously (Jackson, 2015). The system was calibrated with arsenic species, arsenite, dimethylarsinic acid (DMA), monomethylarsonic (MMA) and arsenate and NIST 1568b which is certified for inorganic arsenic (sum of arsenite (3+) and arsenate (5+)), DMA and MMA. Recovery of Inorganic arsenic was $108 (\pm 23) \%$ and organic arsenic (DMA + MMA) was $93 (\pm 6) \%$. Method detection limits for arsenic species were 3 ng/g.

2.5 Mercury speciation

Methylmercury concentrations were determined using an automated methyl mercury analyzer (MERX-M, Brooks Rand Instruments, Seattle, WA) coupled to ICP-MS and quantified by isotope dilution as described previously. Duplicates of SRMs NIST 1566b Oyster Tissue, certified at 0.0132 $\mu\text{g/g}$ MeHg, and NIST 2976 mussel tissue, certified at 0.0272 $\mu\text{g/g}$ MeHg, were analyzed and recovery $\pm \%$ relative difference was $118 \pm 2.6\%$ and $102 \pm 7.1\%$. Detection limits for methylmercury were 0.12 ng/g.

2.6 Statistical Analysis

We conducted one-way analysis of variance (ANOVA) or one-sample t-tests as appropriate on data for the metal concentrations in food products, using $\log_{10}$ transformation to normalize metal concentration data, which tended to be left skewed. For data measurements below the limit of detection, we assigned a value of half of the instrument detection limit.

Means comparisons were conducted using Tukey's statistical tests. We used non-parametric Spearman's statistical test to examine correlations between total arsenic concentrations (all species combined), inorganic arsenic (arsenite + arsenate), organic arsenic (DMA and MMA) and the measured total mercury and methylmercury concentrations.

3.0 RESULTS AND DISCUSSION

Table 1 summarizes the total elemental concentrations and results of one-sample t-tests and ANOVA statistical analysis on the food and product groups analyzed in this study.

3.1 Arsenic

The total arsenic concentration in whole grain rice, rice flour and processed foods containing rice was significantly higher than non-rice flours and processed foods based on other grains [one-sample $t(39.0) = 10.71$, $p < 0.0001$]. As shown in numerous studies, rice and rice products contain elevated arsenic concentrations compared to non-rice foods (Carbonell-Barrachina, Wu, Ramirez-Gandolfo, Norton, Burlo, Deacon, et al., 2012). Our speciation analysis of arsenic found that inorganic arsenic made up on average 63 ($\pm$ SD 15) % of the total arsenic across all rice-containing foods sampled in this study (N=38).

Between the three rice grain types (brown, white and enriched white) total arsenic concentrations were significantly different ($p=0.0335$; $F = 4.2320$), with brown rice and enriched white rice having a higher concentration of arsenic than non-enriched white. The inorganic arsenic concentration ($p=0.0029$, $F = 8.8611$) and the percent of inorganic arsenic ($p=0.0029$, $F=8.8380$) was also significantly higher in brown rice than enriched white or non-enriched white ($p=0.01$; $F = 6.0126$). Inorganic arsenic accumulation in bran layers of the rice grain (Carey, Norton, Deacon, Scheckel, Lombi, Punshon, et al., 2011) is a likely explanation for the higher inorganic arsenic concentrations measured in brown rice. However, the reason why enriched white rice grain contains higher organic arsenic concentrations than brown or non-enriched white rice is not clear, and may be an artifact the small sample size of each grain type (where certain suppliers had more than one grain in each product type).

We found that only one whole grain rice sample measured below detection limits for arsenic (2 ng/g), an upland rice grown under non-flooded conditions. This cultivation technique is an effective method for limiting arsenic uptake in rice because arsenic remains bound to the soil and not available for plant uptake (Hu, Li, Yuan, Ouyang, Zhou, Huang, et al., 2013). However, this rice sample also contained the highest concentration of cadmium of all the grains and food products tested (0.12 μ g/g, compared to a mean of 0.02 μ g/g cadmium for whole grain rice in this study), which again has been shown to be a consequence of growing rice aerobically. Under aerobic soil conditions cadmium is not sequestered as the insoluble sulfide and is more available for uptake (Tai, Li, McBride, & Yang, 2017).

The average inorganic arsenic content of rice and rice products measured in this study was 93.9 ng/g; with and an individual serving of either grain or pasta estimated to be 50 g, a serving could contain an average of 4.6 μ g of inorganic As. An individual following a GF

diet eating 3–4 servings of rice and rice products could conceivably consume 14 – 18 µg of inorganic arsenic per day.

3.2 Mercury

Mercury concentrations were very low, < 4 ng/g for all foods tested. This concentration level is much lower than fish and other seafood, which is generally the major source of mercury to diet. The US FDA reported mean levels of mercury in fish that ranged from 45 ng/g for anchovies to > 1000 ng/g for tuna, halibut and other high trophic level fish (FDA, 2017). While much lower than fish, the mercury concentrations in the foods we tested were of interest because of recent biomonitoring studies and data analysis of the NHANES database, which showed that self-reporting rice eaters and people following a gluten free diet had higher levels of blood mercury than non-rice eaters or those not on a GF diet (Bulka, Davis, Karagas, Ahsan, & Argos, 2017; Raehsler, Choung, Marietta, & Murray, 2017).

When data for all products was grouped by the constituent grain type, food containing rice had significantly higher mercury concentrations than those based on wheat or other grains [one-sample $t(56.3) = 6.49, p < 0.0001$] (Table 1). Within the whole grain rice category, mercury concentrations did not significantly differ between white, brown or enriched white rice, suggesting that mercury does not differentially accumulate in the bran layers in the way that arsenic does.

Methylmercury concentration of rice grains was highly correlated with the independent measure of total mercury and suggested that essentially all the mercury in these rice grains was methylated (Figure 2). Studies of rice grown in Hg-contaminated soils around artisanal gold mining operations have found high levels of total mercury and methylmercury in rice grains (Rothenberg, Windham-Myers, & Creswell, 2014). These studies have mostly focused on rice grown in contaminated soils; however, more recent studies have shown that even in non-contaminated soils, rice is a source of mercury to diet (Hong, Yu, Liu, Cheng, & Rothenberg, 2016). Like arsenic, the presence of methylmercury in rice is due to the practice of growing rice in flooded soils, because methylmercury formation is microbially driven and favored in slightly reducing environments where iron-reducing and sulfate-reducing bacteria predominate (Rothenberg & Feng, 2012). Although levels of mercury and methylmercury in rice are low, rice can be a significant source of these toxicants to the human diet because rice is a staple food (Rothenberg, Windham-Myers, & Creswell, 2014). Considerable concern has been raised about inorganic arsenic levels in infant rice cereal, because very young children are exposed to higher body burdens of inorganic arsenic and early life is a particularly sensitive window of exposure. Similar arguments could be raised for methylmercury exposure through rice cereal during developmentally-sensitive life stages (Hong, Yu, Liu, Cheng, & Rothenberg, 2016), and perhaps additional concern is warranted for combined metalloid exposure from rice consumption. Using an average MeHg concentration of 2.5 ng/g, for an adult (60 kg) on a GF-diet consuming 3–4 rice-based servings (50g) per day, an individual's MeHg exposure would be 0.006 – 0.008 µg/kg body weight. This is over an order of a magnitude lower than JECFA and EPA recommended daily limits and therefore methylmercury exposure from uncontaminated rice to adults is unlikely to pose any risk.

As with arsenic, the whole grain rice sample grown aerobically had undetectable mercury (below 0.5 ng/g), which is unexpected because chemically, mercury availability is believed to be similar to cadmium in its increased availability for uptake when not sequestered as the insoluble sulfide. This suggests that it is actually the methylation of mercury that may be of most importance in plant uptake.

We also found that, among the whole grain rice products tested (N=19), total arsenic and mercury concentrations were strongly correlated (Table 2). Inorganic and organic forms of arsenic were equally correlated with total mercury, whereas the correlation between organic arsenic and methylmercury was marginally stronger than inorganic arsenic and methylmercury. A strong correlation between arsenic and methylmercury concentrations of rice based infant products was also recently found in a recent market basket analysis (Rothenberg, Jackson, Carly McCalla, Donohue, & Emmons, 2017).

3.3 Other contaminant elements

Cadmium concentrations were significantly higher in wheat-based prepared foods compared to rice or other grains and between gluten containing and GF food groupings. Increased uptake of cadmium in wheat compared to other grains is well recognized, and wheat is a main source of cadmium to the human diet (Arduini, Masoni, Mariotti, Pampana, & Ercoli, 2014). As noted earlier, rice can also take up cadmium, especially when grown aerobically, and, indeed, the highest cadmium containing food we measured was an aerobically grown rice which had a concentration of 0.12 µg/g. Due to relatively high uptake of cadmium and high consumption levels the European Food Safety Authority identified cereals and cereal products as the highest source of cadmium to the diet (EFSA, 2009).

Concentrations of lead in all rice containing foods were significantly higher than those based on wheat and, when the spelt flour was omitted from the 'other' grain category, rice was significantly higher than either of the other two grain types. Also, within the food products category, Rice products were higher in Pb than the wheat-based products.

3.4 Nutrient elements

Rice and rice based products were significantly lower in the nutrients iron, nickel, copper, zinc and cobalt compared to wheat flours and food products and the other (rice-free) GF flours. The superior nutritional quality of pseudocereals (e.g. quinoa, buckwheat and amaranth) are well known (Alvarez-Jubete, Arendt, & Gallagher, 2010), having a higher protein content wheat, and constituting a good source of calcium, magnesium and iron. Despite seven of the rice grain products being fortified with ferric orthophosphate, they were comparatively lower than wheat flours and foods containing other rice-free GF flours. This may be due to the fact that only whole grain rice is fortified, whereas ingredients processed from rice are not. We observed that iron concentrations in enriched rice were highly variable in comparison with brown rice. Of the seven enriched white rice grain products analyzed, only three products had iron concentrations that exceeded 12 µg/g – the highest iron concentration measured in non-enriched white rice. One suggestion for this variation would be the specific method use to apply micronutrients to rice grain. Brown rice is known to have a higher nutrient content than white rice due to the intact bran layers, and we found that zinc,

iron and manganese concentrations were significantly higher in brown rice than in white rice, with brown rice also having a higher average copper concentration.

Wheat was significantly higher in selenium than rice or the other GF flours. Wheat grown in the US readily accumulates selenium from Se-rich US soils (Reilly, 2006), whereas rice does not take up comparable concentrations of this micronutrient. Copper was significantly higher in non-rice flours and wheat pasta than rice pasta or grain. In fact, concentrations of Fe, Cr, Co, Ni, Cu and Zn in specialty flours were all significantly higher than rice. When the specialty flours were combined with the rice data to create a GF food category for comparison with the wheat-based non-GF foods then only Se remained significantly higher in the NGF foods; in other words, the nutrient deficiencies of rice were negated by pooling the data with other GF grains, which were generally higher in nutrient content.

A multi-elemental market basket study of 27 GF foods purchased in and around Palermo City (Italy) found that the nutritional quality of the foods, in comparison with recommended dietary intake levels for the essential micronutrient elements, was relatively poor (Orecchio, Amorello, Raso, Barreca, Lino, & Di Gaudio, 2014).

4.0 CONCLUSIONS

Rice and rice products were significantly higher in As, Hg and Pb and lower in Se, Fe, Cu and Zn than foods based either on wheat or alternative grains. Wheat flours and wheat-based foods were higher in Cd than rice and rice-based products. We found a strong correlation between arsenic and mercury in rice and rice based products that warrants further investigation. Assuming that the concentrations of metals measured in these products are indicative of their bioavailable concentrations upon ingestion, our study reinforces nutritional advice to those concerned about dietary arsenic exposure as part of a GF diet, namely to consume a wide variety of grains, which would reduce arsenic, and mercury exposure while supplying essential micronutrients.

Acknowledgments

This research was supported by NIH/NIEHS P42 ES007373, NIH/NCI P30CA023108, NIH/NIEHS 1UG3OD023275-01 and NIH/NIEHS P01ES022832, EPA RD83544201 and by NIGMS P20 GM104416 to TP. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the Environmental Protection Agency.

BIBLIOGRAPHY

- Alvarez-Jubete L, Arendt EK, Gallagher E. Nutritive value of pseudocereals and their increasing use as functional gluten-free ingredients. *Trends in Food Science & Technology*. 2010; 21(2):106–113.
- Arduini I, Masoni A, Mariotti M, Pampana S, Ercoli L. Cadmium uptake and translocation in durum wheat varieties differing in grain-Cd accumulation. *Plant Soil and Environment*. 2014; 60(1):43–49.
- Bulka CM, Davis MA, Karagas MR, Ahsan H, Argos M. The Unintended Consequences of a Gluten-Free Diet. *Epidemiology*. 2017
- Carbonell-Barrachina AA, Wu X, Ramirez-Gandolfo A, Norton GJ, Burlo F, Deacon C, Meharg AA. Inorganic arsenic contents in rice-based infant foods from Spain, UK, China and USA. *Environmental Pollution*. 2012; 163:77–83. [PubMed: 22325434]

- Carey AM, Norton GJ, Deacon C, Scheckel KG, Lombi E, Punshon T, Guerinot ML, Lanzirotti A, Newville M, Choi Y, Price AH, Meharg AA. Phloem transport of arsenic species from flag leaf to grain during grain filling. *New Phytologist*. 2011; 192(1):87–98. [PubMed: 21658183]
- Cubadda F, Jackson BP, Cottingham KL, Van Horne YO, Kurzius-Spencer M. Human exposure to dietary inorganic arsenic and other arsenic species: State of knowledge, gaps and uncertainties. *Sci Total Environ*. 2017; 579:1228–1239. [PubMed: 27914647]
- DiGiacomo DV, Tennyson CA, Green PH, Demmer RT. Prevalence of gluten-free diet adherence among individuals without celiac disease in the USA: results from the Continuous National Health and Nutrition Examination Survey 2009–2010. *Scand J Gastroenterol*. 2013; 48(8):921–925. [PubMed: 23834276]
- EFSA. Scientific Opinion on Contaminants in the Food Chain on a request from the European Commission on cadmium in food: Cadmium in Food. *The EFSA Journal*. 2009; 980:1–139. The European Food Safety Authority.
- EU, EU. Commission Regulation (EU) 2015/1006 of 25 June 2015 amending Regulation No. 1881/2006 as regards maximum levels of inorganic arsenic in foodstuffs. 2016 (pp. Free access to EU law (treaties, directives, regulations, decisions, consolidated legislation), preparatory acts, case-law, international agreements, EFTA documents, summaries of EU legislation and other documents.). 2015. <http://eur-lex.europa.eu/eli/reg/2015/1006/ojEUR-Lex>
- European Food Safety Authority, E. EFSA Panel on Contaminants in the Food Chain (CONTAM); Scientific Opinion on Arsenic in Food. *EFSA Journal*. 2009; 7(10):1351, 1198.
- FAO. Dimensions of need: An atlas of food and agriculture. Banson: 1995.
- FDA. Arsenic in rice and rice products. In: U. FDA. , editor. Foodborne Illnesses and Contaminants. Vol. 2016. US FDA; 2016. <http://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm319870.htm>
- FDA. Mercury Levels in Commercial Fish and Shellfish (1990–2012). In: FDA. , editor. National Marine Fisheries Service Survey of Trace Elements in te Fishery Resource. Food and Drug Administration; 2017.
- FDA, U. FDA proposes limit for inorganic arsenic in infant rice cereal. Sucher, L., editor. US Food and Drug Administration; 2016.
- Grandjean P, Herz KT. Trace elements as paradigms of developmental neurotoxicants: lead, methylmercury and arsenic. *Journal of trace elements in medicine and biology : organ of the Society for Minerals and Trace Elements*. 2015; 31:130–134.
- Group M. Percentage of US adults trying to cut down or avoid gluten in their diets reaches new high in 2013, Reports NPD. In: McLynn, K., editor. National Purchase Diary Press Releases. National Purchase Diary; 2013. 2017
- Harris NS, Taylor GJ. Cadmium uptake and translocation in seedlings of near isogenic lines of durum wheat that differ in grain cadmium accumulation. *BMC plant biology*. 2004; 4:4. [PubMed: 15084224]
- Hong C, Yu XD, Liu JH, Cheng Y, Rothenberg SE. Low-level methylmercury exposure through rice ingestion in a cohort of pregnant mothers in rural China. *Environ Res*. 2016; 150:519–527. [PubMed: 27423706]
- Hu PJ, Li Z, Yuan C, Ouyang YN, Zhou LQ, Huang JX, Huang YJ, Luo YM, Christie P, Wu LH. Effect of water management on cadmium and arsenic accumulation by rice (*Oryza sativa* L.) with different metal accumulation capacities. *Journal of Soils and Sediments*. 2013; 13(5):916–924.
- Jackson BP. Fast ion chromatography-ICP-QQQ for arsenic speciation. *Journal of Analytical Atomic Spectrometry*. 2015; 30(6):1405–1407. [PubMed: 26366032]
- JECFA, J. F. W. E. C. o. F. A.. WHO Food Additives Series. Vol. 58. Geneva, Switzerland: World Health Organization; 2007. Safety evaluation of certain food additives and contaminants.
- Le XC, Ma M, Lu X, Cullen WR, Aposhian HV, Zheng B. Determination of monomethylarsonous acid, a key arsenic methylation intermediate, in human urine. *Environmental Health Perspectives*. 2000; 108(11):1015–1018.
- Meharg AA, Norton G, Deacon C, Williams P, Adomako EE, Price A, Zhu YG, Li G, Zhao FJ, McGrath S, Villada A, Sommella A, De Silva PMCS, Brammer H, Dasgupta T, Islam MR.

- Variation in Rice Cadmium Related to Human Exposure. *Environmental Science & Technology*. 2013; 47(11):5613–5618. [PubMed: 23668419]
- Meng B, Feng X, Qiu G, Anderson CW, Wang J, Zhao L. Localization and speciation of mercury in brown rice with implications for pan-asian public health. *Environmental Science & Technology*. 2014; 48(14):7974–7981. [PubMed: 24925231]
- Munera-Picazo S, Ramirez-Gandolfo A, Burlo F, Carbonell-Barrachina AA. Inorganic and total arsenic contents in rice-based foods for children with celiac disease. *Journal of Food Science*. 2014; 79(1):T122–128. [PubMed: 24313911]
- Norton GJ, Pinson SRM, Alexander J, McKay S, Hansen H, Duan GL, Islam MR, Ismam S, Stroud JL, Zhao FJ, McGrath SP, Zhu YG, Lahner B, Yakubova E, Guerinot ML, Tarpley L, Eizenga GC, Salt DE, Meharg AA, Price AH. Variation in grain arsenic assessed in a diverse panel of rice (*Oryza sativa*) grown in multiple sites. *New Phytologist*. 2012; 193(3):650–664. [PubMed: 22142234]
- Orecchio S, Amorello D, Raso M, Barreca S, Lino C, Di Gaudio F. Determination of trace elements in gluten-free food for celiac people by ICP-MS. *Microchemical Journal*. 2014; 116:163–172.
- Punshon T, Jackson BP, Meharg AA, Warczack T, Scheckel K, Guerinot ML. Understanding arsenic dynamics in agronomic systems to predict and prevent uptake by crop plants. *Science of the Total Environment* 581-582. 2017 Mar 1.:2009–2220. 2017.
- Raehsler SL, Choung RS, Marietta EV, Murray JA. Accumulation of Heavy Metals in People on a Gluten-Free Diet. *Clin Gastroenterol Hepatol*. 2017
- Reilly, C. Selenium in food and health. 2nd ed. New York, N.Y: Springer; 2006.
- Rothenberg SE, Feng XB. Mercury cycling in a flooded rice paddy. *Journal of Geophysical Research-Biogeosciences*. 2012; 117
- Rothenberg SE, Jackson BP, Carly McCalla G, Donohue A, Emmons AM. Co-exposure to methylmercury and inorganic arsenic in baby rice cereals and rice-containing teething biscuits. *Environ Res*. 2017; 159(Supplement C):639–647. [PubMed: 28938205]
- Rothenberg SE, Windham-Myers L, Creswell JE. Rice methylmercury exposure and mitigation: a comprehensive review. *Environ Res*. 2014; 133:407–423. [PubMed: 24972509]
- Rothenberg SE, Yu XD, Liu JH, Biasini FJ, Hong C, Jiang X, Nong YF, Cheng Y, Korrick SA. Maternal methylmercury exposure through rice ingestion and offspring neurodevelopment: A prospective cohort study. *International Journal of Hygiene and Environmental Health*. 2016; 219(8):832–842. [PubMed: 27503636]
- Sun GX, Williams PN, Carey AM, Zhu YG, Deacon C, Raab A, Feldmann J, Islam RM, Meharg AA. Inorganic arsenic in rice bran and its products are an order of magnitude higher than bulk grain. *Environmental Science and Technology*. 2008; 42:7542–7546. [PubMed: 18939599]
- Tai YP, Li ZA, McBride MB, Yang Y. Dry cultivation enhances cadmium solubility in contaminated soils but minimizes cadmium accumulation in a leafy vegetable. *Journal of Soils and Sediments*. 2017; 17(12):2822–2830.
- Taylor VF, Jackson BP, Chen CY. Mercury speciation and total trace element determination of low-biomass biological samples. *Anal Bioanal Chem*. 2008; 392(7–8):1283–1290. [PubMed: 18828006]
- Theethira TG, Dennis M. Celiac Disease and the Gluten-Free Diet: Consequences and Recommendations for Improvement. *Digestive Diseases*. 2015; 33(2):175–182. [PubMed: 25925920]
- USEPA, USEPA. Integrated Risk Information System (IRIS) Chemical Assessment Summary. United States Environmental Protection Agency; 2001. Integrated risk information system. Methylmercury (MeHg).
- Vici G, Belli L, Biondi M, Polzonetti V. Gluten free diet and nutrient deficiencies: A review. *Clin Nutr*. 2016; 35(6):1236–1241. [PubMed: 27211234]

HIGHLIGHTS

- Rice-based foods contained more As, Hg, Pb than others
- Rice-based foods contained less Se, Fe, Cu and Zn than others
- Arsenic and mercury in rice and rice products were strongly correlated
- Virtually all of the mercury in rice and rice based foods was methylmercury

Figure 1. Concentration of arsenic (ng/g) in a range of gluten-free foods and their non gluten-free equivalents, including A: whole grain rice, B: flours and C: pasta and bread.

Figure 2. Correlation between independent measures of total mercury concentration and methylmercury concentrations in rice products.

Table 1

Mean (SD) elemental concentration of food products, grouped by product categories and type of grain as determined by ICP-MS only.

Product	Grain type	N	Mn	Fe	Ni	Cu	Zn	Cr	Co	As	Se	Cd	Sb	Hg	Pb
ng/g															
Method detection limit															
Below detection limit [N (%)]															
% SRM recovery (n=4) Mean (SD)															
Flour															
	Rice	4	22.5 (18.2)	31.2 (19.2) ^b	0.7 (0.8) ^{ab}	2.4 (1.6) ^b	14.9 (6.8) ^b	54 (31)	30.6 (29.9) ^{ab}	112 (92) ^a	117 (57)	26 (35.6)	1.8 (1.0)	2.2 (1.1) ^a	4 (2)
	Wheat	6	25.9 (25.4)	38.0 (15.6) ^{ab}	0.3 (0.3) ^b	3.3 (2.0) ^{ab}	20.1 (13.4) ^b	10 (9)	12.8 (17.0) ^b	11 (4) ^b	375 (164)	45 (23.5)	4.8 (5.8)	0.9 (0.9) ^b	3 (3)
	Other	12	31.7 (21.6)	72.4 (29.5) ^a	2.3 (2.1) ^{ab}	10.3 (9.3) ^a	38.0 (15.5) ^a	52 (84)	108.5 (103.9) ^a	13 (5) ^b	199 (184)	31 (25.3)	5.3 (3.7)	0.6 (0.2) ^b	8 (6)
Whole grain rice															
	White	5	7.8 (3.1) ^b	3.7 (4.4)	0.4 (0.4)	2.3 (1.5)	13.1 (6.0) ^b	48 (100)	9.3 (5.9) ^b	94 (89) ^a	92 (71)	38.0 (42.7)	2.2 (1.2)	1.4 (1.1)	19 (25)
	Brown	6	28.9 (2.7) ^a	11.5 (1.5)	0.4 (0.2)	3.1 (1.0)	22.5 (4.3) ^a	18 (14)	26.4 (15.9) ^a	183 (46) ^a	104 (103)	27.0 (28.6)	1.7 (1.2)	1.8 (1.2)	8 (6)
	Enriched white	7	9.3 (3.3) ^b	19.4 (17.8)	0.2 (0.1)	1.8 (0.4)	12.3 (4.0) ^b	58 (59)	12.1 (4.3) ^b	177 (69) ^a	137 (81)	10.9 (4.1)	6.9 (13.8)	1.9 (1.0)	21 (23)
Prepared Foods & Pasta															
	Rice	18	20.1 (22.4)	13.3 (5.2)	0.4 (0.2)	2.0 (0.9)	14.3 (5.1)	189 (106)	23.5 (16.9)	141 (82)	115 (91)	15.9 (11.3)	2.9 (1.6)	3.4 (2.0)	22 (12)
	Non-rice	8	13.9 (11.0)	26.1 (14.4)	0.2 (0.1)	3.4 (1.6)	20.9 (8.4)	111 (113)	6.7 (5.5)	15 (12)	340 (323)	52.4 (42.7)	1.7 (1.7)	1.0 (0.4)	7 (3)
All products combined															
	Rice	38	17.9 (17.2) ^b	14.0 (11.8) ^c	0.3 (0.3) ^b	2.2 (1.1) ^b	15.0 (5.9) ^b	108 (110)	19.3 (15.0) ^a	147 (78) ^a	114 (84) ^b	21.3 (24.2) ^b	3.3 (5.9)	2.5 (1.7) ^a	18 (16) ^a
	Wheat	15	17.7 (18.1) ^b	32.0 (15.8) ^b	0.2 (0.2) ^b	3.2 (1.7) ^b	20.2 (10.0) ^b	66 (98)	9.1 (11.0) ^b	13 (9) ^b	333 (269) ^a	46.5 (36.2) ^a	2.9 (3.9)	0.9 (0.6) ^b	5 (4) ^b
	Other	13	33.1 (20.1) ^a	70.7 (31.4) ^a	2.4 (2.0) ^a	10.4 (9.2) ^a	38.1 (15.3) ^a	55 (83)	113.8 (99.7) ^c	13 (4) ^b	207 (177) ^{ab}	31.5 (24.5) ^{ab}	5.2 (3.8)	0.6 (0.3) ^b	8 (6) ^{ab}
Gluten Content															
	Contains gluten	13	16.5 (18.9)	28.8 (16.8)	0.2 (0.2)	2.9 (1.7)	18.2 (10.2)	85 (133)	9.5 (11.8)	18 (19)	307 (256)	37.8 (27.9)	3.1 (4.1)	1.0 (0.7)	7 (7)
	GF	55	21.5 (18.5)	27.6 (29.1)	0.8 (1.2)	4.1 (5.4)	20.6 (12.8)	90 (97)	38.8 (60.5)	112 (90)	149 (146)	26.7 (29.3)	3.6 (5.4)	2.0 (1.7)	15 (15)

Results of one-sample t-tests or ANOVA statistical tests are indicated: ns not significant;

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

*
Value < 0.05;
d

> 0.01 and

< 0.0001. Tukey's means comparison test results are indicated by superscript lettering. nc=not certified. Italicized SRM recoveries indicate reference values supplied on SRM certification. The method detection limit was based on the higher value of either the instrument detection limit or 3σ of the digestion blanks. The method detection limit also accounts for the sample digestion dilution and reflects the concentration limit in the food product.

Table 2

Results of Spearman's non-parametric statistical test between measured concentrations of arsenic species (total^a, inorganic^b, organic^b) and total mercury^a and methylmercury^c in whole grain rice (N=19).

Arsenic	Total Hg		MeHg	
	Spearman's ρ	<i>p</i> -Value	Spearman's ρ	<i>p</i> -Value
Total	0.7309	<.0001	0.6368	0.0034
Inorganic	0.5083	0.0001	0.5088	0.0311
Organic	0.5296	<.0001	0.6257	0.0055

^a Measured by ICP-MS

^b Measured by LC-ICP-MS

^c Measured by direct mercury analysis