Published in final edited form as:

Org Lett. 2017 December 01; 19(23): 6356–6359. doi:10.1021/acs.orglett.7b03196.

Catalytic Enantioselective Intermolecular [5+2] Dipolar Cycloadditions of a 3-Hydroxy-4-Pyrone-Derived Oxidopyrylium Ylide§

Katherine N. Fuhr^a, Danielle R. Hirsch^{b,c}, Ryan P. Murelli^{b,c,*}, and Stacey E. Brenner-Moyer^{a,*}

^aDepartment of Chemistry, Rutgers University, Newark, New Jersey, USA 07102

^bDepartment of Chemistry, Brooklyn College, The City University of New York, Brooklyn, New York, USA 11210

^cPhD Program in Chemistry, The Graduate Center, The City University of New York, New York, NY, USA 10016

Abstract

The first catalytic enantioselective [5+2] dipolar cycloaddition of a 3-hydroxy-4-pyrone-derived oxidopyrylium ylide is described. These studies leveraged the recently recognized ability of oxidopyrylium dimers to serve as the source of ylide, which was found to be key to increasing yields and achieving enantiomeric excesses up to 99%. General reaction conditions were identified for an array of α,β -unsaturated aldehyde dipolarophiles. Reaction products possess four stereocenters, and subsequent reduction introduced a fifth contiguous stereocenter with total stereocontrol.

Graphical abstract

The [5+2] dipolar cycloaddition of oxidopyrylium ylides enables facile access to the 8-oxabicyclo[3.2.1]octane scaffold,¹ and has been widely used in target-oriented synthesis.² In addition to being a prevalent substructure in bioactive natural products,³ 8-

Corresponding Authors. seb244@rutgers.edu, rpmurelli@brooklyn.cuny.edu.

§Dedicated to Professor Paul A. Wender on the occasion of his 70th birthday.

ASSOCIATED CONTENT

Supporting Information

Supporting information includes detailed experimental procedures and ¹H and ¹³C NMR spectra. These materials are available free of charge via the Internet at <http://pubs.acs.org>.

oxabicyclo[3.2.1]octanes provide an entrée into functionalized tetrahydrofurans and cycloheptanes.⁴⁻⁵ These two ring systems are of high interest to synthetic chemists, with the latter being relatively challenging to construct due to ring-strain and entropic considerations.⁶

Despite the widespread utility of oxidopyrylium cycloadditions, including asymmetric variants of the reaction,^{4b,7} catalytic enantioselective [5+2] dipolar cycloadditions of oxidopyrylium ylides have eluded the synthetic chemistry community until very recently. The first such reaction was intramolecular and was reported by Jacobsen and coworkers in 2011, and was subsequently adapted as an intermolecular process in 2014.⁸ These reactions used a combination of enamine catalysis and a hydrogen-bonding catalyst to promote the asymmetric cycloaddition (eq 1, Scheme 1). In 2015, Reyes and Vicario reported the first catalytic enantioselective [5+2] dipolar cycloaddition of a benzopyrylium oxide via dienamine catalysis (eq 2).⁹ These remain the only strategies for catalytic asymmetric oxidopyrylium cycloadditions published to date.

Both of these prior studies leveraged carboxylate elimination of acyloxypyran to generate reactive oxidopyrylium ylides, which is the method most commonly associated with oxidopyrylium [5+2] cycloadditions.^{1,2} A less studied intermediate to oxidopyrylium ylides are 3-hydroxy-4-pyrones (**9**, Scheme 2), which are classically converted to the active oxidopyrylium ylides through thermal group transfer (**9** → **10a**) and trapped by a tethered alkene or alkyne.^{5a,10} Intermolecular 3-hydroxy-4-pyrone-based oxidopyrylium cycloadditions, on the other hand, can be performed by first converting the pyrone to 3-hydroxy-4-methoxypyrylium triflate salts (**11**), and then deprotonating them with aniline bases in the presence of the dipolarophile.¹¹ We further adapted this intermolecular reaction through our recent discovery that oxidopyrylium dimers (**12**) can be used as the source of the oxidopyrylium ylide, thereby eliminating conjugate acid and residual base present in the *in situ* deprotonation protocol.¹² 3-Hydroxy-4-pyrone-based oxidopyrylium ylides (**10a/b**) are inherently electron rich due to the presence of an added donor oxygen, and as such are known to react preferentially with electron poor dipolarophiles.^{11a, 12c} Thus, unlike prior catalytic asymmetric intermolecular processes wherein oxidopyrylium ylides reacted with electron rich dipolarophiles (Scheme 1), 3-hydroxy-4-pyrone-derived oxidopyrylium ylides were viewed as an ideal ylide with which to develop an asymmetric oxidopyrylium cycloaddition reaction employing electron poor dipolarophiles accessible through iminium catalysis.¹³

Early studies revealed that commercially available catalyst **14** was capable of promoting an enantioselective reaction between salt **11** and cinnamaldehyde upon addition of *i*-Pr₂NPh (method A, Table 1), generating product in high ee and dr, albeit in prohibitively low yields at rt (entry 1). Efforts to improve yields by raising the temperature from rt to 50 °C led to significant deterioration of both enantio- and diastereoselectivity (entry 2). After extensive optimization,¹⁴ the best results were obtained in acetonitrile by doubling the loading of catalyst, which at room temperature generated the product in excellent ee and outstanding dr, but still in only 45% yield after 7 days (entry 3). We thus turned our attention to dimer **12** as the source of oxidopyrylium ylide (method B, Table 1).

When **12** was subjected to comparable reaction conditions, no conversion was observed, even at 50 °C (entry 4). This was not surprising given that catalytic acid is generally necessary to promote iminium ion formation, and that initial studies revealed no background reaction between **12** and cinnamaldehyde at this temperature (result not shown). What was surprising, however, was that the addition of catalytic benzoic acid led only to trace amounts of products (entry 5). Given the promising results with **11** and base, anilinium triflate was reintroduced into the reaction mixture through the addition of stoichiometric *i*-Pr₂NPh and TfOH (entries 6–9). Adjustment of time and temperature resulted in much higher product yields and comparable enantioselectivities relative to those obtained with salt **11**. In the absence of catalyst, some background reaction was observed (entry 10), demonstrating that the anilinium triflate promotes racemic background reaction. The equivalents of additives were reduced to achieve a concentration of anilinium triflate that promoted iminium activation, while not detrimentally catalyzing the racemic background reaction, which gave nearly enantiopure products (entry 11).¹⁴

It is, at present, unclear why *N,N*-diisopropylanilinium triflate serves as an efficient additive for the reaction, whereas benzoic acid does not. Assuming that both additives are capable of catalyzing iminium ion formation, one hypothesis is that the greater nucleophilicity of the resulting benzoate, relative to both *i*-Pr₂NPh and triflate, may promote side reactions such as demethylation of the oxidopyrylium ylide (i.e., via loss of methoxide or methyl ether cleavage),^{12b–c} thereby inhibiting desired reactivity. Confirming the importance of a non-nucleophilic conjugate base, TfOH alone as an additive did promote the cycloaddition reaction with comparable enantio- and diastereoselectivities, however lower yields were observed due to decomposition (entry 12).

Upon identification of optimal reaction conditions, an investigation into the substrate scope of this transformation ensued (Scheme 3). The reaction exhibited broad substrate compatibility, affording 8-oxabicyclo[3.2.1]octane products in all cases in moderate to high yields and dr, and in high ee. Both electron-rich and –poor benzene rings were tolerated (**13a–g**), as were heteroaromatic and polyaromatic R groups (**13h–i**), although an aromatic R group was not required for this transformation. Adjacent non-aromatic sp² centers, as in ester **13j**, were acceptable. Importantly, aliphatic enals were also suitable substrates (**13k**). Also notable is the tolerance of this cycloaddition reaction for steric bulk. The enal with the most hindered β-position, generated the corresponding product, **13g**, in comparable yield to other substrates, but in higher ee and as a single diastereomer.

Remarkably, despite the diversity of enals evaluated in this reaction, the conditions developed for **13a** proved general and, with only one exception (**13j**), did not require extensive reoptimization. Only a minor adjustment in temperature was necessary for certain substrates to achieve high ee's (**13c–e**, **13h**), and there was no discernable trend as to which substrates would require these slightly modified conditions. While the reaction time, product yield, and ee all increased when these substrates were subjected to a slightly reduced reaction temperature, curiously, the dr of the products decreased in all cases, sometimes dramatically so. For example, **13c** and **13e** were each formed in 13:1 dr at 40 °C and in 7:1 dr at 35 °C, while **13h** was produced in 10:1 dr at 40 °C and in 3:1 dr at 35 °C. With enal **13j**, however, there was a significant uncatalyzed background reaction even at 0 °C, so low

temperature conditions were employed. α -Methyl-*trans*-cinnamaldehyde was unreactive under the standard conditions after three days.

Representative subsequent transformations of the reaction products are shown in Scheme 4. Global reduction of **13** to provide **16** introduced another stereocenter. Diols **16g** and **16a** were produced in quantitative and nearly quantitative yield, respectively, and as a single diastereomer. Acidic hydrolysis of the methyl enol ether in **16a** revealed ketone **17** in 51% unoptimized yield. Finally, selective silyl protection of the primary alcohol afforded **18** in 84% yield. Reaction products, **13**, effectively contain three carbonyl groups (including one masked as an enol ether), which have been differentiated in **18** for further orthogonal elaboration.

The relative and absolute configurations of **13a** and **16g**, respectively, were established by X-ray crystallography.¹⁵ X-ray crystallography also revealed that the major diastereomer is the exo diastereomer. Comparison of ¹H NMR data revealed that the exo diastereomer was the major diastereomer in all cases, and the exclusive diastereomer in the case of **13g**. Based on this data, a model for asymmetric induction using the oxidopyrylium ylide form is depicted in Figure 1. It cannot be ruled out, however, that the conjugated iminium ion reacts directly with dimer **12** or an intermediate.

In conclusion, the first catalytic enantioselective [5+2] dipolar cycloaddition employing 3-hydroxy-4-pyrone-based oxidopyrylium ylides was developed. This also represents the first catalytic enantioselective intermolecular oxidopyrylium cycloaddition of electron-deficient dipolarophiles, which were accessible via iminium catalysis, using a commercially available *L*-proline-derived catalyst. The oxidopyrylium ylide dimer reactant enabled use of substoichiometric quantities of strong acid (TfOH) and base (*N,N*-diisopropylaniline), which was critical to attain the best yields, selectivities, and reaction times. This transformation has a remarkable substrate scope with respect to the enal reactant, yet general reaction conditions that were optimal for nearly all enals were identified. This reaction generated 8-oxabicyclo[3.2.1]octanes with four contiguous stereocenters in 91–99% ee, and the exo adduct was the major diastereomer. Reaction products can be readily elaborated, including introduction of a fifth stereocenter.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Science Foundation (CAREER-1461919; SEB-M), Rutgers University (SEB-M), and the National Institutes of Health (SC1GM111158; RPM). We also thank Dr. William Brennessel (University of Rochester) and Dr. Furong Sun (University of Illinois) for the acquisition of X-ray crystallographic and mass spectrometry data, respectively.

References

1. For reviews of [5+2] cycloadditions of oxidopyrylium species, see: Clavier H, Pellissier H. *Methods and Applications of Cycloaddition Reactions in Organic Syntheses*. Nishiwaki N. John Wiley &

- Sons, Inc. Hoboken 2014: 631–654. Pellissier H. *Adv. Synth. Catal.* 2011; 353:189–218. Singh V, Krishna UM, Vikrant, Trivedi GK. *Tetrahedron*. 2008; 64:3405–3428.
2. For some recent examples, see: Nicolau KC, Kang Q, Ng SY, Chen DY-K. *J. Am. Chem. Soc.* 2010; 132:8219–8222. [PubMed: 20496885] Zhang M, Liu N, Tang W. *J. Am. Chem. Soc.* 2013; 135:12434–12438. [PubMed: 23930656] Strych S, Journot G, Pemberton RP, Wang SC, Tantillo DJ, Trauner D. *Angew. Chem. Int. Ed.* 2015; 54:5079–5083. Chen B, Liu X, Hu Y-J, Zhang D-M, Deng L, Lu J, Min L, Ye W-C, Li C-C. *Chem. Sci.* 2017; 8:4961–4966. [PubMed: 28959419]
 3. See, for example: a) Komaroviquinone: Uchiyama N, Kiuchi F, Ito M, Honda G, Takeda Y, Khodzhimatov OK, Ashurmetov OA. *J. Nat. Prod.* 2003; 66:128–131. [PubMed: 12542361] b) Intricarene: Marrero J, Rodríguez AD, Barnes CL. *Org. Lett.* 2005; 7:1877–1880. [PubMed: 15844929] c) Coristatin: Aoki S, Watanabe Y, Sanagawa M, Setiawan A, Kotoku N, Kobayashi M. *J. Am. Chem. Soc.* 2006; 128:3148–3149. [PubMed: 16522087] d) Englerin A: Ratnayake R, Covell D, Ransom TT, Gustafson KR, Beutler JA. *Org. Lett.* 2008; 11:57–60. e) Lancolide: Shi Y-M, Wang X-B, Li X-N, Luo X, Shen Z-Y, Wang Y-P, Xiao W-L, Sun H-D. *Org. Lett.* 2013; 15:5068–5071. [PubMed: 24059675] f) Waltherione: Jadulco RC, Pond CD, Van Wagoner RM, Koch M, Gideon OG, Matainaho TK, Piskaut P, Barrows LR. *J. Nat. Prod.* 2014; 77:183–187. [PubMed: 24392742] g) Wenyujinin: Yin G-P, Li L-C, Zhang Q-Z, An Y-W, Zhu J-J, Wang Z-M, Chou G-X, Wang Z-t. *J. Nat. Prod.* 2014; 77:2161–2169. [PubMed: 25275213]
 4. For some examples of the synthesis of 7-membered ring-containing natural products from 8-oxabicyclo[3.2.1]octanes: Bromidge SM, Sammes PG, Street LJ. *J. Chem. Soc. Perkin Trans. 1.* 1985:1725–1730. Wender PA, Rice KD, Schnute ME. *J. Am. Chem. Soc.* 1997; 119:7897–7898. Wender PA, Jesudason CD, Nakahira H, Tamura N, Tebbe AL, Ueno Y. *J. Am. Chem. Soc.* 1997; 119:12976–12977. Geng Z, Chen B, Chiu P. *Angew. Chem. Int. Ed.* 2006; 45:6197–6201. Kawasumi M, Kanoh N, Iwabuchi Y. *Org. Lett.* 2011; 13:3620–3623. [PubMed: 21657253] Leung LT, Chiu P. *Chem. Asian J.* 2015; 10:1042–1049. [PubMed: 25556852]
 5. Synthesis of tetrahydrofurans from 8-oxabicyclo[3.2.1]octanes: Rodríguez JR, Rumbo A, Castedo L, Mascareñas JL. *J. Org. Chem.* 1999; 64:4560–4563. Mascareñas JL, Rumbo A, Castedo L. *J. Org. Chem.* 1997; 62:8620–8621. Fishwick CWG, Mitchell G, Pang PFW. *Synlett.* 2005:285–286. Krishna UM. *Tetrahedron Lett.* 2010; 51:2148–2150. Yadav AA, Sarang PS, Trivedi GK, Salunkhe MM. *Synlett.* 2007:989–991. Ali MA, Bhogal N, Findlay JBC, Fishwick CWG. *J. Med. Chem.* 2005; 48:5655–5658. [PubMed: 16134933]
 6. For reviews of 7-membered carbocycle syntheses, see: Battiste MA, Pelphrey PM, Wright DL. *Chem. Eur. J.* 2006; 12:3438–3447. [PubMed: 16402402] Nguyen TV, Hartmann JM, Enders D. *Synthesis.* 2013; 45:845–873.
 7. For some examples of diastereoselective oxidopyrylium cycloadditions, see: López F, Castedo L, Mascareñas JL. *Org. Lett.* 2000; 2:1005–1007. [PubMed: 10768208] Krishna UM, Deodhar KD, Trivedi GK. *Tetrahedron.* 2004; 60:4829–4836. For an example of stoichiometric chiral reagent control, see: Gerard B, Sangji S, O’Leary DJ, Porco JA Jr. *J. Am. Chem. Soc.* 2006; 128:7754–7755. [PubMed: 16771486]
 8. a) Burns NZ, Witten MR, Jacobsen EN. *J. Am. Chem. Soc.* 2011; 133:14578–14581. [PubMed: 21848300] b) Witten MR, Jacobsen EN. *Angew. Chem. Int. Ed.* 2014; 53:5912–5916.
 9. Orue A, Uria U, Reyes E, Carrillo L, Vicario JL. *Angew. Chem. Int. Ed.* 2015; 54:3043–3046.
 10. a) Volkmann RA, Weeks PD, Kuhla DE, Whipple EB, Chmurny GN. *J. Org. Chem.* 1977; 42:3976–3978. b) Garst ME, McBride BJ, Douglass JG. *Tetrahedron Lett.* 1983; 24:1675–1678. c) Wender PA, McDonald FE. *J. Am. Chem. Soc.* 1990; 112:4956–4958. d) Rumbo A, Mourinho A, Castedo L, Mascareñas JL. *J. Org. Chem.* 1996; 61:6114–6120. [PubMed: 11667444] e) Mascareñas JL, Pérez I, Rumbo A, Castedo L. *Synlett.* 1997:81–82. f) Wender PA, D’Angelo N, Elitzin VI, Ernst M, Jackson-Ugueto EE, Kowalski JA, McKendry S, Rehfeuter M, Sun R, Voigtlaender D. *Org. Lett.* 2007; 9:1829–1832. [PubMed: 17408281]
 11. a) Wender PA, Mascareñas JL. *Tetrahedron Lett.* 1992; 33:2115–2118. b) Wender PA, Mascareñas JL. *J. Org. Chem.* 1991; 56:6267–6269. c) Mei G, Yuan H, Gu Y, Chen W, Chung LW, Li C-C. *Angew. Chem. Int. Ed.* 2014; 53:11051–11055.
 12. a) D’Erasmus MP, Masaoka T, Wilson JA, Hunte EM Jr, Beutler JA, Le Grice SFJ, Murelli RP. *Med. Chem. Commun.* 2016:1789–1792. b) D’Erasmus MP, Meck C, Lewis CA, Murelli RP. *J. Org.*

Chem. 2016; 81:3744–3751. [PubMed: 27018974] c) Meck C, Mohd N, Murelli RP. Org. Lett. 2012; 14:5988–5991. [PubMed: 23167954]

13. a) For a review of iminium catalysis, see: Erkkilä A, Majander I, Pihko PM. Chem. Rev. 2007; 107:5416–5470. [PubMed: 18072802] For select recent examples of our development of iminium- and organocatalyzed transformations, see: Guo Q, Fraboni AJ, Brenner-Moyer SE. Org. Lett. 2016; 18:2628–2631. [PubMed: 27186662] Fraboni AJ, Brenner-Moyer SE. Org. Lett. 2016; 18:2146–2149. [PubMed: 27070296]
14. See the Supporting Information.
15. CCDC 1567374 and 1567375 contain the supplementary crystallographic data for this paper. These data are provided free of charge by The Cambridge Crystallographic Data Centre.

Figure 1. Proposed Mode of Asymmetric Induction

Scheme 1. Catalytic Asymmetric Intermolecular Oxidopyrylium Cycloadditions Employing Acyloxypyranes

Scheme 2. Overview of 3-Hydroxy-4-pyrone-Based Oxidopyrylium Cycloaddition

Scheme 3. Substrate Scope Studies.^a

^a Yields reflect isolated yields, with dr determined by ¹H NMR, and ee determined by chiral phase HPLC. ^b Equivalents and concentrations reported relative to monomeric ylide. ^c Reaction run at 35 °C. ^d Single diastereomer. ^e Reaction run for 11 days at –25 °C.

Scheme 4. Elaboration of Oxabicyclo Scaffold

Table 1
Representative Results from Asymmetric Oxidopyrylium Cycloaddition Optimization Studies.

Method A

entry	14 (mol %)	t (d)	temp (°C)	yield (%) ^a	ee (%) ^b	dr ^c	
1 ^d	20	7	rt	9	90	16:1	
2	20	0.1	50	47	67	5:1	
3 ^e	40	7	rt	45	97	33:1	
Method B

entry	additive (equiv) ^f	t (d)	temp (°C)	yield (%) ^a	ee (%) ^b	dr ^c	
4 ^g	-	7	50	nr	--	--	
5 ^g	BzOH (0.2)	4	50	trace	--	--	
6	<i>i</i> -Pr ₂ NPh (1.0)	7	rt	32	95	10:1	
7	<i>i</i> -Pr ₂ NPh (1.0)	2	35	42	93	6:1	
8 ^d	<i>i</i> -Pr ₂ NPh (1.0)	2	40	82	94	7:1	
9	<i>i</i> -Pr ₂ NPh (1.0)	2	45	91	89	6:1

entry	additive (equiv) ^f	t (d)	temp (°C)	yield (%) ^a	ee (%) ^b	dr ^c
10 ^h	<i>i</i> -Pr ₂ NPh (1.0) TfOH (1.0)	4	40	8	--	1.5:1
11	<i>i</i> -Pr ₂ NPh (0.4) TfOH (0.4)	2	40	78	99	9:1
12	TfOH (0.4)	2	40	41	98	8:1

^aIsolated yield.

^bDetermined by chiral phase HPLC.

^cDetermined by ¹H NMR.

^dAverage of 2 experiments.

^eSolvent was MeCN.

^fEquivalents and concentrations reported relative to monomeric ylide for consistency to method A.

^gSolvent was CHCl₃.

^hIn the absence of **14**.