Published in final edited form as:

Curr Opin Psychiatry. 2013 September ; 26(5): 511–519. doi:10.1097/YCO.0b013e3283642cb6.

Assessing prescription stimulant use, misuse and diversion among youth 10 to 18 years of age

Linda B. Cottler, PhD, MPH, Catherine Woodstock Striley, PhD, MPE, and Sonam O. Lasopa, MPhil

University of Florida

Abstract

Purpose—Assessing medical and non-medical use of stimulants and diversion is a challenge, especially among youth, with different methods for recruitment, and definitions of non-medical use and use. The field needs inexpensive, yet effective and reliable, methods of data collection to understand the prescription drug use problem. Most studies of youth are school or web-based, and conducted with teens.

Recent Findings—The National Monitoring of Adolescent Prescription Stimulants Study (N-MAPSS) recruited 11,048 youth 10 to 18 years of age from urban, rural and suburban areas in ten US cities using an entertainment venue intercept study. This review discusses the effectiveness of the method and results from four cross sections as well as the representativeness of the sample. Lifetime prevalence of any stimulant use was 14.8%, with rates highest among rural 16 to 18 year olds. The rate of past 30 day use was 7.3%, with over half (3.9%) non-medical use. Nearly 12% of all youth (whether a user or not) reported lifetime incoming/outgoing diversion of prescription stimulants.

Summary—Because no study has focused on stimulant use among youth as young as 10 and 11, this study is a landmark for future comparisons and offers a unique strategy for sampling and data collection.

Keywords

Intercept venue study; prescription stimulant use; non-medical use; drug use; diversion

INTRODUCTION

Many methodological challenges exist in the assessment of non-medical use (NMU) of prescription medications among adults. This begins with substantial variation in the definition and questions used to assess it (1–3). NMU is defined as “misuse” or “abuse”, and is often referred to as use that deviates from the prescribed dose, route, or frequency of administration, or use without a prescription. Medical use is typically defined as use

Author of correspondence: Linda B. Cottler, **Address:** Department of Epidemiology, University of Florida, 1225 Center Dr. PO Box 100231, Gainesville, Florida, 32611, **Telephone number:** 352-273-5468, lbcottler@ufl.edu.

Conflict of Interest:

Authors declare no conflicts of interest other than those in the acknowledgements.

according to the labeled route, frequency, and dose of drugs obtained with one's own prescription. Any use includes both medical and NMU. Due to the definitional challenges, and because NMU with a prescription is often neglected as a definition of misuse, comparisons among studies are difficult. This paper examines two areas of emerging concern: surveillance of young adults, and misuse and diversion of specific stimulant drugs by illustrating these issues with data from a recent US national survey.

Engaging Youth in Research

Challenges exist in assessing youth who are often thought to be questionable sources of information on their own use and NMU. Thus, many researchers use parents as proxy reporters because children may not be aware of the name or indication of their own prescription(s), often dispensed by parents at home or by teachers at school. Also, parental consent is usually required to obtain data from youth themselves and parents may decline to allow their children to participate in research.

Despite the earlier mentioned limitations, two United States national sources of data on prescription drug use among youth exist: 1) the National Survey on Drug Use and Health (NSDUH) and 2) Monitoring the Future (MTF). NSDUH conducts computer-assisted personal interviews with respondents 12 years of age and older in their homes (4–6); MTF relies on self-administered questionnaires from 8th (13 to 14 year olds), 10th (15 to 16 year olds) and 12th (17 to 18 year olds) graders completed in their classrooms (7). NMU of prescription drugs is assessed in NSDUH by the question: "Have you ever, even once, used [DRUG], that was not prescribed for you or that you took only for the experience or feeling it caused?" In MTF, it is assessed with: "On how many occasions (if any) have you used [DRUG] on your own—that is, without a doctor telling you to take them?" Both studies require parental consent.

Emerging Problem of Prescribed Stimulant Use and Diversion

Sweeney and colleagues (8) reported lifetime NSDUH prevalence of stimulant misuse to be 3.5% among respondents 12+ years of age. MTF estimated stimulant misuse in the past one year (including methamphetamine) to be 4.5% among 8th graders, and 12% among 12th graders (7). Definitional and other methodological issues affect reported prevalence of substance use as well. Absent students and non-enrolled youth are excluded from the MTF sample, which could lead to biased rates (9). Hispanics, who have higher school dropout rates (10), would be under-represented in school-based surveys—this is especially problematic in that non-whites have higher medication diversion rates compared to whites (11). Additionally, a child could be taking prescribed stimulants fraudulently obtained by faking symptoms with the physician.

Another national study—the National Survey of Adolescents-Replication (NSA-R) which is telephone based, reported a 2.7% prevalence of past year stimulant NMU among youth 12 to 17 years (12). For that study, youth were asked: Have you ever taken 'on your own' or non-medically stimulants like Ritalin, speed, Adderall, diet pills? These questions differ from those previously mentioned, further illustrating variability that exists in the assessment of nonmedical prescription stimulant use.

Regional studies have also been conducted; they have assessed stimulant misuse through a web survey with middle and high school students and found lifetime NMU of ADHD stimulants to be 4.5% (9). Kroutil and colleagues (13) identified lifetime prevalence of misuse of stimulants to be 2.6% among 12 to 17 year olds. Another found NMU of prescription stimulants to be 2.4% among youth in Mississippi schools (14). These are the only studies of stimulant misuse that can be found among youth. Others assess “use of prescription drugs to get high”, without looking specifically at rates of each type.

The National Monitoring of Adolescent Prescription Stimulants Study

To resolve these methodological issues to detect current levels of NMU of prescription stimulants among pre-teens and adolescents, the National Monitoring of Adolescent Prescription Stimulants Study (N-MAPSS) was launched with an innovative method to detect signals of use and misuse. An entertainment venue intercept method—used previously to recruit people in popular venues where they congregate or visit—allowed us to economically reach a large and diverse sample of over 11,000 youth (15–20). This method and rate comparisons with other more traditional sources of recruitment are the focus of this review.

METHODS

Our methods are described below.

Research Site Selection

To begin, standard federal regions of the United States (Office of Management and Budget) Circular A-105, used by the National Survey of Children’s Health were selected. Then, the IMS database—which identifies prescriptions dispensed in the US at retail pharmacies, using the ZIPcode of each prescriber—was used to identify States with the highest volume of stimulant prescriptions, and the city within each with the highest volume.

The regions, States, cities, and their 2008 rankings are shown in Table 1; four cities represented eastern US (Boston, New York, Philadelphia, Tampa); three represented central US (St. Louis, Cincinnati, Houston), and western US (Denver, Los Angeles, Seattle).

ZIPCode Enumeration

Areas were characterized as urban, suburban and rural, determined by city limits, proximity to city limits, and population density. All 5-digit ZIPcodes contained in the selected city boundary were defined “urban”. Those contiguous to the urban ZIPcodes were defined suburban and included if a population density was less than urban, but more than rural (the US Census does not define suburban). ZIPcodes contiguous to suburban areas with fewer than 1,000 persons per square mile were considered rural.

Eligibility Criteria and Participant Recruitment

Youth 10 to 18 years of age residing in an urban, suburban or rural ZIPcode from one of the ten cities were eligible. Youth unaware of their ZIPcode, non-English readers, those cognitively impaired, or in college, were excluded. N-MAPSS utilized the entertainment

venue intercept method to comprehensively and inclusively reach youth where they were likely to be, to achieve representativeness. This was accomplished by sending Recruiter-Interviewers (RIs) to carefully selected venues (shopping malls, movie theaters, sports and recreation centers, libraries, arcades, skate parks, and parks) to locate a diverse and representative sample of youth in each region. Criteria for venue selection included being inside the designated ZIPcode boundary, being youth friendly, having a large customer base, being age appropriate and allowing study recruitment. Home-schooled and truant youth were included. The most common venue visited was malls (69%), followed by parks and other shopping areas (13%). Libraries, skate parks, recreation centers, movie theaters and restaurants/food courts accounted for 11% of venues.

RIs approached potential respondents in venues based on the time and day youth most likely attend each venue, similar to the MacKellar et al (19) and Zhao et al (16) studies. RIs documented the city, date, time, gender, race/ethnicity and number of eligible youth in the approached group and age, grade and ZIPcode for each person contacted. Youth were coded as complete, refused, or ineligible (due to college, age, language, out of ZIPcode or over quota). Demographics of non-completers were used to calculate a response rate.

RIs answered respondent's questions about the study and obtained implied assent, indicated by survey completion. Parental permission was not solicited per Washington University and University of Florida Human Protection Research Offices because all survey data were anonymous.

Sampling Goals

The target sample size was based on a power calculation assuming a 4% rate of stimulant use; 270 youth per site was adequate (80% power) to detect a 3% difference in rates across sites. With four data collection periods, 1,080 youth per site would be adequate to show even small differences.

Recruitment goals ensured representation of all ages and urban, suburban and rural status. Our goal was to include 20% 10 to 12 year olds, 40% 13 to 15 year olds and 40% 16 to 18 year olds as well as 50% urban youth, 30% suburban and 20% rural, from each city. When adolescents at the venues were in small groups ("clusters"), RIs surveyed one 10 to 12 year old, two 13 to 15 year olds, and three 16 to 18 year olds per cluster. RIs were notified when urbanicity or age quotas were met for their city.

Assessment

The cost and logistics of supplying computers to 30-40 interviewers in ten cities motivated us to use a full-color paper assessment, which was familiar to youth, easy to use, and eliminated technical complications of computer malfunctions.

The N-MAPSS research team adapted survey questions from the Substance Abuse Module (SAM) (21) and the Washington University Risk Behavior Assessment (RBA) (22) on quantity and frequency, route of administration, reasons and source of use. The 20 minute assessment, conducted in private, was divided into two (Table 2). After Part I was completed, mostly on demographics and dosage and form recognition, respondents completed Part II,

where information on lifetime and past 30 day use of Adderall[®], Concerta[®], Daytrana[®], Ritalin[®] and Vyvanse[®] were assessed with photos of each. (All formulations were queried, including immediate release and extended release. NMU was assessed by: use other than by mouth (except for Daytrana[®]), use of someone else's medications, more than prescribed, or use "to get high", "out of curiosity" or "just because". After completion, participants were given a \$10 electronics store gift card.

Test-Retest Sub-study

Before the study, a Pre-Test Reliability Feasibility study was conducted with a sample of ten each of 10-12 year olds, 13-14 year olds, 15-16 year olds, and 17-18 year olds to ensure the assessment was appropriate for all. Participants were recruited via flyers in St. Louis; to be enrolled youth had to have used any Adderall[®], Concerta[®], Daytrana[®], Ritalin[®] or Vyvanse[®].

Interviewers guided 10 to 12 year olds through the written survey, but were present in the room with 13 to 18 year olds to answer any questions during survey self-administration (Time1). Respondents returned seven days later for the Time2 survey with a different interviewer. Since locating information was needed for the Time2 interview, parental and youth Informed Consent was needed. All participants received a \$40 gift certificate to an area department store.

The data showed good to excellent agreement between Time1 and Time2 interviews (kappas = 0.6-1.0); use of drugs and an ADD diagnosis were in nearly perfect agreement (0.95 to 1.0).

After Time2, the project coordinator conducted a Discrepancy Interview Protocol (DIP) and Debriefing Interview to identify questions needing modification (23). The DIP compares answers from Time1 to Time2 to assess if inconsistent responses were due to misunderstanding questions, inability to remember answers, change in situation since the first interview, or if the youth was not paying attention. A Debriefing Interview was then conducted with 50 10- to 18-year olds to evaluate their understanding of the intent of the questions. Participants received a \$10 gift card for these interviews. Responses indicated that youth had no difficulty understanding the questions; no other inconsistencies were found, confirming that the survey was feasible.

Hiring and Training of Recruiter Interviewers (RIs)

Craigslist was used to find RIs, along with university listservs and career centers. Once hired, RIs received a Training Manual with ZIP Codes, Question-by-Question Specifications, Screeners, Logs, Adverse Event Forms, surveys, and Code of Conduct material. A two-hour training via Skype was held which covered a practice interview and editing session. Additional time was needed to review details before RIs entered the field. The project coordinator was available every day and evening via text messaging to address issues that arose; she also gave immediate feedback and guidance on surveys. Booster training continued throughout field periods to reinforce protocol adherence. For subsequent cross-sections, RIs were chosen from among the excellent performers. St. Louis project staff and investigators also collected surveys for first-hand experience (LBC, CS).

Quality Control

Quality control from St. Louis included a check of handwriting on all self-administered surveys; analyses were conducted to identify variable outliers, suspicious patterns and potentially fabricated data (24). Questionable surveys were reviewed by the team resulting in 164 surveys being excluded.

Surveys were also checked with the SAS random number generator for duplication both within, and across field periods. Suspicious cases were matched on ZIPcode, gender, age, ethnicity, birth order and grade in school. In five replications, the pool of potential repeaters was found to be 1.0% to 2.5%. We examined 7 pairs that matched on all 6 criteria and agreed on cigarette, alcohol, marijuana, and prescription stimulant use, and history of ADD/ADHD. Three of these also had similar family compositions and handwriting; they were excluded from further analysis.

RESULTS

The study was carried out in four cross-sections of data collection with fall and spring selected. In fall of 2008, in 52 days, 2,820 adolescents were surveyed. In spring 2009, in 74 days, 2,878 youth were surveyed. The third period, fall 2010, took 82 days to yield 2,839 surveys. The final period, in spring 2011, concluded 84 days later with 2,931 surveys collected.

Among the 21,444 youth approached (Figure 1), 25% did not stop to hear the RI's introduction. Additionally, 21% were ineligible mostly (84%) due to their age or an ineligible ZIPcode. Another 10%, who stopped to hear about the survey refused participation. Among the 11,468 youth who participated, 420 were eliminated as shown in the figure. The remaining 11,048 youth represented an 86.7% participation rate and a 68% overall response rate (11,048/16,143). Among these, 48%, 37% and 15% were urban, suburban, and rural, respectively, in line with the goal.

As shown in Table 3, data are stratified by urbanicity and age with 55.6% of youth living with both parents and 43.7% reporting excellent health. Other demographic characteristics are shown in the table. The sample, overall, mimicked population rates for race. Reports of drinking increased with age. Regarding stimulant use, 14.8% reported any lifetime use; rates increased with age with highest rates among rural 16 to 18 year olds. Among 10 to 12 year olds, 7.6% to 10.1% reported use. Additionally, 63.2% of youth reported thinking that prescription stimulant use was a moderate to large problem among youth.

Past 30 day use of stimulants was reported by 7.3% of youth (Table 4) with over half (53%) being NMU (3.9%). Diversion was stratified by having received (incoming) or given (outgoing) pills. Older youth were more likely than younger to report being approached to divert medications, or diverting and receiving prescription stimulants. In fact, 2.4% to 4.9% of 10 to 12 year olds reported incoming diversion. About 12% of youth reported lifetime diversion of prescription stimulants; 4.3% reported both incoming and outgoing diversion, with rates increasing with age. The highest rate of any lifetime diversion ($6.6 + 3.1 + 7.6\% = 17.4\%$) was among 16 to 18 year olds rural youth.

DISCUSSION

Assessing stimulant use from teens and pre-teens presents challenges around definitions, sampling and validity. This is the first US national study that monitored pre-teen and teenage use, non-medical use and diversion of prescription stimulants, using photo cards with and without brand names, and multiple questions to assess NMU and medical use. N-MAPSS employed an innovative sampling technique of recruiting respondents and conducting surveys in popular youth-oriented entertainment venues that helped overcome limitations of school based sampling, such as home-schooled adolescents, school drop-outs, and other selective absenteeism (25). By conducting an RI paper-based survey, and not a web-based one, there was the ability to validate the approximate age of our participants, unlike other studies recently completed (9, 26, 27).

Also, unlike other surveys, N-MAPSS focused primarily on prescription stimulants, and collected brand-level information with photos of each drug. Specifically, five stimulants were focused on and 8 questions were asked per drug which allowed for the assessment of medical as well as the nonmedical use of each stimulant along with subtypes of diversion and other risk factors. This enabled N-MAPSS to be uniquely comprehensive as well as exhaustive in assessing prescription stimulant use. Other information, not the focus of this review, such as qualitative data on what teens think could be done to reduce the NMU of prescription drugs, adds to the study.

The lifetime prevalence of use of prescription stimulants assessed in the NMAPSS is 14.8%. Learning from past experience in substance use measurement and prior, albeit limited prescription stimulant research, and as reported earlier, the NMAPPS survey asked specific questions for each of five stimulants that allows for a distinction to be made between medical use and nonmedical use exclusively. Using this distinction, the rate of past 30 day NMU only was found to be 2.7%. A past month prevalence rate of 3.9 % (both medical and nonmedical) was found; most other studies have reported a past twelve month prevalence of under 5% (12, 28–30).

The entertainment venue methodology was found to be highly successful as measured in several ways. First, only 10% of eligible youth declined participation—a mark of success at a time when household surveys are experiencing record low response rates (31). N-MAPSS has a large database of 1,406 10- to 12-year olds, which other studies lack. Second, the rates of self-reported ADD/ADHD, stimulant use, NMU, medical use, school suspension history and other indicators were consistent across four cross-sections. Third, when the age, gender, race and urban/rural composition of the 11,048 youth in the ten N-MAPSS cities were compared to that of the 2010 US Census for each of the cities, the sample was found to be highly representative, though there were slightly less males, Hispanics, and African Americans in several cities. Fourth, this method, modeled on the venue-time sampling method, allowed us to recruit a large number of youth in a relatively short period, without the need for parental approval (15, 19).

Conclusion

The N-MAPSS survey was launched in view of the limitations imposed by variable methodology, definition and measurement in the understanding of prescription stimulant use and misuse. The utilization of innovative methods to best recruit and survey youth based on ZIPcodes across ten cities in the US led to the comprehensive assessment of a nationally representative sample of youth. Additionally, inclusion of youth not reachable through conventional methods of school based or telephone surveys was possible. The past 30 day prevalence rate for NMU as assessed in the N-MAPSS was 3.9% overall (1.2% plus 2.7%).

Acknowledgments

N-MAPSS was implemented by Washington University in St. Louis and University of Florida under contract from Pinney Associates, Inc., with funding provided by Shire Development LLC and Noven Therapeutics.

Abbreviations

NMU Non-medical use

N-MAPSS National Monitoring of Adolescent Prescription Stimulants Study

References

1. Compton WM, Volkow ND. Abuse of prescription drugs and the risk of addiction. *Drug Alc Dependence*. 2006; 83(1):4–7.
2. Barrett SP, Meisner JR, Stewart SH. What constitutes prescription drug misuse? Problems and pitfalls of current conceptualizations. *Curr Drug Abuse Rev*. 2008; 1:255–62. [PubMed: 19630724]
3. Denisco RA, Chandler RK, Compton WM. Addressing the intersecting problems of opioid misuse and chronic pain treatment. *Exp Clinical Psychopharm*. 2008; 16(5):417–28.
4. Substance Abuse and Mental Health Services Administration. National survey on drug use and health: National Findings. Rockville, MD: Department of Health and Human Services, Office of Applied Studies; 2007. Contract No: DHHS publication No. SMA 08-4343DoHaH; 2008
5. Substance Abuse and Mental Health Services Administration. Overview of Findings from the 2003 National survey on drug use and health. Department of Health and Human Services, Office of Applied Studies; Rockville, MD: 2004.
- 6**. Nakawaki B, Crano W. Predicting adolescents' persistence, non-persistence, and recent onset of nonmedical use of opioids and stimulants. *Addictive behaviors*. 2012 Jun.37(6) Epub 2012 Feb 17. The study uses combined data from 2003 to 2009 from the National Survey of Drug Use and Health to examine nonmedical prescription opioids and stimulants and their association with onset of age and duration of use.
- 7**. Monitoring the Future 2012 Survey Results. National Institute on Drug Abuse; Bethesda, MA: <http://www.drugabuse.gov/related-topics/trends-statistics/infographics/monitoring-future-2012-survey-results>. This report indicates one of the most recent rates of prescription drug use in youth.
8. Sweeney CT, Sembower MA, Ertischek MD, Shiffman S, et al. Nonmedical use of prescription ADHD stimulants and preexisting patterns of drug abuse. *J Addict Dis*. 2013 Jan; 32(1):1–10. [PubMed: 23480243]
9. McCabe SE, Teter CJ, Boyd CJ. The use, misuse, and diversion of prescription stimulants among middle and high school students. *Subst Use Misuse*. 2004; 39:1095–116. [PubMed: 15387205]
10. Johnston, LD., O'Malley, PM., Bachman, JG., Schulenberg, JE. Various stimulant drugs show continuing gradual declines among teens in 2008 most illicit drugs hold steady [news release]. Ann Arbor, MI: University of Michigan News Service; Dec. 2008 p. 112008

11. McCabe SE, West BE, Teter CJ, Ross-Durrow P, et al. Characteristics associated with the diversion of controlled medications among adolescents. *Drug Alcohol Depend.* 2011; 118:452–8. [PubMed: 21665384]
12. McCauley JL, Danielson C, Amstadter AB, Ruggiero KJ, et al. The role of traumatic event history in non-medical use of prescription drugs among a nationally representative sample of US adolescents. *J Child Psychol Psychiatry.* 2010 Jan; 51(1):84–93. [PubMed: 19674194]
13. Kroutil LA, Van Brunt DL, Herman-Stahl MA, Heller DC, et al. Nonmedical use of prescription stimulants in the United States. *Drug Alcohol Depend.* 2006; 84(2):135–43. Epub 2006/02/17. [PubMed: 16480836]
- 14*. Viana AG, Trent L, Tull MT, Heiden L, et al. Non-medical use of prescription drugs among Mississippi youth: constitutional, psychological, and family factors. *Addict Behav.* 2012 Dec. 37:1382–8. Epub 2012 Jun 24. This is one of the most recent studies that examines individual factors within domains to increase the knowledge base of non medical prescription drug use. [PubMed: 22781877]
15. Vanden Bergh W, Nostlinger C, Buvé A, Beelaert G, et al. A venue based HIV prevalence and behavioral study among men who have sex with men in Antwerp and Ghent, Flanders, Belgium, October 2209 to March 2010. *Euro Surveill.* 2011 Jul 14.16(28) pii19914.
16. Zhao J, Cai WD, Chen L, Zhao JK, et al. A comparison of HIV infection and related risks among male sex workers in different venues in Shenzhen. *China AIDS Behav.* 2011 Apr; 15(3):635–42. [PubMed: 20711650]
17. Toledo CA, Varangrat A, Wimolsate W, Chemnasiri T, et al. Examining HIV infection among male sex workers in Bangkok, Thailand: A comparison of participants recruited at entertainment and street venues. *AIDS Educ Prev.* 2010 Aug; 22(4):299–311. [PubMed: 20707691]
18. Kendall C, Kerr LR, Gondim RC, Werneck GL, et al. An empirical comparison of respondent-driven sampling, time location sampling, and snowball sampling for behavioral surveillance in men who have sex with men, Fortaleza, Brazil. *AIDS Behav.* 2008 Jul; 12(4 Suppl):S97–104. Epub 2008 Apr 4. [PubMed: 18389357]
19. MacKellar D, Valleroy L, Karon J, Lemp G, et al. The young men's survey: methods for estimating HIV seroprevalence and risk factors among young men who have sex with men. *Public Health Rep.* 1996; 111(Suppl 1):138–44.
- 20*. Liu S, Zhao J, Rou K, Chen L, et al. A survey of condom use behaviors and HIV/STI prevalence among venue-based money boys in Shenzhen, China. *AIDS Behav.* 2012; 16(4):835–46. This study illustrates the effectiveness of time venue sampling to recruit difficult and hard to reach populations. [PubMed: 21800179]
21. Horton J, Compton W, Cottler LB. Reliability of substance use disorder diagnoses among African-Americans and Caucasians. *Drug Alcohol Depend.* 2000; 3:203–9.
22. Shacham E, Cottler LB. Sexual behaviors among club drug users: prevalence and reliability. *Arch Sex Behav.* 2010; 39(6):1331–41. Epub 2009/09/17. [PubMed: 19757011]
23. Cottler LB, Compton WM, Brown L, Shell A. The discrepancy interview protocol: A method for evaluating and interpreting discordant survey responses. *Int J Methods Psychiatr Res.* 1994; 4:173–82.
24. Johnson TP, Parker V, Clements C, Johnson TP, et al. Detection and prevention of data falsification in survey research. *Survey Research.* 2001; 32(3):1–3.
25. Hallfors D, Vevea JL, Iritani B, Cho H, et al. Truancy, grade point average, and sexual activity: A meta-analysis of risk indicators for youth substance use. *J Sch Health.* 2009; 72:205–11.
26. McCabe SE, Boyd CJ, Young A. Medical and nonmedical use of prescription drugs among secondary school students. *J Adolesc Health.* 2007; 40:76–83. Epub 2006/12/23. [PubMed: 17185209]
27. Cassidy TA, Varughese S, Russo L, Budman SH, et al. Nonmedical use and diversion of ADHD stimulants among U.S. adults ages 18–49: A national internet survey. *J Attent Disord.* 2012 Dec.: 26. [Epub ahead of print].
28. Simoni-Wastila L, Yang HW, Lawler J. Correlates of prescription drug nonmedical use and problem use by adolescents. *J Addict Med.* 2008; 2:31–9. [PubMed: 21768970]

29. McCabe SE, Teter CJ, Boyd CJ. The use, misuse, and diversion of prescription stimulants among middle and high school students. *Subst Use Misuse*. 2004; 39:1095–116. [PubMed: 15387205]
30. Schepis TS, Krishnan-Sarin S. Characterizing adolescent prescription misusers: a population-based study. *J Am Acad Child and Adolesc Psychiatry*. 2008 Jul.47:745–54. Epub 2011/08/02. [PubMed: 18520963]
31. Galea S, Tracy M. Participation rates in epidemiologic studies. *Ann Epidemiol*. 2007; 17:643–53. [PubMed: 17553702]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Key points

- First US national study that monitored pre-teen and teenage use, non-medical use and diversion of prescription stimulants with significant details for each topic.
- Showed effectiveness of an entertainment venue intercept method in recruiting a nationally representative sample for the survey.
- Found lifetimes rates of stimulant use to be nearly 15%
- Past 30 day rate of prescription stimulant use was 7.3%, with NMU reported to be more than half that rate (3.9%).

Figure 1.

Accrual of the N-MAPSS sample from 21,444 approached to 11,048 completed

Table 1**N-MAPSS Cities**

OMB Circular A-105 Region	State	State Volume Rank in Nation ¹	City	City Volume Rank in Nation ²
Eastern US:				
Region I	Massachusetts	13	Boston	43
Region II	New York	9	New York City	2
Region III	Pennsylvania	7	Philadelphia	17
Region IV	Florida	2	Tampa	22
Central US:				
Region V	Ohio	3	Cincinnati	6
Region VI	Texas	1	Houston	1
Region VII	Missouri	16	Saint Louis	4
Western US:				
Region VIII	Colorado	31	Denver	49
Region IX	California	4	Los Angeles	50
Region X	Washington	24	Seattle	45

¹ States ranked #5-6, 8, 10-12, 14-15, 17-23, 25-30 are within already selected federal regions and, therefore, were not selected.

² Cities ranked 3, 5, 7-16, 18-21, 23-42, 44, 46-48 are within already selected regions

Table 2

Data Collected by Topic

Topic – Part 1	Specifics
Demographics	Gender, age, race, ZIP code
Family	Who Respondent lived with (in the last 7 days, # of siblings, birth order, # times ate meals with family (in last 7 days)
Education	Grade in school, grades
Income	Employment, # of hours worked, cash on hand, debit card
Activity Level	# of sports teams, # hours watching tv, # hours playing video games, bedtime, waking time
Health	Perceived health
Conduct	# of tickets, arrests, suspensions
Behavioral Problems	Selected conduct disorder, anxiety, depression, anorexia symptoms
ADD/ADHD	Did a doctor diagnose ADD/ADHD, five symptoms
Brand Level Data	# of youth able to identify drugs by name, source of information
Topic – Part 2	Specifics
Prescription Stimulants	Use, misuse, abuse and/or diversion, use of more than one prescription stimulant, route of administration, source, reasons for use, likelihood of taking Rx Stimulant in one and five years, dependence symptoms, perception of how big a problem prescription stimulants are with kids, Respondents' age
Prescription Sedatives	Use and misuse
Prescription Opioids	Use and misuse
Illegal drugs	Use of marijuana, cocaine, crack, heroin, club drugs (like ecstasy), hallucinogens (like LSD or mushrooms), anabolic steroids, cough syrup/ "purple drank" to get high, methamphetamine, inhalants (like gasoline or paint)
Energy drink	Use in last 7 days
Tobacco	Use, onset, # cigarettes smoked on days when Respondent smokes
Alcohol	Use, onset, # of drinks in last 7 days, # days drank in last 30 days, binge drinking, source
Gambling	Internet, poker games and sports bets, ever played Second Life
Friends	# of close friends, # of close friends Respondent thinks has tried Adderall, even once
Poly drugs	Within and between prescription drugs and illicit drugs
Truthfulness	How truthful Respondents were in answering questions on the survey (not at all, somewhat or completely)
Qualitative Data	How should kids your age be told about prescriptions drugs and their effects? If you ran the world, how would you stop kids from taking others people's prescription medicines? Why do people use prescription stimulants without a prescription?
Height & Weight	Self-reported
Accompanied by Parent	Yes or No

Table 3
Descriptive Characteristics of N-MAPSS Sample Residential Area and Age (N = 11,048)

	Rural						Suburban						Urban						Total (N = 11048)	
	10 to 12 (N = 211)	13 to 15 (N = 622)	16 to 18 (N = 836)	10 to 12 (N = 553)	13 to 15 (N = 1635)	16 to 18 (N = 1935)	10 to 12 (N = 642)	13 to 15 (N = 2126)	16 to 18 (N = 2488)	N	%	N	%	N	%	N	%	N		%
Gender (Male)	110	52.1	265	42.6	410	49.0	238	43.0	730	44.7	980	50.7	313	48.8	977	46.0	1260	50.6	5283	47.8
Race/Ethnicity																				
Caucasian	120	57.7	441	71.2	603	72.6	234	42.8	805	49.4	885	45.8	185	29.0	662	31.4	780	31.5	4715	42.9
African American	25	12.0	44	7.1	69	8.3	102	18.6	207	12.7	292	15.1	228	35.7	478	22.7	741	30.0	2186	19.9
Asian	7	3.4	29	4.7	21	2.5	57	10.4	130	8.0	183	9.5	41	6.4	195	9.2	192	7.8	855	7.8
Hispanic/Latino	25	12.0	52	8.4	72	8.7	101	18.5	292	17.9	357	18.5	125	19.6	502	23.8	500	20.2	2026	18.4
Other	31	14.9	53	8.6	66	7.9	53	9.7	195	12.0	214	11.1	60	9.4	273	12.9	260	10.5	1205	11.0
Live w/Mom and Dad	145	68.7	411	66.1	514	61.6	345	62.4	951	58.2	1068	55.2	341	53.1	1167	54.9	1194	48.0	6136	55.6
Level of general health																				
Excellent	109	51.7	286	46.1	378	45.2	259	46.9	736	45.2	782	40.5	322	50.2	953	44.9	991	39.9	4816	43.7
Good	84	39.8	281	45.3	375	44.9	251	45.5	718	44.1	924	47.8	283	44.1	951	44.8	1140	45.9	5007	45.4
Fair/Poor	18	8.5	53	8.5	83	9.9	42	7.6	174	10.7	226	11.7	37	5.8	219	10.3	353	14.2	1205	10.9
Has a job	6	2.8	84	13.5	427	51.1	36	6.5	229	14.0	810	41.9	38	5.9	320	15.1	950	38.3	2900	26.3
Has a debit card	7	3.3	68	11.0	376	45.0	20	3.6	182	11.2	765	39.6	26	4.1	262	12.4	943	38.1	2649	24.1
Think RxStim are moderate to large problem	93	44.7	399	64.8	582	70.4	245	44.7	984	60.7	1341	70.0	320	50.6	1319	62.9	1617	65.7	6900	63.2
Ever drank alcohol	7	3.3	242	39.2	549	66.4	40	7.3	624	38.5	1236	64.3	37	5.8	770	36.6	1461	59.2	4966	45.3
Lifetime Rx Stimulant Use	16	7.6	94	15.1	206	24.8	56	10.1	202	12.4	374	19.5	55	8.6	230	10.9	399	16.1	1632	14.8

Table 4
N-MAPSS Prevalence of Use, Misuse, and Diversion of Rx Stimulants by Residential Area and Age

	Rural						Suburban						Urban						Total (N = 11048)	
	10 to 12 (N = 211)		13 to 15 (N = 622)		16 to 18 (N = 836)		10 to 12 (N = 553)		13 to 15 (N = 1635)		16 to 18 (N = 1935)		10 to 12 (N = 642)		13 to 15 (N = 2126)		16 to 18 (N = 2488)			
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%		
RxStim Use in the Past 30 Days																				
No RxStim Use	194	94.6	527	92.0	626	87.2	497	94.1	1425	93.4	1548	90.2	582	94.3	1884	95.6	2080	92.6	9363	92.7
Nonmedical Use Only	0	0.0	17	3.0	38	5.3	3	0.6	22	1.4	77	4.5	9	1.5	27	1.4	81	3.6	274	2.7
Medical Use Only	10	4.9	24	4.2	41	5.7	27	5.1	69	4.5	57	3.3	23	3.7	45	2.3	50	2.2	346	3.4
Both	1	0.5	5	0.9	13	1.8	1	0.2	10	0.7	35	2.0	3	0.5	14	0.7	36	1.6	118	1.2
Approached to Divert ¹																				
Asked to Sell RxStim	2	1.0	66	10.7	152	18.4	42	7.6	153	9.4	324	16.9	24	3.8	172	8.2	348	14.1	1283	11.7
Asked to Give RxStim	1	0.5	39	6.3	85	10.3	31	5.6	79	4.9	204	10.6	16	2.5	100	4.7	181	7.3	736	6.7
Asked to Trade RxStim	2	1.0	53	8.6	120	14.5	29	5.3	109	6.7	241	12.6	20	3.2	112	5.3	273	11.1	959	8.8
	2	1.0	28	4.5	63	7.6	22	4.0	45	2.8	131	6.8	10	1.6	54	2.6	153	6.2	508	4.6
Outgoing Diversion ¹																				
Sold RxStim	0	0.0	16	2.6	49	5.9	5	0.9	26	1.6	94	4.9	4	0.6	32	1.5	102	4.1	328	3.0
Gave RxStim	1	0.5	25	4.1	61	7.4	15	2.7	42	2.6	137	7.1	10	1.6	61	2.9	147	6.0	499	4.6
Trade RxStim	0	0.0	14	2.3	45	5.4	6	1.1	21	1.3	78	4.1	2	0.3	20	0.9	77	3.1	263	2.4
Had RxStim stolen	1	0.5	17	2.8	24	2.9	11	2.0	18	1.1	39	2.0	5	0.8	22	1.0	43	1.7	180	1.6
Incoming Diversion ¹																				
Stolen RxStim	5	2.4	48	7.8	118	14.2	27	4.9	101	6.2	266	13.8	15	2.4	140	6.6	287	11.6	1007	9.2
Got RxStim for Free	0	0.0	14	2.3	29	3.5	5	0.9	25	1.5	62	3.2	4	0.6	22	1.0	69	2.8	230	2.1
Borrowed RxStim	2	1.0	27	4.4	99	12.0	24	4.3	65	4.0	217	11.3	10	1.6	90	4.3	235	9.5	769	7.0
	3	1.4	27	4.4	47	5.7	10	1.8	48	3.0	115	6.0	11	1.7	80	3.8	123	5.0	464	4.2
Lifetime Diversion ²																				
None	203	97.1	552	89.5	684	82.6	512	92.8	1492	92.0	1601	83.3	604	96.2	1932	91.7	2120	85.9	9700	88.6
Incoming Diversion Only	4	1.9	25	4.1	55	6.6	18	3.3	59	3.6	136	7.1	9	1.4	85	4.0	148	6.0	539	4.9
Outgoing Diversion Only	1	0.5	17	2.8	26	3.1	13	2.4	28	1.7	54	2.8	9	1.4	35	1.7	62	2.5	245	2.2

	Rural						Suburban						Urban						Total (N = 11048)	
	10 to 12 (N = 211)		13 to 15 (N = 622)		16 to 18 (N = 836)		10 to 12 (N = 553)		13 to 15 (N = 1635)		16 to 18 (N = 1935)		10 to 12 (N = 642)		13 to 15 (N = 2126)		16 to 18 (N = 2488)			
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%		
Both Incoming and Outgoing Diversion	1	0.5	23	3.7	63	7.6	9	1.6	42	2.6	130	6.8	6	1.0	55	2.6	139	5.6	468	4.3

Note

¹ Any type;

² Incoming or outgoing