Published in final edited form as:

Front Biosci (Landmark Ed). ; 23: 1552–1563.

Lipid droplet autophagy during energy mobilization, lipid homeostasis and protein quality control

Enrique J. Garcia^{*}, Jason D. Vevea[†], and Liza A. Pon^{*}

^{*}Department of Pathology and Cell Biology, Columbia University, New York, NY, 10032 USA

Abstract

Lipid droplets (LDs) have well-established functions as sites for lipid storage and energy mobilization to meet the metabolic demands of cells. However, recent studies have expanded the roles of LDs to protein quality control. Lipophagy, or LD degradation by autophagy, plays a vital role not only in the mobilization of free fatty acids (FFAs) and lipid homeostasis at LDs, but also in the adaptation of cells to certain forms of stress including lipid imbalance. Recent studies have provided new mechanistic insights about the diverse types of lipophagy, in particular microlipophagy. This review summarizes key findings about the mechanisms and functions of lipophagy and highlights a novel function of LD microlipophagy as a mechanism to maintain endoplasmic reticulum (ER) proteostasis under conditions of lipid imbalance.

Keywords

Lipid Droplets; Autophagy; Macrolipophagy; Microlipophagy; Lipophagy; ER Quality cControl; Proteostasis; ATG core proteins; ESCRT; Lipid Imbalance; Lipid Homeostasis

2. INTRODUCTION

Lipid droplets (LDs) are present from bacteria to metazoans (1–6). In eukaryotes, they are produced at the endoplasmic reticulum (ER) (7, 8). The boundary membrane of LDs is a single phospholipid monolayer, which contains conserved integral and peripheral membrane proteins and envelops a hydrophobic core composed primarily of triacylglycerol (TAG) and sterol esters (SE) (9). LDs are sites of lipid storage and mobilization for energy homeostasis and membrane biogenesis (10). They also store hydrophobic vitamins and lipid signaling molecules, and sequester toxic lipids. Finally, emerging evidence support a role for LDs in protein storage, assembly, and quality control (10, 11).

To use lipids and other compounds stored in LDs, cells have developed mechanisms to transfer LD contents to other compartments. Lipolysis is the best-characterized pathway for release of lipids from LDs. In response to depletion of extracellular free fatty acids (FFA) or the need for mobilization of energy stores, protein kinases including cAMP-dependent protein kinase (PKA) and 5-AMP activated protein kinase (AMPK) are activated by

Correspondence: Liza A. Pon, Ph.D., Department of Pathology and Cell Biology, Columbia University, 630 W. 168th St. P&S 14-442, New York, NY 10032, lap5@cumc.columbia.edu.

[†]current address: HHMI and Dept. of Neuroscience, University of Wisconsin, Madison, WI, 53705 USA

hormonal cues (12). This results in the phosphorylation of cytosolic neutral lipid lipases (adipose triglyceride lipase, ATGL and hormone sensitive lipase, HSL), and of perilipins (PLIN), a conserved family of proteins that serve as scaffolds and regulators on the LD surface (13). ATGL and HSL are then recruited to PLIN proteins on LDs where they catalyze the breakdown of neutral lipids within LDs to generate FFAs that are released from LDs (14).

Other studies indicate that the ubiquitin-proteasome system (UPS) remodels LDs and regulates LD activity. Specifically, when lipid availability is limiting and LD abundance is low, LD proteins including PLIN1 and PLIN2, ATGL and its inhibitor GOS2, and the LD fusion protein FSP27/CIDEA are degraded by the UPS (15–21). Conversely, induction of LD biogenesis (e.g. by growth of cells with oleate) results in stabilization of these proteins. While the UPS can control LD protein composition and regulate TAG lipolysis in LD, the mechanisms that target LD proteins for UPS-mediated degradation are not well understood.

Finally, emerging studies indicate that breakdown of LD constituents and LDs can occur by autophagy in response to cellular and environmental cues, and that this affects LD function in energy, lipid homeostasis, and ER protein quality control. Here, we discuss the process, mechanism, and consequences of LD autophagy, generally called lipophagy.

3. LIPOPHAGY: LD AUTOPHAGY

While the autophagy of various organelles was described as early as the 1960s, it was only recently that autophagy was implicated in the degradation of LDs (22). There are three main types of autophagy: macroautophagy, microautophagy, and chaperone-mediated autophagy (CMA). As described below, all three forms of autophagy function in LD degradation.

3.1. Macrolipophagy

Macroautophagy is defined by the formation of an autophagosome, a double-membrane structure that surrounds and sequesters cytosolic components, and mediates fusion of the enveloped cargo with the lysosome (or vacuole in yeast) for degradation (Fig. 1) (23). LD macrolipophagy was first described in mouse hepatocytes under conditions of starvation and lipid exposure (22). These studies revealed that exposure of hepatocytes to serum starvation results in co-localization of autophagosome and lysosome marker proteins with LDs and conversion of LC3-I to LC3-II, a vital step for initiating and lengthening autophagosomal membranes, at LDs. These studies also revealed that inhibition of macroautophagy, pharmacologically or by silencing of the autophagy machinery (ATG genes), leads to the accumulation of TAGs and LDs in serum-starved hepatocytes (22). Thus, nutrient limitation triggers LD degradation by macroautophagy in hepatocytes.

RAB7 and RAB10, members of the Rab GTPase family, have emerged as mediators of macrolipophagy in hepatocytes. RAB7 was originally identified as a regulator of late endosomes and autophagosome maturation (24–26). RAB10 has multiple roles in vesicular and membrane trafficking in adipocytes, epithelial cells and developing axons (27–29). Recent studies indicate that RAB7 and RAB10 accumulate on LDs in response to nutrient limitations and that activation of RAB7 and RAB10 is required to recruit autophagosome

and lysosomal marker proteins to LDs. Conversely, depletion of either protein leads to LD accumulation in hepatocytes during starvation conditions (30, 31). Overall, these findings support the model that RAB7 and RAB10 play a role in the priming or licensing of LDs for macrolipophagy.

Recent findings also support a role for PLIN1 and PLIN2 in regulating LD macroautophagy. Stimulation of β -adrenergic receptors in 3T3-L1 cells activates PKA, and promotes recruitment of RAB7 to LDs and lipophagy. Knock-down of PLIN1 inhibits association of RAB7 and lysosomal marker proteins with LDs in activated 3T3-L1 cells (32). Similarly, PLIN2 deficiency reduces LD levels and enhances autophagic lipolysis in mouse liver and cultured hepatoma cells (33). In contrast, overexpression of PLIN2 protects hepatic lipid droplets from autophagic lipolysis. Thus, PLIN1 and PLIN2 may function as physical barriers that prevent recruitment of the macroautophagy machinery to LDs. Since PLIN1 and PLIN2 are targets for UPS degradation, it is possible that the UPS regulates LD function through effects on macrolipophagy. For a more extensive recent review of macrolipophagy see Schulze et al. (34).

3.2. CMA and Lipophagy

CMA is a specialized form of autophagy that targets cytosolic proteins bearing the KFERQ degradation pentapeptide signal for lysosomal degradation (Fig. 1) (35). Although CMA exclusively targets proteins and not lipids, several findings support a role for CMA in LD homeostasis. CMA is triggered by starvation, and mice with defects in liver CMA develop marked hepatosteatosis (36, 37). Moreover, PLIN2 and PLIN3 contain the KFERQ degradation pentapeptide and are CMA substrates (37). CMA-dependent degradation of PLIN2 and PLIN3 at the LD surface promotes the recruitment of a neutral lipid lipase (ATGL) and macroautophagy machinery components to the LD. Conversely, impairment of CMA or mutations in the pentapeptide signal of PLIN2 or PLIN3 prevents association of ATG core proteins with LDs and promotes LD accumulation (37). Overall, these findings indicate that CMA-mediated degradation of PLIN2 and PLIN3 stimulates lipolysis and macrolipophagy by promoting the recruitment of lipases and macrolipophagy machinery components to LDs.

3.3. Microlipophagy

In yeast, LDs are also degraded by microautophagy. In microlipophagy, LDs are not enveloped by autophagosomes, but are delivered directly to the vacuole (the yeast lysosome). Microlipophagy has been detected in yeast in response to nitrogen starvation, glucose depletion, survival during stationary phase, and phospholipid imbalance. Under these conditions, LDs are taken up into the vacuole at sites of vacuolar membrane invagination (Fig. 1) (38–41). However, different environmental conditions stimulate different mechanisms for microlipophagy.

3.3.1. LD-vacuolar contact sites during ATG core gene dependent

microlipophagy—In nitrogen starvation- and stationary phase-induced microlipophagy, LDs bind to specific sterol-rich, raft-like liquid ordered (L_o) vacuolar microdomains (38, 39). Recent studies support a role for two essential Niemann-Pick type C (NPC) sterol

transporter proteins, Ncr1p and Npc2p, in L_o microdomain formation under both conditions (42). These proteins are internalized into the vacuolar lumen, where they transfer sterols to the vacuolar membrane. If these proteins are absent or fail to localize to the vacuolar lumen, the formation and expansion of L_o microdomains needed for microlipophagy is severely affected.

Interestingly, the source for sterols in L_o microdomains differs in yeast in stationary phase and undergoing nitrogen starvation. Sterol-rich intraluminal vesicles (ILVs) are the main source of L_o microdomain sterols in nitrogen starved yeast (42). These ILVs are transported to the vacuole by multivesicular bodies (MVBs), membrane bound structures that are generated from late endosomes, contain internal membranes and deliver proteins and lipids to the vacuole (43). In contrast, sterol esters in LD are the sterol source in stationary phase yeast cells (39). Interestingly, there are reciprocal interactions between microlipophagy and these sterol-rich microdomains in stationary phase yeast. Vacuolar microdomains are required for microlipophagy (39). Conversely, microlipophagy is essential to maintain the L_o microdomains, potentially because LDs that are taken up into the vacuole by microlipophagy are a major source for sterol esters for vacuolar L_o microdomains (44).

Finally, core ATG genes, which encode proteins that mediate autophagosome formation, are required for microlipophagy in both stationary phase and nitrogen starved yeast (38, 39, 42). However, the mechanism of ATG gene function varies under these two conditions. ATG genes are required for localization of NPC proteins to the vacuole in stationary phase cells. Specifically, deletion of *ATG1*, 2, 3, 5, 7, 8, or 18 results in accumulation of Ncr1p and Npc2p as punctate, presumably aggregated structures in the cytosol (42). Surprisingly, under conditions of nitrogen starvation, localization of Ncr1p, Npc2p, or ILVs to the vacuole is not affected in ATG mutants. Rather, ATG genes are required for autophagosome-mediated delivery of sphingolipids, another vital component of the L_o microdomains, to the vacuole (45, 46).

Currently, it is not clear whether Ncr1p, Npc2p, or ILV-dependent sterols are required for microlipophagy under conditions of acute glucose depletion. However, microlipophagy in glucose-depleted yeast requires ATG core genes including Atg14p, an autophagy-specific subunit of phosphatidylinositol 3-kinase complex I that localizes to L_o microdomains (41). Moreover, *ATG14* is required for L_o formation and microlipophagy, and localization of Atg14p to L_o sites is dependent upon the AMPK pathway for energy sensing during acute glucose depletion. The defect in microlipophagy observed in glucose-depleted *atg14* cells does not appear to be due to effects on macroautophagy: *ATG14* is required for macroautophagy in fed yeast cells but not in yeast undergoing acute glucose depletion (41). This finding suggests an unexpected plasticity in autophagy in response to different cellular cues and a novel function of Atg14p in microlipophagy.

3.3.2. ATG core gene-independent forms of microlipophagy—Microlipophagy that does not require core ATG genes has been observed in yeast under two conditions: lipid imbalance produced by defects in PC biosynthesis (47), and transition through the diauxic shift, the shift from glycolysis- to respiration-driven growth when glucose becomes limiting (48). Since core ATG genes are required for L_o microdomain formation (42), it is possible

that LD uptake into vacuoles does not occur at L_o microdomains in ATG-independent forms of microautophagy. Alternatively, L_o microdomains may function in all forms of microlipophagy; however, generation of those subdomains may occur by different mechanisms.

Interestingly, both forms of ATG gene-independent microlipophagy require the endosome sorting complexes required for transport (ESCRT) machinery, conserved membrane complexes that mediate invagination of late endosomal membranes to produce MVBs (49). Since MVBs transport hydrolases and other macromolecules to the vacuole, ESCRT affects micro- and macrolipophagy through effects on vacuolar biogenesis. However, emerging evidence supports a direct role for the ESCRT in vacuolar membrane remodeling in ATG gene-independent microlipophagy.

Lipid imbalance-induced microlipophagy requires Vps4p, which mediates ESCRT III disassembly, and Esm1p/Atg39p, a newly identified class V ESCRT protein that has also been implicated in nucleophagy (40, 50). Microautophagy of cargos including LDs in post-diauxic yeast requires multiple ESCRT components, including Vps27p, a clathrin- and ubiquitin-binding protein and component of the ESCRT-0 complex normally responsible for initiating the MVB pathway (49). Indeed, passage through the diauxic shift results in relocalization of Vps27p from endosomes to punctate structures on the vacuolar membrane. The clathrin-binding domain of Vps27p is not required for its relocalization to the vacuole after the diauxic shift. However, it is required for efficient microlipophagy under these conditions. Other studies confirm that ESCRT can function directly on the vacuolar membrane to sort and direct polyubiquitinated protein cargo into the vacuolar lumen for degradation (51). These findings raise the possibility that ESCRT mediates invagination of the vacuolar membrane for LD uptake during microlipophagy in yeast after the diauxic shift, and potentially also in yeast exposed to lipid stress.

4. FUNCTIONS OF LIPOPHAGY

4.1. Energy mobilization

A role for lipophagy in release of free fatty acids (FFAs) from LDs for energy mobilization in response to starvation was first identified in mouse hepatocytes (22). Other studies revealed a role for lipophagy in starvation-induced FFA release in the proximal tubule of the kidney (52), hypothalamic and primary striatal neurons, glial cells, lymphocytes, macrophages, cultured adipocytes, gastrointestinal epithelial cells, and prostate cancer cells (53). Lipophagy in response to nutrient limitation is also observed in the yeast *Saccharomyces cerevisiae*, the nematode *Caenorhabditis elegans*, and the fungus *Fusarium graminearum* (53).

Finally, lipophagy can be activated in brown adipose tissue by cold shock, where it promotes the release of FFAs from brown fat to supply peripheral tissues with energy and promote their survival during cold stress (54). Thus, lipophagy is an established mechanism to supply energy to cells under various metabolic demands.

As described above, starvation-induced LD degradation occurs by macroautophagy in mouse liver (22). In contrast, yeast subjected to nitrogen starvation or acute glucose limitation degrade LDs by microlipophagy (38, 41). Since deletion of cytosolic neutral lipid lipases does not have severe effects on survival of yeast in stationary phase, microlipophagy may also be a major pathway for TAG metabolism during that stage of the yeast life cycle (55–57). Yet to be determined is why different cell types use different mechanisms for microlipophagy.

Although lipolysis can be used to release FFAs from LDs, lipophagy may be activated during starvation in some cell types because it allows for rapid release of FFAs. Moreover, use of lipophagy as opposed to lipolysis for energy mobilization from LDs varies among tissues. Lipolysis is the primary mechanism for FFA release from LDs in white adipose tissue, which expresses high levels of HSL and ATGL (12, 58–61). In contrast, lipophagy appears to be the major mechanism for FFA mobilization in liver, which expresses low levels of HSL and ATGL (13). Elevated FFA results in lipotoxicity, elevated reactive oxygen species production, uncoupling of mitochondria, apoptosis, and insulin resistance (62–65). Therefore, use of the relatively slower process of lipolysis for FFA release in white adipose tissue may protect against release of excessive levels of FFA from the large stores of TAGs in those cells.

4.2. Lipid homeostasis

The function of lipophagy in lipid regulation extends beyond TAG storage and FFA release. Lipophagy is a mechanism to protect cells from excess intracellular lipids and lipotoxicity. Up-regulation of macrolipophagy was first detected in hepatocytes exposed to high levels of oleic acid (22). Subsequent studies showed that lipid overload can increase autophagy in cells including neurons, beta cells, myocytes and epithelial cells (66–70). Conversely, blocking autophagy in cultured myocytes leads to elevated TG levels and cell death as a result of lipid overload (71). Thus, lipophagy protects cells from lipotoxicity by removal of excess intracellular lipids.

Lipophagy also protects yeast from lipid stress produced by inhibition of PC biosynthesis (40). The immediate response to inhibition of PC biosynthesis is altered levels of phospholipids, decreased growth rates, and defects in the morphology and distribution of mitochondria and ER, the two organelles where PC biosynthesis occurs. Surprisingly, yeast adapt to this lipid stress: their growth rates, ER and mitochondrial morphology and distribution, and levels of many phospholipids are restored (40). We found that TAG and sterol ester levels as well as LD biogenesis and degradation by microlipophagy are increased in cells that have adapted to this lipid stress. Moreover, we find that mutations that block TAG and LD biosynthesis or microlipophagy severely inhibit adaptation. These studies support the model that excess lipids are stored in LDs in the form of TAG and degradation of those lipids by microlipophagy protects yeast cells from lipid imbalance.

Finally, lipophagy has also been implicated in protecting cells from excess cholesterol. Reverse cholesterol transport, transport of free cholesterol from macrophage foam cells in atherosclerotic plaques to the liver where it is excreted in bile, is one mechanism to reduce cholesterol levels (72). Recent studies indicate that lysosomes contribute to lipid breakdown

in lipid-loaded macrophage foam cells (73). These studies also revealed co-localization of autophagosome and lysosome marker proteins with LDs, conversion of LC3-I to LC3-II in subcellular fractions enriched in LDs, and a role for lysosomal acid lipases in release of free cholesterol from cholesterol esters in LDs in macrophage foam cells. Thus, LD breakdown in lipid-loaded macrophage foam cells occurs by macroautophagy. Finally, these studies showed that lipophagy is required for release of free cholesterol from cholesterol esters in LDs and reverse cholesterol transport from lipid-loaded macrophage foam cells in culture and in whole animals (73). Thus, lipophagy may contribute to clearance of excess cholesterol from peripheral tissues and be a target for treatment of atherosclerosis (74).

4.3. Protein quality control

LDs have also emerged as cellular “protein sinks.” LDs sequester histones H2A and H2B during oogenesis in *Drosophila* (75). Defects in this process results in toxic accumulation of histones, which leads to mitotic error and apoptosis (76, 77). LDs also sequester misfolded cytosolic proteins and prevent them from forming toxic aggregates under several pathological states and conditions (78, 79). Finally, LDs have been implicated in the assembly of hepatitis C virus (HCV) and in liver steatosis caused by HCV infection. Specifically, two HCV core proteins localize to LDs where they contribute to transfer of lipids from LDs to the developing virion during virus assembly (80–82).

Our group has recently identified a novel role for LD and lipophagy in ER proteostasis under conditions of lipid imbalance. Lipid imbalance produced by defects in PC biosynthesis results in accumulation of unfolded proteins in the ER (ER stress) and activation of the unfolded protein response (UPR) (40), a signal transduction pathway that down-regulates protein synthesis and up-regulates proteostasis mechanisms in response to ER stress (83). One pathway that is activated by the UPR is ER-associated protein degradation (ERAD), a pathway for retrotranslocation of unfolded ER proteins to the surface of the organelle, where they are ubiquitinated and targeted for degradation by the proteasome (84, 85).

LD biogenesis and microautophagy have emerged as an ERAD-independent mechanism for ER proteostasis in yeast undergoing acute lipid imbalance. Specifically, polyubiquitinated proteins and Kar2p, the yeast homologue of the ER chaperone BiP, are enriched in LDs isolated from yeast undergoing lipid stress (40). Furthermore, these LDs are targeted to the vacuole for degradation by microautophagy. As described above, lipid stress induced microlipophagy does not require the core autophagy gene *ATG7*, but does require ESCRT components. Finally, microlipophagy is essential for yeast cell survival during lipid stress (40).

These findings support a novel mechanism for ER proteostasis in response to lipid stress whereby LDs remove unfolded proteins from the ER and target them for degradation by lipophagy. They also provide support for the model that LDs function as “escape hatches” in ER quality control (86). Interestingly, LDs may also remove damaged proteins from mitochondria in yeast (87). Under conditions of stress, specific damaged mitochondrial outer membrane proteins associate with LDs, which in turn are degraded by lipophagy. Therefore, LDs may serve as a vehicle for segregation and degradation of damaged proteins from different compartments.

5. CONCLUSION

It is now clear that LD autophagy occurs and contributes to established and newly identified LD functions in energy and lipid homeostasis and ER proteostasis. However, there are fundamental questions that remain unresolved. Indeed, lysosomes can be 100 to 1000 times smaller than LDs in mammalian cells. In light of this, it is not clear how LDs can be taken up into lysosomes. Are there mechanisms for fragmentation of LDs that are targeted for autophagic degradation?

Studies from multiple groups revealed that macrolipophagy and microlipophagy are used to release of FFAs from LDs. However, why different cell types use different autophagy mechanisms and how those mechanisms are regulated are not well understood. Indeed, although macroautophagy has been studied extensively, our understanding of microautophagy and microlipophagy are limited. Nonetheless, available evidence indicates that microlipophagy is complex. There appears to be multiple forms of microlipophagy in yeast that are induced by different conditions, rely upon or do not rely upon core ATG genes, and utilize different mechanisms for generating docking sites for LDs on the vacuole. Future studies will reveal the precise mechanism underlying these pathways, how they are regulated and whether they are conserved in other cell types.

Finally, recent studies on microlipophagy have revealed novel mechanisms including a possible role for the ESCRT machinery in invagination of the vacuolar membrane for LD uptake and a role for LD biogenesis and microlipophagy in ER proteostasis. Ongoing and future studies will reveal whether ESCRT mediates lysosomal membrane invagination during microlipophagy in metazoans, other possible targets for ESCRT function in membrane curvature, whether LDs regulate ER proteins quality control in metazoans, the targets for ER proteostasis by LDs, and whether LDs control the quality of other organelles and cellular compartments.

Acknowledgments

We thank the members of the Pon laboratory for valuable discussion and critical review of the manuscript. This work was supported by grants from the National Institutes of Health (NIH) (GM45735, GM122589 and AG051047) to LP and (GM007367 and AR070013) to EJG, and from the HHMI (56006760) and the NIH (NS098604) to JDV.

Abbreviations

LD	lipid droplet
FFA	free fatty acid
ER	endoplasmic reticulum
TAG	triacylglycerol
SE	sterol esters
AMPK	5-AMP activated protein kinase
PKA	cAMP-dependent protein kinase

ATGL	adipose triglyceride lipase
HSL	hormone sensitive lipase
UPS	ubiquitin-proteasome system
CMA	chaperone mediated autophagy
PC	phosphatidylcholine
NPC	Niemann-Pick type C
ILV	intraluminal vesicle
MVB	multivesicular bodies
ESCRT	endosome sorting complexes required for transport
HCV	hepatitis C virus
UPR	unfolded protein response
ERAD	endoplasmic reticulum associated degradation

References

1. Waltermann M, Steinbuchel A. Neutral lipid bodies in prokaryotes: recent insights into structure, formation, and relationship to eukaryotic lipid depots. *J Bacteriol.* 2005; 187(11):3607–19. DOI: 10.1128/JB.187.11.3607-3619.2005 [PubMed: 15901682]
2. Tzen J, Cao Y, Laurent P, Ratnayake C, Huang A. Lipids, Proteins, and Structure of Seed Oil Bodies from Diverse Species. *Plant Physiol.* 1993; 101(1):267–276. [PubMed: 12231682]
3. Murphy DJ. The biogenesis and functions of lipid bodies in animals, plants and microorganisms. *Prog Lipid Res.* 2001; 40(5):325–438. [PubMed: 11470496]
4. Kohlwein SD, Veenhuis M, van der Klei IJ. Lipid droplets and peroxisomes: key players in cellular lipid homeostasis or a matter of fat--store 'em up or burn 'em down. *Genetics.* 2013; 193(1):1–50. DOI: 10.1534/genetics.112.143362 [PubMed: 23275493]
5. Murphy DJ, Vance J. Mechanisms of lipid-body formation. *Trends Biochem Sci.* 1999; 24(3):109–15. [PubMed: 10203758]
6. Yang L, Ding Y, Chen Y, Zhang S, Huo C, Wang Y, Yu J, Zhang P, Na H, Zhang H, Ma Y, Liu P. The proteomics of lipid droplets: structure, dynamics, and functions of the organelle conserved from bacteria to humans. *J Lipid Res.* 2012; 53(7):1245–53. DOI: 10.1194/jlr.R024117 [PubMed: 22534641]
7. Jacquier N, Choudhary V, Mari M, Toulmay A, Reggiori F, Schneiter R. Lipid droplets are functionally connected to the endoplasmic reticulum in *Saccharomyces cerevisiae*. *J Cell Sci.* 2011; 124(Pt 14):2424–37. DOI: 10.1242/jcs.076836 [PubMed: 21693588]
8. Ohsaki Y, Cheng J, Suzuki M, Fujita A, Fujimoto T. Lipid droplets are arrested in the ER membrane by tight binding of lipidated apolipoprotein B-100. *J Cell Sci.* 2008; 121(Pt 14):2415–22. DOI: 10.1242/jcs.025452 [PubMed: 18577578]
9. Tauchi-Sato K, Ozeki S, Houjou T, Taguchi R, Fujimoto T. The surface of lipid droplets is a phospholipid monolayer with a unique Fatty Acid composition. *J Biol Chem.* 2002; 277(46):44507–12. DOI: 10.1074/jbc.M207712200 [PubMed: 12221100]
10. Walther TC, Farese RV Jr. Lipid droplets and cellular lipid metabolism. *Annu Rev Biochem.* 2012; 81:687–714. DOI: 10.1146/annurev-biochem-061009-102430 [PubMed: 22524315]
11. Welte MA, Gould AP. Lipid droplet functions beyond energy storage. *Biochim Biophys Acta.* 2017; 1862(10 Pt B):1260–1272. DOI: 10.1016/j.bbalip.2017.07.006 [PubMed: 28735096]

12. Lass A, Zimmermann R, Oberer M, Zechner R. Lipolysis - a highly regulated multi-enzyme complex mediates the catabolism of cellular fat stores. *Prog Lipid Res.* 2011; 50(1):14–27. DOI: 10.1016/j.plipres.2010.10.004 [PubMed: 21087632]
13. Zechner R, Madeo F, Kratky D. Cytosolic lipolysis and lipophagy: two sides of the same coin. *Nat Rev Mol Cell Biol.* 2017; 18(11):671–684. DOI: 10.1038/nrm.2017.76 [PubMed: 28852221]
14. Sztalryd C, Brasaemle DL. The perilipin family of lipid droplet proteins: Gatekeepers of intracellular lipolysis. *Biochim Biophys Acta.* 2017; 1862(10 Pt B):1221–1232. DOI: 10.1016/j.bbali.2017.07.009 [PubMed: 28754637]
15. Takahashi S, Homma K, Zhou Y, Nishimura S, Duan C, Chen J, Ahmad A, Cheatham MA, Zheng J. Susceptibility of outer hair cells to cholesterol chelator 2-hydroxypropyl-beta-cyclodextrine is prestin-dependent. *Sci Rep.* 2016; 6:21973.doi: 10.1038/srep21973 [PubMed: 26903308]
16. Olzmann JA, Richter CM, Kopito RR. Spatial regulation of UBXD8 and p97/VCP controls ATGL-mediated lipid droplet turnover. *Proc Natl Acad Sci U S A.* 2013; 110(4):1345–50. DOI: 10.1073/pnas.1213738110 [PubMed: 23297223]
17. Xu G, Sztalryd C, Lu X, Tansey JT, Gan J, Dorward H, Kimmel AR, Londos C. Post-translational regulation of adipose differentiation-related protein by the ubiquitin/proteasome pathway. *J Biol Chem.* 2005; 280(52):42841–7. DOI: 10.1074/jbc.M506569200 [PubMed: 16115879]
18. Xu G, Sztalryd C, Londos C. Degradation of perilipin is mediated through ubiquitination-proteasome pathway. *Biochim Biophys Acta.* 2006; 1761(1):83–90. DOI: 10.1016/j.bbali.2005.12.005 [PubMed: 16448845]
19. Ghosh M, Niyogi S, Bhattacharyya M, Adak M, Nayak DK, Chakrabarti S, Chakrabarti P. Ubiquitin Ligase COP1 Controls Hepatic Fat Metabolism by Targeting ATGL for Degradation. *Diabetes.* 2016; 65(12):3561–3572. DOI: 10.2337/db16-0506 [PubMed: 27658392]
20. Nian Z, Sun Z, Yu L, Toh SY, Sang J, Li P. Fat-specific protein 27 undergoes ubiquitin-dependent degradation regulated by triacylglycerol synthesis and lipid droplet formation. *J Biol Chem.* 2010; 285(13):9604–15. DOI: 10.1074/jbc.M109.043786 [PubMed: 20089860]
21. Zhang X, Heckmann BL, Xie X, Saarinen AM, Liu J. Regulation of FSP27 protein stability by AMPK and HSC70. *Am J Physiol Endocrinol Metab.* 2014; 307(11):E1047–56. DOI: 10.1152/ajpendo.00125.2014 [PubMed: 25315694]
22. Singh R, Kaushik S, Wang Y, Xiang Y, Novak I, Komatsu M, Tanaka K, Cuervo AM, Czaja MJ. Autophagy regulates lipid metabolism. *Nature.* 2009; 458(7242):1131–5. DOI: 10.1038/nature07976 [PubMed: 19339967]
23. Feng Y, He D, Yao Z, Klionsky DJ. The machinery of macroautophagy. *Cell Res.* 2014; 24(1):24–41. DOI: 10.1038/cr.2013.168 [PubMed: 24366339]
24. Feng Y, Press B, Wandinger-Ness A. Rab 7: an important regulator of late endocytic membrane traffic. *The Journal of Cell Biology.* 1995; 131(6):1435. [PubMed: 8522602]
25. Press B, Feng Y, Hoflack B, Wandinger-Ness A. Mutant Rab7 Causes the Accumulation of Cathepsin D and Cation-independent Mannose 6-Phosphate Receptor in an Early Endocytic Compartment. *The Journal of Cell Biology.* 1998; 140(5):1075. [PubMed: 9490721]
26. Gutierrez MG, Munafó DB, Berón W, Colombo MI. Rab7 is required for the normal progression of the autophagic pathway in mammalian cells. *Journal of Cell Science.* 2004; 117(13):2687.doi: 10.1242/jcs.01114 [PubMed: 15138286]
27. Sano H, Roach William G, Peck Grantley R, Fukuda M, Lienhard Gustav E. Rab10 in insulin-stimulated GLUT4 translocation. *Biochemical Journal.* 2008; 411(1):89.doi: 10.1042/BJ20071318 [PubMed: 18076383]
28. Wang T, Liu Y, Xu XH, Deng CY, Wu KY, Zhu J, Fu XQ, He M, Luo ZG. Lgl1 activation of rab10 promotes axonal membrane trafficking underlying neuronal polarization. *Dev Cell.* 2011; 21(3): 431–44. DOI: 10.1016/j.devcel.2011.07.007 [PubMed: 21856246]
29. Schuck S, Gerl MJ, Ang A, Manninen A, Keller P, Mellman I, Simons K. Rab10 is involved in basolateral transport in polarized Madin-Darby canine kidney cells. *Traffic.* 2007; 8(1):47–60. DOI: 10.1111/j.1600-0854.2006.00506.x [PubMed: 17132146]
30. Schroeder B, Schulze RJ, Weller SG, Sletten AC, Casey CA, McNiven MA. The small GTPase Rab7 as a central regulator of hepatocellular lipophagy. *Hepatology.* 2015; 61(6):1896–907. DOI: 10.1002/hep.27667 [PubMed: 25565581]

31. Li Z, Schulze RJ, Weller SG, Krueger EW, Schott MB, Zhang X, Casey CA, Liu J, Stöckli J, James DE, McNiven MA. A novel Rab10-EHBP1-EHD2 complex essential for the autophagic engulfment of lipid droplets. *Science Advances*. 2016; 2(12):e1601470.doi: 10.1126/sciadv.1601470 [PubMed: 28028537]
32. Lizaso A, Tan KT, Lee YH. beta-adrenergic receptor-stimulated lipolysis requires the RAB7-mediated autolysosomal lipid degradation. *Autophagy*. 2013; 9(8):1228–43. DOI: 10.4161/auto.24893 [PubMed: 23708524]
33. Tsai TH, Chen E, Li L, Saha P, Lee HJ, Huang LS, Shelness GS, Chan L, Chang BH. The constitutive lipid droplet protein PLIN2 regulates autophagy in liver. *Autophagy*. 2017; 13(7): 1130–1144. DOI: 10.1080/15548627.2017.1319544 [PubMed: 28548876]
34. Schulze RJ, Sathyanarayan A, Mashek DG. Breaking fat: The regulation and mechanisms of lipophagy. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids*. 2017; 1862(10 Part B):1178–1187. doi:<https://doi.org/10.1016/j.bbalip.2017.06.008>. [PubMed: 28642194]
35. Kaushik S, Cuervo AM. Chaperone-mediated autophagy: a unique way to enter the lysosome world. *Trends in Cell Biology*. 2012; 22(8):407–417. doi:<https://doi.org/10.1016/j.tcb.2012.05.006>. [PubMed: 22748206]
36. Schneider JL, Suh Y, Cuervo AM. Deficient chaperone-mediated autophagy in liver leads to metabolic dysregulation. *Cell metabolism*. 2014; 20(3):417–432. DOI: 10.1016/j.cmet.2014.06.009 [PubMed: 25043815]
37. Kaushik S, Cuervo AM. Degradation of lipid droplet-associated proteins by chaperone-mediated autophagy facilitates lipolysis. *Nat Cell Biol*. 2015; 17(6):759–70. DOI: 10.1038/ncb3166 [PubMed: 25961502]
38. van Zutphen T, Todde V, de Boer R, Kreim M, Hofbauer HF, Wolinski H, Veenhuis M, van der Klei IJ, Kohlwein SD. Lipid droplet autophagy in the yeast *Saccharomyces cerevisiae*. *Mol Biol Cell*. 2014; 25(2):290–301. DOI: 10.1091/mbc.E13-08-0448 [PubMed: 24258026]
39. Wang CW, Miao YH, Chang YS. A sterol-enriched vacuolar microdomain mediates stationary phase lipophagy in budding yeast. *J Cell Biol*. 2014; 206(3):357–66. DOI: 10.1083/jcb.201404115 [PubMed: 25070953]
40. Vevea JD, Garcia EJ, Chan RB, Zhou B, Schultz M, Di Paolo G, McCaffery JM, Pon LA. Role for Lipid Droplet Biogenesis and Microlipophagy in Adaptation to Lipid Imbalance in Yeast. *Dev Cell*. 2015; 35(5):584–99. DOI: 10.1016/j.devcel.2015.11.010 [PubMed: 26651293]
41. Seo AY, Lau P-W, Feliciano D, Sengupta P, Gros MAL, Cinquin B, Larabell CA, Lippincott-Schwartz J. AMPK and vacuole-associated Atg14p orchestrate μ -lipophagy for energy production and long-term survival under glucose starvation. *eLife*. 2017; 6:e21690.doi: 10.7554/eLife.21690 [PubMed: 28394250]
42. Tsuji T, Fujimoto M, Tatematsu T, Cheng J, Orii M, Takatori S, Fujimoto T. Niemann-Pick type C proteins promote microautophagy by expanding raft-like membrane domains in the yeast vacuole. *Elife*. 2017; 6doi: 10.7554/eLife.25960
43. Piper RC, Katzmann DJ. Biogenesis and Function of Multivesicular Bodies. *Annual review of cell and developmental biology*. 2007; 23:519–547. DOI: 10.1146/annurev.cellbio.23.090506.123319
44. Chen XJ, Wang X, Butow RA. Yeast aconitase binds and provides metabolically coupled protection to mitochondrial DNA. *Proc Natl Acad Sci U S A*. 2007; 104(34):13738–43. DOI: 10.1073/pnas.0703078104 [PubMed: 17698960]
45. Lingwood D, Simons K. Lipid rafts as a membrane-organizing principle. *Science*. 2010; 327(5961):46–50. DOI: 10.1126/science.1174621 [PubMed: 20044567]
46. Hechtberger P, Zinser E, Saf R, Hummel K, Paltauf F, Daum G. Characterization, quantification and subcellular localization of inositol-containing sphingolipids of the yeast, *Saccharomyces cerevisiae*. *Eur J Biochem*. 1994; 225(2):641–9. [PubMed: 7957179]
47. Vevea JD, Wolken DM, Swayne TC, White AB, Pon LA. Ratiometric biosensors that measure mitochondrial redox state and ATP in living yeast cells. *J Vis Exp*. 2013; (77):50633.doi: 10.3791/50633 [PubMed: 23912244]

48. Oku M, Maeda Y, Kagohashi Y, Kondo T, Yamada M, Fujimoto T, Sakai Y. Evidence for ESCRT- and clathrin-dependent microautophagy. *J Cell Biol.* 2017; 216(10):3263–3274. DOI: 10.1083/jcb.201611029 [PubMed: 28838958]
49. Schmidt O, Teis D. The ESCRT machinery. *Current Biology.* 2012; 22(4):R116–R120. DOI: 10.1016/j.cub.2012.01.028 [PubMed: 22361144]
50. Mochida K, Oikawa Y, Kimura Y, Kirisako H, Hirano H, Ohsumi Y, Nakatogawa H. Receptor-mediated selective autophagy degrades the endoplasmic reticulum and the nucleus. *Nature.* 2015; 522(7556):359–62. DOI: 10.1038/nature14506 [PubMed: 26040717]
51. Zhu L, Jorgensen JR, Li M, Chuang YS, Emr SD. ESCRTs function directly on the lysosome membrane to downregulate ubiquitinated lysosomal membrane proteins. *Elife.* 2017; 6doi: 10.7554/eLife.26403
52. Minami S, Yamamoto T, Takabatake Y, Takahashi A, Namba T, Matsuda J, Kimura T, Kaimori JY, Matsui I, Hamano T, Takeda H, Takahashi M, Izumi Y, Bamba T, Matsusaka T, Niimura F, Isaka Y. Lipophagy maintains energy homeostasis in the kidney proximal tubule during prolonged starvation. *Autophagy.* 2017; 13(10):1629–1647. DOI: 10.1080/15548627.2017.1341464 [PubMed: 28813167]
53. Martinez-Lopez N, Singh R. Autophagy and Lipid Droplets in the Liver. *Annual Review of Nutrition.* 2015; 35(1):215–237. DOI: 10.1146/annurev-nutr-071813-105336
54. Martinez-Lopez N, Garcia-Macia M, Sahu S, Athonvarangkul D, Liebling E, Merlo P, Cecconi F, Schwartz Gary J, Singh R. Autophagy in the CNS and Periphery Coordinate Lipophagy and Lipolysis in the Brown Adipose Tissue and Liver. *Cell Metabolism.* 2016; 23(1):113–127. doi:<https://doi.org/10.1016/j.cmet.2015.10.008>. [PubMed: 26698918]
55. Athenstaedt K, Daum G. Tgl4p and Tgl5p, two triacylglycerol lipases of the yeast *Saccharomyces cerevisiae* are localized to lipid particles. *J Biol Chem.* 2005; 280(45):37301–9. DOI: 10.1074/jbc.M507261200 [PubMed: 16135509]
56. Kohlwein SD. Triacylglycerol homeostasis: insights from yeast. *J Biol Chem.* 2010; 285(21):15663–7. DOI: 10.1074/jbc.R110.118356 [PubMed: 20231294]
57. Koffel R, Tiwari R, Falquet L, Schneiter R. The *Saccharomyces cerevisiae* YLL012/YEH1, YLR020/YEH2, and TGL1 genes encode a novel family of membrane-anchored lipases that are required for steryl ester hydrolysis. *Mol Cell Biol.* 2005; 25(5):1655–68. DOI: 10.1128/mcb.25.5.1655-1668.2005 [PubMed: 15713625]
58. Langin D, Dicker A, Tavernier G, Hoffstedt J, Mairal A, Ryden M, Arner E, Sicard A, Jenkins CM, Viguerie N, van Harmelen V, Gross RW, Holm C, Arner P. Adipocyte lipases and defect of lipolysis in human obesity. *Diabetes.* 2005; 54(11):3190–7. [PubMed: 16249444]
59. Schweiger M, Schreiber R, Haemmerle G, Lass A, Fledelius C, Jacobsen P, Tornqvist H, Zechner R, Zimmermann R. Adipose triglyceride lipase and hormone-sensitive lipase are the major enzymes in adipose tissue triacylglycerol catabolism. *J Biol Chem.* 2006; 281(52):40236–41. DOI: 10.1074/jbc.M608048200 [PubMed: 17074755]
60. Nielsen TS, Jessen N, Jorgensen JO, Moller N, Lund S. Dissecting adipose tissue lipolysis: molecular regulation and implications for metabolic disease. *J Mol Endocrinol.* 2014; 52(3):R199–222. DOI: 10.1530/jme-13-0277 [PubMed: 24577718]
61. Zimmermann R, Strauss JG, Haemmerle G, Schoiswohl G, Birner-Gruenberger R, Riederer M, Lass A, Neuberger G, Eisenhaber F, Hermetter A, Zechner R. Fat Mobilization in Adipose Tissue Is Promoted by Adipose Triglyceride Lipase. *Science.* 2004; 306(5700):1383. [PubMed: 15550674]
62. Brookheart RT, Michel CI, Schaffer JE. As a matter of fat. *Cell Metab.* 2009; 10(1):9–12. DOI: 10.1016/j.cmet.2009.03.011 [PubMed: 19583949]
63. Yao H-R, Liu J, Plumeri D, Cao Y-B, He T, Lin L, Li Y, Jiang Y-Y, Li J, Shang J. Lipotoxicity in HepG2 cells triggered by free fatty acids. *American Journal of Translational Research.* 2011; 3(3):284–291. [PubMed: 21654881]
64. Goldberg IJ, Trent CM, Schulze PC. Lipid metabolism and toxicity in the heart. *Cell Metab.* 2012; 15(6):805–12. DOI: 10.1016/j.cmet.2012.04.006 [PubMed: 22682221]
65. Bobulescu IA. Renal lipid metabolism and lipotoxicity. *Current opinion in nephrology and hypertension.* 2010; 19(4):393–402. DOI: 10.1097/MNH.0b013e32833aa4ac [PubMed: 20489613]

66. Jiang L, Wang W, He Q, Wu Y, Lu Z, Sun J, Liu Z, Shao Y, Wang A. Oleic acid induces apoptosis and autophagy in the treatment of Tongue Squamous cell carcinomas. *Sci Rep.* 2017; 7(1): 11277.doi: 10.1038/s41598-017-11842-5 [PubMed: 28900281]
67. Tang Y, Chen Y, Jiang H, Nie D. Short-chain fatty acids induced autophagy serves as an adaptive strategy for retarding mitochondria-mediated apoptotic cell death. *Cell Death Differ.* 2011; 18(4): 602–18. DOI: 10.1038/cdd.2010.117 [PubMed: 20930850]
68. Komiya K, Uchida T, Ueno T, Koike M, Abe H, Hirose T, Kawamori R, Uchiyama Y, Kominami E, Fujitani Y, Watada H. Free fatty acids stimulate autophagy in pancreatic beta-cells via JNK pathway. *Biochem Biophys Res Commun.* 2010; 401(4):561–7. DOI: 10.1016/j.bbrc.2010.09.101 [PubMed: 20888798]
69. Nowicki M, Serke H, Kosacka J, Muller K, Spanel-Borowski K. Oxidized low-density lipoprotein (oxLDL) induces cell death in neuroblastoma and survival autophagy in schwannoma cells. *Exp Mol Pathol.* 2010; 89(3):276–83. DOI: 10.1016/j.yexmp.2010.07.009 [PubMed: 20692253]
70. Pauloin A, Chat S, Pechoux C, Hue-Beauvais C, Droineau S, Galio L, Devinoy E, Chanut E. Oleate and linoleate stimulate degradation of beta-casein in prolactin-treated HC11 mouse mammary epithelial cells. *Cell Tissue Res.* 2010; 340(1):91–102. DOI: 10.1007/s00441-009-0926-3 [PubMed: 20182891]
71. Lam T, Harmancey R, Vasquez H, Gilbert B, Patel N, Hariharan V, Lee A, Covey M, Taegtmeier H. Reversal of intramyocellular lipid accumulation by lipophagy and a p62-mediated pathway. *Cell Death Discovery.* 2016; 2:16061.doi: 10.1038/cddiscovery.2016.61 [PubMed: 27625792]
72. Ono K. Current concept of reverse cholesterol transport and novel strategy for atheroprotection. *Journal of Cardiology.* 2012; 60(5):339–343. doi:<https://doi.org/10.1016/j.jjcc.2012.07.014>. [PubMed: 23063661]
73. Ouimet M, Franklin V, Mak E, Liao X, Tabas I, Marcel YL. Autophagy regulates cholesterol efflux from macrophage foam cells via lysosomal acid lipase. *Cell Metab.* 2011; 13(6):655–67. DOI: 10.1016/j.cmet.2011.03.023 [PubMed: 21641547]
74. Liao X, Sluimer JC, Wang Y, Subramanian M, Brown K, Pattison JS, Robbins J, Martinez J, Tabas I. Macrophage autophagy plays a protective role in advanced atherosclerosis. *Cell Metab.* 2012; 15(4):545–53. DOI: 10.1016/j.cmet.2012.01.022 [PubMed: 22445600]
75. Cermelli S, Guo Y, Gross SP, Welte MA. The lipid-droplet proteome reveals that droplets are a protein-storage depot. *Curr Biol.* 2006; 16(18):1783–95. DOI: 10.1016/j.cub.2006.07.062 [PubMed: 16979555]
76. Li Z, Johnson MR, Ke Z, Chen L, Welte MA. Drosophila lipid droplets buffer the H2Av supply to protect early embryonic development. *Curr Biol.* 2014; 24(13):1485–91. DOI: 10.1016/j.cub.2014.05.022 [PubMed: 24930966]
77. Li Z, Thiel K, Thul Peter J, Beller M, Kühnlein Ronald P, Welte Michael A. Lipid Droplets Control the Maternal Histone Supply of Drosophila Embryos. *Current Biology.* 2012; 22(22):2104–2113. doi:<https://doi.org/10.1016/j.cub.2012.09.018>. [PubMed: 23084995]
78. Cole NB, Murphy DD, Grider T, Rueter S, Brasaemle D, Nussbaum RL. Lipid droplet binding and oligomerization properties of the Parkinson's disease protein alpha-synuclein. *J Biol Chem.* 2002; 277(8):6344–52. DOI: 10.1074/jbc.M108414200 [PubMed: 11744721]
79. Ohsaki Y, Cheng J, Fujita A, Tokumoto T, Fujimoto T. Cytoplasmic lipid droplets are sites of convergence of proteasomal and autophagic degradation of apolipoprotein B. *Mol Biol Cell.* 2006; 17(6):2674–83. DOI: 10.1091/mbc.E05-07-0659 [PubMed: 16597703]
80. Barba G, Harper F, Harada T, Kohara M, Goulinet S, Matsuura Y, Eder G, Schaff Z, Chapman MJ, Miyamura T, Brechot C. Hepatitis C virus core protein shows a cytoplasmic localization and associates to cellular lipid storage droplets. *Proc Natl Acad Sci U S A.* 1997; 94(4):1200–5. [PubMed: 9037030]
81. McLauchlan J, Lemberg MK, Hope G, Martoglio B. Intramembrane proteolysis promotes trafficking of hepatitis C virus core protein to lipid droplets. *EMBO J.* 2002; 21(15):3980–8. DOI: 10.1093/emboj/cdf414 [PubMed: 12145199]
82. Shi ST, Polyak SJ, Tu H, Taylor DR, Gretch DR, Lai MM. Hepatitis C virus NS5A colocalizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology.* 2002; 292(2): 198–210. DOI: 10.1006/viro.2001.1225 [PubMed: 11878923]

83. Thibault G, Shui G, Kim W, McAlister GC, Ismail N, Gygi SP, Wenk MR, Ng DT. The membrane stress response buffers lethal effects of lipid disequilibrium by reprogramming the protein homeostasis network. *Mol Cell*. 2012; 48(1):16–27. DOI: 10.1016/j.molcel.2012.08.016 [PubMed: 23000174]
84. Brodsky JL. Cleaning up: ER-associated degradation to the rescue. *Cell*. 2012; 151(6):1163–7. DOI: 10.1016/j.cell.2012.11.012 [PubMed: 23217703]
85. Wu H, Ng BS, Thibault G. Endoplasmic reticulum stress response in yeast and humans. *Biosci Rep*. 2014; 34(4)doi: 10.1042/BSR20140058
86. Ploegh HL. A lipid-based model for the creation of an escape hatch from the endoplasmic reticulum. *Nature*. 2007; 448(7152):435–8. DOI: 10.1038/nature06004 [PubMed: 17653186]
87. Bischof J, Salzmann M, Streubel MK, Hasek J, Geltinger F, Duschl J, Bresgen N, Briza P, Haskova D, Lejskova R, Sopjani M, Richter K, Rinnerthaler M. Clearing the outer mitochondrial membrane from harmful proteins via lipid droplets. *Cell Death Discovery*. 2017; 3:17016.doi: 10.1038/cddiscovery.2017.16 [PubMed: 28386457]

Figure 1. A schematic illustrating different forms of lipid droplet (LD) autophagy
Please refer to the text for explanation.