Published in final edited form as:

J Pediatr Surg. 2018 July ; 53(7): 1399–1402. doi:10.1016/j.jpedsurg.2017.08.049.

The Extent of Intestinal Failure-Associated Liver Disease in Patients Referred for Intestinal Rehabilitation is Associated with Increased Mortality: An Analysis of the Pediatric Intestinal Failure Consortium Database

Patrick J. Javid, M.D.^a, Assaf P. Oron, Ph.D.^a, Christopher Duggan, M.D., M.P.H.^b, Robert H. Squires, M.D.^c, and Simon P. Horslen, M.B., Ch.B.^a on behalf of On behalf of the Pediatric Intestinal Failure Consortium

^aSeattle Children's Hospital and University of Washington School of Medicine

^bBoston Children's Hospital

^cChildren's Hospital of Pittsburgh of UPMC

Abstract

Background—The advent of regional multidisciplinary intestinal rehabilitation programs has been associated with improved survival in pediatric intestinal failure. Yet, the optimal timing of referral for intestinal rehabilitation remains unknown. We hypothesized that the degree of intestinal failure-associated liver disease (IFALD) at initiation of intestinal rehabilitation would be associated with overall outcome.

Methods—The multicenter, retrospective Pediatric Intestinal Failure Consortium (PIFCon) database was used to identify all subjects with baseline bilirubin data. Conjugated bilirubin (CBili) was used as a marker for IFALD, and we stratified baseline bilirubin values as CBili <2 mg/dL, CBili 2–4 mg/dL, and CBili >4 mg/dL. The association between baseline CBili and mortality was examined using Cox proportional hazards regression.

Results—Of 272 subjects in the database, 191 (70%) children had baseline bilirubin data collected. 38% and 28% of patients had CBili >4 mg/dL and CBili <2 mg/dL, respectively, at baseline. All cause mortality was 23%. On univariate analysis, mortality was associated with CBili 2–4 mg/dL, CBili >4 mg/dL, prematurity, race, and small bowel atresia. On regression analysis controlling for age, prematurity, and diagnosis, the risk of mortality was increased by 3-fold for baseline CBili 2–4 mg/dL (HR 3.25 [1.07 – 9.92], $p=0.04$) and 4-fold for baseline CBili >4 mg/dL (HR 4.24 [1.51–11.92], $p=0.006$). On secondary analysis, CBili >4 mg/dL at baseline was associated with a lower chance of attaining enteral autonomy.

Please address correspondence to: Patrick J. Javid, M.D., Seattle Children's Hospital, 4800 Sand Point Way NE, Ocean A.9.220, Seattle, WA 98105, Phone: (206) 987-6129, Fax: (206) 987-3925, patrick.javid@seattlechildrens.org.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusion—In children with intestinal failure treated at intestinal rehabilitation programs, more advanced IFALD at referral is associated with increased mortality and decreased prospect of attaining enteral autonomy. Early referral of children with intestinal failure to intestinal rehabilitation programs should be strongly encouraged.

Keywords

Intestinal failure; short bowel syndrome; intestinal failure-associated liver disease; mortality; bilirubin; parenteral nutrition

Introduction

Long-term outcomes for the child with intestinal failure have improved significantly over the past decade. Many factors have contributed to this improved survival including the use of alternative lipid emulsions, innovative forms of bowel lengthening, and better prevention of sepsis (1–3). It has been proposed that the evolution of multi-disciplinary intestinal failure programs has made the greatest contribution to improved outcomes (4–6). Such programs emphasize long range care supervised by a team of specialists including pediatric surgeons, gastroenterologists, hepatologists, transplant surgeons, pharmacists, nutritionists, nurses and social workers.

While the role of multidisciplinary specialized care in these patients has been well described, the optimal timing of referral to an intestinal rehabilitation program is controversial. It remains common practice for some infants to be referred only when complications, such as intestinal failure-associated liver disease (IFALD) or line sepsis, arise or at the time of initial hospital discharge. Previous single center data have suggested that early referral may be beneficial when infants with intestinal failure have progressive IFALD (7).

We hypothesized that referral to an intestinal rehabilitation program early in the evolution of liver disease would be associated with improved long-term outcomes in children with intestinal failure. We chose to query the multicenter Pediatric Intestinal Failure Consortium (PIFCon) database to evaluate whether the degree of IFALD at the time of enrollment in an intestinal failure center impacted overall patient survival in a large cohort of infants with intestinal failure.

Methods

PIFCon was initiated through funding by the National Institutes of Health and National Institute of Diabetes and Digestive and Kidney Diseases in 2006 and consists of 14 centers with established multidisciplinary pediatric intestinal failure programs (8). A list of participating programs is included in Appendix A. PIFCon performed a multicenter retrospective cohort study that included all infants with intestinal failure who had received prolonged parenteral nutrition (PN) defined as the administration of PN in 60 of 74 consecutive days prior to 12 months of age. Subject study entry took place from January 1, 2000 to December 31, 2004 with study entry at one institution extended to December 31,

2005. Patient data were collected retrospectively to accrue follow-up data for at least 2 years from study entry. Data from the PIFCon database have been analyzed previously (8, 9).

After IRB approval (Seattle Children's IRB #15456), the PIFCon database was queried to identify all subjects with baseline serum bilirubin data at time of enrollment. Conjugated bilirubin (CBili) was chosen as a marker for IFALD as it represents an accurate measure of cholestatic liver disease, is not an approximated value like direct bilirubin, and has been previously studied in this population (7). In addition, the use of direct bilirubin may overestimate conjugated hyperbilirubinemia as the measurement also includes delta bilirubin. In the PIFCon dataset, baseline CBili values were available for 77 (40%) patients. For patients in which both conjugated and total bilirubin were recorded, correlation was high ($r=0.93$). A robust linear conversion formula from total to substitute-conjugated bilirubin was estimated on the square-root scale where correlation was even higher. By contrast, correlation with direct bilirubin was moderate ($r=0.56$) and offered almost no additional cases after the incorporation of total bilirubin data. Therefore only conjugated bilirubin and converted-total-bilirubin data were used in the analyses. A separate sensitivity model found that the conjugated bilirubin data generated from this conversion had no statistically significant effect on our survival analysis. Baseline bilirubin data was defined as the first serum bilirubin measurement obtained within 45 days of enrollment into the PIFCon database.

We stratified baseline bilirubin values at enrollment as CBili <2 mg/dL, CBili 2–4 mg/dL, and CBili >4 mg/dL. Association between baseline CBili and mortality was examined using a Cox proportional hazard regression model adjusting for diagnosis, prematurity, race, and clustering by hospital as well as censoring at enteral autonomy. Deaths after intestinal transplant were included in the original analysis, and an ancillary analysis was performed that censored patients at time of intestinal transplantation. A separate Cox proportional hazard regression model was created to examine the association of baseline CBili at enrollment with the overall prospect of attaining enteral autonomy; this model censored at patient death.

A final multivariable analysis was performed to explore the relationship between subjects' CBili over time after enrollment to mortality. This analysis was constructed as a time-varying Cox proportional hazard model and employed additional stratification of conjugated bilirubin values given the increased quantity of bilirubin data available over the follow-up period ($n=1164$ bilirubin values). Bilirubin values measured within 30 days of patient death were excluded from analysis.

$P < 0.05$ was used for statistical significance. Data analysis was performed using R 3.2 software.

Results

Of 272 subjects in the PIFCon database, baseline serum bilirubin data at study entry was recorded in 191 (70%) patients at mean age 63 (range 60–365) days. Baseline bilirubin data consisted of total bilirubin in 114 (60%) subjects and CBili in 77 (40%) subjects. Of

children with baseline bilirubin data, 137 (72%) presented with a CBili > 2 mg/dL, and 73 subjects (38%) presented with a CBili of 4 mg/dL or greater.

The most common diagnoses in this cohort included gastroschisis (31%), necrotizing enterocolitis (29%), small bowel atresia (13%) and midgut volvulus (13%). Gestational age data demonstrated that 146 (77%) subjects were born at less than 37 weeks, and 78 (41%) subjects were born at 32 weeks or earlier. Forty-three (23%) subjects died during the study period, and enteral autonomy was achieved in 98 (51%) of patients; these data are similar to previously published data from the entire PIFCon cohort (8). Sixty (31%) patients received an intestinal transplant, and 10 patients died following transplant. Of the post-transplant deaths, 7 patients died within the first 120 days of transplant.

All cause mortality was directly correlated with the level of CBili at time of enrollment in the intestinal failure program (Figure 1, $p=0.003$). On univariate analysis, mortality was associated with earlier gestational age, baseline bilirubin greater than 2 mg/dL, non-white race, and the diagnosis of intestinal atresia (Table I). On multivariable regression controlling for chronologic age, gestational age, and intestinal failure diagnosis, mortality was independently associated with baseline bilirubin levels between 2 mg/dL and 3.9 mg/dL (HR 3.68 [1.89 – 7.16], $p<0.001$) and greater than 4 mg/dL (HR 4.11 [1.78 – 9.48], $p=0.001$, Table II). An additional regression analysis that censored patients at time of intestinal transplantation was also performed; these data demonstrated that mortality was again independently associated with baseline CBili levels of 2 – 3.9 mg/dL (HR 3.59 [1.24, 10.42], $p=0.02$) and greater than 4 mg/dL (HR 4.80 [1.55, 14.86], $p=0.007$). The multivariable analysis of follow-up CBili data after initiation of intestinal rehabilitation demonstrated a direct relationship between higher bilirubin levels and mortality. After controlling for diagnosis and prematurity, each step-wise increase in CBili during the study period was associated with an increased risk of mortality ($p<0.05$, Table III).

In a secondary analysis using enteral autonomy as the outcome variable, a CBili value greater than 4 mg/dL at study entry was associated with a significant reduction in the overall chance of attaining enteral autonomy (HR 0.45 [0.27 – 0.74], $p<0.01$). In this analysis, the diagnosis of necrotizing enterocolitis was associated with an increased prospect of achieving enteral autonomy (HR 1.66 [1.03 – 2.68], $p<0.05$) as described previously (9).

Discussion

The field of pediatric intestinal failure has seen significant progress over the past two decades. Long-term survival is now standard due to clinical advances carried out through regional multidisciplinary intestinal failure programs. Such programs bring together medical and surgical specialists with the goal of safely rehabilitating the intestinal remnant and weaning PN over time. Data have shown that intestinal rehabilitation centers are a driving factor behind the improved outcomes in pediatric intestinal failure care (4–6).

The benefits associated with multidisciplinary, regionalized care in intestinal failure are multiple. Intestinal failure programs have access to and experience with an assortment of treatment options including alternative lipid emulsions to reverse or prevent IFALD, bowel

lengthening techniques to increase functional intestinal capacity, ethanol locks to decrease the incidence of central line-associated blood stream infection, and options for transplantation if intestinal rehabilitation is not successful. These treatment options are not readily available or utilized consistently at smaller centers without dedicated clinical programs to treat intestinal failure. In a similar fashion, programs specializing in intestinal failure also have access to research trials investigating the next generation of therapies for this condition (10). Finally, providers in intestinal rehabilitation centers have experience with the complications of intestinal failure including bacterial overgrowth, central line infections, and IFALD.

While the value of intestinal failure programs has been established, the optimal timing of referral to these specialized programs has not been defined. It remains common practice for centers without dedicated intestinal rehabilitation programs to continue to care for these young children despite progressive liver disease and the need for long-term PN. Such patients may be cared for by a single specialty or by a dedicated but solo practitioner without the benefit of a team approach. It is established that an early and sustained increase in serum bilirubin is a poor prognostic indicator in children with short bowel syndrome (11, 12). A previous single institution study found that referral to an intestinal rehabilitation program with a CBili greater than 7.2 mg/dL was associated with a 15-fold increased risk of mortality (7). Despite these data, patients are still referred for intestinal rehabilitation weeks to months after the initial diagnosis of intestinal failure.

The present analysis of multicenter data reinforces the finding that the extent of liver disease at the initiation of intestinal rehabilitation is independently associated with mortality in this population. Children enrolled in intestinal failure programs earlier in their course of liver disease had a significant survival benefit and were more likely to attain enteral autonomy. These findings are especially relevant since only 28% of children in our study did not have IFALD at the time of study entry. In our data, mortality was increased even in infants who presented with CBili levels between 2 and 4 mg/dL, often considered "mild" liver disease. This is associated remained significant even when subjects were censored at time of intestinal transplantation in a secondary analysis. In addition, the data demonstrate that CBili represents an on-going marker for disease severity and risk of mortality in this population. Even after enrollment in an intestinal failure program, higher bilirubin levels over time were independently associated with an increased risk of death. In this way, advanced IFALD may not be reversible and can continue to progress despite dedicated multi-disciplinary care. Earlier treatment of IFALD has been shown to improve its long-term prognosis (13).

The primary limitation to this study is the fact that these data, while still the largest set of multi-center data to track a cohort of children with intestinal failure, are dated. Since the completion of the PIFCon data collection, new treatment options for pediatric intestinal failure have evolved. Indeed, most of the data presented here were recorded prior to the widespread use of fish-oil based lipid emulsion to treat IFALD. There is strong evidence that Omegaven® (Fresenius Kabi, Bad Homburg, Germany) can reverse IFALD and improve overall survival in this population if administered in an appropriate and timely fashion (1, 14).

Nonetheless, these data are still relevant to the contemporary care of children with intestinal failure and can help to define the optimal time to refer these children for intestinal rehabilitation. Recent data have demonstrated that the failure of Omegaven to reverse IFALD is associated with patient age and the extent of liver disease at the time of Omegaven initiation (15). Thus, while the widespread use of Omegaven has contributed to improved outcomes in intestinal failure, it still requires timely administration early in the course of IFALD. In addition, the availability of Omegaven remains primarily on a research basis, and its usage requires approval of an Investigational New Drug (IND) application by the Food and Drug Administration and Institutional Review Board of each hospital. It can be argued, then, that intravenous fish oil therapy should be administered primarily by centers with advanced parenteral nutrition experience and infrastructure. In the current era, the optimal administration of intravenous lipid is even more important as the use of SMOF lipid (soy, medium chain triglyceride, olive, and fish oils) has been recently approved for use in the adult population and clinical trials are ongoing in children.

In a similar fashion, the use of lipid restriction has been shown to both reverse and prevent IFALD in children on long-term PN (16–18). This treatment option uses reduced allotments of soy-based lipid emulsion to improve liver function while maintaining growth patterns. Lipid restriction has only recently been studied, and a randomized, prospective trial is currently ongoing at select intestinal failure centers. Lipid restriction is utilized routinely by intestinal failure programs as an alternative to fish oil-based lipid emulsions.

Thus, while the specific strategies to mitigate complications in intestinal failure have evolved since these data were collected, the basic premise that these advanced treatment options are best utilized only in select centers has not changed. From the data presented here, the timely implementation of these strategies may have a direct effect on survival in this population of chronically ill children. For these reasons, early referral of infants with intestinal failure to a multidisciplinary intestinal rehabilitation center remains paramount to optimizing outcomes in IFALD. Liver disease in patients with intestinal failure should be diagnosed promptly and considered a sign of progressive disease requiring more comprehensive care. Since even mild elevations in CBili were associated with an increased risk of mortality in this study, we recommend transfer to a multidisciplinary intestinal failure center for patients who develop a consistently elevated CBili > 2 mg/dL. Strategies to educate referring institutions that care for infants with intestinal failure about these survival patterns should be strongly considered. In a similar fashion, long-distance care partnerships between regional intestinal failure centers and outlying pediatric providers should be encouraged.

Acknowledgments

Support for this manuscript was provided by NIH grants K24 DK104676 (PI: C Duggan), 2P30 DK040561 (PI: C Duggan), 1R21 DK081059-01 (PI: R. Squires), and the National Center For Advancing Translational Sciences UL1 TR000423.

References

1. Puder M, Valim C, Meisel JA, Le HD, de Meijer VE, Robinson EM, Zhou J, Duggan C, Gura JM. Parenteral fish oil improves outcomes in patients with parenteral nutrition-associated liver injury. *Ann Surg.* 2009; 250:395–402. [PubMed: 19661785]
2. Ardura MI, Lewis J, Harp PL, Dienhart MC, Balint JP. Central catheter-associated bloodstream infection reduction with ethanol lock prophylaxis in pediatric intestinal failure: broadening quality improvement initiatives from hospital to home. *JAMA Pediatr.* 2015; 169:324–31. [PubMed: 25642912]
3. Pakarinen MP, Kurvinen A, Koivulsalo AI, Iber T, Rintala RJ. Long-term controlled outcomes after autologous intestinal reconstruction surgery in treatment of severe short bowel syndrome. *J Pediatr Surg.* 2013; 48:339–44. [PubMed: 23414862]
4. Stanger JD, Oliveira C, Blackmore C, Avitzur Y, Wales PW. The impact of multidisciplinary intestinal rehabilitation programs on the outcome of pediatric patients with intestinal failure: a systematic review and meta-analysis. *J Pediatr Surg.* 2013; 48:983–92. [PubMed: 23701771]
5. Hess RA, Welch KB, Brown PI, Teitelbaum DH. Survival outcomes of pediatric intestinal failure patients: analysis of factors contributing to improved survival over the past two decades. *J Surg Res.* 2011; 170:27–31. [PubMed: 21601876]
6. Modi BP, Langer M, Ching YA, Valim C, Waterford SD, Iglesias J, Duro D, Lo C, Jaksic T, Duggan C. Improved survival in a multidisciplinary short bowel syndrome program. *J Pediatr Surg.* 2008; 43:20–4. [PubMed: 18206449]
7. Javid PJ, Malone FR, Bittner R, Healey PJ, Horslen SP. The optimal timing of referral to an intestinal failure program: the relationship between hyperbilirubinemia and mortality. *J Pediatr Surg.* 2011; 46:1052–6. [PubMed: 21683197]
8. Squires RH, Duggan C, Teitelbaum DH, et al. Natural history of pediatric intestinal failure: initial report from the pediatric intestinal failure consortium. *J Pediatr.* 2012; 161:732–8.
9. Khan FA, Squires RH, Litman HJ, et al. Predictors of enteral autonomy in children with intestinal failure: a multicenter cohort study. *J Pediatr.* 2015; 167:29–34. [PubMed: 25917765]
10. Carter BA, Cohran VC, Cole CR, et al. Outcomes from a 12-week, open-label, multicenter trial of Teduglutide in pediatric short bowel syndrome. *J Pediatr.* 2017; 181:102–11. [PubMed: 27855998]
11. Teitelbaum DH, Drongowski R, Spivak D. Rapid development of hyperbilirubinemia in infants with the short bowel syndrome as a correlate to mortality: possible indication for early small bowel transplantation. *Transplant Proc.* 1996; 28:2699–2700. [PubMed: 8908015]
12. Bueno J, Ohwada S, Kocoshis S, et al. Factors impacting the survival of children with intestinal failure referred for intestinal transplantation. *J Pediatr Surg.* 1999; 34:27–33. [PubMed: 10022138]
13. Cowles RA, Ventura KA, Martinez M, et al. Reversal of intestinal failure-associated liver disease in infants and children on parenteral nutrition: experience with 93 patients at a referral center for intestinal rehabilitation. *J Pediatr Surg.* 2010; 45:84–7. [PubMed: 20105585]
14. Diamond IR, Sterescu A, Pencharz PB, Kim JH, Wales PW. Changing the paradigm: omegaven for the treatment of liver failure in pediatric short bowel syndrome. *J Pediatr Gastroenterol Nutr.* 2009; 209–15. [PubMed: 19179884]
15. Nandivada P, Baker MA, Mitchell PD, O'Loughlin AA, Potemkin AK, Anez-Bustillos L, Carlson SJ, Dao DT, Fell JM, Gura JM, Puder M. Predictors of failure of fish-oil therapy for intestinal failure-associated liver disease in children. *Am J Clin Nutr.* 2016; 104:663–70. [PubMed: 27510535]
16. Rollins MD, Ward RM, Jackson WD, Mulroy CW, Spencer CP, Ying J, Greene T, Book LS. Effect of decreased parenteral soybean lipid emulsion on hepatic function in infants at risk for parenteral nutrition-associated liver disease: a pilot study. *J Pediatr Surg.* 2013; 48:1348–56. [PubMed: 23845629]
17. Sanchez SE, Braun LP, Mercer LD, Sherrill M, Stevens J, Javid PJ. The effect of lipid restriction on the prevention of parenteral nutrition-associated cholestasis in surgical infants. *J Pediatr Surg.* 2013; 48:573–8. [PubMed: 23480915]

18. Cober MP, Killu G, Brattain A, Welch KB, Kunisaki SM, Teitelbaum DH. Intravenous fat emulsions reduction for patients with parenteral nutrition-associated liver disease. *J Pediatr*. 2012; 160:421–7. [PubMed: 21982303]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

All cause mortality in the PIFCon dataset was directly related to the level of conjugated bilirubin at study entry ($p=0.003$).

Table I

Univariate analysis of factors associated with mortality in infants with intestinal failure.

Effect	Hazard Ratio	Confidence Interval	P-value
Female (vs male)	0.86	0.47, 1.58	0.63
Gestational age (per week)	0.92	0.86, 0.98	0.008
Achieved enteral autonomy	0.00	0.00, INF	0.99
Age at study entry	1.01	0.98, 1.05	0.43
Baseline CBili 2.0 –3.9 mg/dL	3.54	1.17, 10.66	0.03
Baseline CBili 4.0 mg/dL	5.20	1.80, 14.98	0.002
NEC	1.38	0.75, 2.53	0.30
Gastroschisis	0.61	0.30, 1.25	0.18
Small bowel atresia	1.93	1.02, 3.66	0.04
Midgut volvulus	0.76	0.34, 1.72	0.51
Race (Non-white vs White)	2.46	1.24, 4.86	0.01

Table II

Multivariable regression analysis of factors associated with mortality in infants with intestinal failure.

Effect	Hazard Ratio	Confidence Interval	P-value
Baseline CBili 2.0 – 3.9 mg/dL	3.68	1.89, 7.16	<0.001
Baseline CBili 4.0 mg/dL	4.11	1.78, 9.48	0.001
NEC	0.67	0.25, 1.84	0.44
Gastroschisis	0.68	0.37, 1.27	0.23
SBA	1.76	1.14, 2.71	0.01
Volvulus	0.86	0.50, 1.46	0.57
Race (Non-white vs White)	2.32	0.88, 6.12	0.09

Table III

Multivariable regression data evaluating the association of follow-up bilirubin levels with mortality over time in an intestinal failure program.

Effect	Hazard Ratio	Confidence Interval	P-value
CBili 2.0 – 3.9 mg/dL	4.62	1.06, 20.20	0.04
CBili 4.0 – 6.9 mg/dL	5.60	1.54, 20.37	0.009
CBili 7.0 – 9.9 mg/dL	5.85	1.64, 20.90	0.007
CBili 10.0 mg/dL	7.20	1.80, 28.84	0.005
NEC	1.21	0.38, 3.83	0.74
Gastroschisis	0.89	0.86, 1.16	0.40
Small bowel atresia	1.38	0.73, 2.62	0.32
Volvulus	1.28	0.91, 1.79	0.15
Race (Non-white vs White)	2.58	1.10, 6.05	0.03