Figure 1 Tibial nerve stimulator.

pedal Doppler ultrasound. The tibial nerve stimulator can be used repeatedly during the operation to aid surgical resection. Surgery continues until provocation with the nerve stimulator no longer causes flow disturbance in the pedal vessels.

DISCUSSION

This technique is a relatively simple way of ensuring that the structures compressing the popliteal artery have been satisfactorily resected during surgery. Nerve stimulation is used routinely in anaesthesia without complication.³ Its application in this setting hypothetically reduces the need for a more extensive dissection or resection and potentially reduces the need for repeat surgery.

References

1. Sinha S, Houghton J, Holt PJ *et al*. Popliteal entrapment syndrome. *J Vasc Surg* 2012; **55**(1): 252–262.e30.
2. Peach, G; Sinha, S; Hinchliffe, RJ. Arterial pathologies in athletes. *J Cardiovasc Surg* 2017; **58**(2): 360–368.
3. Association of Anaesthetists of Great Britain and Ireland. Recommendations for standards of monitoring during anaesthesia and recovery 2015. *Anaesthesia* 2016; **71**: 85–93.

A simple endoscopic method for removal of a knotted urethral catheter

B Rampersad

University of the West Indies, Trinidad and Tobago

CORRESPONDENCE TO

Barbara Rampersad, E: barbara.rampersad@sta.uwi.edu
doi 10.1308/rcsann.2017.0108

BACKGROUND

Small infant feeding tubes (5Fr and 6Fr) are commonly used in many institutions for catheterisation of neonates and infants. This is most likely because feeding tubes are cheaper, more flexible and more readily available than infant urinary catheters.¹ Spontaneous intravesicular knotting of a feeding tube used for urethral catheterisation is a rare but documented complication.² Removal of these intravesicular knots has been attempted in a number of ways such as use of a guidewire to

Figure 1 Cystoscope in knot of catheter

untangle the knot, use of excess lubrication and traction with or without anaesthetic, and open or endoscopic removal via cystostomy. This method of removal using a cystoscope has not been described previously.

TECHNIQUE

Our technique uses a well lubricated 7Fr cystoscope inserted urethrally alongside the feeding tube. Introduction of the cystoscope into the knot is possible as the knot is usually stuck at the bladder neck or posterior urethra. The knot is then further tightened by pulling on the distal end of the catheter. It is subsequently possible to remove the catheter on withdrawal of the cystoscope (Fig 1).

DISCUSSION

If standard methods of removal such as gentle traction or use of a guidewire are not successful in cases of a knotted catheter, this is a simple technique that can be employed. This method might avoid the use of more invasive procedures such as cystostomy.

References

1. Arena B, McGillivray D, Dougherty G. Urethral catheter knotting: be aware and minimize the risk. *CJEM* 2002; **4**: 108–110.
2. Foster H, Ritchey M, Bloom D. Adventitious knots in urethral catheters: report of 5 cases. *J Urol* 1992; **148**: 1,496–1,498.

Use of sponge heel boots as a temporary splint for paediatric elbow supracondylar fractures

M Gandhi

Robert Jones and Agnes Hunt Orthopaedic Hospital NHS Foundation Trust, UK

CORRESPONDENCE TO

Maulik Gandhi, E: maulikgandhi@yahoo.co.uk
doi 10.1308/rcsann.2017.0118

Splints are not mentioned as initial management in the British Orthopaedic Association guidelines on supracondylar fractures of the humerus in children.¹ In order to minimise fracture movement (causing pain and/or worsening neurovascular function), plaster casts can be applied. Application is time consuming, and the casts can be challenging to remove and reapply (for x-rays or clinical reassessment) without causing distress to the child.

Figure 1 Heel boot applied to the elbow of a non-injured four-year old boy

Figure 1 Femoral head topped and being reamed for bone harvest

Figure 2 Heel boot applied to the elbow of a non-injured seven-year old girl

Figure 2 Residual shelled out femoral head

Using an adult sized heel boot (designed to reduce the risk of pressure ulcers on the heel area) means the paediatric elbow is adequately splinted and can be reassessed in seconds. This reduces the need for extra analgesia, staff and equipment for casting (Figs 1 and 2).

References

1. British Orthopaedic Association. *BOAST 11: Supracondylar Fractures of the Humerus in Children*. <http://www.boa.ac.uk/wp-content/uploads/2015/01/BOAST-11.pdf> (cited May 2017).

The 'egg topping' technique for bone graft harvesting in hip arthroplasty

AS Awad, MG Williams, GA Higgins
Torbay and South Devon NHS Foundation Trust, UK

CORRESPONDENCE TO

Ashraf Awad, E: ashrafawad@nhs.net
doi 10.1308/rcsann.2017.0125

BACKGROUND

During hip arthroplasty, bone graft from the femoral head is usually acquired either using a bone mill (with its associated costs) or it is done by hand with a rongeur. Further described techniques include reaming the head with an assistant holding it in a clamp or with a swab. These methods carry a risk of dropping the femoral head and injuring the assistant. We describe a simple technique to safely and efficiently obtain cancellous bone graft from the femoral head during hip arthroplasty procedures.

TECHNIQUE

After femoral head dislocation, a saw is used to mark the definitive neck osteotomy. A second osteotomy removes the top of the femoral head (analogous to topping an egg), presenting a flat surface to ream directly on to (Fig 1). Employing a small acetabular reamer, the cancellous bone is obtained from the head, leaving a residual shell of the head proximal to the neck osteotomy (Fig 2). The neck osteotomy is then completed with the oscillating saw.

DISCUSSION

This in vivo technique offers a safe, quick and cost effective alternative to other reported techniques.