CORRESPONDENCE

The Treatment of Illnesses Arising in Pregnancy

by Dr. med. Michael Bolz, Dr. med. Sabine Körber, Prof. Dr. med. Toralf Reimer, Prof. Dr. med. Johannes Buchmann, Prof. Dr. med. Hans-Christof Schober, and Prof. Dr. med. Volker Briese in issue 37/2017

German Pregnancy Register „Rhekiss“

With respect to the very interesting article from Bolz and colleagues (1), it is important to note that numerous inflammatory and systemic autoimmune rheumatic diseases are frequently concomitant with pregnancy, and that pregnancy can substantially alter these diseases. Conversely, activity of the underlying rheumatic disorder can also influence the course of pregnancy, either positively or negatively. Classic examples, which together affect up to 5% of pregnant women (and thus are significantly more frequently than hyperthyroidism, at 1%), are rheumatoid arthritis, spondyloarthritides, psoriatic arthritis, and of course all connective tissue diseases (e.g. systemic lupus erythematosus, systemic sclerosis, myositides) and vasculitides. Furthermore, patients with underlying rheumatic diseases are as a rule on long-term immunosuppressive drugs, the effectiveness of which is crucial for allowing these women to become pregnant and successfully complete pregnancies. As numerous new findings have been associated specifically with these medications in recent years, every woman with an inflammatory rheumatic disease who becomes pregnant should be monitored by a rheumatologist (2). It should also be noted that the presence of connective tissue disease-associated antiphospholipid antibodies defines a high-risk group precisely for venous and arterial thrombosis, and being informed about this and treating it appropriately can be decisive for the outcome of pregnancy (3). To allow us to make more precise statements in the future, the nationwide pregnancy register „Rhekiss“ was established for all medical colleagues in Germany, in which prospective data on the course of pregnancies in women with inflammatory rheumatic diseases can and should be documented (4). DOI: 10.3238/arztebl.2018.0112a

References

1. Bolz M, Körber S, Reimer T, Buchmann J, Schober HC, Briese V: The treatment of illnesses arising in pregnancy. *Dtsch Arztebl Int* 2017; 114: 616–26.
2. Götestam Skorpen C, Hoeltzenbein M, Tincani A, et al.: The EULAR points to consider for use of antirheumatic drugs before pregnancy, and during pregnancy and lactation. *Ann Rheum Dis* 2016; 75: 795–810.
3. Andreoli L, Bertias GK, Agmon-Levin N, et al.: EULAR recommendations for women's health and the management of family planning, assisted reproduction, pregnancy and menopause in patients with systemic lupus erythematosus and/or antiphospholipid syndrome. *Ann Rheum Dis* 2017; 76: 476–85.
4. Rheuma, Kinderwunsch und Schwangerschaft. www.rhekiss.de (last accessed on 15 January 2018).

Prof. Dr. med. Ulf Müller-Ladner

Abteilung Rheumatologie und Klinische Immunologie, Justus-Liebig-Universität Gießen, Campus Kerckhoff, Bad Nauheim
u.mueller-ladner@kerckhoff-klinik.de

PD Dr. med. Rebecca Fischer-Betz

Abteilung Rheumatologie, Universitätsklinikum Düsseldorf

Conflict of interest statement

PD Dr. Fischer-Betz has received speaking honoraria from Abbvie, Chugai, MSD, Roche, UCB, Pfizer, Cellegue, Janssen, and Lilly.

Prof. Müller-Ladner declares that no conflict of interest exists.

Anatomically Incorrect

With interest we read the article by Bolz et al. (1). As coloproctologists we sometimes treat patients who develop proctologic complaints during pregnancy or after delivery. Unfortunately, under the heading „Varicose veins (chronic venous insufficiency)“, the authors also mention „hemorrhoids“ („The following sites of involvement are important in pregnancy because of the possibility of hemorrhagic complications: ... and hemorrhoids [27–30]“). Anatomically, physiologically, and functionally, this is incorrect: Hemorrhoids are arteriovenous, cavernous cushions that have no direct relationship to veins and venous insufficiency. Women who are pregnant or have given birth often present with perianal (venous) thrombosis or hematoma. These occur in the (very sensitive) area of the anocutaneous line, under which extends a subcutaneous venous plexus.

As coloproctologists, and from an anatomical perspective, we find it important to clarify (also for this readership) that hemorrhoids have nothing to do with veins and certainly are not „varicose veins“.

DOI: 10.3238/arztebl.2018.0112b

References

1. Bolz M, Körber S, Reimer T, Buchmann J, Schober HC, Briese V: The treatment of illnesses arising in pregnancy. *Dtsch Arztebl Int* 2017; 114: 616–26.

Dr. med. Johannes Jongen, Prof. Dr. med. Volker Kahlke

Proktologische Praxis Kiel, Germany
j.jongen@kielnet.net

Prof. Dr. med. Thilo Wedel

Anatomisches Institut, Christian-Albrechts-Universität Kiel, Germany

Conflict of interest statement

The authors declare that no conflict of interest exists.

Independent Measurement of PEF Values

The article calls for an optimal control of asthma when caring for pregnant women. However, the only parameters of success given are monitoring of the lung

function and of the exhaled nitric oxide (NO) (1). As an internist working as a general practitioner, I do consider neither the proposed monthly control of the lung function nor NO measurements (which are not covered by the Statutory Health Insurance) to be efficient, since both are snapshots and thus not necessarily representative of the clinical picture.

In contrast, I rely on frequent, independent measurement checks of the peak expiratory flow (PEF) when caring for pregnant women with asthma. PEF meters are inexpensive (at 15–22 €) and their benefits easily explained (comparable to blood sugar controls for pregnant women with type I diabetes). In case of viral respiratory tract infections, which occur frequently, therapy can be escalated very rapidly (for example, by increasing the dose of the inhaled budesonide)—often before typical asthma symptoms appear. This minimizes the risks associated with asthma exacerbation in pregnant women.

DOI: 10.3238/arztebl.2018.0113a

Reference

1. Bolz M, Körber S, Reimer T, Buchmann J, Schober HC, Briese V: The treatment of illnesses arising in pregnancy. *Dtsch Arztebl Int* 2017; 114: 616–26.

Dr. med. Heidrun Höppner
Forchheim, Germany
heidrun.hoeppner@gmx.com

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank all authors of the correspondences for their interest and comments on our article (1).

To address the correspondence of Dr. Höppner: Thank you for your commentary about follow-up of bronchial asthma during pregnancy. Since peak expiratory flow (PEF) variability correlates well with the extent of bronchial hyper-responsiveness, it is used for monitoring and therapy control. While the PEF value can be directly read from the flow-volume loop on spi-

rometry, this requires patient collaboration. Alternatively, the patient can measure PEF at home several times a day using a peak flow meter. Although unlike spirometry, the peak flow meter is not a calibrated measuring method, differences between the two methods are irrelevant as the peak flow meter records intra-individual peak flow variability (2).

From our point of view, it is important in general to require regular monitoring and therapy control of the disorder throughout pregnancy. Which type of pulmonary function examination is preferred is then of secondary importance. Furthermore, no distinction is currently made between controlled and uncontrolled lung function in bronchial asthma. Here, the parameters of symptomatology, limitation of everyday activity, nocturnal symptoms, use of on-demand medication, and exacerbation play the decisive role.

We thank Dr. Jongen, Prof. Kahlke, and Prof. Wedel for their corrective supplement to our article.

To address the correspondence of Prof. Müller-Ladner and Dr. Fischer-Betz: We also believe that inflammatory and systemic autoimmune rheumatic diseases are of great importance in pregnancy. In this respect, we agree with the authors. However, as a concomitant disease, the incidence of thyroid diseases during pregnancy is more frequent overall and therefore was taken into account in our article.

DOI: 10.3238/arztebl.2018.0113b

References

1. Bolz M, Körber S, Reimer T, Buchmann J, Schober HC, Briese V: The treatment of illnesses arising in pregnancy. *Dtsch Arztebl Int* 2017; 114: 616–26.
2. AWMF: gültige S2k-Leitlinie Spirometrie 020–017. www.awmf.org/leitlinien/detail/II/020-017.html (last accessed on 15 January 2018).

On behalf of the authors:

Dr. med. Michael Bolz
Universitätsfrauenklinik
Klinikum Südstadt Rostock, Germany
michael.bolz@kliniksued-rostock.de

Conflict of interest statement

The author declares that no conflict of interest exists.