Published in final edited form as:

J Chem Theory Comput. 2017 May 09; 13(5): 1874–1882. doi:10.1021/acs.jctc.7b00194.

Reparameterization of Solute—Solute Interactions for Amino Acid-Sugar Systems Using Isopiestic Osmotic Pressure Molecular Dynamics Simulations

Wesley K. Lay, Mark S. Miller, and Adrian H. Elcock

Department of Biochemistry, University of Iowa, Iowa City, IA 52242

Abstract

AMBER/GLYCAM and CHARMM are popular force fields for simulations of amino acids and sugars. Here we report excessively attractive amino acid-sugar interactions in both force fields, and corrections to nonbonded interactions that match experimental osmotic pressures of mixed aqueous solutions of diglycine and sucrose. The modified parameters also improve the G_{trans} of diglycine from water to aqueous sucrose and, with AMBERff99SB/GLYCAM06, eliminate a caging effect seen in previous simulations of the protein ubiquitin with glucose.

Protein-carbohydrate interactions are critical to a number of important cellular functions such as protein signaling and energy storage,^{1–2} as well as to the construction and maintenance of the peptidoglycan component of the bacterial cell wall.^{3–4} In addition, sugars are known to stabilize the native states of proteins,^{5–6} and this feature is commonly exploited by osmotically-stressed and freeze-resistant organisms.⁷ Properly accounting for the interactions of sugars with amino acids, therefore, is likely to be important for developing a deeper understanding of their roles in biology. To this end, a number of recent studies have used molecular dynamics (MD) simulations, often in connection with traditional experiments, in an attempt to model complex systems containing both carbohydrates and amino acids. Examples include: (a) the use of MD simulations to understand the stabilization of proteins by glycosylation,^{8–10} (b) insightful attempts to model and simulate the peptidoglycan layers of Gram-positive¹¹ and Gram-negative¹² bacteria, and (c) the use of MD to simulate lipid-linked lipopolysaccharides,^{13–14} whose carbohydrate chains are known to interact with outer membrane proteins.¹⁵

The predictive utility of these models and simulations relies heavily on the accuracy of the force fields that are used. A number of recent studies, however, have identified potential problems in the description of nonbonded interactions in two popular force fields that are

Correspondence to: Adrian H. Elcock.

Supporting Information is available free of charge on the ACS Publications website at DOI: to-be-determined.

Scatter plot of the compositions of diglycine and sucrose that have been shown experimentally to be isopiestic; parameterization of nonbonded parameters to reproduce osmotic pressures of diglycine, sucrose, and diglycine-sucrose mixtures; comparison of osmotic coefficients obtained from simulations using modified glucose-glucose parameters with the TIP3P and TIP4P-Ew water models with experiment; plot of osmotic pressure of sucrose as a function of the σ value of oxygen-oxygen interactions in GLYCAM06; list of the diglycine/sucrose compositions that were simulated; values for the optimized nonbonded interaction parameters for AMBERff99SB/GLYCAM06, and CHARMM36.

otherwise well equipped to describe both carbohydrate and amino acid-based systems. These include AMBERff99SB^{16–17} and its variants (widely used to model proteins and amino acids), GLYCAM06¹⁸ (the AMBER-compatible carbohydrate force field), and CHARMM36.^{19–22} With regard to carbohydrates, non-physical aggregation of glucose molecules has been shown to occur^{23–24} at experimentally-soluble concentrations using the GLYCAM06 force field, and the simulated osmotic coefficients of aqueous solutions of glucose and sucrose have been shown to be much too low, relative to experiment, with both the GLYCAM06 and CHARMM36 force fields.²⁴ With regard to amino acids, on the other hand, both the Aksimentiev group²⁵ and our group^{26–27} have reported osmotic pressure simulations that indicate that excessively attractive solute-solute interactions can occur with both force fields depending on the water model used.

Excessively favorable interactions between solutes in aqueous solution reflect an imbalance in the energetic descriptions of solute-solute, solute-solvent, and solvent-solvent interactions in the force fields. One way to quantify the effective strengths of solute-solute interactions in simulations and to compare them to experiment, is to perform osmotic pressure simulations using procedures similar to those outlined originally by Murad²⁸ and by Luo and Roux.²⁹ In such simulations, virtual walls are used to mimic a semi-permeable membrane that restricts solute molecules to a central region of a simulation cell, while leaving the water molecules free to explore the entire cell. The simulated osmotic pressure of the solution is obtained from the forces exerted by the solute molecules on the virtual walls. Carefully selected modifications can then be made to the nonbonded parameters describing solute-solute interactions in an attempt to bring the simulated osmotic pressure into line with experiment; importantly, this can be done while leaving other simulated properties of the solutes, such as their free energies of hydration, unchanged.

A number of recent studies have used this kind of approach to improve the description of solute-solute interactions in a variety of force fields. For example: Luo and Roux have optimized parameters for NaCl and KCl in CHARMM27;²⁹ Yoo and Aksimentiev have optimized parameters for cation-acetate, cation-phosphate,³⁰ amine-carboxylate and amine-phosphate²⁵ interactions in both the CHARMM36 and AMBERff99SB force fields; more recently, the same authors have parameterized a hydrated model of the Ca²⁺ ion³¹ and have shown that their previously proposed modifications improved the realism of protein folding simulations;³² our group has optimized nonbonded parameters for sugar-sugar interactions in GLYCAM06 and CHARMM36,²⁴ and amino acid-amino acid interactions in a number of force fields;²⁷ the Grafmüller group has optimized nonbonded parameters for sugar-sugar interactions in GLYCAM06 using the TIP5P water model;³³ Saxena and Garcia have improved the behavior of multisite models of divalent cations (Mg²⁺ and Ca²⁺) by calibrating against the experimental osmotic pressure of concentrated salt solutions;³⁴ finally, the MacKerell group has used osmotic pressure measurements to verify improved nonbonded parameters for guanidinium-acetate interactions in the newly-released CHARMM36m force field.³⁵

In each of these previous studies, the experimental osmotic pressure data that have been targeted have referred to what are effectively binary solutions, i.e. solutions involving a single type of solute molecule dissolved in water. But many important biomolecular systems

involve combinations of molecule types, with the protein-carbohydrate systems noted at the beginning of this manuscript being good examples. To begin the process of parameterizing such mixed interactions, therefore, we report here the use of osmotic pressure simulations to validate and optimize nonbonded parameters describing amino acid-sugar interactions in both the AMBERff99SB/GLYCAM06 and CHARMM36 force fields. In all of the simulations reported here we have used the TIP4P-Ew water model³⁶ in conjunction with the AMBERff99SB/GLYCAM06 force field, and the CHARMM-modified TIP3P water model^{37–38} in conjunction with the CHARMM36 force field; the TIP4P-Ew water model was selected for use with the AMBER force field as it has been shown to modestly outperform the TIP3P model when used to simulate amino acid and peptide systems.³⁹

Experimental osmotic pressure data for ternary solutions (which, in the present context, means mixtures of amino acids and sugars in water) are often available in the literature in the form of isopiestic osmotic or activity coefficient data. In a typical experimental isopiestic setup, ternary aqueous solutions with different compositions (e.g. 1 m diglycine + 0.5 m sucrose, etc.) are stored together with a reference solution (e.g. aqueous KCl) and water is allowed to equilibrate between the various solutions by evaporation. At equilibrium, the osmotic pressures of all solutions will be identical and their absolute values can be determined from the molal concentration of the solute in the reference solution, for which high quality osmotic pressure data as a function of concentration are available. Importantly for this work, isopiestic data for aqueous solutions of mixtures of diglycine and sucrose were reported many years ago by Uedaira.⁴⁰ The availability of these data enable the MD-simulated strengths of amino acid-sugar interactions to be tested in the following way. Using the concentration data provided by Uedaira, we generate a range of mixtures of diglycine and sucrose in water, all of which should produce identical simulated osmotic pressures if the simulation force fields are realistic. We first measure the simulated osmotic pressures of supposedly-isopiestic binary solutions (diglycine + water, and sucrose + water) in order to validate and, if necessary, reparameterize amino acid-amino acid, and sugar-sugar interactions. We then measure the simulated osmotic pressures of the supposedly-isopiestic ternary solutions (diglycine + sucrose + water) and, if necessary, reparameterize the amino acid-sugar interactions in an attempt to match the experimental osmotic pressure over the entire range of mixtures from diglycine + water, through diglycine + sucrose + water mixtures, to sucrose + water.

In the present study, we chose Uedaira's reference KCl concentration of 0.81372 m, for which five isopiestic mixtures, ranging from 0.585 m diglycine + 0.869 m sucrose to 1.394 m diglycine + 0.250 m sucrose, have been reported (see Table S1 in the Supporting Information). Based on the reference data for KCl reported by Robinson and Stokes,⁴¹ all of these compositions should produce an osmotic pressure of 36.17 bar. When the diglycine and sucrose concentrations that are isopiestic are plotted against each other (Figure S1), they follow a clear trend that can be fit extremely well to either a simple linear or quadratic function; since experimental data for other systems can deviate from linearity (see, e.g.⁴²) we chose here to use the more general quadratic fit, but very similar results would be obtained with either fit function. We extrapolate the fit function to both axes in order to determine the concentrations of the binary solutions that should be isopiestic with Uedaira's ternary solutions. The extrapolated concentrations are 1.69 m and 1.29 m for diglycine +

water and sucrose + water, respectively. Importantly, in the present case, we can explicitly check both of these values against experimental osmotic pressure data that have been reported independently for binary solutions of the same solutes. According to the data reported by Ellerton *et al.*,⁴³ a 1.69 m solution of diglycine in water should have an osmotic pressure of 36.57 bar; according to the data reported by Stokes and Robinson,⁴⁴ a 1.29 m solution of sucrose in water should have an osmotic pressure of 36.39 bar. Encouragingly, both of these values are in close agreement with, and are therefore isopiestic with, the osmotic pressure data for diglycine + sucrose + water ternary solutions. In addition to these two extreme conditions, we also extrapolated Uedaira's data in an attempt to obtain a more uniformly sampled range of compositions, thereby identifying a seventh mixture composition for simulation (see data point second from the left in Figure S1).

Having obtained a set of seven mixture compositions that should all be isopiestic, we tested the ability of the AMBERff99SB/GLYCAM06 and CHARMM36 force fields to reproduce these data. The osmotic pressures obtained with the original versions of the two force fields are shown in Figure 1A. With AMBERff99SB/GLYCAM06 (blue circles), the simulated osmotic pressures of the mixtures are all too low, and the level of agreement with experiment becomes progressively worse as the mole fraction of sucrose increases. This behavior, while somewhat disappointing, is at least consistent with our previous studies, which suggested that while amino acid-amino acid interactions described by AMBERff99SB^{26–27} are reasonably accurate, the sugar-sugar interactions described by GLYCAM06²⁴ are far too attractive. Interestingly, this is the case despite the fact that GLYCAM's nonbonded parameters and atom types are largely borrowed from AMBER. This suggests, therefore, that amino acid-amino acid and sugar-sugar interactions might require unique and independent sets of atom types in AMBER in order to produce equally realistic descriptions of behavior in solution.

With the CHARMM36 force field (Figure 1A, red upward triangles) the simulated osmotic pressures are again somewhat too low, but the level of agreement with experiment is effectively independent of the mole fraction of sucrose. The generally lower values obtained with CHARMM36 are consistent with our group's previous work on carbohydrates,²⁴ and with work from the Aksimentiev group on glycine.²⁵ But the generally flat nature of the plot for CHARMM36 (red line) is more encouraging. In particular, it suggests that the force field's descriptions of amino acid-amino acid, sugar-sugar, and amino acid-sugar interactions are likely to be well balanced, making it better suited to dealing with mixed solutions, and potentially better suited to modeling transfer between different solutions, than the less well-balanced AMBERff99SB/GLYCAM06 force field combination.

The results for the two binary solutions (left-most and right-most data points in Figure 1A) make it clear that there are problems with both force fields' descriptions of amino acid-amino acid and sugar-sugar interactions. The ternary solutions, on the other hand, involve not only amino acid-amino acid and sugar-sugar interactions, but also amino acid-sugar interactions. It is not clear at this stage, therefore, whether the poor results obtained for the ternary solutions (intermediate data points in Figure 1A) reflect additional problems with the force fields' descriptions of amino acid-sugar interactions, or whether they can be attributed

entirely to the errors that we already know exist in the descriptions of amino acid-amino acid and sugar-sugar interactions.

To distinguish between these two possibilities, we must first modify both force fields so that they reproduce the experimental osmotic pressure of the two binary solutions: diglycine + water, and sucrose + water. To do so, we implemented force field modifications similar to those used previously to correct the osmotic coefficients of glycine²⁵ and glucose.²⁴ Specifically, for diglycine, we followed the approach developed by Yoo and Aksimentiev for glycine,²⁵ and increased the Lennard-Jones σ value describing intermolecular interactions of the amino group nitrogen and the carboxylate oxygen atoms; we have recently shown that this approach can correct amino acid-amino acid interactions with a variety of other force fields.²⁷ Figure S2 shows that diglycine's osmotic pressure can be brought into close agreement with experiment by modestly increasing σ with both the AMBERff99SB/GLYCAM06 (Figure S2A) and CHARMM36 (Figure S2B) force fields; in percentage terms these modifications amount to changes of ~5 and ~2.5%, respectively.

For sucrose with AMBERff99SB/GLYCAM06, we followed the approach used in our recent work for glucose,²⁴ and decreased the Lennard-Jones ϵ values describing inter-molecular carbon-carbon and carbon-oxygen interactions while leaving intra-molecular interactions unaltered. We note that we also tried the alternative approach of increasing the σ values for intermolecular oxygen-oxygen interactions, but, as was also the case in our recent work on glucose,²⁴ were unable to reproduce experiment (Figure S3). For sucrose with CHARMM36, on the other hand, we were successful in following a similar strategy to that used for the amino acids, since we found that experiment could be reproduced by increasing the σ value describing intermolecular oxygen-oxygen interactions. The success of both types of modifications in matching the experimental osmotic pressure of sucrose can be seen in Figure S2C for AMBERff99SB/GLYCAM06 and Figure S2D for CHARMM36. The full set of modifications – which, it should be stressed, are almost certainly not the only way by which agreement with experiment can be achieved – are detailed in Tables S2A and S2B for AMBERff99SB/GLYCAM06 and CHARMM36, respectively.

The osmotic pressures obtained with these modified versions of the two force fields are shown in Figure 1B. By construction, the osmotic pressures for the binary solutions of diglycine + water (left-most data points) and sucrose + water (right-most data points) are now within error of experiment with both AMBERff99SB/GLYCAM06 (blue circles) and CHARMM36 (red upward triangles). But with AMBERff99SB/GLYCAM06, the osmotic pressures for the ternary solutions (intermediate blue data points) are still found to be much too low, which strongly suggests that amino acid-sugar interactions are excessively attractive with this force field combination. With CHARMM36, the osmotic pressures for the ternary solutions are in much better agreement with experiment (again reflecting the well-balanced nature of the original force field), but there remains a tendency for the simulated values to be too low.

In an attempt to eliminate the remaining deviations from experiment, we implemented force field modifications to the diglycine-sucrose interactions in both force fields. With AMBERff99SB/GLYCAM06, we decreased the Lennard-Jones ϵ values describing

intermolecular carbon-carbon interactions; with CHARMM36, we very slightly increased the σ value describing intermolecular nitrogen-oxygen and oxygen-oxygen interactions so as to weaken the hydrogen bonding interactions. With both force fields, this parameterization work was carried out only at a single composition (1.158 m diglycine + 0.432 m sucrose; mole fraction of sucrose = 0.272). The success of both types of modifications in matching the experimental osmotic pressure of the mixed diglycine-sucrose solution can be seen in Figures S2E and S2F for AMBERff99SB/GLYCAM06 and CHARMM36, respectively. The full sets of adjusted parameter values are again detailed in Tables S2A and S2B for AMBERff99SB/GLYCAM06 and CHARMM36, respectively. The osmotic pressures obtained when the fully modified versions of the two force fields are applied to the entire range of simulated compositions are shown in Figure 1C. Encouragingly, with both AMBERff99SB/GLYCAM06 (blue circles) and CHARMM36 (red upward triangles), the computed osmotic pressures are found to be within error of experiment over the entire range of compositions tested.

At this point, therefore, we have optimized the nonbonded interaction parameters for diglycine-sucrose interactions so that they reproduce isopiestic osmotic pressure data. In order to validate these parameters further, we attempted to reproduce the experimental transfer free energy (G_{trans}) of +0.39 kcal/mol associated with transferring diglycine from water to a 15% aqueous sucrose solution.⁴⁵ Since the experimental G_{trans} value was reported by a different group (that of Chatterjee and Basumallick⁴⁵), using a quite different experimental approach (solubility measurements), attempting to reproduce this G_{trans} represents an important, independent test of the modified parameters. To this end, we performed a series of umbrella sampling simulations in which a single diglycine molecule was restrained to 96 different positions along a reaction coordinate that transfers it from water, into a 15 wt% sucrose solution, and then back into water (Figure 2A; see Computational Methods for details). The resulting potentials of mean force (PMF), computed using the weighted histogram analysis method (WHAM),⁴⁶ are plotted in Figure 2 for each of the force field combinations described earlier: (a) the original force field, (blue circles), (b) the original force field with modified sugar-sugar interactions (green upward triangles), and (c) the original force field with modified sugar-sugar and amino acid-sugar interactions (red downward triangles). Note that the modifications made to amino acid-amino acid interactions are irrelevant to this particular analysis as the simulations involve only a single molecule of diglycine.

The PMFs obtained from simulations using AMBERff99SB/GLYCAM06 are shown in Figure 2B. From this it can be seen that the original force field parameters (blue circles) predict a large negative G_{trans} value (~ -1.3 kcal/mol) which is in qualitative disagreement with the positive G_{trans} observed experimentally. Modifying only the sucrose-sucrose interactions (green upward triangles) decreases the magnitude of the negative G_{trans} to ~ -0.4 kcal/mol, but the result remains qualitatively wrong, suggesting (as was seen previously in the osmotic pressure simulations) that diglycine-sucrose interactions are also excessively attractive using the original force field parameters. Finally, modifying both the sucrose-sucrose and diglycine-sucrose interactions so that they match the isopiestic data of Uedaira⁴⁰ (red downward triangles), results in a PMF that, while still too low, is more in line with

experiment: the diglycine molecule now shows no obvious preference for either the water or the aqueous sucrose phases.

The PMFs obtained from corresponding simulations using CHARMM36 are shown in Figure 2C. The original CHARMM36 parameters predict an essentially flat PMF that is in much closer agreement with experiment than was obtained with the AMBER force field, and that again suggests that while the solute-solute interactions in CHARMM36 are sometimes too attractive, they are, on the whole, very well balanced. Modifying only the sucrose-sucrose interactions (green downward triangles) does not result in any significant change to the PMF, but the addition of modifications to the diglycine-sucrose interactions (red upward triangles) produces a slight upward shift in the PMF that again brings it into somewhat better agreement with experiment.

Taken together, the results shown in Figure 2 indicate that force field modifications introduced in order to improve the modeling of osmotic pressure behavior also improve the modeling of transfer free energies involving the same molecule types. Finally, for a more ambitious test of our force field modifications, we looked to apply them to a model protein-sugar system. An interesting recent MD simulation study from the Dal Peraro group has reported a ‘caging’ of the small protein ubiquitin by glucose molecules when simulated with the AMBERff99SB/GLYCAM06 force field.⁴⁷ Simulation snapshots shown in that work appear to indicate that the glucose molecules aggregate around the protein at a glucose concentration that would normally be considered soluble (i.e. 325 g/L, or 1.8 M). Given the behavior seen in our previous study of sugar solutions,²⁴ and given the behavior seen in the isopiestic osmotic pressure simulations reported here (see earlier), we thought it worthwhile to determine whether the interesting behavior seen in the simulations of Spiga *et al.* might be an unfortunate consequence of excessively attractive sugar-sugar and amino acid-sugar interactions in the AMBERff99SB/GLYCAM06 force field.

We therefore attempted to roughly mimic the simulation conditions reported in the work of Spiga *et al.*, but using each of the three AMBER force field iterations described here, namely: (a) the original AMBERff99SB/GLYCAM06 force field, (b) the original force field but with modified sugar-sugar interactions, and (c) the original force field with modified sugar-sugar and amino acid-sugar parameters. For the sugar-sugar interactions, we used the parameters optimized for 1M glucose with the TIP3P water model in our previous study.²⁴ Here, they were used together with the TIP4P-Ew water model since this is our preferred model for use in simulations of amino acids and proteins (see above); additional simulations, however, showed that while the parameters were originally derived for use with TIP3P, they also work well for describing glucose-glucose interactions in TIP4P-Ew water (see Figure S4). For the amino acid-sugar interactions, we used the parameters optimized here to reproduce the osmotic pressure of diglycine-sucrose mixtures (see Table S2A). Although these parameters were optimized only for sugar-interactions of diglycine, we chose to apply the same modifications to the sugar-interactions of all non-aromatic carbon atoms in ubiquitin (i.e. atom types CT and C). We chose not to modify the sugar-interactions of aromatic carbon atoms (i.e. those in the sidechains of Phe, Tyr, Trp or His) since, at this stage, we have no direct evidence that these interactions are incorrectly described by the original force field. In recognition of the above issues, therefore, we should consider the

modified parameters used in the following simulations to be a preliminary set of parameters for use with protein-carbohydrate systems, not a finalized force field.

Snapshots showing the state of the ubiquitin + glucose systems after a 100 ns simulation with each of the three force fields are displayed in Figure 3. As hoped, the behavior seen with the original force field (Figure 3A) appears to match closely that described by Spiga *et al.*⁴⁷: the ubiquitin is surrounded by a cage of aggregated glucose molecules. This indicates that the behavior reported by those authors is a consistent prediction of the AMBERff99SB/GLYCAM06 force field and is robust to changes in the water model (since TIP4P-Ew was used here, whereas TIP3P was used in the Spiga *et al.* study). In contrast, the snapshot obtained at the end of the simulation that used modified (i.e. more realistic) glucose-glucose interactions (Figure 3B), shows an obvious difference in that the glucose molecules remain unaggregated. Less obvious from the figure, however, is the fact that the glucose molecules in this simulation show a similarly high affinity for the surface of the ubiquitin to that seen with the original force field.

This can be seen more clearly in Figure 4, which shows a histogram of the minimum heavy-atom distance between each glucose molecule and the ubiquitin, sampled over the entire 100 ns of each simulation. With both the original force field (blue line) and the glucose-glucose-only modified force field (green line), a large peak is obtained at a distance (~ 2.8 Å) indicative of strong hydrogen bonding between glucose molecules and polar atoms on the ubiquitin surface. The differences between the two force fields only become apparent at longer distances, where the original force field (blue line) produces a diffuse peak from ~ 6 to ~ 8 Å that reflects binding of secondary glucose molecules to the primary glucose molecules that are in direct contact with the ubiquitin. This peak disappears in the simulations using the modified glucose-glucose parameters (green line) since their interactions have been significantly weakened in order to match the experimental osmotic pressure data.

Finally, the snapshot obtained at the end of the simulation that used modified glucose-glucose and amino acid-glucose interactions (Figure 3C), shows no aggregation of the glucose molecules and a much lower level of association of glucose molecules to the ubiquitin. The latter aspect is especially noticeable in the histogram of minimum distances (see red line in Figure 4), where the large peak indicative of direct hydrogen bonding is significantly diminished. The overall effect of this weakened association between glucose molecules and the ubiquitin is that the protein tumbles and diffuses much more rapidly than is seen with the original force field parameters (data not shown). We think that the fundamentally different views of glucose's effects on ubiquitin that emerge from simulations using the original AMBERff99SB/GLYCAM06 force field, and from the modified version proposed here, reiterates the importance of developing fully validated force fields for modeling biomolecular systems.

In closing, we reiterate that the present report is simply the latest in a number of recent studies highlighting the potential power of osmotic pressure simulations as a means of validating and refining the nonbonded parameters used to describe biomolecular interactions in aqueous solution. Previous studies have focused on improving the behavior of a range of

chose to optimize agreement with experiment for sucrose by diminishing the well-depth, ϵ , of carbon-carbon and carbon-oxygen interactions, and we chose to apply a similar approach in attempting to match the osmotic data for the mixed diglycine-sucrose solutions. With CHARMM36, on the other hand, we found here that adjusting σ values for oxygen-oxygen interactions was sufficient to match the osmotic data for sucrose and we chose, therefore, to follow a similar strategy in attempting to match the data for the mixed solutions. Although it would seem preferable to have a more unified approach to modifying force fields, it seems quite possible that different force fields will be deficient in different ways so fundamentally different solutions to the problems might be required. Future applications to other systems will hopefully clarify this issue.

Finally, it should not be thought that the osmotic-pressure centered approach outlined here is the only way to proceed in the development of improved force field parameters. The Maginn⁴⁸ and Hünenberger⁴⁹ groups, for example, have recently chosen to focus on modifying the electrostatic interactions of carbohydrates and organic solutes in GROMOS in order to improve their behavior. The Smith group has, for a number of years, taken a similar approach in developing parameters for their influential Kirkwood-Buff force field.^{50–52} Finally, the Nerenberg group^{53–54} has chosen to modify the van der Waals parameters describing solute-solute and solute-solvent interactions in AMBERff99SB in order to reproduce experimental enthalpies of vaporization and free energies of hydration, respectively. Given the Chong group's recent demonstration that a new protein force field can be derived essentially 'from scratch' in the space of a few months, it is likely that still other routes to force field optimization will be explored in the near future.⁵⁵

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH R01 GM087290 awarded to A.H.E.

References

1. Madigan, MT., Martinko, JM., Parker, J., Brock, TD. *Biology of Microorganisms*. Vol. 985. Prentice Hall; Upper Saddle River, NJ: 1997.
2. Varki, A. *Essentials of Glycobiology*. Cold Spring Harbor Laboratory Press; Cold Spring Harbor, N.Y.: 2009.
3. Vollmer W, Blanot D, de Pedro MA. Peptidoglycan Structure and Architecture. *FEMS Microbiol Rev*. 2008; 32:149–167. [PubMed: 18194336]
4. Vollmer W, Bertsche U. Murein (Peptidoglycan) Structure, Architecture and Biosynthesis in *Escherichia Coli*. *Biochim Biophys Acta*. 2008; 1778:1714–1734. [PubMed: 17658458]
5. Lee JC, Timasheff SN. The Stabilization of Proteins by Sucrose. *J Biol Chem*. 1981; 256:7193–7201. [PubMed: 7251592]
6. Kim Y-S, Jones LS, Dong A, Kendrick BS, Chang BS, Manning MC, Randolph TW, Carpenter JF. Effects of Sucrose on Conformational Equilibria and Fluctuations Within the Native-State Ensemble of Proteins. *Protein Sci*. 2003; 12:1252–1261. [PubMed: 12761396]
7. Yancey P, Clark M, Hand S, Bowlus R, Somero G. Living With Water Stress: Evolution of Osmolyte Systems. *Science*. 1982; 217:1214–1222. [PubMed: 7112124]

8. Zhong L, Xie J. Investigation of the Effect of Glycosylation on Human Prion Protein by Molecular Dynamics. *J Biomol Struct Dyn*. 2009; 26:525–533. [PubMed: 19236103]
9. Mallajosyula SS, Jo S, Im W, MacKerell AD Jr. Molecular Dynamics Simulations of Glycoproteins using CHARMM. *Methods Mol Biol*. 2015; 1273:407–429. [PubMed: 25753723]
10. Lee HS, Qi Y, Im W. Effects of N-Glycosylation on Protein Conformation and Dynamics: Protein Data Bank Analysis and Molecular Dynamics Simulation Study. *Sci Rep*. 2015; 5:8926. [PubMed: 25748215]
11. Beeby M, Gumbart JC, Roux B, Jensen GJ. Architecture and Assembly of the Gram-positive Cell Wall. *Mol Microbiol*. 2013; 88:664–672. [PubMed: 23600697]
12. Gumbart JC, Beeby M, Jensen GJ, Roux B. Escherichia coli Peptidoglycan Structure and Mechanics as Predicted by Atomic-Scale Simulations. *PLoS Comput Biol*. 2014; 10:e1003475. [PubMed: 24586129]
13. Kirschner KN, Lins RD, Maass A, Soares TA. A Glycam-Based Force Field for Simulations of Lipopolysaccharide Membranes: Parametrization and Validation. *J Chem Theory Comput*. 2012; 8:4719–4731. [PubMed: 26605626]
14. Wu EL, Engström O, Jo S, Stuhlsatz D, Yeom MS, Klauda JB, Widmalm G, Im W. Molecular Dynamics and NMR Spectroscopy Studies of E. coli Lipopolysaccharide Structure and Dynamics. *Biophys J*. 2013; 105:1444–1455. [PubMed: 24047996]
15. Pavlova A, Hwang H, Lundquist K, Balusek C, Gumbart JC. Living on the Edge: Simulations of Bacterial Outer-Membrane Proteins. *Biochim Biophys Acta - Biomembranes*. 2016; 1858:1753–1759.
16. Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C. Comparison of Multiple AMBER Force Fields and Development of Improved Protein Backbone Parameters. *Proteins*. 2006; 65:712–725. [PubMed: 16981200]
17. Lindorff-Larsen K, Piana S, Palmo K, Maragakis P, Klepeis JL, Dror RO, Shaw DE. Improved Side-Chain Torsion Potentials for the Amber ff99SB Protein Force Field. *Proteins*. 2010; 78:1950–1958. [PubMed: 20408171]
18. Kirschner KN, Yongye AB, Tschampel SM, González-Outeiriño J, Daniels CR, Foley BL, Woods RJ. GLYCAM06: a generalizable biomolecular force field. *Carbohydrates. J Comput Chem*. 2008; 29:622–655. [PubMed: 17849372]
19. Guvench O, Hatcher E, Venable RM, Pastor RW, MacKerell AD Jr. CHARMM Additive All-Atom Force Field for Glycosidic Linkages Between Hexopyranoses. *J Chem Theory Comput*. 2009; 5:2353–2370. [PubMed: 20161005]
20. Raman EP, Guvench O, MacKerell AD Jr. CHARMM Additive All-Atom Force Field for Glycosidic Linkages in Carbohydrates Involving Furanoses. *J Phys Chem B*. 2010; 114:12981–12994. [PubMed: 20845956]
21. Guvench O, Mallajosyula SS, Raman EP, Hatcher E, Vanommeslaeghe K, Foster TJ, Jamison FW, MacKerell AD Jr. CHARMM Additive All-Atom Force field for Carbohydrate Derivatives and Its Utility in Polysaccharide and Carbohydrate–Protein Modeling. *J Chem Theory Comput*. 2011; 7:3162–3180. [PubMed: 22125473]
22. Huang J, MacKerell AD Jr. CHARMM36 All-Atom Additive Protein Force Field: Validation Based on Comparison to NMR Data. *J Comput Chem*. 2013; 34:2135–2145. [PubMed: 23832629]
23. Sauter J, Grafmüller A. Solution Properties of Hemicellulose Polysaccharides with Four Common Carbohydrate Force Fields. *J Chem Theory Comput*. 2015; 11:1765–1774. [PubMed: 26574386]
24. Lay WK, Miller MS, Elcock AH. Optimizing Solute–Solute Interactions in the GLYCAM06 and CHARMM36 Carbohydrate Force Fields Using Osmotic Pressure Measurements. *J Chem Theory Comput*. 2016; 12:1401–1407. [PubMed: 26967542]
25. Yoo J, Aksimentiev A. Improved Parameterization of Amine–Carboxylate and Amine–Phosphate Interactions for Molecular Dynamics Simulations Using the CHARMM and AMBER Force Fields. *J Chem Theory Comput*. 2016; 12:430–443. [PubMed: 26632962]
26. Miller MS, Lay WK, Elcock AH. Osmotic Pressure Simulations of Amino Acids and Peptides Highlight Potential Routes to Protein Force Field Parameterization. *J Phys Chem B*. 2016; 120:8217–8229. [PubMed: 27052117]

27. Miller MS, Lay WK, Li S, Hacker WC, An J, Ren J, Elcock AH. Reparametrization of Protein Force Field Nonbonded Interactions Guided by Osmotic Coefficient Measurements from Molecular Dynamics Simulations. *J Chem Theory Comput.* 2017; 13:1812–1826. [PubMed: 28296391]
28. Murad S, Powles J. A Computer Simulation of the Classic Experiment on Osmosis and Osmotic Pressure. *J Chem Phys.* 1993; 99:7271–7272.
29. Luo Y, Roux B. Simulation of Osmotic Pressure in Concentrated Aqueous Salt Solutions. *J Phys Chem Lett.* 2009; 1:183–189.
30. Yoo J, Aksimentiev A. Improved Parametrization of Li^+ , Na^+ , K^+ , and Mg^{2+} Ions for All-Atom Molecular Dynamics Simulations of Nucleic Acid Systems. *J Phys Chem Lett.* 2012; 3:45–50.
31. Yoo J, Wilson J, Aksimentiev A. Improved model of hydrated calcium ion for molecular dynamics simulations using classical biomolecular force fields. *Biopolymers.* 2016; 105:752–763. [PubMed: 27144470]
32. Yoo J, Aksimentiev A. Refined Parameterization of Nonbonded Interactions Improves Conformational Sampling and Kinetics of Protein Folding Simulations. *J Phys Chem Lett.* 2016; 7:3812–3818. [PubMed: 27617340]
33. Sauter J, Grafmüller A. Predicting the Chemical Potential and Osmotic Pressure of Polysaccharide Solutions by Molecular Simulations. *J Chem Theory Comput.* 2016; 12:4375–4384. [PubMed: 27529356]
34. Saxena A, García AE. Multisite Ion Model in Concentrated Solutions of Divalent Cations (MgCl_2 and CaCl_2): Osmotic Pressure Calculations. *J Phys Chem B.* 2015; 119:219–227. [PubMed: 25482831]
35. Huang J, Rauscher S, Nawrocki G, Ran T, Feig M, de Groot BL, Grubmüller H, MacKerell AD Jr. CHARMM36m: an improved force field for folded and intrinsically disordered proteins. *Nat Methods.* 2017; 14:71–73. [PubMed: 27819658]
36. Horn HW, Swope WC, Pitera JW, Madura JD, Dick TJ, Hura GL, Head-Gordon T. Development of an Improved Four-site Water Model for Biomolecular Simulations: TIP4P-Ew. *J Chem Phys.* 2004; 120:9665–9678. [PubMed: 15267980]
37. Reiher, W. E. I. Theoretical Studies of Hydrogen Bonding Ph D Thesis. Harvard University; Cambridge, MA, USA: 1985.
38. MacKerell AD Jr, Bashford D, Bellott M, Dunbrack RL, Evanseck JD, Field MJ, Fischer S, Gao J, Guo H, Ha S, Joseph-McCarthy D, Kuchnir L, Kuczera K, Lau FT, Mattos C, Michnick S, Ngo T, Nguyen DT, Prodhom B, Reiher WE, Roux B, Schlenkrich M, Smith JC, Stote R, Straub J, Watanabe M, Wiorkiewicz-Kuczera J, Yin D, Karplus M. All-Atom Empirical Potential for Molecular Modeling and Dynamics Studies of Proteins. *J Phys Chem B.* 1998; 102:3586–3616. [PubMed: 24889800]
39. Beauchamp KA, Lin Y-S, Das R, Pande VS. Are Protein Force Fields Getting Better? A Systematic Benchmark on 524 Diverse NMR Measurements. *J Chem Theory Comput.* 2012; 8:1409–1414. [PubMed: 22754404]
40. Uedaira H. Activity Coefficients of Glycylglycine and α -Aminobutyric Acid in Aqueous Sucrose Solutions. *Bull Chem Soc Jpn.* 1977; 50:1298–1304.
41. Robinson, RA., Stokes, RH. *Electrolyte Solutions: The Measurement and Interpretation of Conductance, Chemical Potential and Diffusion in Solutions of Simple Electrolytes.* Butterworths Scientific Publications; 1959.
42. Han H, Li D, Guo L, Yao Y, Yang H, Zeng D. Isopiestic Measurements of Water Activity for the $\text{NaCl-KCl-MgCl}_2\text{-H}_2\text{O}$ Systems at 323.15 K. *J Chem Eng Data.* 2015; 60:1139–1145.
43. Ellerton HD, Reinfelds G, Mulcahy DE, Dunlop PJ. Activity, Density, and Relative Viscosity Data for Several Amino Acids, Lactamide, and Raffinose in Aqueous Solution at 25°. *J Phys Chem.* 1964; 68:398–402.
44. Stokes R, Robinson R. Interactions in Aqueous Nonelectrolyte Solutions. I. Solute-Solvent Equilibria. *J Phys Chem.* 1966; 70:2126–2131.
45. Chatterjee JP, Basumallick IN. Thermodynamics of Transfer of Glycine, Diglycine, p-Nitroaniline and Benzoic Acid from Water to Aqueous Solutions of Polyhydroxy Compounds. *J Chem Soc Faraday Trans 1.* 1988; 84:2697–2701.

46. Kumar S, Rosenberg JM, Bouzida D, Swendsen RH, Kollman PA. The Weighted Histogram Analysis Method for Free-Energy Calculations on Biomolecules. I. The Method. *J Comput Chem.* 1992; 13:1011–1021.
47. Spiga E, Abriata LA, Piazza F, Dal Peraro M. Dissecting the Effects of Concentrated Carbohydrate Solutions on Protein Diffusion, Hydration, and Internal Dynamics. *J Phys Chem B.* 2014; 118:5310–5321. [PubMed: 24773474]
48. Batista ML, Perez-Sanchez G, Gomes JR, Coutinho JA, Maginn EJ. Evaluation of the GROMOS 56ACARBO Force Field for the Calculation of Structural, Volumetric, and Dynamic Properties of Aqueous Glucose Systems. *J Phys Chem B.* 2015; 119:15310–15319. [PubMed: 26572856]
49. Horta BAC, Merz PT, Fuchs PFJ, Dolenc J, Riniker S, Hünenberger PH. A GROMOS-Compatible Force Field for Small Organic Molecules in the Condensed Phase: The 2016H66 Parameter Set. *J Chem Theory Comput.* 2016; 12:3825–3850. [PubMed: 27248705]
50. Kang M, Smith PE. A Kirkwood-Buff Derived Force Field for Amides. *J Comput Chem.* 2006; 27:1477–1485. [PubMed: 16823811]
51. Benteñitis N, Cox NR, Smith PE. A Kirkwood-Buff Derived Force Field for Thiols, Sulfides, and Disulfides. *J Phys Chem B.* 2009; 113:12306–12315. [PubMed: 19681588]
52. Ploetz EA, Benteñitis N, Smith PE. Developing Force Fields from the Microscopic Structure of Solutions. *Fluid phase equilib.* 2010; 290:43. [PubMed: 20161692]
53. Chapman DE, Steck JK, Nerenberg PS. Optimizing Protein–Protein van der Waals Interactions for the AMBER ff9x/ff12 Force Field. *J Chem Theory Comput.* 2014; 10:273–281. [PubMed: 26579910]
54. Nerenberg PS, Jo B, So C, Tripathy A, Head-Gordon T. Optimizing Solute–Water van der Waals Interactions To Reproduce Solvation Free Energies. *J Phys Chem B.* 2012; 116:4524–4534. [PubMed: 22443635]
55. Debiec KT, Cerutti DS, Baker LR, Gronenborn AM, Case DA, Chong LT. Further along the Road Less Traveled: AMBER ff15ipq, an Original Protein Force Field Built on a Self-Consistent Physical Model. *J Chem Theory Comput.* 2016; 12:3926–3947. [PubMed: 27399642]

Figure 1. Comparison of osmotic pressures obtained from simulations with experiment
A. Osmotic pressures of diglycine/sucrose solutions plotted versus mole fraction of sucrose. The expected osmotic pressure exerted based on experimental data is 36.17 bar^{40, 44} and depicted by the dashed horizontal line; data were obtained from simulations using the original AMBERff99SB/GLYCAM06 and CHARMM36 force fields. Error bars were obtained by averaging the osmotic pressures in 20 ns blocks and calculating the standard deviation of these averages. **B.** Same as **A** but showing results obtained with modified sugar-sugar nonbonded parameters in the AMBERff99SB/GLYCAM06 and CHARMM36 force

fields. **C.** Same as **A** but showing results obtained with modified sugar-sugar + amino acid-sugar parameters. Osmotic pressures are given in bar units and diglycine/sucrose compositions are expressed relative to mole fraction sucrose (i.e. 0 mol fraction sucrose indicates diglycine only in water and *vice versa*).

Figure 2. Potentials of mean force (PMF) for the transfer of diglycine from water to an aqueous sucrose solution

A. Setup for umbrella sampling simulations. Sucrose molecules are confined to the region between the two gray walls while water molecules (not shown) can move freely, effectively creating a 15 wt% sucrose solution at the center region of the unit cell. A series of umbrella sampling simulations were performed where a single diglycine molecule is pulled across the entire length of the cell (see Computational Methods in the SI for details). **B.** Free energy change, ΔG , as a function of the z-position of the central nitrogen atom of diglycine. Dashed vertical lines at 24 and 72 Å indicate the left and right boundaries of the aqueous sucrose

solution. Umbrella sampling simulations were performed with the original (blue circles), modified sugar-sugar (green upward triangles), and modified sugar-sugar + amino acid-sugar parameters (red downward triangles). Results obtained with the AMBERff99SB/GLYCAM06 force field are shown on this panel. **C.** Same as **B** but showing results for the CHARMM36 force field.

Figure 3. Final snapshots of simulations of ubiquitin in aqueous glucose solutions using original and modified AMBERff99SB/GLYCAM06 force fields

A. snapshot obtained after 100 ns of MD using original parameters. **B.** same as **A** but showing result obtained with modified sugar-sugar parameters. **C.** same as **A** but showing result obtained with modified sugar-sugar + amino acid-sugar parameters.

Figure 4. Minimum distance histograms of glucose molecules to the ubiquitin with original and adjusted parameters

Histograms depicting the sampling of the minimum distance of each glucose molecule to the ubiquitin using original (blue line), modified sugar—sugar (green line) and modified sugar—sugar + sugar—amino acid (red line) parameters for AMBERff99SB/GLYCAM06. Each 100 ns trajectory was analyzed at every 1 ps and the histogram binned at 0.1 Å.