Published in final edited form as:

J Allergy Clin Immunol Pract. 2018 ; 6(2): 677–679. doi:10.1016/j.jaip.2017.09.023.

Food allergen component proteins are not detected in early-childhood vaccines

Alice E. W. Hoyt, M.D.^{1,2}, Martin D. Chapman, Ph.D.⁴, Eva M. King, M.Sc., Ph.D.⁴, Thomas A. E. Platts-Mills, M.D., Ph.D., F.R.S.^{1,3}, and John W. Steinke, Ph.D.^{1,3,**}

¹Department of Medicine, University of Virginia Health System, Charlottesville, VA 22908 USA

²Department of Pediatrics, University of Virginia Health System, Charlottesville, VA 22908 USA

³Carter Immunology Center, University of Virginia Health System, Charlottesville, VA 22908 USA

⁴Indoor Biotechnologies, Charlottesville, VA 22903 USA

Keywords

Food allergy; vaccines; immunizations; component proteins; critical window; hepatitis B vaccine

To the Editor

Since the 1990's, a substantial increase has occurred in the incidence of food allergy, particularly in early childhood.¹ Although some vaccines contain potential allergens,^{2,3} social media and internet websites has promulgated the concern that vaccines contain food proteins, which then induce sensitization. Parental unease regarding the immunologic impact of vaccines appears to be quite common as 72.5% of pediatricians report that parents are concerned that the number of vaccines recommended stress their child's immune system and may be the cause of a variety of diseases.⁴ Studies that address a potential association between vaccines and Th2 immunity are limited. The likelihood is low that the early-childhood vaccines would contain food allergens. However, casein derivatives are used in the manufacture of many vaccines, egg protein can be found in influenza and yellow fever vaccines and parents may not understand that only these two vaccines are made with chicken eggs, and there is always the worry of contamination or cross-reactivity with allergen components. This study sought to address parental – and physician – concern that early-childhood vaccines given within the first 6–18 months contain food allergens.

The immune response that underlies allergic disease is thought to begin during a critical window in early childhood.⁵ Given that some children already have manifestations of allergic disease as early as two months of age, this investigation initially targeted the earliest

^{**}To whom correspondence should be addressed at: Asthma and Allergic Disease Center, Box 801355, Charlottesville, VA 22903
phone #: (434) 243-6570; fax #: (434) 982-3270; js3ch@virginia.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

vaccine administered to children: the hepatitis B (HepB) vaccine. The CDC recommends that the HepB vaccine be administered within 24 hours of birth and 2–3 subsequent times in the first year of life.⁶ Because Vitamin K (VitK) is also administered to newborns at birth (to prevent hemorrhagic disease of the newborn), it too may be considered a candidate for early-life induction of sensitization and was included in the study. As early as six weeks of age, children may receive Pediarix, which protects against HepB as well as DTaP and poliovirus. Given Pediarix contains HepB and may be administered within the potential critical window, we also included it in our investigation. As the time-frame for early sensitization is not known, we included two other 1st year vaccines in our analysis: Prevnar 13 and ActHIB. The aim was to determine whether component proteins of the common food allergens peanut, egg, and cow's milk were detectable in these early life injections. These three allergens were selected based on the prevalence of these allergies in children.

To investigate the possibility that early-childhood vaccines and injectable VitK contain food proteins, we used ELISA to detect specific food allergens. Engerix-B (one of two brands of HepB-only containing vaccines administered at birth), Pediarix (which contains HepB, DTaP, and poliovirus), Prevnar 13, ActHIB and VitK were purchased through the University of Virginia Pharmacy. Vaccine samples were analyzed using highly sensitive allergen-specific ELISA methods for five peanut allergens, three milk allergens, and one egg allergen (Indoor Biotechnologies, Inc.). The antibody pairs used in each assay are specified in brackets as follows:

- Peanut – Ara h 1, [2C12/2F7]; Ara h 2, [1C4/pAb]; Ara h 3, [1E8/4G9]; Ara h 6, [3B8/3E12]; and Ara h 8, [4G6/pAb];
- Milk – beta-lactoglobulin, as native Bos d 5, [NBD5-1/NBD5-2]; as denatured Bos d 5, [DBD5-1/DBD5-2]; and as beta casein Bos d 11 [CC11/VB1C];
- Egg – Gal d 1 [CB-1/CB-2].

The Limits of Detection (LOD) of the ELISA were: Ara h 1, 156.25 ng/ml; Ara h 2, 2.44 ng/ml; Ara h 3, 4.88 ng/ml; Ara h 6, 0.98 ng/ml; Ara h 8, 4.88 ng/ml; nBos d 5, 0.78 ng/ml; dBos d 5, 7.81 ng/ml; Bos d 11, 31.25 ng/ml; and Gal d 1, 13.28 ng/ml. Samples from two lots of Engerix-B, four lots of Pediarix, two lots of ActHIB, one lot of Prevnar 13 and one lot of injectable VitK were tested.

The results demonstrate that none of the nine specific food allergens from peanut, cow's milk, and hen's egg were detected in Engerix-B, Pediarix, Prevnar 13, ActHIB or injectable VitK (Table I). These findings do not support the concept that early-childhood vaccines or VitK injections contain allergy-causing food allergens. Without evidence of significant food allergen within the vaccines, the aforementioned hypothesis of vaccine-associated sensitization is unsupported. These findings also suggest that there are no cross-reacting components in the ingredients listed on the packaging (Table II) with any of the food allergens tested. While the ELISA detection limits are quite low, the amount of allergen required to cause sensitization is poorly understood. Theoretically, it could be argued that the ELISA limits of detection may not be low enough, though this seems unlikely as these are the most sensitive assays available with detection limits in the nanogram range and have been used before to detect these antigens.⁷ What is reassuring is that children with peanut,

cow's milk, and egg allergy continue to receive these vaccines in the first year of life without having allergic reactions. This implies that even if allergen was present (in sub-picogram amounts), the quantities are too low for the immune system to mount an IgE-mediated allergic response.

This study does not negate the concept that other components of the vaccine, such as aluminum adjuvant, could alter the immune response in a developing child. It is well known that aluminum adjuvant induces a robust immune response that results in great protection by many vaccines. While the effect on IgE production in animal models is somewhat understood, the effect of aluminum adjuvant in humans is uncertain. The impact on IgE production is particularly unclear during early childhood, when the immune system is developing away from the tolerant, fetal phenotype and into a more mature, adult immune system. It has been suggested that allergen exposure at the time of vaccination may direct the immune response towards a Th2 bias leading to allergen sensitization. This is unlikely for the HepB vaccine and VitK as they are given at the time of birth before allergen exposure would occur. What is highly unlikely – and refuted in our study – is that early-life vaccines and injectable VitK contain food allergens and thereby contribute to the increase in food allergy. Some vaccines do contain food proteins, such as gelatin, and people with gelatin allergy, including patients with the galactose-alpha-1,3-galactose (alpha-gal) allergy, should consult with their allergy-immunology specialist regarding questions about vaccines.⁸ In the majority of cases, the benefits of the vaccine outweigh the risk of a reaction, and the vaccine can be administered in the specialist's office. Overall, the concept that food proteins in early-life vaccines are the cause of food allergy is unlikely and unsupported and, thus, should not be a concern when considering vaccination.

Acknowledgments

Funding

This work was supported by National Institutes of Health grant AI20565 and by the financial support of local donor Panera Bread of Charlottesville.

Abbreviations

DTaP	diphtheria, tetanus, acellular pertussis
HepB	hepatitis B
VitK	vitamin K

References

1. Branum AM, Lukacs SL. Food allergy among U.S. children: trends in prevalence and hospitalizations. NCHS Data Brief. 2008; (10):1–8.
2. Kattan JD, Konstantinou GN, Cox AL, et al. Anaphylaxis to diphtheria, tetanus, and pertussis vaccines among children with cow's milk allergy. J Allergy Clin Immunol. 2011; 128(1):215–218. [PubMed: 21624648]
3. Stone CA, Hemler JA, Commins SP, et al. Anaphylaxis after zoster vaccine: Implicating alpha-gal allergy as a possible mechanism. J Allergy Clin Immunol. 2016

4. Hough-Telford C, Kimberlin DW, Aban I, et al. Vaccine Delays, Refusals, and Patient Dismissals: A Survey of Pediatricians. *Pediatrics*. 2016; 138(3)
5. Prescott SL, Smith P, Tang M, et al. The importance of early complementary feeding in the development of oral tolerance: concerns and controversies. *Pediatr Allergy Immunol*. 2008; 19(5): 375–380. [PubMed: 18266825]
6. Prevention CDC. Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger, UNITED STATES, 2017. 2017 [Accessed March 14, 2017, 2017]
7. Filep S, Block DS, Smith BRE, et al. Specific allergen profiles of peanut foods and diagnostic or therapeutic products. *J Allergy Clin Immunol*. 2017 Jul 12. pii: S0091-6749(17)31091-6 [Epub ahead of print]. doi: 10.1016/j.jaci.2017.05.049
8. Mullins RJ, James H, Platts-Mills TA, Commins S. Relationship between red meat allergy and sensitization to gelatin and galactose- α -1,3-galactose. *J Allergy Clin Immunol*. 2012; 129(5):1334–1342. e1331. [PubMed: 22480538]
9. Prevention CDC. [Accessed March 14, 2017, 2017] The Pink Book. 2017. <https://www.cdc.gov/vaccines/pubs/pinkbook/downloads/appendices/B/excipient-table-2.pdf>

Clinical Implications

Limited studies have explored the potential impact of vaccines on food allergy development. This study investigates and does not support the theory that early-childhood vaccines contain specific food allergens that induce sensitization.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

TABLE I
Absence of food allergens in early-childhood vaccines and in Vitamin K injection

Medication	Lot #	ng/mL								
		Ara h 1	Ara h 2	Ara h 3	Ara h 6	Ara h 8	nBos d 5	dBos d 5	Bos d 11	Gal d 1
Engerix-B	EG9Y2	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Engerix-B	EG9Y2	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Engerix-B	P7XF7	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
ActHIB	UI787AAA	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
ActHIB	UI735AAA	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Pediarix	3J9R3	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Pediarix	437M5	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Pediarix	529PP	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Pediarix	FY7FK	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Prevnar 13	S31255	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3
Vitamin K	58-272-EV	<156	<2.44	<4.88	<0.98	<4.88	<0.78	<7.81	<31.3	<13.3

Engerix-B contains hepatitis B virus surface antigen (recombinant). *Pediarix* contains hepatitis B virus surface antigen (recombinant); diphtheria and tetanus toxoids and acellular pertussis (DTaP); and inactivated poliovirus vaccine. *Prevnar 13* contains pneumococcal 13-valent conjugate. *Vitamin K* contains phytonadione. Complete list of product ingredients can be found in Table II. ELISA LOD is based on testing vaccines at 1/5 dilution.

TABLE II

Ingredients in early-childhood vaccines⁹ and in Vitamin K injection

Medication	Vaccine Antigens	Ingredients
Engerix-B	Hepatitis B	Aluminum hydroxide, yeast protein, sodium chloride, disodium phosphate dihydrate, sodium dihydrogen phosphate dihydrate
ActHIB	Haemophilus b	sodium chloride, modified Mueller and Miller medium (the culture medium contains milk-derived raw materials [casein derivatives]), formaldehyde, sucrose
Pediarix	Hepatitis B Diphtheria, Tetanus, acellular Pertussis (DTaP) Poliovirus	Fenton medium containing a bovine extract, modified Latham medium derived from bovine casein, formaldehyde, modified Stainer-Scholte liquid medium, VERO cells, a continuous line of monkey kidney cells, calf serum and lactalbumin hydrolysate, aluminum hydroxide, aluminum phosphate, aluminum salts, sodium chloride, polysorbate 80 (Tween 80), neomycin sulfate, polymyxin B, yeast protein
Prevnar 13	Pneumococcal 13-valent	soy peptone broth, casamino acids and yeast extract-based medium, CRM197 carrier protein, polysorbate 80, succinate buffer, aluminum phosphate
Vitamin K	N/A	Phytonadione, polyoxyethylated fatty acid derivative, hydrous dextrose, benzyl alcohol; may contain hydrochloric acid

N/A, not applicable.