Published in final edited form as:

Physiol Behav. 2018 May 01; 188: 11–17. doi:10.1016/j.physbeh.2018.01.019.

Apolipoprotein A-IV enhances cholecystokinin secretion

Jesse Zhan^{1,2}, Jonathan Weng^{1,3}, Brian G Hunt², W. Sean Davidson², Min Liu², and Chunmin C Lo¹

¹Department of Biomedical Sciences, Diabetes Institute and Heritage College of Osteopathic Medicine, Ohio University, Athens, OH, USA

²Department of Pathology and Laboratory Medicine, Metabolic Diseases Institute, University of Cincinnati, Cincinnati, OH, USA

³Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY, USA

Abstract

Cholecystokinin (CCK) and apolipoprotein A-IV (ApoA-IV) are gastrointestinal peptides that play an important role in controlling energy homeostasis. Lymphatic ApoA-IV and plasma CCK secretion are mediated via a chylomicron formation-dependent pathway during a dietary lipid infusion. Given their similar roles as satiating proteins, the present study examines how the two peptides interact in their function. Specifically, this study sought to understand how ApoA-IV regulates CCK secretion. For this purpose, *Cck* gene expression in the small intestines of ApoA-IV knockout (ApoA-IV-KO) and wild-type (WT) mice were compared under an array of feeding conditions. When fed with a chow or high-fat diet (HFD), basal levels of *Cck* transcripts were significantly reduced in the duodenum of ApoA-IV-KO mice compared to WT mice. Furthermore, after an oral gavage of a lipid mixture, *Cck* gene expression in the duodenum was significantly reduced in ApoA-IV-KO mice relative to the change seen in WT mice. To determine the mechanism by which ApoA-IV modulates *Cck* gene expression, STC-1 cells were transfected with predesigned mouse lysophosphatidic acid receptor 5 (LPAR5) small interfering RNA (siRNA) to knockdown *Lpar5* gene expression. In this *in-vitro* study, mouse recombinant ApoA-IV protein increased *Cck* gene expression in enteroendocrine STC-1 cells and stimulated CCK release from the STC-1 cells. However, the levels of CCK protein and *Cck* expression were attenuated when *Lpar5* was knocked down in the STC-1 cells. Together these observations suggest that dietary lipid-induced ApoA-IV is associated with *Cck* synthesis in the duodenum and that ApoA-IV protein directly enhances CCK release through the activation of a LPAR5-dependent pathway.

Keywords

Apolipoprotein A-IV; Cholecystokinin; lipids; STC-1 cells

Address correspondence to: Chunmin Lo, Departments of Biomedical Sciences, Irvine Hall 228, 1 Ohio University, Athens, OH 45237-0507, USA, Phone : 740-593-2328, Fax : 740-597-2778, loc1@ohio.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Obesity is a global epidemic that increases the risk of type II diabetes mellitus, cardiovascular disease, and many cancers (2, 8, 15, 46). Consumption of a high-fat diet (HFD) promotes positive energy balance, which is stored as fat and leads to subsequent development of obesity and insulin resistance (19, 48). Gastrointestinal peptides such as cholecystokinin (CCK) and apolipoprotein A-IV (ApoA-IV) are important satiating proteins for controlling energy homeostasis (21, 47). Both ApoA-IV and CCK inhibit gastric emptying and food intake (11, 12, 18, 35, 45). There are two types of CCK receptors that exist in humans and rodents: CCK A receptors (CCK-AR), which are more abundant in peripheral tissues than in the brain, and CCK B receptors (CCK-BR), which are considered the gastric receptor and are abundantly present in both peripheral tissues and the brain (40). Peripherally administered ApoA-IV and CCK do not cross the blood-brain barrier (31, 39). CCK acts by directly binding to CCK-AR on vagal afferent neurons in the intestinal enteric plexuses to relay satiation signals to the hypothalamus via vagal nerve endings in the nucleus of the solitary tract in the hindbrain (31, 32). ApoA-IV is involved in lipid-induced activation of the vagal afferent pathway and requires vagal nerve transmission to activate neurons in the hindbrain to suppress food intake (13, 28, 45). Together, ApoA-IV and CCK work synergistically to reduce food intake, specifically, ApoA-IV requires a CCK-dependent pathway to induce neuronal activation in the NTS and suppress food intake (29, 50). Thus, the interaction of ApoA-IV and CCK are important in controlling energy homeostasis through the stimulation of vagal activity.

Dietary lipids and proteins stimulate CCK release in the small intestine (24, 37). In addition, different species have different gastrointestinal physiologies in response to nutrients. In humans, dietary lipids are the major physiological stimuli of CCK secretion, whereas in rats dietary proteins are the strongest stimulant of CCK secretion (9, 25). This response occurs in a dose-dependent manner, which CCK secretion by endocrine I cells in the small intestine is elevated with higher doses of lipids (24, 37). ApoA-IV, apolipoprotein A-I (ApoA-I) and apolipoprotein B (ApoB) are major protein constituents of lymphatic triglycerides (TG)-rich lipoproteins and are produced by enterocytes (14, 17). The ApoA-IV protein is the only apolipoprotein that responds to varying amounts of dietary lipids, whereas lymphatic secretion of ApoA-I and ApoB proteins remains consistent (1, 16, 17, 38, 43). During a dietary lipid infusion, lymphatic ApoA-IV and plasma CCK secretion are mediated via a chylomicron formation-dependent pathway (17, 34). These findings imply that dietary lipids are important regulators of ApoA-IV and CCK production in the small intestine and induce secretion of CCK and ApoA-IV in a co-dependent manner. However, the effect of ApoA-IV on the regulation of intestinal CCK remains unknown. Protein and protein hydrolysates (peptones) directly interact with a lysophosphatidic acid receptor 5 (LPAR5 is also known as G protein-coupled receptor 93)-dependent pathway to produce a marked elevation in CCK release (5, 10, 25). It was hypothesized that ApoA-IV enhances CCK levels through the activation of a LPAR5-dependent pathway as well. The present study was designed to determine whether intestinal *Cck* gene expression in ApoA-IV-KO and WT mice would be altered after diets with different fat contents, and whether ApoA-IV-induced CCK secretion by mouse enteroendocrine STC-1 cells is mediated via a LPAR5-dependent pathway.

Material and Methods

Animals

ApoA-IV -KO mice were kindly provided by Dr. Jan Breslow (Rockefeller University, New York, NY) and were backcrossed for >15 generations onto a C57BL/6J genetic background. All mice were genotyped by polymerase chain reaction (PCR) analysis of tail deoxyribonucleic acid (DNA) (6, 20). ApoA-IV-KO and wild-type (WT) mice (C57BL/6J) were housed in an AAALAC-accredited facility under conditions of controlled illumination (12:12-h light-dark cycle, lights from 0600 to 1800 h). All animal protocols were approved by the Institutional Animal Care and Use Committee in Ohio University and the University of Cincinnati.

Animal feeding and small intestine collection

Starting at 10 weeks of age, male ApoA-IV -KO and WT mice received free access to water and chow diet (6% fat, LM-485 Mouse/Rat Sterilizable Diet, Envigo, Madison, WI). To confirm that the duodenum is a major site for CCK synthesis in ApoA-IV-KO and WT mice, the small intestines of 5 h-fasted, 18 weeks old WT and ApoA-IV-KO mice maintained on a chow diet were collected. The basal *Cck* gene expression levels of these samples were then determined for experiment 1. For the study of lipid-induced CCK synthesis in ApoA-IV-KO mice in experiment 2, chow-fed WT and ApoA-IV-KO mice at 15 weeks of age received a lipid mixture containing 2.4 mg triolein, 0.2 mg cholesterol (CHOL) and 0.2 mg phospholipids (200 µl) by gavage after a 5-h fast. Two h later, the small intestine was collected on dry ice for the determination of duodenal CCK levels. To examine the influence of HFD on *Cck* gene expression in the duodenum in experiment 3, WT and ApoA-IV-KO mice at 12 weeks of age were fed with either a semi-purified high-fat diet (HFD, 20% butter fat by weight) or a matched low-fat diet (LFD, 5% butter fat content, Research Diets, Inc, New Brunswick, NJ) for 8 weeks (20). After a 5-h fast, the duodenum was collected on dry ice for the determination of *Cck* gene expression using a real-time quantitative reverse transcription (qRT-PCR) analysis.

STC-1 cells

STC-1 cells, a mouse-intestinal cell-line, were provided by Dr. Douglas Hanahan (UCSF) (3, 5). Briefly, STC-1 cells were grown in 6-well plate with Dulbecco's modified Eagles medium, 10% fetal bovine serum, penicillin (100IU/ml) and streptomycin (100 µg/ml). To determine basal level of CCK, the STC-1 cells (80 % confluency) were incubated with Hanks' balanced salt solution (HBSS) as a vehicle for 0.5, 1 and 2 h. Then the media was collected for measuring the level of CCK. To determine whether CCK release by ApoA-IV increased in a dose-dependent manner, STC-1 cells were incubated with different doses of mouse recombinant ApoA-IV proteins (50, 100, 200 and 400 µg/ml) for 0.5, 1 or 2 h. After incubation the media was collected for CCK determination. To compare CCK secretion by STC-1 cells induced by bovine serum albumin (BSA) and ApoA-IV, STC-1 cells was incubated with BSA (Sigma-Aldrich, St. Louis, MO) or mouse recombinant ApoA-IV protein at 200 µg/ml.

To determine whether the stimulation of CCK by ApoA-IV is mediated via the LPAR5 pathway, *Lpar5*-expressing STC-1 cells were transfected with predesigned mouse *Lpar5* small interfering RNA (siRNA) to knockdown *Lpar5* gene expression. In a 35-mm tissue culture plate, 2×10^5 STC-1 cells per well were seeded in 2 ml-antibiotic-free normal growth medium supplemented with fetal bovine serum 24 h before transfection. Specific siRNAs targeting mouse *Lpar5* gene (catalog number sc-75195) and control siRNAs (catalog number sc-37007) were obtained from Santa Cruz Biotechnology (Santa Cruz Biotechnology, Inc., Dallas, TX) and utilized according to the manufacturer's instruction. Five groups for each treatment were used in this experiment, including five untransfected groups without either siRNA transfection medium or siRNA treatment as the negative controls, five groups incubated with a mixture of siRNA transfection medium plus *Lpar5* siRNA (60 pmols), and five groups incubated with mixture of siRNA transfection medium plus control siRNA (0.9 μ mol). Briefly, 6 μ l of control siRNAs (0.1 μ moles) were diluted in 200 μ l siRNA transfection medium and 6 μ l of siRNA transfection reagent (Santa Cruz) in each well of a 6-well culture plate. In parallel, 6 μ l of the siRNA duplex (60 pmols siRNA) were diluted in 200 μ l of the siRNA transfection Medium and 6 μ l of the siRNA transfection reagent. The mixtures were incubated for 45 min at room temperature for complex formation and then added to the medium with STC-1 cells (80% confluency) without antibiotics at 37 °C. After a 7-h incubation, STC-1 cells were incubated with normal growth medium for an additional 18-h incubation. After 3 washes with ice-cold phosphate buffered saline (PBS, Gibco™, Fisher Scientific), STC-1 cells were incubated with ApoA-IV (200 μ g/ml) and vehicle (HBSS). After, the cells were washed with ice-cold PBS and harvested for qRT-PCR analysis, while the medium was assessed for CCK level using a CCK enzyme immunoassay kit assay.

qRT-PCR for *Cck* mRNA measurement

Total RNA was isolated from the duodenum, jejunum and ileum using Tri-Reagent (Ambion, Austin, TX). Total RNA was isolated, and first-strand cDNA was synthesized from 1 μ g of total RNA (50). qRT-PCR was performed using SYBR green RT-PCR master mixes at a 25- μ l final reaction volume with the Bio-Rad iCycler iQ instrument (Hercules, CA) according to the manufacturer's instructions. Cyclophilin mRNA levels from each sample were used as internal controls to normalize mRNA levels. The sequences of the primers (Integrated DNA Technologies, Coralville, IA) were as follows: mouse *Cck*, 5'-CTA GCGCGATACATCCAGCAGGTC-3' (forward) and 5'-ACTTAATAAATAGATACTCAAACC-3' (reverse); mouse *Lpar5*, 5'-GCTCTGCCTGGGCGTGTGGGCTCTCATCCTGC-3' (forward)(5) and 5'-GCGTCGGGCGCTCGCCAGTGTCCAGAAGAC-3' (reverse); and mouse *cyclophilin*, 5'-TTCATGTGCCAGGGTGGTGACT-3' (forward) and 5'-TCAGTCTTGGCAGTGCAGAT-3' (reverse). Threshold cycle readings for each of the unknown samples were used, and the results were analyzed in Excel using the Ct method (27).

Measurement of the CCK release

CCK immunoreactivity was measured using an enzyme immunoassay kit (Phoenix Pharmaceuticals, Inc, Burlingame, CA) that cross-reacts at 100% with CCK-33 and CCK-8,

and at 100% with sulfated gastrin-17(5). 50 µl of samples or standards were added to a 96-well microtiter plate pre-coated with anti-CCK. The detection antibody was added to the captured peptide molecules according to the manufacture's specifications. After an incubation period, 100 µl of a streptavidin-horseradish peroxidase conjugate and 3,3',5,5'-tetramethylbenzidine were added to the mixture. The absorbance was read at 450 nm using a microplate reader (Synergy HT, BioTek Instruments, Inc, Richmond, VA). According to the manufacture's specifications, the O.D. absorbance of each sample was determined at 450nm using a microplate reader. The concentration of CCK was determined by locating the sample's optical density (O.D.) absorbance on the Y-axis, then drawing a horizontal line to intersect with the standard curve.

To verify CCK sensitivity and recovery by the enzyme immune assay kit, a series of mouse CCK-33 standard solutions (#06911, Phoenix Pharmaceuticals, Inc) were prepared according to the manufacture's specifications. Varying concentrations of CCK33 were dissolved in 500µl Dulbecco's modified Eagles medium (GE Healthcare Life Science, #SH3002101) and 50 µl of standards and samples were added to each well of the 96-well microtiter plate. The R^2 for the CCK-33 standard curve was determined to be 0.9637 and the sample quality control was calculated to be 0.34 ng/ml (sample quality control range: 0.15-0.35ng/ml provided by the manufacture's protocol). The CCK levels were then converted to pmol/mL CCK (i.e.CCK-33 equivalents). To determine mouse CCK-33 recovery using this assay, CCK-33 at 0.025 ng or 0.25 ng was added to 50 µl of medium. CCK recovery (%) was calculated by dividing the concentration of CCK-33 obtained from the standard curve by the known concentration of CCK-33 added (determined by using the kit). The recovery for CCK-33 at 0.5 ng was $48.0 \pm 1.0\%$ (n=2) and the recovery for CCK-33 at 5 ng was $90.5 \pm 10.5\%$ (n=2).

Plasma parameters

Plasma TG and cholesterol (CHOL) were determined using Randox triglyceride kits (Antrim, UK) and Infinity cholesterol kits (Thermo Electron, Noble Park, Victoria, Australia), respectively.

Data analysis and statistical analysis

All values are presented as mean $\pm$ SE. Parametric statistical analyses, one-way analysis of variance (one-way ANOVA), and two-way analysis of variance (two-way ANOVA) were performed using GraphPad™ Prism (version 6.0, San Diego, CA), followed by a *post hoc* Sidak correction. All differences were considered significant if the P value was < 0.05 .

Results

Cck gene expression in ApoA-IV-KO mice

To confirm that the duodenum is a major site for CCK synthesis in ApoA-IV-KO and WT mice, *Cck* gene expression in the small intestine of 5 h-fasted WT and ApoA-IV-KO mice were determined when maintained on a chow diet in experiment 1. In the WT mice, the *Cck* mRNA level in the duodenum was 4-fold higher than that in the jejunum ($P < 0.05$, Figure 1). Relative to the WT mice, ApoA-IV-KO mice had significantly lower levels of *Cck* gene

expression in the duodenum, but comparable levels of *Cck* expression in the jejunum and ileum ($P < 0.05$, Figure 1). These findings indicate that the duodenum is a major site for CCK synthesis at basal levels in both genotype mouse models. In addition, the absence of ApoA-IV reduces basal CCK synthesis in the duodenum when animals are maintained on a chow diet.

Dietary lipids stimulate ApoA-IV synthesis in the small intestine (43). To investigate whether dietary lipid-induced ApoA-IV enhances CCK synthesis in the duodenum, duodenal *Cck* gene levels in WT and ApoA-IV-KO mice were determined following an acute oral gavage of lipid mixtures in experiment 2. Since TG levels are maximized at 2 h in response to a gastric gavage of dietary lipids (49), the duodenum was collected 2 h after infusion of lipid mixtures. When ApoA-IV-KO and WT mice were maintained on chow diets, no differences in body weight, fat mass, or plasma levels of TG and CHOL at 0 h were found between the two genotypes (Table 1). In the WT mice, plasma TG and CHOL were elevated 2 h after infusion of dietary lipids relative to the fasted group ($P < 0.05$, Table 1). Compared to the WT mice, ApoA-IV-KO mice had significantly reduced plasma TG ($P < 0.05$, Table 1), but comparable levels of plasma CHOL 2 h after infusion of lipid mixtures. ApoA-IV-KO mice had comparable basal levels of *Cck* gene expression in the duodenum relative to WT mice (Figure 2). Relative to basal levels of duodenal *Cck* gene expression in the WT mice, dietary lipids induced a 17-fold increase in duodenal *Cck* gene expression in WT mice and a 8-fold increase in duodenal *Cck* gene expression in ApoA-IV-KO mice ($P < 0.05$, Figure 2). These observations indicate that dietary lipids enhance CCK synthesis in the duodenum, and lipid-induced CCK synthesis is associated with ApoA-IV.

To determine the influence of a HFD on *Cck* gene expression in the duodenum, WT and ApoA-IV-KO mice were fed with either a LFD or a HFD for 8 weeks in experiment 3. HFD-fed ApoA-IV-KO mice have been reported to have comparable body weight and reduced levels of TG and CHOL in the plasma relative to WT mice (20). In current experiment, LFD-fed ApoA-IV-KO mice had lower duodenal *Cck* gene expression than the LFD-fed WT mice, however, the difference in *Cck* levels was not statistically significant (Figure 3). In the WT mice, a HFD increased basal levels of *Cck* gene expression relative to a LFD treatment ($P < 0.05$, Figure 3). Duodenal *Cck* gene expression in the HFD-fed ApoA-IV -KO mice was significantly lower than that in the HFD-fed WT mice ($P < 0.05$, Figure 3). These findings suggest that ApoA-IV is partially involved in the control of duodenal *Cck* gene expression during a chronic feeding of dietary lipids.

ApoA-IV induced CCK release by STC-1 cells

To determine if ApoA-IV would dose-dependently increase the levels of CCK protein, STC-1 cells were incubated with different doses of mouse recombinant ApoA-IV (50 to 400 $\mu\text{g/ml}$)(3, 5). After a 0.5-h incubation, the CCK levels in the media of STC-1 cells were comparable between ApoA-IV- and vehicle-treated cells; or above 100 $\mu\text{g/ml}$ significantly elevated CCK levels in the media of STC-1 cells after a 1-h and 2-h incubation of STC-1 cells ($P < 0.05$, Figure 4B & 4C). To determine the specificity of ApoA-IV's effect on CCK secretion, STC-1 cells were incubated with BSA or ApoA-IV (both at a dose of 200 $\mu\text{g/ml}$) for 2 h. Relative to vehicle treatment (0.059 ± 0.0 pmol/mL equivalent to CCK-33), ApoA-

IV significantly elevated CCK release (0.16 ± 0.03 pmol/ml) in the media of STC-1 cells ($P < 0.05$, Figure 5). In contrast, BSA induced comparable CCK release in the medium of STC-1 cells (Figure 5). ApoA-IV stimulated a 2-fold increase in CCK levels in the medium of STC-1 cells relative to BSA treatment at the same dose. These observations suggest that BSA is unable to stimulate CCK release by STC-1 cells.

ApoA-IV-induced CCK secretion from STC-1 cells through a LPAR5-dependent pathway

Protein hydrolysates interact with LPAR5 in the membrane of the enteroendocrine STC-1 cells to induce CCK release (5). To determine if the LPAR5-signaling pathway also mediates the action of ApoA-IV-induced CCK release, STC-1 cells treated with *Lpar5* siRNA (knockdown) or scramble siRNA (negative control) were incubated with ApoA-IV at 200 μ g/ml or vehicle. Relative to *Lpar5* gene expression in the vehicle-treated STC-1 cells without knockdown of *Lpar5*, there was an 89% reduction of *Lpar5* expression in vehicle-treated STC-1 cells with knockdown of *Lpar5* using a SiRNA treatment ($P < 0.05$, Figure 6A). After a 2-h incubation, there was no difference in vehicle-induced *Cck* gene expression between STC-1 cells treated with or without *Lpar5* siRNA treatment (Figure 6B), suggesting that knockdown of *Lpar5* gene did not alter basal levels of CCK synthesis in the STC-1 cells. ApoA-IV at 200 μ g/ml increased *Cck* gene expression in STC-1 cells treated with scramble siRNA ($P < 0.05$, Figure 6B). In contrast, ApoA-IV failed to elevate *Cck* expression in the STC-1 cells treated with *Lpar5* siRNA. Consistent with the induction of *Cck* gene expression by ApoA-IV in Figure 6B, the CCK protein levels were also increased by ApoA-IV in the culture medium of STC-1 cells treated with scrambled siRNA, relative to vehicle treatment ($P < 0.05$, Figure 6C). And assuredly, ApoA-IV-induced CCK levels in the medium of STC-1 cells transfected with *Lpar5* siRNA were attenuated. These observations suggest that ApoA-IV interacts with membrane LPAR5 to stimulate *Cck* synthesis and CCK protein secretion in STC-1 cells.

Discussion

CCK is a well-known satiating protein that binds with CCK-AR on intestinal vagal afferents to relay neural signals to the brain (40). Dietary lipids and proteins are major stimulants of CCK secretion in humans and rats (24, 40). The basal concentration of CCK in human plasma is around 1 pmol/l (24, 37). After nutrients consumption, CCK levels in human plasma are increased within 20 min and then gradually reduced to reach a second peak after 1.5-2 h (24, 40). Longer fatty acids with chains greater than or equal to C12 stimulate CCK secretion much more than fatty acids with chains shorter than C12 in human and STC-1 cells (30, 40). Thus, dietary lipids play an important role in the stimulation of CCK secretion in humans and rats (24, 40).

Apolipoprotein A-IV (ApoA-IV) is one of the most abundant proteins (as much as 3% of the proteins) made by the enterocytes and the hypothalamus in the presence of dietary lipids, and it functions to modulate lipid metabolism as well as glucose homeostasis in mice (21, 42). Both ApoA-IV and CCK are secreted in response to dietary lipids in human, rats and mice (9, 14, 21, 25, 26, 37). Thus, the present study investigated whether dietary lipid-induced ApoA-IV elevates CCK levels in the small intestine using ApoA-IV-KO and WT

mice. ApoA-IV-KO mice fed a chow diet have normal body weight, fat mass, and *Cckar* gene expression in the duodenum (21). In response to dietary lipids, CCK is mainly secreted by the duodenum and jejunum in humans (37). The present experiment confirmed that duodenum was a major site for CCK synthesis in ApoA-IV-KO and WT mice. Furthermore, ApoA-IV-KO mice on a chow diet had reduced basal levels of duodenal CCK synthesis, compared with the control group (33). Thus, mice without ApoA-IV have impaired CCK synthesis in the duodenum and normal *Cck* expression in the jejunum and ileum.

A short-term feeding of dietary lipids enhances ApoA-IV levels in the jejunum of human and rats (21, 43). In the current study, dietary lipids elevated *Cck* gene expression in the duodenum of WT mice, consistent with the observations of increased CCK protein secretion stimulated by dietary lipids in rats and human (24, 34, 37). After an acute gavage of dietary lipids, the ApoA-IV-KO mice had lower levels of *Cck* mRNA level in the duodenum compared to that of the WT mice. Thus, the elevation of intestinal CCK synthesis induced by an acute feeding of dietary lipids is associated with the presence of ApoA-IV. Chronic feeding of a HFD elevated basal levels of duodenal CCK synthesis in WT mice, consistent with the observations of increased CCK protein secretion stimulated by a HFD in rats and human (24, 37). A HFD enhances the efficiency of triglyceride digestion and luminal long-chain fatty acids trigger CCK secretion by endocrine cells in the small intestine of rats and human (24, 34, 37). Thus, HFD enhances plasma CCK responses to dietary lipids. During the HFD feeding, the ApoA-IV-KO mice exhibited lower levels of *Cck* expression in the duodenum relative to WT mice *Cck* expression. Thus, these findings suggest that the elevation of duodenal *Cck* gene expression in the HFD-fed WT mice is associated with the presence of ApoA-IV. ApoA-IV and CCK are synthesized by different cells on the villi (enterocytes versus endocrine cells) and the secretion of ApoA-IV and CCK are chylomicron-dependent (17, 34). The working hypothesis for the current *in-vivo* study is that chylomicrons with ApoA-IV secreted by enterocytes in the villi pass through the lamina propria equipped with capillaries and nerve fibers to enter the lacteal (lymphatic vessels) located centrally in the core of the lamina propria (41). In the lamina propria, ApoA-IV on the surface of chylomicrons may stimulate CCK release by endocrine cells via a paracrine mode before entering the lacteal (40). However, further investigations for the site of ApoA-IV and CCK interaction through chylomicron circulation is required.

The ApoA-IV-KO mice fed a chow diet had increased body weight, comparable fat mass, and reduced levels of TG and CHOL in the fasted plasma (20). ApoA-IV-KO mice have comparable fat absorption in the small intestine and normal transport of TG from the small intestine into the plasma (21). ApoA-IV-KO mice produce decreased secretion of ApoA-I and ApoC-III in the lymph after an intraduodenal infusion of dietary lipids (20, 44) because ApoA-IV-KO mice have a global deficiency of a gene cluster containing ApoA-IV, ApoA-I, and ApoC-III (21). The ApoA-IV-KO mice had reduced levels of plasma lipids following an acute oral gavage of lipid mixtures, possibly due to decreased ApoC-III-induced lipolysis (20, 21). Since the role of ApoA-I and ApoC-III in the regulation of CCK levels remains unknown, further study of increased CCK release induced by ApoA-I or ApoC-III is required. Thus, these observations suggest that the downregulation of intestinal CCK synthesis in the ApoA-IV-KO mice induced by an acute feeding of dietary lipids or chronic feeding of HFD may be associated with the deficiency of ApoA-IV, ApoA-I and ApoC-III.

Since STC-1 cells do not produce detectable gastrin (30), CCK-producing STC-1 cells were used to study a direct effect of ApoA-IV on CCK released by STC-1 cells in the present experiment. Dietary proteins are the major stimulus of CCK release in rats (9, 26). The present study showed that mouse recombinant ApoA-IV proteins at or above 100 µg/ml directly induced CCK release in a dose-dependent manner after 1-h and 2-h incubations. CCK-8 release by STC-1 cells into the culture medium is induced by various secretagogues (3). Thus, CCK-8 may be the major form of the ApoA-IV-induced CCK in the medium of STC-1 cells. In contrast, CCK-58 should be most secreted CCK by ApoA-IV in WT and ApoA-IV-KO mice because of CCK-58 is the major form in rodent plasma (36, 37). Since the ApoA-IV receptor has not yet been identified, the mechanism by which ApoA-IV directly stimulates CCK release by STC-1 cells remains unknown. Albumin is the major protein in the plasma and acts as a carrier protein for fatty acids (23). BSA in the current experiment was used to mimic the action of albumin in the capillaries in the stimulation of CCK release in the lamina propria (41). BSA at 1% has been reported to induce a moderate increase in CCK release by STC-1 cells (7). Thus, BSA at the same dose of ApoA-IV was used to determine if ApoA-IV's effect on CCK secretion by STC-1 cells is specific. The present study demonstrated that BSA at the same dose of ApoA-IV failed to enhance CCK release by STC-1 cells, suggesting that ApoA-IV is a more potent stimulant than BSA in elevating CCK levels in the medium of STC-1 cells. LAPR5 is highly expressed in the mucosa of small intestine and is also present in STC-1 cells (4, 5, 22). Protein hydrolysates (peptones) have been reported to produce a marked elevation in CCK release by STC-1 cells through the activation of LAPR5-dependent pathway (7, 10). To examine whether ApoA-IV-induced CCK release by STC-1 cells through the activation of LPAR5-dependent pathway, CCK levels in the media of STC-1 cells with knockdown of *Lpar5* and treatment of ApoA-IV were determined. In the current study, the blockade of a LPAR5-dependent pathway with siRNA significantly attenuated ApoA-IV's stimulatory effect on *Cck* gene expression and CCK secretion in STC-1 cells. Thus, the culture studies are fundamental for the connection between ApoA-IV and CCK via a LPAR5-dependent pathway.

Conclusion

In light of the observations in the current study, ApoA-IV-induced CCK response to dietary lipids in mice and in STC-1 cells may provide important information about shared roles in the modulation of gastric emptying and satiation through the activation of vagal activity. Furthermore, ApoA-IV-induced CCK may be involved in lipid absorption in the lumen of the small intestine, because CCK release by enteroendocrine cells is dependent on the presence of ApoA-IV in chylomicrons during a lipid infusion. Consequently, CCK stimulates gall bladder contraction and pancreatic acini to evoke the secretions of bile acids and pancreatic lipase, which both assist in the digestion and absorption of cholesterol and fatty acid in the lumen of small intestine. Thus, ApoA-IV-induced CCK secretion plays an important role in the regulation of lipid metabolism and energy homeostasis.

Acknowledgments

We would like to thank the Mouse Metabolic Phenotyping Center at the University of Cincinnati for their excellent assistance with the generation of ApoA-IV-KO mice. This work was supported by National Institutes of Health Grants DK83550, DK97436, DK 92779 and 95440.

References

1. Apfelbaum TF, Davidson NO, Glickman RM. Apolipoprotein A-IV synthesis in rat intestine: regulation by dietary triglyceride. *The American journal of physiology*. 1987; 252:G662–666. [PubMed: 3555115]
2. Callow AD. Cardiovascular disease 2005--the global picture. *Vascular pharmacology*. 2006; 45:302–307. [PubMed: 17074537]
3. Chang CH, Chey WY, Sun Q, Leiter A, Chang TM. Characterization of the release of cholecystokinin from a murine neuroendocrine tumor cell line, STC-1. *Biochimica et biophysica acta*. 1994; 1221:339–347. [PubMed: 8167157]
4. Choi S, Lee M, Shiu AL, Yo SJ, Aponte GW. Identification of a protein hydrolysate responsive G protein-coupled receptor in enterocytes. *American journal of physiology Gastrointestinal and liver physiology*. 2007; 292:G98–G112. [PubMed: 16935853]
5. Choi S, Lee M, Shiu AL, Yo SJ, Hallden G, Aponte GW. GPR93 activation by protein hydrolysate induces CCK transcription and secretion in STC-1 cells. *American journal of physiology Gastrointestinal and liver physiology*. 2007; 292:G1366–1375. [PubMed: 17290006]
6. Cohen RD, Castellani LW, Qiao JH, Van Lenten BJ, Lusis AJ, Reue K. Reduced aortic lesions and elevated high density lipoprotein levels in transgenic mice overexpressing mouse apolipoprotein A-IV. *The Journal of clinical investigation*. 1997; 99:1906–1916. [PubMed: 9109435]
7. Cordier-Bussat M, Bernard C, Haouche S, Roche C, Abello J, Chayvialle JA, Cuber JC. Peptones stimulate cholecystokinin secretion and gene transcription in the intestinal cell line STC-1. *Endocrinology*. 1997; 138:1137–1144. [PubMed: 9048620]
8. DeFronzo RA. Current issues in the treatment of type 2 diabetes. Overview of newer agents: where treatment is going. *The American journal of medicine*. 2010; 123:S38–48. [PubMed: 20206731]
9. Douglas BR, Woutersen RA, Jansen JB, de Jong AJ, Lamers CB. The influence of different nutrients on plasma cholecystokinin levels in the rat. *Experientia*. 1988; 44:21–23. [PubMed: 3350114]
10. Foltz M, Ansems P, Schwarz J, Tasker MC, Loubakos A, Gerhardt CC. Protein hydrolysates induce CCK release from enteroendocrine cells and act as partial agonists of the CCK1 receptor. *Journal of agricultural and food chemistry*. 2008; 56:837–843. [PubMed: 18211011]
11. Fujimoto K, Cardelli JA, Tso P. Increased apolipoprotein A-IV in rat mesenteric lymph after lipid meal acts as a physiological signal for satiation. *The American journal of physiology*. 1992; 262:G1002–1006. [PubMed: 1616031]
12. Gibbs J, Young RC, Smith GP. Cholecystokinin decreases food intake in rats. *Journal of comparative and physiological psychology*. 1973; 84:488–495. [PubMed: 4745816]
13. Glatzle J, Darcel N, Rechs AJ, Kalogeris TJ, Tso P, Raybould HE. Apolipoprotein A-IV stimulates duodenal vagal afferent activity to inhibit gastric motility via a CCK1 pathway. *American journal of physiology Regulatory, integrative and comparative physiology*. 2004; 287:R354–359.
14. Green PH, Glickman RM, Riley JW, Quinet E. Human apolipoprotein A-IV. Intestinal origin and distribution in plasma. *The Journal of clinical investigation*. 1980; 65:911–919. [PubMed: 6987270]
15. Guo F, Moellering DR, Garvey WT. The progression of cardiometabolic disease: validation of a new cardiometabolic disease staging system applicable to obesity. *Obesity*. 2014; 22:110–118. [PubMed: 23894121]
16. Hayashi H, Fujimoto K, Cardelli JA, Nutting DF, Bergstedt S, Tso P. Fat feeding increases size, but not number, of chylomicrons produced by small intestine. *The American journal of physiology*. 1990; 259:G709–719. [PubMed: 2240215]
17. Hayashi H, Nutting DF, Fujimoto K, Cardelli JA, Black D, Tso P. Transport of lipid and apolipoproteins A-I and A-IV in intestinal lymph of the rat. *Journal of lipid research*. 1990; 31:1613–1625. [PubMed: 2123231]
18. Holzer HH, Turkelson CM, Solomon TE, Raybould HE. Intestinal lipid inhibits gastric emptying via CCK and a vagal capsaicin-sensitive afferent pathway in rats. *The American journal of physiology*. 1994; 267:G625–629. [PubMed: 7943327]

19. Kahn BB, Flier JS. Obesity and insulin resistance. *The Journal of clinical investigation*. 2000; 106:473–481. [PubMed: 10953022]
20. Kohan AB, Wang F, Li X, Bradshaw S, Yang Q, Caldwell JL, Bullock TM, Tso P. Apolipoprotein A-IV regulates chylomicron metabolism-mechanism and function. *American journal of physiology Gastrointestinal and liver physiology*. 2012; 302:G628–636. [PubMed: 22207575]
21. Kohan AB, Wang F, Lo CM, Liu M, Tso P. ApoA-IV: current and emerging roles in intestinal lipid metabolism, glucose homeostasis, and satiety. *American journal of physiology Gastrointestinal and liver physiology*. 2015; 308:G472–481. [PubMed: 25591862]
22. Kotarsky K, Boketoft A, Bristulf J, Nilsson NE, Norberg A, Hansson S, Owman C, Sillard R, Leeb-Lundberg LM, Olde B. Lysophosphatidic acid binds to and activates GPR92, a G protein-coupled receptor highly expressed in gastrointestinal lymphocytes. *The Journal of pharmacology and experimental therapeutics*. 2006; 318:619–628. [PubMed: 16651401]
23. Levitt DG, Levitt MD. Human serum albumin homeostasis: a new look at the roles of synthesis, catabolism, renal and gastrointestinal excretion, and the clinical value of serum albumin measurements. *Int J Gen Med*. 2016; 9:229–255. [PubMed: 27486341]
24. Liddle RA. Cholecystokinin cells. *Annual review of physiology*. 1997; 59:221–242.
25. Liddle RA, Goldfine ID, Rosen MS, Taplitz RA, Williams JA. Cholecystokinin bioactivity in human plasma. Molecular forms, responses to feeding, and relationship to gallbladder contraction. *The Journal of clinical investigation*. 1985; 75:1144–1152. [PubMed: 2580857]
26. Liddle RA, Green GM, Conrad CK, Williams JA. Proteins but not amino acids, carbohydrates, or fats stimulate cholecystokinin secretion in the rat. *The American journal of physiology*. 1986; 251:G243–248. [PubMed: 3740265]
27. Liu M, Shen L, Liu Y, Woods SC, Seeley RJ, D'Alessio D, Tso P. Obesity induced by a high-fat diet downregulates apolipoprotein A-IV gene expression in rat hypothalamus. *American journal of physiology Endocrinology and metabolism*. 2004; 287:E366–370. [PubMed: 15082422]
28. Lo CC, Langhans W, Georgievsky M, Arnold M, Caldwell JL, Cheng S, Liu M, Woods SC, Tso P. Apolipoprotein AIV requires cholecystokinin and vagal nerves to suppress food intake. *Endocrinology*. 2012; 153:5857–5865. [PubMed: 23027805]
29. Lo CM, Zhang DM, Pearson K, Ma L, Sun W, Sakai RR, Davidson WS, Liu M, Raybould HE, Woods SC, Tso P. Interaction of apolipoprotein AIV with cholecystokinin on the control of food intake. *American journal of physiology Regulatory, integrative and comparative physiology*. 2007; 293:R1490–1494.
30. McLaughlin JT, Lomax RB, Hall L, Dockray GJ, Thompson DG, Warhurst G. Fatty acids stimulate cholecystokinin secretion via an acyl chain length-specific, Ca²⁺-dependent mechanism in the enteroendocrine cell line STC-1. *The Journal of physiology*. 1998; 513(Pt 1):11–18. [PubMed: 9782155]
31. Moran TH, Baldessarini AR, Salorio CF, Lowery T, Schwartz GJ. Vagal afferent and efferent contributions to the inhibition of food intake by cholecystokinin. *The American journal of physiology*. 1997; 272:R1245–1251. [PubMed: 9140026]
32. Moran TH, Norgren R, Crosby RJ, McHugh PR. Central and peripheral vagal transport of cholecystokinin binding sites occurs in afferent fibers. *Brain research*. 1990; 526:95–102. [PubMed: 2078822]
33. Pressler JW, Haller A, Sorrell J, Wang F, Seeley RJ, Tso P, Sandoval DA. Vertical sleeve gastrectomy restores glucose homeostasis in apolipoprotein A-IV KO mice. *Diabetes*. 2015; 64:498–507. [PubMed: 25157093]
34. Raybould HE, Meyer JH, Tabrizi Y, Liddle RA, Tso P. Inhibition of gastric emptying in response to intestinal lipid is dependent on chylomicron formation. *The American journal of physiology*. 1998; 274:R1834–1838. [PubMed: 9841489]
35. Raybould HE, Zittel TT, Holzer HH, Lloyd KC, Meyer JH. Gastrointestinal sensory mechanisms and CCK in inhibition of gastric emptying in response to a meal. *Digestive diseases and sciences*. 1994; 39:41S–43S. [PubMed: 7995213]
36. Reeve JR Jr, Green GM, Chew P, Eysselein VE, Keire DA. CCK-58 is the only detectable endocrine form of cholecystokinin in rat. *American journal of physiology Gastrointestinal and liver physiology*. 2003; 285:G255–265. [PubMed: 12686511]

37. Rehfeld JF. Cholecystokinin-From Local Gut Hormone to Ubiquitous Messenger. *Frontiers in endocrinology*. 2017; 8:47. [PubMed: 28450850]
38. Rodriguez MD, Kalogeris TJ, Wang XL, Wolf R, Tso P. Rapid synthesis and secretion of intestinal apolipoprotein A-IV after gastric fat loading in rats. *The American journal of physiology*. 1997; 272:R1170–1177. [PubMed: 9140017]
39. Shen L, Pearson KJ, Xiong Y, Lo CM, Tso P, Woods SC, Davidson WS, Liu M. Characterization of apolipoprotein A-IV in brain areas involved in energy homeostasis. *Physiology & behavior*. 2008; 95:161–167. [PubMed: 18577393]
40. Steinert RE, Feinle-Bisset C, Asarian L, Horowitz M, Beglinger C, Geary N. Ghrelin, CCK, GLP-1, and PYY(3-36): Secretory Controls and Physiological Roles in Eating and Glycemia in Health, Obesity, and After RYGB. *Physiological reviews*. 2017; 97:411–463. [PubMed: 28003328]
41. Tso P, Balint JA. Formation and transport of chylomicrons by enterocytes to the lymphatics. *The American journal of physiology*. 1986; 250:G715–726. [PubMed: 3521320]
42. Wang F, Kohan AB, Lo CM, Liu M, Howles P, Tso P. Apolipoprotein A-IV: a protein intimately involved in metabolism. *Journal of lipid research*. 2015; 56:1403–1418. [PubMed: 25640749]
43. Weinberg RB, Dantzker C, Patton CS. Sensitivity of serum apolipoprotein A-IV levels to changes in dietary fat content. *Gastroenterology*. 1990; 98:17–24. [PubMed: 2104541]
44. Weinstock PH, Bisgaier CL, Hayek T, Aalto-Setälä K, Sehayek E, Wu L, Sheffele P, Merkel M, Essenburg AD, Breslow JL. Decreased HDL cholesterol levels but normal lipid absorption, growth, and feeding behavior in apolipoprotein A-IV knockout mice. *Journal of lipid research*. 1997; 38:1782–1794. [PubMed: 9323588]
45. Whited KL, Lu D, Tso P, Kent Lloyd KC, Raybould HE. Apolipoprotein A-IV is involved in detection of lipid in the rat intestine. *The Journal of physiology*. 2005; 569:949–958. [PubMed: 16239275]
46. WHO. Obesity and Overweight World Health Organization. Fact sheet No 311. 2015
47. Woods SC. The control of food intake: behavioral versus molecular perspectives. *Cell metabolism*. 2009; 9:489–498. [PubMed: 19490904]
48. Woods SC, Seeley RJ, Rushing PA, D'Alessio D, Tso P. A controlled high-fat diet induces an obese syndrome in rats. *The Journal of nutrition*. 2003; 133:1081–1087. [PubMed: 12672923]
49. Yoder SM, Yang Q, Kindel TL, Tso P. Stimulation of incretin secretion by dietary lipid: is it dose dependent? *American journal of physiology Gastrointestinal and liver physiology*. 2009; 297:G299–305. [PubMed: 19520739]
50. Yoshimichi G, Lo CC, Tamashiro KL, Ma L, Lee DM, Begg DP, Liu M, Sakai RR, Woods SC, Yoshimatsu H, Tso P. Effect of peripheral administration of cholecystokinin on food intake in apolipoprotein AIV knockout mice. *American journal of physiology Gastrointestinal and liver physiology*. 2012; 302:G1336–1342. [PubMed: 22461023]

Highlights

- Dietary lipid-induced ApoA-IV is associated with the increased levels of *Cck* gene expression in the duodenum.
- Recombinant ApoA-IV protein directly increases CCK protein released from enteroendocrine STC-1 cells.
- Recombinant ApoA-IV protein acts on the LPAR5-dependent pathway to stimulate CCK release from STC-1 cells.

Figure 1.

Comparison of intestinal *Cck* gene expression in the duodenum, jejunum and ileum in ApoA-IV-KO and WT mice on a chow diet. Following a 5-h fast, small intestines of ApoA-IV-KO and WT were collected and *Cck* gene expression in the small intestine were determined using qPCR. Data are expressed as mean $\pm$ SEM (n= 6-7 animals per group), and *Cck* mRNA was normalized to cyclophilin housekeeping gene expression. Values with asterisks represent significant differences between different groups ($P < 0.05$).

Figure 2.

The effect of lipid on duodenal *Cck* gene expression in ApoA-IV-KO and WT mice. Animals on a chow diet received a bolus of lipid mixture (200 μ l) by oral gavage following a 5-h fast. The duodenum was collected prior to lipid treatment (0 h) and 2 h after infusion of lipid mixtures (2 h). Data are expressed as mean $\pm$ SEM (n= 5-6 animals per group), and *Cck* mRNA was normalized to cyclophilin housekeeping gene expression. Values with asterisks represent significant differences between different groups ($P < 0.05$).

Figure 3.

The effect of chronic HFD-feeding on duodenal *Cck* gene expression in ApoA-IV-KO and WT mice. Animals at 10 weeks of age received either a LFD or a HFD for 8 weeks. Following a 5-h fast, duodenum was collected and *Cck* gene expressions were determined. *Cck* mRNA was normalized to cyclophilin housekeeping gene expression. Data are expressed as mean $\pm$ SEM, $n = 6-7$ animals per group and values with asterisks represent significant differences between different groups ($P < 0.05$).

Figure 4.

ApoA-IV increased CCK levels in the media of STC-1 cells. ApoA-IV-induced CCK level after incubation of a 0.5-h (A), 1-h (B), or 2-h (C). STC-1 cells were incubated with vehicle or mouse recombinant ApoA-IV proteins, and the media were collected at different time points for the determination of CCK level (pmol/mL equivalent to CCK-33). Data are expressed as mean $\pm$ SEM for the medium of STC-1 cells from 5-6 wells per group. Values with asterisks represent significant differences relative to the vehicle treatment ($P < 0.05$).

Figure 5.

Comparison of CCK levels in the media of STC-1 cells after treated with Apo-AIV and BSA. STC-1 cells were incubated with vehicle, ApoA-IV proteins or BSA at 200 $\mu\text{g/ml}$, and the media were collected after a 2-h incubation for the determination of CCK level (pmol/mL equivalent to CCK-33). Data are expressed as mean $\pm$ SEM for the medium of STC-1 cells from 5-6 wells per group. Values with asterisks represent significant differences relative to the vehicle treatment ($P < 0.05$).

Figure 6.

ApoA-IV increased CCK level through a LPAR5-dependent pathway. *Lpar5* gene expression in STC-1 cells (A), *Cck* gene expression in STC-1 cells (B) and CCK level (pmol/mL equivalent to CCK-33) in the medium of STC-1 cells (C). STC-1 cells transfected with or without *Lpar5* siRNA were incubated with ApoA-IV or vehicle. After a 2-h incubation, the media and cells of the STC-1 cells were collected for CCK level and gene expression. Data are expressed as mean $\pm$ SEM for the cells of STC-1 cells from 7-8 wells per group and *Cck* mRNA was normalized to cyclophilin housekeeping gene expression. Data are expressed as mean $\pm$ SEM for the medium of STC-1 cells from 6-7 wells per group. Values with asterisks represent significant differences between different treatments ($P < 0.05$).

Table 1
Body weight, adipose tissue weights, and plasma parameters in animals

	WT	ApoA-IV-KO
BW (g)	31.08 ± 0.9	29.92 ± 1.7
BAT (g)	0.12 ± 0.0	0.12 ± 0.0
Epididymal Fat (g)	0.37 ± 0.1	0.63 ± 0.2
Inguinal Fat (g)	0.12 ± 0.0	0.17 ± 0.1
Triglyceride (mg/dl) 0-h	56.40 ± 4.1	33.23 ± 3.3
Triglyceride (mg/dl) 2-h	398.22 ± 61.5[#]	163.12 ± 42.6^{*,#}
Cholesterol (mg/dl) 0-h	68.40 ± 9.1	62.85 ± 8.7
Cholesterol (mg/dl) 2-h	155.18 ± 30.1[#]	139.6 ± 32.1

Plasma and tissues from ApoA-IV -KO and WT mice (n =6 per group) on a chow diet were collected either 0 h or 2 h after infusion of lipid mixtures. Values represent mean ± SEM, and

* asterisk indicates a significant difference ($P < 0.05$) compared to WT controls at the same treatment and the same time point.

[#] Number sign indicates a significant difference ($P < 0.05$) compared to WT controls 0 h after infusion of dietary lipids.