Antimicrobial resistance and putative virulence genes of *Pseudomonas aeruginosa* isolates from patients with respiratory tract infection

Heba Y. Al Dawodeyah¹, Nathir Obeidat², Luay F. Abu-Qatouseh³, Asem A. Shehabi^{4,*}

Abstract

Introduction *Pseudomonas aeruginosa* is a common agent causing community acquired and nosocomial respiratory tract infections, with particularly life-threatening manifestations in patients who are immunocompromised or who have cystic fibrosis. This study investigated the occurrence of extended-spectrum β -lactamases (ESBLs) and metallo β -lactamase (MBL) in association with important putative virulence genes and genotypes variation among *P. aeruginosa* isolates from respiratory tract infection of Jordanian patients.

Methods Over a period of 8-month, a total of 284 respiratory tract samples were obtained from patients diagnosed with respiratory tract infection while attending the Pulmonary Clinic/Intensive Care Unit, Jordan University Hospital (JUH). At the time of sampling most were inpatients (86.9%). Samples were cultured specifically for *P. aeruginosa*.

Results A total of 61/284 (21.5%) *P. aeruginosa* isolates were recovered from respiratory samples of patients. The percentage of MDR *P. aeruginosa* isolates was 52.5%, and all isolates were susceptible to colistin with lower rates of susceptibility to other tested antibiotics. Positive genes of *bla*_{CTX-M}, *bla*_{VEB}, *bla*_{TEM}, *bla*_{GES} and *bla*_{SHV} were detected in 68.9%, 18.9%, 18.9%, 15.6% and 12.5% of isolates, respectively. Genotyping revealed no significant genetic relationship among MDR *P. aeruginosa* isolates from hospitalized patients as judged by the constructed dendrogram and the presence of 14 genotypic groups. The percentages of the virulence genes *algD*, *lasB*, *toxA*, *exoS*, and *exoU* among *P. aeruginosa* isolates were 98%, 98%, 80%, 33% and 33%, respectively, and 87% of isolates produced pyocyanin.

Conclusion The present study demonstrates high occurrence of MDR *P. aeruginosa* isolates carrying *bla*_{CTX-M} genes. No specific associations were found between antibiotic resistance, virulence genes and genotypes among MDR isolates.

Keywords *P. aeruginosa*, antimicrobial resistance, virulence factors, respiratory infections, Jordan patients.

Introduction

Pseudomonas aeruginosa respiratory infections are causing high morbidity and mortality due to the germ's capacity to rapidly develop antibiotic resistance, especially during antibiotic treatment of patients. The organism can carry a variety of putative virulence factors, which are highly controlled by cell-to-cell signaling systems.^{1,2}

P. aeruginosa can secrete an exopolysaccharide called alginate (*algD*) and an elastase B (*lasB*) enzyme in response to environmental conditions. Both these virulence factors can influence pathogenesis by enhancing adhesion, colonization, and invasion of tissues, causing chronic pulmonary inflammation.³ Additionally, elastase B is an important protease of *P.*

Received: 05 December 2017; revised: 16 February 2018; accepted: 19 February 2018.

¹MSc, Department of Pathology-Microbiology, School of Medicine, The Jordan University, Queen Rania str. 100, Amman, Jordan; ²MD, Department of Internal Medicine/Division of Pulmonary Diseases, The Jordan University Hospital, Queen Rania str. 100, Amman, Jordan; ³PhD, Faculty of Pharmacy, Department of Biomedical Sciences, University of Petra, Airport str. 317, Amman, Jordan; ⁴DSc, Department of Pathology-Microbiology,

School of Medicine, The Jordan University, Queen Rania str. 100, Amman, Jordan.

*Corresponding author: Asem A. Shehabi, DSc, Department of Pathology-Microbiology, School of Medicine, The Jordan University, Queen Rania str. 100, Amman, Jordan. ashehabi@ju.edu.jo

Article downloaded from www.germs.ro

Published March 2018

© GERMS 2018

ISSN 2248 - 2997

ISSN - L = 2248 - 2997

aeruginosa. This enzyme has a tissue-damaging activity, it can degrade a number of plasma proteins and it contributes to the survival of *P. aeruginosa* in infected tissues.⁴

Other potential virulence factors secreted by *P. aeruginosa* that are important in its pathogenicity include exotoxin A (toxA), which is the most toxic virulence factor detected in this organism. It inhibits protein biosynthesis, it has a necrotizing activity on tissues and contributes to the colonization process.⁵ Exoenzyme S (exoS) is secreted into the cytosol of epithelial cells and it contributes to increased pneumonia severity.⁶

Infections caused by multidrug-resistant (MDR) *P. aeruginosa* are difficult to treat and can be associated with high mortality especially in patients who are immunocompromised or who present chronic lung diseases. *P. aeruginosa* acquires resistance primarily through the production of plasmid mediated AmpC β -lactamase, different types of extended-spectrum β -lactamases (ESBLs) and metallo- β -lactamases (MBLs).^{7,8} The recent emergence of bla-KPC enzymes in *P. aeruginosa* strains, currently KPC-2 and KPC-5, is alarming since these strains show activity against most β -lactams including monobactams, carbapenems, and oxymino-cephalosporins.⁸

Recent studies report that ESBLs and MBLs are increasing among *P. aeruginosa* in the Middle East Arab countries, and this development could result in prolonged hospital stay, increased cost and fatal outcome among patients.^{9,10}

This study investigated the occurrence and relationship of MDR *P. aeruginosa* recovered from Jordanian patients with respiratory tract infection, in association with putative virulence genes and genotypes.

Methods

Patients

This prospective convenience sampling study included a total of 284 respiratory tract samples obtained from adult patients attending the Pulmonary Clinic, The Jordan University Hospital (JUH), over the period from November 2014 to June 2015. JUH includes 550 beds, and is a tertiary care teaching hospital with 24 of the

beds distributed in 3 adult Intensive Care Units (ICUs).

Only one sample was collected from each patient with signs and symptoms of respiratory tract infection. The majority of specimens (247; 86.9%) were obtained from hospitalized patients and only 37 (13.1%) were collected from outpatients. Bio-data of each patient was obtained and registered on a special sheet and included the following: patient's identification number, gender, age, clinical diagnosis, specific admission conditions, type of specimen, type of current or previous antibiotic treatment within the past 4 weeks prior to specimen sampling.

The study was approved by the School of Medicine and the School of Graduate studies, The Jordan. A permission was also obtained from the Institutional Ethical Review Board (IERB) at the Jordan University Hospital (JUH, Permission No. 9/2015). Verbal consent was obtained from all patients after explaining the aim of the study. The identity of patients was protected by applying numbers instead of names for all collected specimens.

Culture, isolation and bacterial identification

All collected respiratory samples (sputum, bronchoalveolar lavage [BAL], pleural fluid) were inoculated directly using cotton swab onto *Pseudomonas* CN agar plates (Oxoid, Basingstoke, United Kingdom) and incubated for 24-48 hours at 37 °C. The culture plates were examined to detect the presence of straw-colored colonies with yellow or blue-green pigmentation, and colonies were confirmed later as being *P. aeruginosa* using the following preliminary characteristics: positive oxidase test, growth at 42 °C, negative lactose, and glucose fermentation in tube of Kligler iron agar. Five colonies of pure *P. aeruginosa* were subcultured into cysteine lactose electrolyte deficient agar plates (CLED, Oxoid), and all isolates were stored in cryotubes containing brain-heart infusion agar with 15% glycerol at -70 °C for further investigation.

Table 1. List of all primers, their product sizes and references used for detection of ESBLs, MBLs and virulence genes

Target genes	Primer	Primer sequence (5' to 3')	Product size (bp)	Reference
<i>bla</i> _{CTX-M1}	CTX-M(F)	GACGATGTCACCTGGCTGAGC	499	13
	CTX-M(R)	AGCCGCCGACGCTAATACA		
<i>bla</i> _{TEM}	TEM-A	ATAAAATTCTTGAAGAC	1,075	14
	TEM-B	TTACCAATGCTTAATCA		
<i>bla</i> _{VEB-1}	VEB-FOR	CGACTTCCATTTCCCGATGC	642	15
	VEB-REV	GGACTCTGCAACAAATACGC		
<i>bla</i> _{SHV-1}	SHV-FOR	TGGTTATGCGTTATATTGCGC	867	21
	SHV-REV	GCTTAGCGTTGCCAGTGCT		
<i>bla</i> _{GES} type	GES-1(F)	CTATTACTGGCAGGGATCG	594	22
	GES-1(R)	CCTCTCAATGGTGTGGGT		
<i>bla</i> _{IMP}	IMP-A	GAAGGCGTTTATGTTTCATAC	587	14
	IMB-B	GTACGTTTCAAGAGTGATGC		
<i>bla</i> _{VIM}	VIM2004A	GTTTGGTCGCATATCGCAAC	382	14
	VIM2004B	AATGCGCAGCACCAGGATAG		
<i>bla</i> _{KPC}	KPC(F)	ATGTCACTGTATCGCCGTCT	880	18
	KPC(R)	TTACTGCCCCGTTGACGCCCA		
<i>algD</i>	algDF	CGTCTGCCGCGAGATCGGCT	313	19
	algDR	GACCTCGACGGTCTTGCGGA		
<i>lasB</i>	lasBF	GGAATGAACGAAGCGTTCTCCGAC	284	19
	lasBR	TTGGCGTCGACGAACACCTCG		
<i>toxA</i>	toxAF	CTGCGCGGGTCTATGTGCC	270	23
	toxAR	GATGCTGGACGGGTCGAG-3		
<i>exoS</i>	exoS1-F	TCTGAATTCTTCCAGGCGGGTGAACATCA	504	6
	exoS1-R	TTTAGATCTCACCCTGGTATCCAAGGCGA		
<i>exoU</i>	exoU-F	GGG AAT ACT TTC CGG GAA GTT	428	20
	exoU-R	CGA TCT CGC TGC TAA TGT GTT		
<i>P. aeruginosa</i> 16S rDNA	PA-SS-F	(GGGGGATCTTTCGGACCTCA	956	12
	PA-SS-R	(TCCTTAGAGTGCCCAACCCG		

Antimicrobial susceptibility

Antimicrobial susceptibility of *P. aeruginosa* isolates was assessed using disc diffusion test according to the recommendation of the Clinical Laboratory and Standards Institute (CLSI, 2015).¹¹ *P. aeruginosa* isolates which showed resistance to three or more antibiotic classes (MDR) were tested for minimum inhibitory concentrations (MICs) by E-test strips for ceftazidime, ciprofloxacin, colistin, imipenem and meropenem (Oxoid). The results of susceptibility tests were interpreted according to the breakpoints of tested antibiotics as recommended in CLSI guidelines (2015). *P. aeruginosa* ATCC 27853 was used as control during all tests.

Plasmid DNA extraction and PCR procedures

After thawing, stored *P. aeruginosa* isolates were subcultured onto blood agar, and incubated at 37 °C for 24 hours. A few colonies were picked from the agar, and inoculated into 3 mL Luria Broth (LB) (Oxoid) and incubated for 24 hours at 37 °C. The bacterial plasmid was extracted using the Zyppy™ Plasmid Miniprep Kit, Zymo (QIAGEN, Germantown, MD, USA) according to manufacturer's instructions. The quantity of DNA from each isolate extract was tested to ensure that the DNA concentration was more than 50 µg/mL⁻¹ with a purity at A₂₆₀/A₂₈₀=1.6-1.8, using a spectrophotometer (Bio-Rad, Hercules, CA, USA).

Identification of *P. aeruginosa* by PCR

Specific identification of *P. aeruginosa* was performed by nested PCR using a first primer set for identification of the genus *Pseudomonas*, followed by a second primer set specific for *P. aeruginosa*. The 16 ribosomal DNA (rDNA) was detected as described by Spilker et al. as shown in Table 1.¹²

Detection of virulence genes by PCR

The following genes: *algD*, *lasB*, *toxA*, *exoS* and *exoU* were detected using specific primers for each gene as described in references listed in Table 1.^{6,13-20} *P. aeruginosa* (PAO-1 strain) was used as a positive control for *lasB* and *toxA* genes and *P. aeruginosa* ATCC 27853 was used as a

positive control for *algD* gene, and nuclease free water was used as a negative control.

Detection of genes of ESBLs, MBLs and *bla*_{KPC}

A total of 32 MDR *P. aeruginosa* isolated were investigated for genes of *bla*_{CTX-M}, *bla*_{TEM}, *bla*_{VEB-1}, *bla*_{SHV}, *bla*_{GES-1} and MBLs (*IMP-1*, *VIM-2*, *bla*_{KPC}), as presented in Table 1. A positive control of *E. coli* (ATCC 51446) was used for detection of *bla*_{CTX-M} genes, *P. aeruginosa* (CL 120 & CL 162, Université Libanaise, Tripoli, Lebanon) for *IMP-1*, *VIM-2* genes, and *K. pneumoniae* (ATCC BAA-1705) for *bla*_{KPC} genes. Nuclease free water was used as a negative control in all tests.

Table 2. Demographic characteristics of the 61 investigated patients and their clinical features in association with positive *P. aeruginosa*

Patient characteristics	Hospitalized patients* (n=45)	Non-hospitalized patients (n=16)	Total	P-value	OR (95%CI)
	No. (%)	No. (%)	No. (%)		
Male	29 (64.4)	6 (37.5)	35 (57.4)	0.061	3.021(0.926-9.845)
mean age±SD, years	36.48±14.56	47.5±18.09	38.37±15.51		
Female	16 (35.5)	10 (62.5)	26 (42.6)		
mean age±SD, years	33.63±13.13	32.1±10.19	33.04±11.89	0.984	1.012 (0.312-3.286)
Fever (≥ 38 °C)	17 (37.8)	6 (37.5)	23 (37.7)		
Cough	43 (95.6)	16 (100)	59 (96.7)	0.541	Undefined
Purulent sputum	45 (100)	15 (93.8)	60 (98.4)	0.262	Undefined
Dyspnea	22 (48.9)	4 (25.0)	26 (42.6)	0.085	2.87 (0.803-10.25)
Chest pain	25 (55.6)	10 (22.2)	35 (57.4)	0.428	0.75 (0.233-2.418)
Hemoptysis	16 (35.6)	2 (12.5)	18 (29.5)	0.074	1.16 (5.89-29.91)
COPD	3 (6.7)	5 (31.3)	8 (13.1)	0.421	0.778 (3.862-19.17)
Cystic fibrosis	4 (8.9)	6 (37.5)	10 (16.4)	0.477	0.038 (0.163-0.687)
Received antibiotics during the past 4 weeks prior to admission	45 (100)	14 (87.5)	59 (96.7)	0.066	Undefined

COPD – chronic obstructive pulmonary disease; OR – odds ratio; SD – standard deviation; 95%CI – 95% confidence interval.

*Fisher's exact test and chi-squared test were used to compare significant characteristics between hospitalized and non-hospitalized patients.

Table 3. Demographic characteristics of the 45 hospitalized patients in clinical wards and intensive care units and their clinical features in association with positive *P. aeruginosa*

Patient characteristics	Patients hospitalized in clinical wards (n=27)	Patients hospitalized in intensive care units (n=18)	Total (n=45)	P-value	OR (95%CI)
	No. (%)	No. (%)	No. (%)		
Fever ($\geq 38^{\circ}\text{C}$)	8 (47.1%)	9 (50.0)	17 (37.8)	0.167	2.375 (0.688-8.202)
Cough	25 (92.6)	18 (100)	43 (95.6)	0.509	Undefined
Purulent sputum	27 (100)	18 (100)	45 (100)	-	Undefined
Dyspnea	12 (44.4)	10 (55.6)	22 (48.9)	0.465	1.562 (0.471-5.189)
Chest pain	17 (63.0)	8 (44.4)	25 (55.6)	0.221	0.471 (0.140-1.585)
Hemoptysis	11 (40.7)	5 (27.8)	16 (35.6)	0.373	0.559 (0.155-2.024)
COPD	0	3 (16.7)	3 (13.1)	0.028	Undefined
Cystic fibrosis	8 (29.6)	2 (11.1)	10 (16.4)	0.143	3.368 (0.624-18.186)
Received antibiotics during the past 4 weeks prior to admission	27 (100)	18 (100)	45 (100)	-	Undefined

COPD – chronic obstructive pulmonary disease; OR – odds ratio; SD – standard deviation; 95%CI – 95% confidence interval.

Gel electrophoresis

In all tests, 6 μL of each amplified PCR product, controls and 100 bp ladder (molecular size marker) were analyzed in 1.5% agarose gel (Promega, Madison, WI, USA) in 1X TBE buffer containing ethidium bromide, and run for 50 minutes at 120 volts using horizontal apparatus. After electrophoresis, the DNA bands in gel were visualized under ultraviolet light (UVP system, Upland, CA, USA).

RAPD-PCR to detect genotypic variations among *P. aeruginosa* isolates

Two short (10 mers) single primers of 272 (5-AGCGGGCCAA-3) and 208 (5-ACGGCCGACC-3) were used in two separated PCR reactions to create a repetitive electrophoresis patterns and to detect the genotypic variation according to the number and weight of detected band differences among the isolates.²¹ The results were analyzed by computer with PyElph software (USA).

Statistical analysis

Data generated from the study were tabulated as Microsoft Excel sheet and uploaded to Statistical Package for Social Sciences version 20 (IBM Corp, Armonk, NY, USA). Frequencies and percentages were calculated for categorical data and Pearson's chi-squared test or Fisher's exact test were applied to determine potential factors associated with *P. aeruginosa* and to determine whether there are any statistical differences between groups. The level of significance was set at a p value of 0.05 to test the hypothesis of no association. Fisher's exact test replaces chi-squared test when the minimum expected count is less than five.

Results

Patients

A total of 61/284 (21.5%) *P. aeruginosa* isolates were recovered from respiratory samples of patients. These included 247 (86.9%) hospitalized patients and 37 (13.1%) non-hospitalized patients. Recovery of *P. aeruginosa*

was significantly more frequent ($p=0.036$) in hospitalized (73.8%) versus non-hospitalized patients (26.2%). Table 2 presents the demographic and clinical characteristics of the 61 investigated patients. Table 3 shows demographic characteristics and clinical features of all patients infected with *P. aeruginosa* and hospitalized in wards (60%) and ICUs (40%). All clinical features were almost similar in both groups. Only one female patient hospitalized in the ICU died during the study period.

Table 4. Antimicrobial susceptibility pattern of 61 *P. aeruginosa* isolates*

Antimicrobial agents	Susceptible isolates No. (%)	MIC ₅₀ (µg/mL)	MIC ₉₀ (µg/mL)	MIC range (µg/mL)
Amikacin	30 (49.1)	ND	ND	
Ciprofloxacin	30 (49.1)	2.3	4.1	0.012-32
Gentamicin	23 (37.7)	ND	ND	
Ceftazidime	50 (82.0)	1.8	3.3	0.38-64
Aztreonam	35 (57.3)	ND	ND	
Meropenem	48 (78.7)	2.8	4.9	0.016-32
Imipenem	49 (80.3)	3.6	6.5	0.19-32
Piperacillin - tazobactam	38 (62.2)	ND	ND	
Cefepime	50 (82.0)	ND	ND	
Cefotaxime	15 (24.6)	ND	ND	
Colistin	61 (100)	0.18	0.32	0.032-3.0

*32 of the isolates were multidrug resistant, recovered from hospitalized patients.

MIC – minimum inhibitory concentration; ND – not done.

Occurrence of *P. aeruginosa* and antimicrobial susceptibility profiles

The identity of 61 *P. aeruginosa* isolates was confirmed by biochemical testing and PCR. A total of 53 (87%) isolates showed clear blue-green pigment production (pyocyanin) on *Pseudomonas* CN agar. The antimicrobial susceptibility patterns and MICs of *P. aeruginosa* isolates for important used antibiotics are shown in Table 4.

A total of 32 (52.5%) isolates were completely resistant to three or more antibiotic classes (MDR) and all were recovered from hospitalized patients (Table 4).

MDR and putative virulence genes

All 32 MDR *P. aeruginosa* isolates carried one to four of the investigated ESBL genes and all were positive for genes of *algD*, *lasB*, *toxA* and pyocyanin, as shown in Table 5. Most MDR isolates (22/32; 68.9%) harbored *bla*_{CTX-M} genes, whereas other ESBL genes were much less common among the isolates, and none of the MDR *P. aeruginosa* isolates was positive for the *bla*_{IMP-1} or *bla*_{KPC} genes (Table 5).

Table 5. Distribution of ESBL and MBL genes in association with potential virulence genes among 32 MDR *P. aeruginosa* isolates

ESBL genes/ MBL genes	<i>P. aeruginosa</i> isolates, no. (%)*	Putative virulence genes / factors (%)**
<i>bla</i> _{CTX-M}	22 (68.8)	<i>algD</i> , <i>lasB</i> , <i>toxA</i> , pyocyanin (100)
<i>bla</i> _{VEB}	6 (18.9)	<i>algD</i> , <i>lasB</i> , <i>toxA</i> , pyocyanin (100)
<i>bla</i> _{TEM}	6 (18.9)	<i>algD</i> , <i>lasB</i> , <i>toxA</i> , pyocyanin (100)
<i>bla</i> _{GES}	5 (15.6)	<i>algD</i> , <i>lasB</i> , <i>toxA</i> , pyocyanin (100)
<i>bla</i> _{SHV}	4 (12.5)	<i>algD</i> , <i>lasB</i> , <i>toxA</i> , pyocyanin (100)
<i>bla</i> _{VIM-2}	3 (9.4)	<i>algD</i> , <i>lasB</i> , <i>toxA</i> , pyocyanin (100)
<i>bla</i> _{IMP-1}	0	-
<i>bla</i> _{KPC}	0	-

*All 10 isolates from cystic fibrosis cases were positive for the genes *bla*_{CTX-M}, *algD*, *lasB*, *toxA* and 4 of them secreted pyocyanin.

**Some of these isolates were simultaneously positive to two to four ESBL genes or one MBL gene.

The most common virulence genes detected among the all 61 isolates were *algD* and *lasB* (98%), followed by *toxA* (80%), *exoS* and *exoU* (33), and 87% were positive for pyocyanin (Table 6).

Table 6. Distribution of potential virulence genes among 61 *P. aeruginosa* isolates

Virulence factor	Gene	Positive virulence genes No. isolates (%)	
Alginate	<i>algD</i>	60	(98)
Elastase B	<i>lasB</i>	60	(98)
Exotoxin A	<i>toxA</i>	49	(80)
Exoenzyme S	<i>exoS</i>	20	(33)
Exoenzyme U	<i>exoU</i>	20	(33)
Pyocyanin		53	(87)

Genotyping

The RAPD profile showed high diversity among the isolates. Each isolate generated 2-6 fragments with band sizes ranging from 180 to 1200 bp. The dendrogram of the isolates is presented in Figure 1; by applying 70% similarity index of dendrogram, the isolates were clustered into 14 different groups. The major group was presented by 5 isolates and all other groups were presented with less than 4 isolates. There was no significant association between genotype groups and antimicrobial resistance phenotypes using RAPD cluster groups.

Discussion

The general rate of *P. aeruginosa* isolates recovery in this study (21.5%) is relatively in the middle of other recently reported studies from neighboring countries such as Egypt, Saudi Arabia, United Arab Emirates, where the isolation rates of *P. aeruginosa* from sputum samples of patients with lower respiratory tract infection ranged between 16.7% and 50%.^{9,21-23} A study from the United Arab Emirates reported that the most common pathogen isolated from the sputum of patients with hospital-acquired pneumonia was *P. aeruginosa* (50%), and that it caused a mortality rate of 24%.²³ A prevalence study conducted in ICU patients for one day in 72 countries around the world has also demonstrated that *P. aeruginosa* was one of the most common organisms, and that it accounted

for 19.9% of all organisms isolated from respiratory tract cultures.²⁴

Numerous studies reported that nosocomially-acquired *P. aeruginosa* isolates tend to be often more MDR and commonly identified ICU patients.^{9,24-26} The rate of MDR *P. aeruginosa* isolates in our study was 52.2%, and this result is similar to recently reported studies from Jordan and neighboring Arab countries (Egypt, Saudi Arabia, Lebanon).^{23,25-27}

The present study showed that hospitalized patients were significantly ($p=0.036$) more frequently infected with *P. aeruginosa* than non-hospitalized patients in association with lower respiratory tract infection. This study also showed that almost all 61 patients with positive *P. aeruginosa* isolates had purulent sputum, cough and were treated with antibiotics (96.7%), whereas other significant clinical respiratory infection features such as chest pain, dyspnea and hemoptysis were less frequently observed in our patients. There was only one death in patients with lower respiratory infection during the study period.

The present study demonstrates that MDR *P. aeruginosa* isolates (68.9%) commonly are ESBL-producers and *bla*_{CTX-M} was the most frequent phenotype detected in these isolates, whereas the other types of *TEM*, *VEB*, *GES*, and *SHV* genes were less often detected (Table 5). The spread of ESBL-producing Gram-negative bacteria is considered an important driver for the use of carbapenems, which can increase the selection of carbapenemase-producing organisms causing infections, particularly in ICU patients.²⁸

Most recent studies reported that hospital outbreaks caused by MDR *P. aeruginosa* clones with MBLs such as VIM and IMP types can be developed by either chromosomal mutations or horizontal gene transfer.²⁹ VIMs are the most frequent MBL type found among members of the Enterobacteriaceae family, as well as among *Pseudomonas* spp. and *Acinetobacter* spp.²⁹ The present study indicated that MBLs were only detected as type VIM-2 in a few of our MDR *P. aeruginosa* isolates (9.4%), while the other types, *bla*_{IMP} and *bla*_{KPC}, were absent (Table 5). In Jordan, a recent study has also reported that all

Figure 1. Dendrogram based on RAPD results showing relatedness between 32 MDR *P. aeruginosa* isolates; the dendrogram was generated by the neighbor joining method with PyElph software (USA).

Acinetobacter baumannii isolates from the hospital, environment, and patients' respiratory tract were negative for *bla*_{VIM} and *bla*_{IMP-1} genes.³⁰

The pathogenic potential of *P. aeruginosa* is complex and multifactorial, and virulence genes, especially *algD*, *lasB*, *toxA* and *exoS* can induce biofilm formation, infection and cytotoxicity in the human body, especially in the respiratory tract.^{6,31} Additionally, *P. aeruginosa* strains carrying *exoU* genes can cause a more virulent form of pneumonia than other strains.³² The prevalences of the virulence genes *algD*, *lasB*, *toxA* and secretion of pyocyanin in this study were very high among *P. aeruginosa* isolates (98%-80%), and all MDR isolates carried those 4 virulence factors, while both *exoS* and *exoU* genes were found less frequently and only in 33% of the isolates (Table 6). A recent study indicated that pyocyanin production was higher in earlier *P. aeruginosa* isolates from the respiratory tract compared to the isolates obtained later in the course of infection.³³ It has been also reported that pyocyanin induces various antagonistic effects on the host, especially pro-inflammatory

and free radical effects which cause cellular damage and death.³³

The relationship between antimicrobial resistance markers and distribution of common virulence factors among *P. aeruginosa* isolates in this study was not significant, since all MDR isolates and most of the susceptible isolates carried 3 potential virulence genes (*algD*, *lasB* and *toxA*) and secreted pyocyanin (Table 5). A French study detected the virulence genes *algD*, *lasB* and *toxA* in all *P. aeruginosa* isolates from the sputa of patients with cystic fibrosis and, to a lesser rate, from other non-cystic fibrosis isolates.³ A Bulgarian study reported that *P. aeruginosa* isolates from all clinical samples including the respiratory tract carried 100% *lasB* genes and a high rate of *algD* genes (95.3%) in cystic fibrosis cases.²³ A study from Iran demonstrated also that all *P. aeruginosa* isolates from the pulmonary tract harbored significantly higher rates of *toxA* genes than wound and burn isolates ($p < 0.004$), and other virulence genes such as *algD* and *lasB* were less common in respiratory tract specimens.⁴

It can be concluded from our study and other studies that virulence genes *lasB*, *algD*, and *toxA* are mostly associated with *P. aeruginosa* isolates from the pulmonary tract. However, none of the mentioned studies has previously reported on the association between antimicrobial resistance and virulence potential of their *P. aeruginosa* isolates as it has been done in this study.

RAPD fingerprinting of 32 MDR *P. aeruginosa* isolates from hospitalized patients and the construction of the dendrogram of the isolates demonstrated high diversity among *P. aeruginosa* isolates, with 14 genotype groups, by applying 70% similarity index (Figure 1). Association between the antimicrobial resistance phenotypes and genotype profiles of these 32 isolates revealed a limited similarity of DNA bands patterns between phenotypes and genotypes. Based on these results, it can be suggested that most isolates originated from various clones and there was no single genotype prevalent among more than 5 isolates.

Conclusions

In conclusion, this study has found no significant genetic relationship among MDR *P. aeruginosa* isolates from the respiratory tract of hospitalized patients or their carriage of virulence genes.

Authors' contributions statement: HYAD performed all laboratory tests, NO supervised all clinical issues of patients, LFAQ helped in genotyping of the isolates, AAS supervised all laboratory tests and writing of the manuscript. All authors read and approved the final version of the manuscript.

Conflicts of interest: All authors – none to disclose.

Funding: This work was supported by a grant from The Jordan University, Dean of Research (No. 74/2014-2015).

Acknowledgement: The authors are thankful to the Jordan University Hospital for allowing them to collect clinical specimens and data from the admitted patients. Also, the authors thank Mrs. Ilham Abu-Khader for performing the statistical analysis of the study.

References

- Nathwani D, Raman G, Sulham K, Gavaghan M, Menon V. Clinical and economic consequences of hospital-acquired resistant and multidrug-resistant *Pseudomonas aeruginosa* infections: a systematic review and meta-analysis. *Antimicrob Resist Infect Control* 2014;3:32. [\[Crossref\]](#)
- Mitov I, Strateva T, Markova B. Prevalence of virulence genes among Bulgarian nosocomial and cystic fibrosis isolates of *Pseudomonas aeruginosa*. *Braz J Microbiol* 2010;41:588-95. [\[Crossref\]](#)
- Lanotte P, Watt S, Mereghetti L, et al. Genetic features of *Pseudomonas aeruginosa* isolates from cystic fibrosis patients compared with those of isolates from other origins. *J Med Microbiol* 2004;53:73-81. [\[Crossref\]](#)
- Nikbin VS, Aslani MM, Sharafi Z, Hashemipour M, Shahcheraghi F, Ebrahimipour GH. Molecular identification and detection of virulence genes among *Pseudomonas aeruginosa* isolated from different infectious origins. *Iran J Microbiol* 2012;4:118-23.
- Michalska M, Wolf P. *Pseudomonas* exotoxin A: optimized by evolution for effective killing. *Front Microbiol* 2015;6:963. [\[Crossref\]](#)
- Shaver CM, Hauser AR. Relative contributions of *Pseudomonas aeruginosa* ExoU, ExoS, and ExoT to virulence in the lung. *Infect Immun* 2004;72:6969-77. [\[Crossref\]](#)
- Farshadzadeh Z, Khosravi AD, Alavi SM, Parhizgari N, Hoveizavi H. Spread of extended-spectrum β -lactamase genes of *bla*_{OXA-10}, *bla*_{PER-1} and *bla*_{CTX-M} in *Pseudomonas aeruginosa* strains isolated from burn patients. *Burns* 2014;40:1575-80. [\[Crossref\]](#)
- Poirel L, Nordmann P, Lagrutta E, Cleary T, Munoz-Price LS. Emergence of KPC-producing *Pseudomonas aeruginosa* in the United States. *Antimicrob Agents Chemother* 2010;54:3072. [\[Crossref\]](#)
- Mansour SA, Eldaly O, Jiman-Fatani A, Mohamed ML, Ibrahim EM. Epidemiological characterization of *P. aeruginosa* isolates of intensive care units in Egypt and Saudi Arabia. *East Mediterr Health J* 2013;19:71-80.
- Hamze M, Mallat H, Dabboussi F, Achkar M. Antibiotic susceptibility and serotyping of clinical *Pseudomonas aeruginosa* isolates in northern Lebanon. *Int Arabic J Antimicrob Agents* 2012;2:1-6.
- Clinical Laboratory and Standards Institute (CLSI). Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; approved standard – tenth edition. CLSI document M07-A10 Villanova, PA, USA: CLSI, 2015.
- Spilker T, Coenye T, Vandamme P, LiPuma JJ. PCR-based assay for differentiation of *Pseudomonas aeruginosa* from other *Pseudomonas* species recovered from cystic fibrosis patients. *J Clin Microbiol* 2004;42:2074-9. [\[Crossref\]](#)
- Bokaeian M, Shahraki Zahedani S, Soltanian Bajgiran M, Ansari Moghaddam A. Frequency of PER, VEB, SHV, TEM and CTX-M genes in resistant strains of *Pseudomonas aeruginosa* producing extended spectrum β -lactamases. *Jundishapur J Microbiol* 2014;8:e13783. [\[Crossref\]](#)
- Pitout J DD, Gregson DB, Poirel L, McClure JA, Le P, Church DL. Detection of *Pseudomonas aeruginosa* producing metallo- β -lactamases in a large centralized laboratory. *J Clin Microbiol* 2005;43:3129-35. [\[Crossref\]](#)
- Jiang X, Zhang Z, Li M, Zhou D, Ruan F, Lu Y. Detection of extended-spectrum β -lactamases in clinical isolates of *Pseudomonas aeruginosa*. *Antimicrob Agents Chemother* 2006;50:2990-5. [\[Crossref\]](#)
- Fazeli N, Momtaz H. Virulence gene profiles of multidrug resistant *Pseudomonas aeruginosa* isolated from Iranian hospital infections. *Iran Red Crescent Med J* 2014;16:e15722. [\[Crossref\]](#)
- Monteiro J, Widen RH, Pignatari AC, Kubasek C, Silbert S. Rapid detection of carbapenemase genes by multiplex real-time PCR. *J Antimicrob Chemother* 2012;67:906-9. [\[Crossref\]](#)
- Akpaka PE, Swanston WH, Ihemere HN, et al. Emergence of KPC-producing *Pseudomonas aeruginosa* in Trinidad and Tobago. *J Clin Microbiol* 2009;47:2670-1. [\[Crossref\]](#)
- Wolska K, Szveda P. Genetic features of clinical *Pseudomonas aeruginosa* strains. *Pol J Microbiol* 2009;58:255-60.
- Mitov I, Strateva T, Markova B. Prevalence of virulence genes among Bulgarian nosocomial and cystic fibrosis isolates of *Pseudomonas aeruginosa*. *Braz J Microbiol* 2010;41:588-95. [\[Crossref\]](#)
- Nanvazadeh F, Khosravi AD, Zolfaghari MR, Parhizgari N. Genotyping of *Pseudomonas aeruginosa* strains isolated from burn patients by RAPD-PCR. *Burns* 2013;39:1409-13. [\[Crossref\]](#)

22. Fatima A, Naqvi SB, Khaliq SA, Perveen S, Jabeen S. Antimicrobial susceptibility pattern of clinical isolates of *Pseudomonas aeruginosa* isolated from patients of lower respiratory tract infections. Springerplus 2012;1:70. [\[Crossref\]](#)
23. Al-Muhairi S, Zoubeydi T, Ellis M, Nicholls MG, Safa W, Joseph J. Demographics and microbiological profile of pneumonia in United Arab Emirates. Monaldi Arch Chest Dis 2006;65:13-8.
24. Vincent JL, Rello J, Marshall J, et al. International study of the prevalence and outcomes of infection in intensive care units. JAMA 2009;302:2323-9. [\[Crossref\]](#)
25. Al-Hajje A, Ezedine M, Hammoud H, et al. [Current status of nosocomial infections in the Lebanese Hospital Center, Beirut]. East Mediterr Health J 2012;18:495-500.
26. Wassef M, El Mahallawy H, Zafer MM, Ghaith D, Abdel Hamid R. Lab based surveillance of multidrug resistant *Pseudomonas aeruginosa* in Cairo University Hospitals, Egypt. J Microbiol Exp 2015;2:00039. [\[Crossref\]](#)
27. Masaadeh HA, Jaran AS. Incident of *Pseudomonas aeruginosa* in post-operative wound infection. Am J Infect Dis 2009;5:1-6. [\[Crossref\]](#)
28. Voor In 't Holt AF, Severin JA, Lesaffre EM, Vos MC. A systematic review and meta-analyses show that carbapenem use and medical devices are the leading risk factors for carbapenem-resistant *Pseudomonas aeruginosa*. Antimicrob Agents Chemother 2014;58:2626-37. [\[Crossref\]](#)
29. Zhao WH, Hu ZQ. Acquired metallo- β -lactamases and their genetic association with class 1 integrons and ISCR elements in Gram-negative bacteria. Future Microbiol 2015;10:873-87. [\[Crossref\]](#)
30. Obeidat N, Jawdat F, Al-Bakri GA, Shehabi AA. Major biologic characteristics of *Acinetobacter baumannii* isolates from hospital environmental and patients' respiratory tract sources. Am J Infect Control 2014;42:401-4. [\[Crossref\]](#)
31. Moreau-Marquis S, Stanton BA, O'Toole GA. *Pseudomonas aeruginosa* biofilm formation in the cystic fibrosis airway. Pulm Pharmacol Ther 2008;21:595-9. [\[Crossref\]](#)
32. Schulert GS, Feltman H, Rabin SD, Martin CG, Battle SE, Rello J, Hauser AR. Secretion of the toxin ExoU is a marker for highly virulent *Pseudomonas aeruginosa* isolates obtained from patients with hospital-acquired pneumonia. J Infect Dis 2003;188:1695-706. [\[Crossref\]](#)
33. Caldwell CC, Chen Y, Goetzmann HS, et al. *Pseudomonas aeruginosa* exotoxin pyocyanin causes cystic fibrosis airway pathogenesis. Am J Pathol 2009;175:2473-88. [\[Crossref\]](#)

Please cite this article as:

Al Dawodeyah HY, Obeidat N, Abu-Qatouseh LF, Shehabi AA. Antimicrobial resistance and putative virulence genes of *Pseudomonas aeruginosa* isolates from patients with respiratory tract infection. GERMS 2018;8(1):31-40. doi: 10.18683/germs.2018.1130