Bacterial Surface Spreading Is More Efficient on Nematically Aligned Polysaccharide Substrates

David J. Lemon,^a Derek A. Schutzman,^a Anthony G. Garza^a

^aDepartment of Biology, Syracuse University, Syracuse, New York, USA

ABSTRACT Biofilm-forming bacteria typically deposit layers of polysaccharides on the surfaces they inhabit; hence, polysaccharides are their immediate environment on such surfaces. Previously, we showed that many biofilm-forming bacteria preferentially spread in the direction of aligned and densely packed polysaccharide fibers in compressed substrates, a behavior we referred to as polymertropism. This arrangement of polysaccharide fibers is likely to be similar to that found in the “slime” trails deposited by many biofilm-forming bacteria and would explain previous observations that bacteria tend to follow these trails of polysaccharides. Here, we show that groups of cells or flares spread more rapidly on substrates containing aligned and densely packed polysaccharide fibers. Flares also persist longer, tend to hold their trajectories parallel to the long axes of polysaccharide fibers longer, and ultimately show an increase in displacement away from their origin. On the basis of these findings and others, we propose a model for polymertropism. Namely, we suggest that the packing of the aligned polymers increases the efficiency of surface spreading in the direction of the polymer’s long axes; therefore, bacteria tend to spread more rapidly in this direction. Additional work suggests that bacteria can leverage polymertropism, and presumably more efficient surface spreading, for a survival advantage. In particular, when two bacterial species were placed in close proximity and in competition with each other, the ability of one species to move rapidly and directly away from the other by utilizing the aligned polymers of compressed agar substrates led to a clear survival benefit.

IMPORTANCE The directed movement of bacteria on compressed substrates was first described in the 1940s and referred to as elasticotaxis (R. Y. Stanier, *J Bacteriol* 44:405–412, 1942). More recently, this behavior was referred to as polymertropism, as it seems to be a response to the nematic alignment and tight packing of polymers in the substrate (D. J. Lemon, X. Yang, P. Srivastava, Y. Y. Luk, A. G. Garza, *Sci Rep* 7:7643, 2017, <https://doi.org/10.1038/s41598-017-07486-0>). The data presented here suggest that bacteria are more efficient at surface spreading when the polymers in the substrate are arranged in this manner. These data also suggest that bacteria can leverage polymertropism, and presumably more efficient surface spreading, for a survival advantage. Namely, one bacterial species was able to use its strong polymertropism response to escape from and survive competition with another species that normally outcompetes it.

KEYWORDS biofilms, elasticotaxis, polymertropism

Bacteria often change their behavior in response to cues from the environment. Because chemical signals have received the most attention from researchers (1–3), much less is known about the physiological and behavioral responses of bacteria to physical or mechanical signals. Despite the focus on chemical signals, responses to physical signals have been identified in some groups of bacteria such as the myxobacteria. For example, an exposure to light stimulates the process of fruiting body

Received 11 October 2017 Accepted 18 December 2017

Accepted manuscript posted online 8 January 2018

Citation Lemon DJ, Schutzman DA, Garza AG. 2018. Bacterial surface spreading is more efficient on nematically aligned polysaccharide substrates. *J Bacteriol* 200:e00610-17. <https://doi.org/10.1128/JB.00610-17>.

Editor George O’Toole, Geisel School of Medicine at Dartmouth

Copyright © 2018 American Society for Microbiology. All Rights Reserved.

Address correspondence to Anthony G. Garza, agarza@syr.edu.

formation in *Stigmatella aurantiaca* (4–6), and placing grooves in the substrate of *Myxococcus xanthus* stimulates its cellular movements (7).

In 1999, Fontes and Kaiser (8) showed that *M. xanthus* responds to compression of its agar substrate at the individual cell, group, and colony levels. In particular, individual *M. xanthus* cells, which are rod shaped and move in the direction of their long axes, orient their long axes perpendicular to the direction of compression of a polysaccharide substrate. Furthermore, the outward movements of groups of cells from the edges of colonies and the overall expansion of these colonies are primarily oriented perpendicular to the direction of compression, which results in elliptically shaped colonies. The directed group movements and colony spreading are abolished when the cells are defective for a type of surface motility called adventurous motility (A-motility), indicating that these *M. xanthus* behaviors are dependent on the active movement of cells and are not passive events (8).

In a recent study of *M. xanthus*, we showed that the directed group movements and colony expansion on compressed polysaccharide substrates are unlikely to be responses to the water that accumulates on the surface after compression or to compression-induced changes in surface topography (9). Because the directed surface movements consistently match the orientation of the long axes of nematically aligned and tightly packed polysaccharide fibers in compressed substrates, we proposed that *M. xanthus* responds to the compression-induced physical arrangement of the polysaccharide fibers. We also suggested that the focal adhesion complex, which is fixed to the substrate and helps generate the traction force for A-motility (10), is better at gripping tightly packed polysaccharide fibers; hence, *M. xanthus* tends to move along these fibers, which are aligned perpendicular to the direction of compression.

This preferential spreading perpendicular to the axis of substrate compression is not unique to *M. xanthus*; we detected this response in many rod-shaped bacteria with motility systems for surface spreading and in one bacterial species (*Bacillus mycoides*) that rapidly spreads across surfaces via an unknown mechanism (9). However, we did not detect this response in bacteria that have been classified as nonmotile and lack the ability to spread extensively on the surface of a semisolid substrate. These findings suggest a strong connection between surface motility and the preferential spreading perpendicular to the axis of substrate compression.

Because the bacteria that respond to substrate compression appear to be responding to the orientation of the polymers in the substrate and are not sensing and moving up or down a gradient as indicated by the previous term (elasticotaxis) given to this behavior (11), we renamed this behavior polymertropism (9); tropism is the directional movement of an organism in response to a particular stimuli.

The work presented here suggests that the robust surface spreading and polymertropism response of *B. mycoides* are due at least in part to surface motility and not strictly to growth; the surface spreading and polymertropism response continue for more than 1.5 days in the absence of nutrients for growth. The strong polymertropism response of *B. mycoides* on agar substrates helps it survive in competition assays with *Serratia marcescens*, which has no detectable polymertropism response on agar (9, 12). Namely, when *B. mycoides* and *S. marcescens* were combined in a single inoculum on compressed agar substrates, *B. mycoides*' ability to survive competition with *S. marcescens* was significantly improved compared to that on uncompressed agar substrates. Evidently, *S. marcescens* outcompetes *B. mycoides* on uncompressed substrates, but on compressed substrates with aligned and tightly packed polysaccharide fibers, *B. mycoides* is able escape the *S. marcescens* footprint, and a greater number of *B. mycoides* cells then survive. This is the first time an advantage derived from polymertropism has been experimentally observed and quantified, although such an advantage has been the subject of speculation (8, 13).

To examine how aligned and tightly packed polysaccharide fibers in compressed substrates affect surface spreading, we placed four bacterial species on compressed and uncompressed agar substrates and tracked flares or groups of cells emerging from the edges of colonies. Interestingly, the general trends for each bacterial species were

similar, even though one species (*M. xanthus*) uses gliding motility to spread across the surface of the substrate (14–16), two species (*Bacillus cereus* and *Bacillus subtilis*) use flagellar motility (17–20), and one species (*B. mycoides*) has an unknown mechanism of surface motility (12, 21). In particular, the vast majority of flares on compressed substrates moved in the direction of the long axes of the tightly packed polysaccharide fibers. Furthermore, these flares maintained the same general trajectory for substantially longer than flares on uncompressed substrates and had greater surface spreading speeds and overall displacements than flares on uncompressed substrates. It is unclear how these collective changes in flare spreading dynamics affect whole-colony spreading in the direction of polysaccharide fiber alignment; however, we were able to identify a strong relationship between flare displacement and this directed colony spreading. A number of pieces of data suggest that polymertropism can be explained by a simple model in which the arrangement of the polymers in the substrate promotes more efficient surface spreading in a particular direction. Examples (in addition to the one described here) of how bacteria might leverage polymertropism and more efficient spreading to their benefit are discussed.

RESULTS

Polymertropism helps *B. mycoides* compete. We previously suggested that polymertropism might be advantageous to bacteria in certain situations (8, 9, 11, 12). For example, many bacteria produce trails of polysaccharides, which are predicted to be aligned and densely packed, as they move across the surfaces of their substrates (22–27). Because of the arrangement of the polysaccharides in the trails, polymertropism could result in the alignment and coordinated movement of large groups of bacterial cells and ultimately help the bacteria organize over large distances without the need for a chemical signal.

To explore whether polymertropism provides a survival advantage, we focused on a bacterium (*B. mycoides*) that spreads rapidly on the surfaces of nutrient-rich agar substrates and has a robust polymertropism response when such substrates are compressed (see Fig. S1 in the supplemental material). Since it is unclear whether *B. mycoides* spreads on a substrate's surface via growth or motility, we examined whether surface spreading and polymertropism occur in the absence of nutrients for growth. As shown in Fig. S1, these behaviors do not require nutrients for growth and continue over the course of 1.5 days in the absence of nutrients. These results suggest that motility plays an important role in surface spreading and polymertropism in *B. mycoides*. However, *B. mycoides* spreads faster and has a stronger polymertropism response on agar substrates that are rich in nutrients (Fig. S1). On the basis of these findings, we propose that motility, as well as growth and/or a nutrient supply for energy, is important for optimal surface spreading and polymertropism in *B. mycoides*.

In subsequent experiments, we subjected *B. mycoides* and *S. marcescens*, which have a strong polymertropism response and no polymertropism response on agar substrates, respectively, to a survival assay. This assay enabled us to directly quantify a fitness or survival benefit derived from the polymertropism response. In particular, various ratios of *B. mycoides* and *S. marcescens* cultures were mixed and spotted on compressed and uncompressed agar substrates, and the cells were incubated for 24 h (Fig. 1A and B). The cells were then harvested and diluted in CTTYE broth (10 g/liter Casitone, 2 g/liter yeast extract, 1 mM KH_2PO_4 , 10 mM Tris-HCl [pH 8.0], and 8 mM MgSO_4) and plated to determine the numbers of white colonies produced by *B. mycoides* cells and red colonies produced by *S. marcescens* cells (Fig. 1C).

When mixed at a 1:3 or 1:1 ratio with *S. marcescens* and placed on uncompressed substrates, *B. mycoides* yielded no CFU. When the ratio of the mixture was changed to 3:1, we were able to detect *B. mycoides* CFU, but at a relatively low level (approximately 5%) compared with that of *S. marcescens*. *B. mycoides* yielded substantially more CFU when the same cell mixtures were placed on compressed substrates; however, *B. mycoides* still yielded far fewer CFU than *S. marcescens* (approximately 11%, 20%, and 11% of that of *S. marcescens* at 1:3, 1:1, and 3:1 ratios, respectively). These findings

FIG 1 Competition assays with *S. marcescens* and *B. mycoides*. Various ratios of *B. mycoides* and *S. marcescens* cultures were mixed, spotted onto compressed and uncompressed agar substrates, and incubated for 24 h. (A) Results from an assay in which *S. marcescens* cells (red) and *B. mycoides* cells (white) were mixed at a ratio of 3 to 1 and spotted onto uncompressed agar. (B) Results when the same ratio of *S. marcescens* and *B. mycoides* cells was spotted onto a compressed agar substrate. The white arrows show the orientation of the axis of compression. (C) Mean numbers of CFU produced by *B. mycoides* and *S. marcescens* cells in three competition assays on uncompressed (UC) and compressed (C) agar. Error bars represent standard errors of the means and asterisks denote significant differences between the indicated means. (D) Some *B. mycoides* colonies that arose from cells exposed to compressed substrates displayed a rough phenotype (left) that is different from the expected filamentous s-shaped phenotype (right). Black arrows indicate *S. marcescens* colonies.

indicate that *B. mycoides*' ability to survive after being placed in mixed cultures with *S. marcescens* substantially improves when the substrate is compressed. Indeed, it seems that a substantial number of *B. mycoides* cells are able to escape the footprint of *S. marcescens* on compressed but not uncompressed substrates (Fig. 1A and B). Interestingly, *S. marcescens* also yielded a larger number of CFU when the cell mixtures were placed on compressed substrates instead of uncompressed substrates, although the increase in cell survival was much less dramatic than that for *B. mycoides* (Fig. 1C). In addition, we found that *B. mycoides* cells harvested from mixed cultures on compressed substrates produced colonies with a rough morphology instead of the typical s-shaped filamentous morphology (Fig. 1D). The latter points are further addressed in Discussion.

***B. mycoides*' flare speed, duration, and displacement increase on compressed substrates.** The results of the survival assays suggest that under certain conditions (i.e., with a high density of aligned polysaccharide already present), a bacterial species might be able to leverage polymertropism to better survive competition with another species

FIG 2 *B. mycoides*' response to substrate compression. Eight 4- μ l drops of mid-log-phase *B. mycoides* liquid cultures were placed in a column down the centerline of uncompressed or compressed agar substrates using a multichannel pipettor. After the spots dried, the plates were incubated for 30 h, and images of each plate were taken. The diameters of the inserted tubes, which held the agar substrates under various degrees of compression, are shown. Arrows show the orientations of the axes of compression. Scale bars, 1 cm.

that does not have a similarly strong polymertropism response. *B. mycoides* has the strongest whole-colony polymertropism response that we have observed to date (Fig. 2) (9, 12). To explore the potential advantages that the polymertropism response confers to *B. mycoides*, we examined how compression changes the surface spreading dynamics of its colony flares. Briefly, by comparing flare tip positions from one image to the next in ImageJ (28) and tracking these changes over time, we generated trajectories for each flare, from which we extracted the minute-to-minute displacement (speed), the flare duration, and the total displacement of the flare at the end of tracking (Fig. 3).

On uncompressed substrates, *B. mycoides* cells formed flares (see Video S1) that tended to gently curve back to the edges of the colonies from which they emerged (Fig. 4A and B). Furthermore, the flares on uncompressed substrates did not have any clear overarching directionality. In contrast, the flares that formed on compressed substrates were much straighter (Fig. 4C and D), moving nearly parallel to the x axis in response to compression in the direction of the y axis. The mean relative (x,y) values were calculated, as shown in Fig. 3, for both conditions, and the plot of the mean positions in both axes revealed that flares on uncompressed substrates generally remained closer to the colony edges from which they emerged, while flares on compressed substrates moved directly away from the colony edges and continued to do so over time (Fig. 4E; see also Fig. S2). In addition to flare direction or trajectory, flare speed was altered. Namely, on uncompressed substrates, *B. mycoides* flares moved at an average speed of $\sim 2.2 \mu\text{m}$ per min, while on compressed substrates, flares moved more than $3.9 \mu\text{m}$ per min, which is an increase in flare speed of approximately 78% on compressed substrates (Table 1). The flare duration also increased on compressed substrates, as shown in Table 1, but only by approximately 20%. On the basis of the graph of the mean flare displacement over time (Fig. 4F; Fig. S2), it is clear that flares on compressed substrates moved away from their origins more quickly and continued to do so, as the mean displacement continued to increase, while flares on uncompressed substrates moved away from their origins initially, but the displacement appeared to plateau after approximately 6 h (Video S1). Together, these findings indicate that compression-induced alignment and tight packing of polysaccharide fibers change the surface

FIG 3 Schematic diagrams of flare tracking data analyses. (A) Flare tracks were extracted from time-lapse video results in raw (x,y) position data. The initial (x,y) value is subtracted from each track's raw (x,y) values such that all flare tracks start at (0,0). (B) The new (x,y) values are multiplied by -1 as needed, such that the flare tracks are predominantly located in quadrant I. (C) Relative (x,y) data were averaged for each time interval to generate mean relative (x,y) values ($\pm$ standard errors of the means). (D) The minute-to-minute distances that flares moved were calculated and pooled to calculate mean flare speed. Flare displacements at the end of tracks (calculated as shown) were pooled to calculate mean displacement at the end of flare tracking.

spreading dynamics of groups of *B. mycoides* cells, and some of these changes, such as an increase in speed and overall displacement, are likely to be beneficial in certain situations.

Spreading dynamics of *B. subtilis*, *B. cereus*, and *M. xanthus* flares are similarly altered on compressed substrates. In addition to that in *B. mycoides*, flare behavior was monitored in *B. subtilis* (Fig. 5; Fig. S2 and Video S2), *B. cereus* (Fig. 6; Fig. S2 and Video S3), and *M. xanthus* (Fig. 7; Fig. S2 and Video S4). These bacterial species were chosen because their polymertropism responses are variable but weaker than that of *B. mycoides* (9). In addition, *M. xanthus* uses gliding motility to spread across the surface of its substrate (14–16, 25), *B. cereus* and *B. subtilis* use flagellar motility (17–21) for surface spreading, and *B. mycoides* has an uncharacterized motility system for surface spreading. Therefore, at least two and perhaps three different mechanisms of surface motility are represented in these bacteria.

Although we observed some variations in flare dynamics between species, the general trends were similar to that of *B. mycoides* flares (Fig. 5 to 7). Namely, on uncompressed substrates, the flares for all species appeared to have no overarching directionality (Fig. 5A and B, 6A and B, and 7A and B), while on compressed substrates, the dominant flare orientation was parallel to the x axis (perpendicular to the axis of compression) (Fig. 5C and D, 6C and D, and 7C and D). Furthermore, on compressed substrates, we observed increases in flare speed, duration, and the overall displacement at the end of tracking (Fig. 5E and F, 6E and F, and 7E and F; Fig. S2). The largest fold increase in flare speed on compressed substrates was observed in *B. subtilis*; its mean flare speed increased 2.07-fold (from approximately 0.88 to 1.82 μm per min) on compressed substrates (Table 1). The speed of *B. cereus* flares increased 1.79-fold on compressed substrates. This increase in flare speed was slightly less than that of *B. subtilis* and nearly identical to that of *B. mycoides* (1.78-fold increase). The smallest increase in flare speed on compressed substrates was observed in *M. xanthus*. Specifically, the speed of *M. xanthus* flares increased only slightly, from approximately 1.28 μm per min on uncompressed substrates to 1.47 μm per min on compressed substrates (Table 1). However, *M. xanthus* had the largest (4-fold) increase in the mean duration of

FIG 4 *B. mycoides* tracking data and analysis. (A, B, E, and F) The data for flare spreading dynamics on uncompressed substrates were extracted from 32 flare tracks on 3 videos. (C, D, E, and F) The data for flare spreading dynamics on compressed substrates were extracted from 21 flare tracks from 3 videos. The graphs in panels E and F with standard errors of the means are shown in Fig. S2A and B in the supplemental material.

its flares on compressed substrates (Table 1). In addition, *M. xanthus* had the largest fold increase in mean flare displacement at the end of tracking on compressed substrates; *M. xanthus* flares moved 7.8 times farther away from their origins on compressed substrates than comparable flares on uncompressed substrates (Table 1). *B. subtilis* showed a 2.4-fold increase in flare duration on compressed substrates (Table 1), while the increases in the durations for *B. cereus* and *B. mycoides* flares on compressed substrates were more modest (1.6- and 1.2-fold, respectively) (Table 1).

There is a strong relationship between flare displacement and whole-colony polymertropism. Given that compression-induced polysaccharide fiber alignment/

TABLE 1 Mean flare speed, duration, and displacement for each bacterial species^a

Species	Flare speed ($\mu\text{m}/\text{min}$)		Flare duration (min)		Final displacement (μm)	
	Uncompressed	Compressed	Uncompressed	Compressed	Uncompressed	Compressed
<i>B. mycoides</i>	2.196 ± 0.04795	3.911 ± 0.09417	272.0 ± 29.71	327.8 ± 23.94	354.9 ± 43.62	$1,045 \pm 77.54$
<i>B. subtilis</i>	0.8769 ± 0.03998	1.819 ± 0.04842	96.90 ± 7.279	236.2 ± 24.30	47.19 ± 4.079	307.8 ± 50.29
<i>B. cereus</i>	2.917 ± 0.07240	5.223 ± 0.1322	112.7 ± 8.455	184.9 ± 16.00	215.5 ± 29.59	720.2 ± 87.83
<i>M. xanthus</i>	1.282 ± 0.03237	1.471 ± 0.02290	179.0 ± 16.77	745.5 ± 51.24	74.54 ± 6.403	584.4 ± 44.03

^aMean values ($\pm$ standard errors of the means) for flare speed, flare duration, and displacement at track ends were derived from the data presented in Fig. 4 to 7.

packing changes the surface spreading dynamics of groups of bacterial cells, we asked whether we could identify a relationship between any of these changes in group surface spreading dynamics and whole-colony polymertropism (see Fig. 8A for images of the polymertropism response). We first quantified whole-colony polymertropism rates by monitoring colony aspect ratios over time (Fig. 8B), rather than at a single time point, which would represent only a snapshot of the colony spreading pattern. For each colony observed, we applied a line of best fit for the colony aspect ratios over time and interpreted the slope of this line to represent the mean rate of polymertropism. It should be noted that polymertropism in *B. subtilis* was not conducive to this type of analysis, as we found a significant amount of variation in the aspect ratios of its colonies (data not shown). Of the three remaining bacterial species, *B. mycoides* had the greatest mean rate of polymertropism, followed by *B. cereus* and *M. xanthus*. The mean polymertropism rates of the three species varied widely: *B. mycoides*' polymertropism rate was approximately 8 times that of *B. cereus*, and *B. cereus*' polymertropism rate was approximately 6 times that of *M. xanthus*. Hence, the colonies of *B. mycoides* become elongated parallel to the *x* axis (perpendicular to the axis of compression) significantly faster than those of *B. cereus* or *M. xanthus*, and *B. cereus* colonies become elongated parallel to the *x* axis significantly faster than those of *M. xanthus*.

We subsequently graphed the mean flare speeds, durations, and displacement as a function of the mean rates of polymertropism for *B. mycoides*, *B. cereus*, and *M. xanthus* and used R^2 values for linear best-fit lines of ≥ 0.9 to indicate a strong relationship between a particular aspect of flare spreading dynamics and polymertropism (see Fig. S3). The only combination that met our threshold was the mean flare displacement on compressed substrates and mean polymertropism rate, which yielded an R^2 value of approximately 0.96. These findings suggest a strong relationship between flare displacement and the directional spreading that is the hallmark of polymertropism.

There are strong relationships between the spreading dynamics on compressed and uncompressed substrates. In additional experiments, we examined the relationship between the rate of polymertropism (rate of directional surface spreading on compressed substrates) and the rate of surface spreading on uncompressed substrates. For this analysis, we used *Paenibacillus dendritiformis* as a substitute for *B. subtilis*, as this bacterium also utilizes flagellar motility for spreading on a surface (29) and has a consistent polymertropism response on compressed agar substrates. As shown in Table 1, the mean rate of polymertropism for *P. dendritiformis* was greater than those for *B. cereus* and *M. xanthus*, but less than that for *B. mycoides*. We determined the mean rates of surface spreading (change in colony diameter over time) of *P. dendritiformis*, *B. cereus*, *M. xanthus*, and *B. mycoides* on uncompressed substrates using the same strategy described above for calculating the mean rate of polymertropism (Fig. 8B to D). The results summarized in Fig. 8C indicate that *B. mycoides* had the highest rate of surface spreading on uncompressed agar, followed by *P. dendritiformis*, *M. xanthus*, and *B. cereus*. When we analyzed the mean polymertropism rates on compressed agar as a function of the mean surface spreading rates on uncompressed agar and calculated the R^2 value as described above, we uncovered a strong relationship between the two whole-colony behaviors: the R^2 value was approximately 0.94 (Fig. 8E). Similarly, we identified a strong relationship between mean flare speeds on uncompressed and compressed substrates ($R^2 = 0.90$) and between mean flare dis-

FIG 5 *B. subtilis* tracking data and analysis. (A, B, E, and F) The data for flare spreading dynamics on uncompressed substrates were extracted from 20 flare tracks on 2 videos. (C, D, E, and F) The data for flare spreading dynamics on compressed substrates were extracted from 20 flare tracks from 1 video. The graphs in panels E and F with standard errors of the means are shown in Fig. S2C and D.

placement on uncompressed and compressed substrates ($R^2 = 0.91$), as shown in Fig. S3. Thus, there is a strong relationship between whole-colony surface spreading rates on compressed and uncompressed substrates, a strong relationship between flare displacement on compressed and uncompressed substrates, and a strong relationship between flare speeds on compressed and uncompressed substrates.

DISCUSSION

Previous studies of *M. xanthus* showed that single-cell orientation (8), the movements of groups of cells advancing from colony edges (also called flares) (8, 9), the

FIG 6 *B. cereus* tracking data and analysis. (A, B, E, and F) The data for flare spreading dynamics on uncompressed substrates were extracted from 23 flare tracks on 2 videos. (C, D, E, and F) The data for flare spreading dynamics on compressed substrates were extracted from 24 flare tracks from 2 videos. The graphs in panels E and F with standard errors of the means are shown in Fig. S2E and F.

colony spreading patterns (8, 12, 21), and the distribution of fruiting bodies during development (11) are all affected by substrate compression. Recently, we presented data suggesting that compression-induced changes in flare movements and colony spreading patterns, historically referred to as elasticotaxis (11), are likely due to the altered arrangement of polymer fibers in the substrate (9). Namely, the preferred orientation of flare movements and colony spreading are parallel to the long axes of the polysaccharide fibers, which have become aligned and densely packed due to compression. Therefore, we referred to the response to substrate compression as

FIG 7 *M. xanthus* tracking data and analysis. (A, B, E, and F) The data for flare spreading dynamics on uncompressed substrates were extracted from 22 flare tracks on 4 videos. (C, D, E, and F) The data for flare spreading dynamics on compressed substrates were extracted from 21 flare tracks from 2 videos. The graphs in panels E and F with standard errors of the means are shown in Fig. S2G and H.

polymertropism. We suggested that *M. xanthus* movements are biased after substrate compression because the focal adhesion complex, which helps generate the traction force for A-type gliding motility (10), is able to grip the clusters of aligned polysaccharide fibers more tightly because of their relatively high density.

Here, we observed increases in flare speeds of *M. xanthus* and three other bacterial species when the cells moved in the direction of polysaccharide fiber alignment on compressed substrates (Table 1). These findings are consistent with our proposal that

FIG 8 Dynamics of surface spreading on compressed and uncompressed agar substrates. (A) Images of *M. xanthus* (120 h), *B. cereus* (72 h), *B. mycoides* (30 h), and *P. dendritiformis* (72 h) on compressed agar substrates. Scale bars, 1 cm. (B) Colony aspect ratio (AR) versus time data from each bacterial species spotted on compressed substrates. (C) Colony diameter versus time data from each bacterial species spotted on uncompressed substrates. Lines of best fit for each individual colony have been drawn in panels B and C. The data for each bacterial species in panels B and C are differentiated by color. For each species, the slopes of the best fit lines for aspect ratio versus time and colony diameter versus time were pooled, and the means were calculated. (D) The mean values for surface spreading on uncompressed agar substrates and compressed agar substrates. (E) Each species' mean surface spreading ($\pm$ standard error of the mean) on compressed substrates was graphed as a function of spreading on uncompressed substrates.

the *M. xanthus* A-motility machinery is more efficient when the cells move on fibers that are aligned and densely packed and suggest that the flagellar motility systems of *B. cereus* and *B. subtilis* (two of the bacterial species tested here) are likely to be more efficient when they move on fibers arranged in this way.

B. mycoides flares also moved faster in the direction of polysaccharide fiber alignment, but unlike the other three bacterial species examined here, the mechanism of surface spreading in this bacterium has remained elusive. A few models for surface spreading have been proposed, but they are based on growth and not on the apparatus for surface motility (30, 31). Our data suggest that motility does indeed play roles in *B. mycoides* surface spreading and in the movements in the direction of polysaccharide fiber alignment. Namely, we found that these behaviors continue in the absence of nutrients for growth (see Fig. S1 in the supplemental material).

In addition to moving at greater speeds on compressed substrates (9), the flares persist longer, tend to hold their trajectories parallel to the long axes of the polysaccharide fibers longer, and ultimately show an increase in displacement away from their

origin (Table 1). While it is unclear how these collective changes together affect the dynamics of whole-colony spreading in the direction of polysaccharide fiber alignment, there does appear to be a strong relationship between flare displacement and directed colony spreading on compressed substrates on the basis of the R^2 values of lines of best fit of data plotted for displacement and spreading (Fig. S3 and Table S2). Using the same method of analysis, we found a strong relationship between flare displacement on compressed and uncompressed substrates (Fig. S3 and Table S2), between flare speeds on compressed and uncompressed substrates (Fig. S3 and Table S2), and between whole-colony surface spreading rates on compressed and uncompressed substrates (Fig. 8). These findings are consistent with the idea that the underlying mechanism driving a bacterium's surface spreading on the two substrates is the same, but the efficiencies of surface spreading are different. These findings are also consistent with our model for the polymertropism response (9). Namely, that polymer packing increases the efficiency of surface spreading, while the polymers' alignment dictates the direction of spreading.

Because polymertropism can be explained by a simple model in which the arrangement of the polymers in the substrate promotes more efficient surface spreading in a particular direction, we suggest that polymertropism might not require a mechanism for detecting a specific stimulus and responding to that stimulus. However, polymertropism might require a mechanism for surface motility, as surface motility is known to be required for polymertropism in *M. xanthus*, all of the polymertropism-positive bacteria to date possess surface motility, and we have yet to find a polymertropism-positive bacterium that lacks surface motility (8, 9). We believe that the question of whether all or the vast majority of surface-motile bacteria are polymertropism positive is still open because we have tested only a small selection of bacterial species for this behavior and used a limited number of conditions when we performed these tests (9). As discussed below, we believe that bacteria are likely to encounter aligned and tightly packed polysaccharide fibers in natural settings and, in certain circumstances, can leverage polymertropism to their advantage.

Many bacteria are likely to encounter aligned and densely packed polysaccharide fibers in nature in the form of "slime" trails. In particular, it has been shown that a number of rod-shaped biofilm-forming bacteria secrete exopolysaccharides or slime trails as they move across their substrate's surface (22–27, 32, 33). When these bacteria encounter slime trails, they alter their behavior such that their movements are oriented parallel to the long axes of these trails (34–36). Slime is likely to be composed largely of aligned and densely packed polymers, as slime is thought to be extruded through fine nozzles in the cell surface and similarly extruded materials are inherently aligned (23, 24, 33, 37, 38). Thus, slime-trail following may be a special case of polymertropism.

How might polymertropism on slime trails be leveraged by bacteria? As discussed previously (9), there is experimental and computational data suggesting that the alignment and directed movements of large groups of bacterial cells on slime trails facilitate the formation of cell aggregates and perhaps the formation of highly structured biofilms (36, 39). Therefore, in this case, the aligned and tightly packed polysaccharides would be acting as a long-range organizational signal via polymertropism.

In addition to its potential role in the organization of bacteria, polymertropism on slime trails might confer a fitness advantage in certain situations. For example, predatory bacteria such as *M. xanthus* might benefit from polymertropism when hunting prey bacteria; when they encounter the slime trails produced by prey bacteria, the change in direction and increased speed (Fig. 4 to 7 and Table 1) might help them locate their prey. Polymertropism on preexisting slime trails might also be useful when bacteria are trying to escape from a competitor or predator. For example, when a bacterial biofilm is invaded by a new bacterial species, the original species may be able to utilize its own slime trails to move quickly and directly away from the invading bacterial cells. This example may be the most relevant to this study, as *B. mycoides* and *S. marcescens* were placed in close proximity and in competition with each other, and the ability of *B. mycoides* to move rapidly and directly away from *S. marcescens* by utilizing the aligned

polymers of the compressed agar substrate led to a clear survival benefit (Fig. 1). It is notable that *S. marcescens* benefited from the ability of *B. mycoides* to escape its footprint; this might be due to a decrease in the need for competition. It is also notable that some *B. mycoides* cells that survived the competition assays on compressed substrates yielded colonies with rough morphologies and others yielded colonies with typical s-shaped filamentous morphologies (Fig. 1D). Both morphologies were also observed when *B. mycoides* cells were harvested from monocultures on compressed substrates with various amounts of nutrients or no nutrients (data not shown). These findings suggest that the rough-colony morphology phenotype is not due a chemical produced by *S. marcescens* or by *S. marcescens*' depletion of the nutrient supply. We initially speculated that compression might yield "superspreader" variants that gain access to more nutrients by moving on the surface of the substrate more rapidly than the general population of *B. mycoides* cells, and that these variants produce colonies with rough morphologies. However, cells isolated from the rough colonies and the filamentous colonies spread across the surface at approximately the same rates and have similar polymertropism responses (Fig. S4). Hence, at this point, we do not have specific details as to how the physiology of the *B. mycoides* cells that form rough colonies is altered, although strains of *B. mycoides* that yield similar rough colonies have been reported (30).

A simple, rapid, strong, and easy-to-produce response such as polymertropism may well be ripe for exploitation. In laboratory settings, polymertropism might be a useful means of elucidating the exact nature of propulsion of surface-motile bacteria; speed increases and changes of direction are easily and immediately elicited by simply compressing the substrate. The polymertropism response also alters rippling and fruiting body morphology, *M. xanthus* collective behaviors not discussed in detail here, but which might be exploited to better understand emergent behavior with the help of soft- or active-matter physics (40). Furthermore, there is evidence that substrate polymer alignment affects the behavior of eukaryotic cells (41–43), and similar techniques of substrate compression and manipulation may be useful in studies of the motility of eukaryotic cells and perhaps in organizing cells for tissue engineering as well as for medical device and implant design.

MATERIALS AND METHODS

Bacterial strains and media for growth. Our wild-type strain of *M. xanthus* (DK1622) was grown at 30°C in CTTYE broth (44) containing 10 g/liter Casitone, 2 g/liter yeast extract, 1 mM KH_2PO_4 , 10 mM Tris-HCl (pH 8.0), and 8 mM MgSO_4 or on plates containing CTTYE broth and 1.5% agar. *B. cereus* (154870A; Carolina Biological Supply), *B. subtilis* (NRRL B-354; USDA Agriculture Research Service), *Paenibacillus dendritiformis* (NRRL B-23299), and *S. marcescens* (NRRL B-23389) were also grown at 30°C in CTTYE broth or on CTTYE plates. *B. mycoides* (NRRL B-14811) was grown at 30°C in CTTYE broth, 75% CTTYE broth (7.5 g/liter Casitone, 1.5 g/liter yeast extract, 1 mM KH_2PO_4 , 10 mM Tris-HCl [pH 8.0], and 8 mM MgSO_4), 50% CTTYE broth (5 g/liter Casitone, 1 g/liter yeast extract, 1 mM KH_2PO_4 , 10 mM Tris-HCl [pH 8.0], and 8 mM MgSO_4), 25% CTTYE broth (2.5 g/liter Casitone, 0.5 g/liter yeast, 1 mM KH_2PO_4 , 10 mM Tris-HCl [pH 8.0], and 8 mM MgSO_4), or placed in TPM buffer (1 mM KH_2PO_4 , 10 mM Tris-HCl [pH 8.0], and 8 mM MgSO_4) and incubated at 30°C. *B. mycoides* was also grown at 30°C on plates containing CTTYE, 75% CTTYE, 50% CTTYE, or 25% CTTYE broth and 1.5% agar or placed on plates containing TPM buffer and 1.5% agar.

Mixed-culture assays. *B. mycoides* and *S. marcescens* were individually grown at 30°C in CTTYE broth to densities of approximately 5×10^8 cells/ml. The cells were pelleted by centrifugation, the supernatants were removed, and the cell pellets were resuspended in CTTYE broth to a final concentration 1×10^8 cells/ml. *B. mycoides* and *S. marcescens* cells were mixed together at ratios of 3:1, 1:1, and 1:3 and spotted on plates containing uncompressed CTTYE agar or on plates containing CTTYE agar compressed with 5.56-mm Tygon tubing (9). The spots were allowed to dry and the plates were incubated at 30°C. After 24 h of incubation, the colonies were examined visually (see Fig. 1), harvested using a spatula, and placed into 1 ml of CTTYE broth. CTTYE broth containing the various mixtures of *B. mycoides* and *S. marcescens* cells was briefly sonicated, diluted, spread onto plates containing uncompressed CTTYE agar, and incubated at 30°C. After 12 to 18 h of incubation, the number of white colonies produced by *B. mycoides* and the number of red colonies produced by *S. marcescens* were determined. Two sample *t* tests were used to detect significant differences in mean CFU. A stereomicroscope, 3.0-megapixel AmScope USB ocular camera, and Touptview image capture software were used to obtain mixed-culture colony still images. Similar experiments were performed with monocultures of *B. mycoides* or *S. marcescens* to show that neither species derived a benefit from growing on compressed CTTYE agar (see Table S1 in the supplemental material).

Aspect ratio versus time and colony diameter versus time assays. Bacterial cells were grown at 30°C in CTTYE broth to a density of approximately 5×10^8 cells/ml. The cells were pelleted using centrifugation, the supernatants were removed, and the cell pellets were resuspended in CTTYE broth to a final concentration of 1×10^8 cells/ml. Four-microliter aliquots of the concentrated cells were spotted down the centerlines of square plates containing compressed or uncompressed CTTYE agar. The spots were allowed to dry, and the plates were incubated at 30°C for 24 h. The aspect ratios of colonies (i.e., colony width divided by the colony height) on compressed CTTYE agar and the diameters of colonies on uncompressed CTTYE agar were determined after 24 h of incubation and at regular intervals thereafter. For each colony, we plotted the aspect ratio as a function of time and calculated the slope of the line of best fit, which provided the mean rate of increase in aspect ratios over time. The mean polymertropism rate of each bacterial species on compressed agar, which we defined as the mean rate of aspect ratio increase over time, was calculated from 5 to 10 slope values for 5 to 10 colonies. The mean surface spreading rate of each bacterial species on uncompressed agar was calculated using a similar strategy; in this case, increases in colony diameters over time were used in the calculations instead of increases in aspect ratios over time. It should be noted that *P. dendritiformis* was substituted for *B. subtilis* in these studies, as the changes in the aspect ratios of *B. subtilis* colonies (but not flares) varied widely on compressed agar substrates.

Analysis of *B. mycoides* surface spreading, polymertropism, and growth. To assay surface spreading, uncompressed 1.5% agar plates containing TPM buffer, 25% CTTYE broth, 50% CTTYE broth, 75% CTTYE broth, or CTTYE broth were inoculated with 4- μ l drops of 1×10^8 cells/ml *B. mycoides*, and the diameter of each colony at its widest point was measured periodically. For each colony, the slope of a line of best fit for colony diameter versus time was calculated, and these values were pooled to generate mean surface spreading rates as described above. Polymertropism assays were performed as described above on compressed 1.5% agar plates containing TPM buffer, 25% CTTYE broth, 50% CTTYE broth, 75% CTTYE broth, or CTTYE broth. To assay growth, 4- μ l drops of 1×10^8 cells/ml *B. mycoides* cells growing in liquid CTTYE were inoculated into three 12-ml replicates each of 25% CTTYE broth, 50% CTTYE broth, 75% CTTYE broth, or CTTYE broth, while three 12-ml replicates of TPM buffer cultures were started at 20 Klett units to ensure that a lack of growth could be adequately detected. The liquid cultures were incubated in flasks at 30°C with vigorous shaking, and the cell density was determined at various time points. The readings taken between 8 and 12 h, when the growth over time was nearly linear, were used to calculate the growth rate for each culture. A line of best fit was applied to each culture's function of cell density versus time, and the slope of the line was used to indicate the growth rate of the cells. The slope values for each distinct medium composition were pooled and means calculated to determine the growth rate under those conditions.

Images of the *B. mycoides* polymertropism response. CTTYE broth containing 1.5% agar was poured on round plates, allowed to solidify, and compressed using 1.5-cm pieces of Tygon tubing with diameters of 3.18, 3.96, 4.76, 5.56, or 6.35 mm. The plates were placed at 30°C to allow the water that was squeezed out of the agar by compression to evaporate, and 4- μ l spots of overnight *B. mycoides* liquid cultures were placed down the centerline of the compressed agar using a multichannel pipettor. After the spots of *B. mycoides* dried, the plates were incubated for 48 h at 30°C and images were taken using a stereomicroscope, 3.0-megapixel AmScope USB ocular camera, and Toupview image capture software.

Microscopy, time-lapse imaging, and analysis of images. Bacterial cells were grown and concentrated as described above, and 4- μ l aliquots of the concentrated cells were spotted onto compressed and uncompressed CTTYE agar. After the spots dried, the agar plates were placed in a 30°C incubator, and time-lapse videos of flares or groups of cells moving out from the colony edges were obtained using an Olympus CK2 inverted microscope with 3.0-megapixel AmScope USB ocular camera and Toupview image capture software. One image was captured each minute for 1,440 min. Flares were tracked using ImageJ's (28) manual tracking plug-in from their initial emergence from the colony edges until they merged with another flare, merged with the colony, advanced past the field of view, or the time-lapse video ended. After the tracking was completed, the *x* and *y* coordinates of the flares in each image were obtained and analyzed as described below.

From each flare tip that was tracked, the initial coordinate (*x,y*) values (Fig. 3A) were subtracted from the raw (*x,y*) values such that each track began at (0,0). These adjusted *x* or *y* values were then multiplied by -1 as needed, such that each track was predominantly located in quadrant I (Fig. 3B). We called these new (*x,y*) values which started at (0,0) and spread into quadrant I relative (*x,y*) (Fig. 3B). Using the relative (*x,y*) values of all the tracked flares of a bacterial species, we were able to calculate the mean relative (*x,y*) values of the species' flares at each time point in the tracks on compressed and uncompressed agar (Fig. 3C). These calculations revealed the mean shape or trajectory of each species' flares on compressed and uncompressed agar. Flare displacement was defined as the straight-line distance between a flare tip's starting and ending positions on a video (Fig. 3D). Using calculations similar to those described above, we were able to determine the mean final displacement of a species' flares. A flare's duration was inferred after counting the 1-min intervals that could be tracked. The mean flare duration was determined using all of each species' flares on compressed or uncompressed agar substrates. For each bacterial species, the distance that each flare tip traveled between each pair of successive images in a video was calculated (Fig. 3D) and used to determine the mean distance that flares of that species traveled in 1 min, which we refer to as the mean flare speed. The mean flare speeds on compressed and uncompressed agar were determined.

SUPPLEMENTAL MATERIAL

Supplemental material for this article may be found at <https://doi.org/10.1128/JB.00610-17>.

SUPPLEMENTAL FILE 1, AVI file, 1.8 MB.

SUPPLEMENTAL FILE 2, AVI file, 2.2 MB.

SUPPLEMENTAL FILE 3, AVI file, 2.9 MB.

SUPPLEMENTAL FILE 4, AVI file, 3.7 MB.

SUPPLEMENTAL FILE 5, PDF file, 1.7 MB.

ACKNOWLEDGMENTS

This work was supported by NSF grant IOS-1354779 (A.G.G.) and a fellowship from NSF IGERT grant DGE-1068780 (D.J.L.).

REFERENCES

- Skotnicka D, Smaldone GT, Petters T, Trampari E, Liang J, Kaever V, Malone JG, Singer M, Sogaard-Andersen L. 2016. A minimal threshold of c-di-GMP is essential for fruiting body formation and sporulation in *Myxococcus xanthus*. *PLoS Genet* 12:e1006080. <https://doi.org/10.1371/journal.pgen.1006080>.
- López D, Vlamakis H, Losick R, Kolter R. 2009. Paracrine signaling in a bacterium. *Genes Dev* 23:1631–1638. <https://doi.org/10.1101/gad.1813709>.
- Waters CM, Bassler BL. 2005. Quorum sensing: cell-to-cell communication in bacteria. *Annu Rev Cell Dev Biol* 21:319–346. <https://doi.org/10.1146/annurev.cellbio.21.012704.131001>.
- White D, Shropshire W, Jr, Stephens K. 1980. Photocontrol of development by *Stigmatella aurantiaca*. *J Bacteriol* 142:1023–1024.
- Inouye T, White D, Inouye M. 1980. Development of *Stigmatella aurantiaca*: effects of light and gene expression. *J Bacteriol* 141:1360–1365.
- Qualls GT, Stephens K, White D. 1978. Light-stimulated morphogenesis in the fruiting myxobacterium *Stigmatella aurantiaca*. *Science* 201:444–445. <https://doi.org/10.1126/science.96528>.
- Meel C, Kouzel N, Oldewurtel ER, Maier B. 2012. Three-dimensional obstacles for bacterial surface motility. *Small* 8:530–534. <https://doi.org/10.1002/sml.201101362>.
- Fontes M, Kaiser D. 1999. *Myxococcus* cells respond to elastic forces in their substrate. *Proc Natl Acad Sci U S A* 96:8052–8057. <https://doi.org/10.1073/pnas.96.14.8052>.
- Lemon DJ, Yang X, Srivastava P, Luk YY, Garza AG. 2017. Polymertropism of rod-shaped bacteria: movement along aligned polysaccharide fibers. *Sci Rep* 7:7643. <https://doi.org/10.1038/s41598-017-07486-0>.
- Faure L, Fiche J-B, Espinosa L, Ducret A, Anantharaman V, Luciano J, Lhospice S, Islam ST, Treguire J, Sotes M, Kuru E, Van Nieuwenhze MS, Brun YV, Theodoly O, Aravind L, Nollman M, Mignot T. 2016. The mechanism of force transmission at bacterial focal adhesion complexes. *Nature* 539:530–535. <https://doi.org/10.1038/nature20121>.
- Stanier RY. 1942. A note on elasticotaxis in myxobacteria. *J Bacteriol* 44:405–412.
- Stratford JP, Woodley MA, Park S. 2013. Variation in the morphology of *Bacillus mycoides* due to applied force and substrate structure. *PLoS One* 8:e81549. <https://doi.org/10.1371/journal.pone.0081549>.
- Dworkin M. 1983. Tactic behavior of *Myxococcus xanthus*. *J Bacteriol* 154:452–459.
- Shi W, Zusman DR. 1993. The two motility systems of *Myxococcus xanthus* show different selective advantages on various surfaces. *Proc Natl Acad Sci U S A* 90:3378–3382. <https://doi.org/10.1073/pnas.90.8.3378>.
- Mignot T, Shaevitz JW, Hartzell PL, Zusman DR. 2007. Evidence that focal adhesion complexes power bacterial gliding motility. *Science* 315:853–856. <https://doi.org/10.1126/science.1137223>.
- Islam ST, Mignot T. 2015. The mysterious nature of bacterial surface (gliding) motility: a focal adhesion-based mechanism in *Myxococcus xanthus*. *Semin Cell Dev Biol* 46:143–154. <https://doi.org/10.1016/j.semcdb.2015.10.033>.
- Calvio C, Celandroni F, Ghelardi E, Amati G, Salvetti S, Cecilian F, Galizzi A, Senesi S. 2005. Swarming differentiation and swimming motility in *Bacillus subtilis* are controlled by swrA, a newly identified dicistronic operon. *J Bacteriol* 187:5356–5366. <https://doi.org/10.1128/JB.187.15.5356-5366.2005>.
- Houry A, Briandet R, Aymerich S, Gohar M. 2010. Involvement of motility and flagella in *Bacillus cereus* biofilm formation. *Microbiology* 156:1009–1018. <https://doi.org/10.1099/mic.0.034827-0>.
- Kearns DB, Losick R. 2003. Swarming motility in undomesticated *Bacillus subtilis*. *Mol Microbiol* 49:581–590. <https://doi.org/10.1046/j.1365-2958.2003.03584.x>.
- Senesi S, Ghelardi E, Celandroni F, Salvetti S, Parisio E, Galizzi A. 2004. Surface-associated flagellum formation and swarming differentiation in *Bacillus subtilis* are controlled by the ifm locus. *J Bacteriol* 186:1158–1164. <https://doi.org/10.1128/JB.186.4.1158-1164.2004>.
- Polka JK, Silver PA. 2014. Induced sensitivity of *Bacillus subtilis* colony morphology to mechanical media compression. *PeerJ* 2:e597. <https://doi.org/10.7717/peerj.597>.
- Pate JL. 1985. Gliding motility in *Cytophaga*. *Microbiol Sci* 2:289–290.
- Hoiczky E, Baumeister W. 1998. The junctional pore complex, a prokaryotic secretion organelle, is the molecular motor underlying gliding motility in cyanobacteria. *Curr Biol* 8:1161–1168. [https://doi.org/10.1016/S0960-9822\(07\)00487-3](https://doi.org/10.1016/S0960-9822(07)00487-3).
- Wolgemuth CW, Hoiczky E, Kaiser D, Oster G. 2002. How myxobacteria glide. *Curr Biol* 12:369–377. [https://doi.org/10.1016/S0960-9822\(02\)00716-9](https://doi.org/10.1016/S0960-9822(02)00716-9).
- Yu R, Kaiser D. 2007. Gliding motility and polarized slime secretion. *Mol Microbiol* 63:454–467. <https://doi.org/10.1111/j.1365-2958.2006.05536.x>.
- Ducret A, Valignat M-P, Mouhamar F, Mignot T, Theodoly O. 2012. Wet-surface-enhanced ellipsometric contrast microscopy identifies slime as a major adhesion factor during bacterial surface motility. *Proc Natl Acad Sci U S A* 109:10036–10041. <https://doi.org/10.1073/pnas.1120979109>.
- Berleman JE, Zemla M, Remis JP, Liu H, Davis AE, Worth AN, West Z, Zhang A, Park H, Bosneaga E, van Leer B, Tsai W, Zusman DR, Auer M. 2016. Exopolysaccharide microchannels direct bacterial motility and organize multicellular behavior. *ISME J* 10:2620–2632. <https://doi.org/10.1038/ismej.2016.60>.
- Rasband W. 1997–2016. ImageJ, U.S. National Institutes of Health, Bethesda, MD.
- Be'er A, Strain SK, Hernandez RA, Ben Jacob E, Florin EL. 2013. Periodic reversals in *Paenibacillus dendritiformis* swarming. *J Bacteriol* 195:2709–2717. <https://doi.org/10.1128/JB.00080-13>.
- Di Franco C, Beccari E, Santini T, Pisaneschi G, Tecce G. 2002. Colony shape as a genetic trait in the pattern-forming *Bacillus mycoides*. *BMC Microbiol* 2:33. <https://doi.org/10.1186/1471-2180-2-33>.
- Gause G. 1939. Some physiological properties of dextral and of sinistral forms in *Bacillus mycoides* flügge. *Biol Bull* 76:448–465. <https://doi.org/10.2307/1537751>.
- Shimkets LJ. 1986. Correlation of energy-dependent cell cohesion with social motility in *Myxococcus xanthus*. *J Bacteriol* 166:837–841. <https://doi.org/10.1128/jb.166.3.837-841.1986>.
- Kaiser D. 2009. Are there lateral as well as polar engines for A-motile gliding in myxobacteria? *J Bacteriol* 191:5336–5341. <https://doi.org/10.1128/JB.00486-09>.
- Burchard RP. 1982. Trail following by gliding bacteria. *J Bacteriol* 152:495–501.
- Kranz WT, Gelimson A, Zhao K, Wong GC, Golestanian R. 2016. Effective dynamics of microorganisms that interact with their own trail. *Phys Rev Lett* 117:038101. <https://doi.org/10.1103/PhysRevLett.117.038101>.

36. Zhao K, Tseng BS, Beckerman B, Jin F, Gibiansky ML, Harrison JJ, Luijten E, Parsek MR, Wong GC. 2013. Psl trails guide exploration and micro-colony formation in *Pseudomonas aeruginosa* biofilms. *Nature* 497: 388–391. <https://doi.org/10.1038/nature12155>.
37. Tan J, Rouse SL, Li D, Pye VE, Vogeley L, Brinth AR, El Arnaout T, Whitney JC, Howell PL, Sansom MS, Caffrey M. 2014. A conformational landscape for alginate secretion across the outer membrane of *Pseudomonas aeruginosa*. *Acta Crystallogr D Biol Crystallogr* 70:2054–2068. <https://doi.org/10.1107/S1399004714001850>.
38. Wang Y, Andole Pannuri A, Ni D, Zhou H, Cao X, Lu X, Romeo T, Huang Y. 2016. Structural basis for translocation of a biofilm-supporting exopolysaccharide across the bacterial outer membrane. *J Biol Chem* 291: 10046–10057. <https://doi.org/10.1074/jbc.M115.711762>.
39. Balagam R, Igoshin OA. 2015. Mechanism for collective cell alignment in *Myxococcus xanthus* bacteria. *PLoS Comput Biol* 11:e1004474. <https://doi.org/10.1371/journal.pcbi.1004474>.
40. Marchetti M, Joanny J-F, Ramaswamy S, Liverpool T, Prost J, Rao M, Simha RA. 2013. Hydrodynamics of soft active matter. *Rev Mod Phys* 85:1143–1189. <https://doi.org/10.1103/RevModPhys.85.1143>.
41. Harris AK, Stopak D, Wild P. 1981. Fibroblast traction as a mechanism for collagen morphogenesis. *Nature* 290:249–251. <https://doi.org/10.1038/290249a0>.
42. Miron-Mendoza M, Seemann J, Grinnel F. 2008. Collagen fibril flow and tissue translocation coupled to fibroblast migration in 3d collagen matrices. *Mol Biol Cell* 19:2051–2058. <https://doi.org/10.1091/mbc.E07-09-0930>.
43. Feng Z, Wagatsuma Y, Kobayashi S, Kosawada T, Sato D, Nakamura T, Kitajima T, Umez M. 2013. Analysis of the contraction of fibroblast collagen gels and the traction force of individual cells by a novel elementary structural model. *Conf Proc IEEE Eng Med Biol Soc* 2013: 6232–6235. <https://doi.org/10.1109/EMBC.2013.6610977>.
44. Giglio KM, Zhu C, Klunder C, Kummer S, Garza AG. 2015. The enhancer binding protein Nla6 regulates developmental genes that are important for *Myxococcus xanthus* sporulation. *J Bacteriol* 197:1276–1287. <https://doi.org/10.1128/JB.02408-14>.