Sublingual microcirculation reveals fluid overload and leukocytosis in a post-cardiac surgery patient

Zühre Uz,¹ Bastianus AJM de Mol,² Thomas M van Gulik,¹ Can Ince³

¹Department of Surgery, AMC, Amsterdam, North Holland, The Netherlands

²Department of Cardio-Thoracic Surgery, AMC, Amsterdam, North Holland, The Netherlands

³Department of Translational Physiology, AMC, Amsterdam, North Holland, The Netherlands

Correspondence to

Professor Thomas M van Gulik, t.m.vangulik@amc.nl

Accepted 30 January 2018

DESCRIPTION

Positive cumulative fluid balance caused by excessive fluid administration during the perioperative phase is known as one of the most common clinical complications in post-intensive care unit cardiac surgery patients.¹ Mostly, administration of diuretics is initiated to correct the fluid overload. However, there is no clinical parameter showing the changes in the fluid balance, while daily measurements of weight and fluid balance are initiated to help to manage diuretic therapy. Herein, we present a case report, in which we show that sublingual microcirculation monitoring (SMiMo) may provide an additional check on optimal filling status in these patients apart from recording fluid balance and weight control.

A 75-year-old woman with a history of B cell chronic lymphocytic leukaemia diagnosed with aortic valve stenosis was admitted for transcatheter aortic valve replacement (TAVR). Patient was discharged from the intensive care unit with a cumulative positive fluid balance of 1800 cm³ and transferred to the cardiac ward. The patient's body weight increased after surgery and was followed by a gradual decrease in the subsequent days (preoperative weight: 64.5 kg, 2 days after surgery: 66.7 kg, 3 and 4 days after surgery: 65.2 and 64.7 kg, respectively). Patients with a severe aortic stenosis suffer from left ventricular hypertrophy requiring careful fluid management.

Therefore, the patient received oral administration of 40 mg furosemide and 25 mg spironolactone per day for the following 3 days to correct fluid overload.

Fluid overload and its microcirculatory impact secondary to cardiac surgery was assessed by SMiMo, using incident dark-field imaging (CytoCam-IDF, Braedius, Huizen, The Netherlands). The CytoCam is a handheld microscope emitting green light with a 548 nm wavelength. The green light ensures optimal absorption by oxyhaemoglobin and deoxyhaemoglobin, through which the red blood cells become visible.²

Images were recorded on day 1 (T0) (figure 1) and day 3 (T1) (figure 2) after TAVR, respectively, showing a substantial increase in vessel density. This significant change can be assessed visually at the same anatomical sublingual site. This is the first report showing a visual effect of diuretic therapy on peripheral microcirculation in a fluid overloaded patient. SMiMo allows comparison of both images without offline analysis, which is usually required. In addition, leukocytosis was also observed at both time points. Subsequent decrease in leucocyte count detected in blood samples ($16.3 \times 10^9/L$ at T0 decreased to $14.5 \times 10^9/L$ at T1) could also be observed on CytoCam images at the same microvascular site.

Offline analysis using dedicated software (Automated Vascular Analysis, AVA, Microvision Medical, Amsterdam, The Netherlands)³ showed an increase in microcirculatory density parameters (total vessel density and perfused vessel density). Microcirculatory flow parameters (proportion of perfused vessels and microvascular mean flow index) showed a minor change indicating a stable flow (see legends of figures 1 and 2).

Figure 1 Sublingual microcirculation 1 day after surgery. Screenshot of video recorded on day 1 (T0) after TAVR. Arrowheads show decreased diameter of vessels shown in figure 2. Crosses indicate disappeared vessels which can be seen in figure 2. Black arrows indicate leukocytes sticking and rolling in a postcapillary venule. Microcirculatory parameters: total vessel density 22.6 mm/mm², perfused vessel density 21.6 mm/mm², proportion of perfused vessels 96.0% and microvascular mean flow index 3.0.

Figure 2 Sublingual microcirculation 3 days after surgery. Screenshot of video recorded 3 days (T1) after TAVR. This is the first time that a patient follow-up of the sublingual microcirculation is obtained at one and the same microvascular bed. Compared with figure 1, the vessel density has increased, and black arrows indicate leukocytes that are sticking and rolling in a postcapillary venule. Microcirculatory parameters: total vessel density 32.2 mm/mm², perfused vessel density 31.36 mm/mm², proportion of perfused vessels 97.3% and microvascular mean flow index 2.9.

To cite: Uz Z, de Mol BAJM, van Gulik TM, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-223681

Learning points

- ▶ Fluid overload is a common complication after cardiac surgery caused by haemodynamic resuscitation performed during the surgical procedure.
- ▶ Patients with a severe aortic stenosis suffer from left ventricular hypertrophy requiring careful fluid management.
- ▶ Sublingual microcirculation monitoring may provide an additional check on optimal filling status in critically ill patients apart from recording fluid balance and weight control.

The patient was discharged from the cardiac surgery ward on postoperative day 5 once a normal weight and fluid balance were obtained.

Contributors ZU participated in the design of the case report, data collection (sublingual microcirculation measurements), the Automated Vascular Analysis of the sublingual microcirculation measurements, interpreted, drafted and revised the

manuscript. CI and TvG helped to draft and revise the manuscript. BAdM participated in the design of the case report, contributed by providing the facilities and drafted and revised the manuscript.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Haase-Fielitz A, Haase M, Bellomo R, *et al.* Perioperative Hemodynamic Instability and Fluid Overload are Associated with Increasing Acute Kidney Injury Severity and Worse Outcome after Cardiac Surgery. *Blood Purif* 2017;43:298–308.
- 2 van Elteren HA, Ince C, Tibboel D, *et al.* Cutaneous microcirculation in preterm neonates: comparison between sidestream dark field (SDF) and incident dark field (IDF) imaging. *J Clin Monit Comput* 2015;29:543–8.
- 3 Dobbe JG, Streekstra GJ, Atasever B, *et al.* Measurement of functional microcirculatory geometry and velocity distributions using automated image analysis. *Med Biol Eng Comput* 2008;46:659–70.

Copyright 2018 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow