Published in final edited form as:

Crit Care Med. 2018 April ; 46(4): 500–505. doi:10.1097/CCM.0000000000002949.

Delay within the 3-Hour Surviving Sepsis Campaign Guideline on Mortality for Patients with Severe Sepsis and Septic Shock

Lisiane Pruinelli, PhD, RN¹, Bonnie L. Westra, PhD, RN, FAAN, FACMI^{1,2}, Pranjul Yadav, PhD³, Alexander Hoff³, Michael Steinbach, PhD³, Vipin Kumar, PhD³, Connie W. Delaney, PhD, RN, FAAN, FACMI^{1,2}, and Gyorgy Simon, PhD^{2,4}

¹University of Minnesota, School of Nursing. 308 Harvard St SE, WDH 5-140, Minneapolis, MN 55455

²University of Minnesota, Institute for Health Informatics. MMC 912 Mayo, 420 Delaware St. SE, Minneapolis, MN 55455

³University of Minnesota, Department of Computer Science and Engineering. Room 4-192 EE/CSci, 200 Union Street SE, Minneapolis, MN 55455

⁴University of Minnesota, School of Medicine. Mayo Building, 420 Delaware Street SE, Minneapolis, MN 55455

Abstract

Objective—To specify when delays of specific 3-hour bundle Surviving Sepsis Campaign (SSC) guideline recommendations applied to severe sepsis or septic shock become harmful and impact mortality.

Design—Retrospective cohort study.

Setting—One health system composed of six hospitals and 45 clinics in a Midwest state from January 01, 2011 and July 31, 2015.

Patients—All adult patients hospitalized with billing diagnosis of severe sepsis or septic shock.

Interventions—Four 3-hour SSC guideline recommendations: 1) obtain blood culture before antibiotics, 2) obtain lactate level, 3) administer broad-spectrum antibiotics, and 4) administer 30 mL/kg of crystalloid fluid for hypotension (defined as mean arterial pressure (MAP) < 65) or lactate (> 4).

Measurements and Main Results—To determine the effect of t minutes of delay in carrying out each intervention, propensity score matching of *baseline* characteristics compensated for

Corresponding author: Lisiane Pruinelli, PhD, MS, RN. pruin001@umn.edu. 6-183 Weaver-Densford Hall, 308 Harvard Street SE, Minneapolis, MN 55455. Phone: 612-626-5144. Fax: 612-625-7091.

Conflict of Interest: There is no conflict of interest to report for any of the authors.

Copyright form disclosure: Drs. Westra, Kumar, and Simon received support for article research from the National Institutes of Health (NIH). Dr. Westra disclosed that this study is supported by National Science Foundation (NSF) grant IIS-1344135 and partially supported by Grant Number 1UL1RR033183 from the National Center for Research Resources (NCRR) of the NIH to the University of Minnesota Clinical and Translational Science Institute (CTSI). Dr. Steinbach disclosed that he is supported in part by NSF and NIH grants and by the University of Minnesota, although no grants were received for this specific work. For this paper, he received NSF related funding. The remaining authors have disclosed that they do not have any potential conflicts of interest.

differences in health status. The Average Treatment effect in the Treated (ATT) computed as the average difference in outcomes between those treated after shorter versus longer delay. To estimate the uncertainty associated with the ATT metric and to construct 95% confidence intervals, bootstrap estimation with 1,000 replications was performed. From 5,072 patients with severe sepsis or septic shock, 1,412 (27.8%) had in-hospital mortality. The majority of patients had the four 3-hour bundle recommendations initiated within three hours. The statistically significant time in minutes after which a delay increased the risk of death for each recommendation was: lactate, 20.0 minutes; blood culture, 50.0 minutes; crystalloids, 100.0 minutes; and antibiotic therapy, 125.0 minutes.

Conclusions—The guideline recommendations showed that shorter delays indicates better outcomes. There was no evidence that 3 hours is safe; even very short delays adversely impact outcomes. Findings demonstrated a new approach to incorporate time *t* when analyzing the impact on outcomes and provide new evidence for clinical practice and research.

Keywords

Severe sepsis; Septic shock; Surviving Sepsis Campaign guideline; Predictive modeling; Mortality; Electronic Health Record data

Introduction

Sepsis is a major health problem with increasing prevalence, high costs, and poor outcomes. In the United States, sepsis accounts for about 5.2% of hospital expenditures, more than \$20 billion per year (1), and increased hospitalizations for more than 1 million people in 2008 (2). Between 1999 and 2014, there were 2,470,666 deaths with sepsis on the death certificate and annual sepsis related deaths increased by 31% (2, 3). Severe sepsis and septic shock are a subset of sepsis conditions with at least one organ dysfunction (2). This subset of conditions accounts for 25–50% of in-hospital mortality (2), ranging from 45% to 65% after six months of discharge. If patients with these conditions survive, they have increasing chronic complications, morbidity, high costs of care, and decreasing quality of life (4, 5). Due to the seriousness of these outcomes, the focus of this study is on severe sepsis and septic shock.

The Surviving Sepsis Campaign (SSC) international guidelines (6) were developed for early detection and treatment of severe sepsis and septic shock. While there is agreement that applying the SSC guidelines improves outcomes, there are several challenges including varying strength of evidence associated with the recommendations (6). Since 2001, the definitions and guidelines for sepsis have continuously evolved (6). In 2004 there were 46 recommendations and pediatric management considerations which were subsequently divided into 6 and 24 hour bundles (7). The guidelines were revised in 2008, 2012 (6), and most recently updated in 2015 (8). The recommendations that repeatedly proved effective were simplified and grouped into 3- and 6-hour bundles. The next question is whether there is more specific timing that would improve outcomes. The 3-hour recommendations, which must be carried out within 3 hours from the first time sepsis is suspected, are: 1) obtain a blood culture before antibiotics, 2) obtain a lactate level, 3) administer broad-spectrum

antibiotics, and 4) administer 30 mL/kg of crystalloid fluid for hypotension (defined as a mean arterial pressure (MAP) < 65) or lactate (> 4).

Previous studies examined the effect of the SSC guideline on various outcomes, such as hospital and intensive care unit length of stays, discharge disposition, hospital readmission and mortality (5, 9–12). There are mixed results regarding a decrease in mortality, which is likely related to outcome definitions, types of interventions evaluated, population heterogeneity, and timing (13–21). It is unclear if the timing of specific SSC guideline recommendations within the 3-hour bundle improves the probability of survival. The availability of electronic health record (EHR) data and newer machine learning analytics may provide new insights for disease management and clinical improvement. Sepsis studies using EHR data have focused on trends in sepsis (22) and early identification of sepsis and/or the impact on outcomes (23–27). A few studies used machine learning, a family of techniques that use artificial intelligence to learn from large collections of data, to develop predictive models for sepsis outcomes (28–31). None focused on evaluating the timing of SSC guideline recommendations within the 3-hour bundle to determine if a delay in these recommendations results in harm. We previously used machine learning and found the SSC guidelines to be effective on outcomes of mortality and complications (32); however, we did not look at the timing within the 3-hour bundle to determine if there is a harmful effect in less than 3 hours. The purpose of this study was to specify the time frame when the delay of specific 3-hour bundle guideline recommendations applied to severe sepsis or septic shock becomes harmful and impacts mortality.

Materials and Methods

Study Design and Data Selection

A retrospective cohort study was conducted using data from all adult patients who were hospitalized with a billing diagnosis of severe sepsis or septic shock between January 01, 2011 and July 31, 2015. After IRB approval (#1311E46042), a de-identified data set was obtained from a clinical data repository which contained over 2.4 million patients. The inclusion criteria were patients 18 years and older, both medical and surgical patients, who were hospitalized with a billing diagnosis of severe sepsis or septic shock (ICD-9 codes 995.92 and 785.5* excluding 785.51). The initial data set contained 5,374 patients. The first hospitalization was selected as the index encounter. Exclusion criteria were: did not meet two of the criteria for sepsis suspicion (n=114), or no data for antibiotic therapy (n=142), respiratory rate (n=3) or white blood cells count (WBC) (n=43). This resulted in a final sample of 5,072 patients. The onset of sepsis (baseline) was estimated as the earliest date and time when the patient met at least two of the following six criteria: MAP < 65, heart rate (HR) >100, respiratory rate (RR) >20, temperature < 95 or >100.94, WBC < 4 or > 12, and lactate > 2.0.(33)

Variables of Interest

Variables of interest included predictors, *delay* in administering guideline recommendations, and outcomes. The 16 predictor variables were: age, gender, race, ethnicity, lactate, WBC, HR, RR, temperature, MAP, Charlson Comorbidity Index (CCI), and five comorbidity

severity scores, which describe the severity of preexisting diseases in each of the following five health categories: respiratory, cardiac, liver, kidney, and vascular system. Using domain experts, comorbidity severity scores were computed as the weighted sum of the diagnoses, identified from the problem list and billing diagnoses prior to the index encounter, where the weights were assigned to reflect severity and acuteness: 0= chronic, 1= mild, 2= moderate to severe. These weights were summed for each comorbidity severity score and each patient for analysis, resulting in a continuous scale. *Delay* of obtaining a lactate level and blood culture or administering a broad-spectrum antibiotic was measured from baseline to the time these actions occurred (up to six hours); for the recommendation of ‘administer crystalloids when lactate is > 4 mmol/L’, *delay* is the time between discovering that lactate is high (time of the laboratory result) and the time of administering crystalloids. The quantity of interest is the average causal effect of *delay* on all-cause in-hospital mortality. The outcome is mortality occurring at any time during that hospital stay.

Data Analysis

The purpose of the analysis was to determine the effect of t minutes of delay in carrying out a recommendation, for any amount of delay t between 15 and 360 minutes. This effect was defined as the Average Treatment effect in the Treated (ATT) and is interpreted as the reduction in the probability of in-hospital mortality as a result of carrying out a particular guideline recommendation with less than t minutes of delay. We performed sequential propensity score matching (PSM) at $t = 15, 30, \dots, 360$ minutes with the “exposed” group consisting of patients who received a particular guideline recommendation in less than t minutes, versus the “control” group who received the recommendation with more than t minutes of delay. Patients who did not receive the guideline recommended treatment before discharge were excluded. The sequential PSM procedure resulted in 24 point estimates of ATT at 24 discrete values of delay. We applied a local polynomial smoother (loess) to the 24 point estimates to obtain a smooth ATT curve from which the effect of delay can be determined for any t between 15 minutes and 6 hours in a non-parametric fashion. The same procedure was repeated for all four guideline recommendations.

The predictors of the propensity score model included predictors of the delay as well as predictors of the outcome. These were age, sex, the baseline lab results and vitals, and the severity score of the pre-existing comorbidities. The MatchIt R package (version 3.0.1) was used for propensity matching, using nearest neighbor matching with a caliper of .25 standard deviations. We used the paired t -test as well as the standardized mean difference (SMD) between the exposed and control groups to assess the achieved balance.⁽³⁴⁾ A p -value of .05 and a SMD of less than .2 were required to achieve significance. All 96 propensity matched populations (24 for each of the 4 recommendations) had a SMD of less than .2, but some matched populations (for Crystalloid beyond 3 hours delay) failed the t -test. Furthermore, this did not impact the ATT, because the SMD was still less than 0.2. In the matched population, ATT was computed as the difference in the proportions of patients who died in the hospital between the exposed and control groups.

The 95% confidence interval for the ATT curve was obtained through bootstrap simulation with 1,000 replications. Continuous variables are reported as mean and interquartile range

(IQR), and categorical variables as a count and percentage. To determine statistical significance, a 5% empirical significance level was adopted.

Results

There were 5,072 patients with severe sepsis or septic shock during the study period. Table 1 describes the baseline characteristics of the sample and variables used for each PSM. There were 1,412 (27.8%) patients who died in the hospital. The majority of patients had the four 3-hour bundle recommendations initiated within three hours. The percentage of patients who did not receive the recommendation within 3 hours included: lactate (14.3%), blood culture collected before antibiotic therapy (8.2%), broad-spectrum antibiotics initiated (2.7%), and crystalloids initiated inside the recommended 3-hours (0.3%). From these 214 (4.2%) patients who did not receive any of the above recommended actions within 3 hours, 88 (41.1%) died.

Figure 1(a) illustrates the casual effect of delays in blood culture draw on in-hospital mortality. The horizontal axis corresponds to *delay* in minutes and the vertical axis corresponds to the ATT. The solid line is the (smoothed) ATT at each time point and the dashed lines correspond to the 95% CI. The arrow indicates the t minutes delay when the recommendation becomes statistically significant. For example, the ATT at 30 minutes is .006 (CI = -.026, .034), meaning that a delay of half an hour increases the risk of mortality by .006 (6 additional mortalities per 1000 patients) but it is not statistically significant. A delay of 50 minutes, however, increases the patient's risk of mortality by .028 (CI = .004, .058) suggesting that a delay of 50 minutes significantly increases the risk of mortality. The distance between these two lines is the effect of the delay and is the solid line in Figure 1(a). The solid line in Figure 1(b) shows the survival probability for treated (patients with less than t minutes of delay in drawing a blood culture) and the dashed line shows for the control patients (those with t or greater minutes of delay). Figure 1(b) helps gauge how big the effect of the delay is relative to the actual survival probability. Similarly, Figure 2(a–b), 3 (a–b) and 4 (a–b) illustrates results for the lactate, antibiotic therapy and crystalloids recommendations, respectively.

Discussion

The effect of *delay* for each SSC 3-hour bundle recommendation on in-hospital mortality for patients with severe sepsis and septic shock was evaluated. A non-parametric methodology of sequential PSM for calculating the ATT at a sequence of time points was used to estimate the effect of implementation delay in the SSC guideline recommendations on all-cause in-hospital mortality. We found that delays in administering all four guideline recommendations, even when they did not exceed 3 hours, was associated with a significant increase in in-hospital mortality.

Overall, this study showed that no delay is safe, though modest delays may not be harmful in a clinically meaningful way. For most outcomes, the ATT curves (solid curve in Figures 1-a, 2-a, 3-a, and 4-a) are uniformly above zero and are strictly monotonically increasing. The longer the delay, the more harm. Up to a certain delay, the effect is not statistically

significant. This does not mean that this much delay is not harmful; it simply means that we do not have sufficient evidence to show the (potential) harm. Statistical significance depends on sample sizes and the delay that is associated with statistically significant harm could be different across hospitals. Thus, the recommended actions can and should be carried out as soon as possible to prevent harm.

The shape of the curves provides further support for our recommendation. The curves indicate that harm accumulates very quickly in the beginning (short delays) but slows down for longer delays and flattens out by about 3 hours. This means that delays exceeding the 3 hours are associated with little additional harm on top of the harm the patient has experienced within 3 hours. We acknowledge that the high compliance rate with the 3-hour bundle may influence our results: the minimal additional harm after 3 hours of delay may be partly artefactual, attributable in part to the fact that most patients have already received the recommended intervention within three hours and the sample size is too small beyond 3 hours of delay to determine further harm. However, if the observed effect was indeed artefactual, we would see the confidence interval widen substantially (as it does for crystalloid use) and also propensity matching would fail. During our analysis, we always observed sufficient support (overlap between the treatment and control groups) for propensity matching and adequate balance of the variables (except Crystalloid beyond 3 hours).

Results showed an in-hospital mortality of 27.8%. This is consistent with other reports that showed similar rates worldwide, specifically in the US and Europe, although rates may vary greatly according to different geographic regions (17, 21). Variability in outcomes may be due to the high heterogeneity of sepsis populations; this study considered incorporating a large variety of health conditions into account when performing PSM. Furthermore, diagnosis differences and, consequently, earlier initiation of treatment vary greatly as a result of several human, institutional, and financial practices (21). Previous studies reported evidence that timing of antimicrobial therapy in the presence of septic shock is associated with higher mortality (6, 8). Our findings agree that mortality increases with delay in antibiotics, but only becomes significant at nearly 3 hours after baseline (8).

There are several strengths in this study. The adopted methodology of sequential PSM in conjunction with bootstraps is novel and necessary to minimize the confounding that occurs in observational studies. We were able to incorporate several covariates in our model, making the matched pairs very alike. This combined methodology provided the estimated effect of treatment considering matched pairs with similar propensity score values, thus, decreasing bias estimation. Another strength of the study is the use of EHR data. While results from EHR-based observational studies are often considered preliminary that requires validation through clinical trials, studying the effect of delaying treatment for research purpose would be unethical and thus designing a trial for this purpose would be difficult.

Limitations exist in this study, such as the reuse of EHR for research from one single clinical source of data. Our data set only provided in-hospital mortality, and we were not able to take into account mortality that occurred outside the hospital or long-term outcomes. We acknowledge that we did not adjust for baseline vasopressors and mechanical ventilation

use, and both are important predictors of mortality among sepsis patients. Another limitation is the identification of a consistent list of broad-spectrum antibiotics. Our approach may have overestimated this, as a single list of intravenous broad-spectrum antibiotics was not found using national data standards nor in the current literature or consulting with two pharmacists. Although efforts to address this gap, a clear listing of intravenous broad-spectrum antibiotics is needed for comparison across studies; particularly in phenotyping EHR data and mapping to sepsis guideline recommendations (8). Likewise, the compliance with crystalloid recommendation could be overestimated, as this study was not able to extract precise data that could account for volume administered and/or the type of crystalloids. Future efforts are needed to standardize non-structured data, such as flowsheet data where crystalloids volume is documented; the researchers could capture and reuse this type of information for comparative effectiveness research across organizations (34, 25). The definition of *delay* does not depend on the length of exposure to sepsis. The actual onset time of sepsis is unknown; it could have occurred outside the hospital. While using actual onset time reflects physiology better; using “first suspicion” reflects clinical reality better: patient’s lactate cannot be measured before they come to the hospital.

Conclusions

The guideline recommendations should be applied as soon as sepsis is suspected; applying the guidelines with shorter delays is associated with better outcomes. Our work found no evidence that 3 hours is safe; we found evidence that even very short delays can adversely impact outcomes. Our study demonstrated a new approach to incorporate time (*delay*) when analyzing the impact on outcomes and provide new evidence for clinical practice and research. It adopted a novel methodology using EHR data and bootstrapping in conjunction with PSM to estimate the ATT of SSC guideline recommendations *delay* on in-hospital mortality. The use of EHR data is important because a clinical trial that delays urgent treatment would be unethical. Based on these findings, future studies should also investigate the impact on long-term outcomes, including national or international data set of clinical data to account for geographic variability and heterogeneity.

Acknowledgments

This study is supported by National Science Foundation (NSF) Grant #1602394 and by the National Institutes of Health (NIH) grant #GM120079. Contents of this document are the sole responsibility of the authors and do not necessarily represent official views of the NSF/NIH.

References

1. Torio, CM., Andrews, RM. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD): 2006. National Inpatient Hospital Costs: The Most Expensive Conditions by Payer, 2011: Statistical Brief #160.
2. Epstein L, Dantes R, Magill S, et al. Varying Estimates of Sepsis Mortality Using Death Certificates and Administrative Codes--United States, 1999–2014. *MMWR Morbidity Mortal Wkly Rep.* 2016; 65:342–345.
3. Verdonk F, Blet A, Mebazaa A. The new sepsis definition [Internet]. *Curr Opin Anaesthesiol.* 2017; 30:200–204. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28207566>. [PubMed: 28207566]

4. Novosad SA, Sapiano MR, Grigg C, et al. Vital Signs: Epidemiology of Sepsis: Prevalence of Health Care Factors and Opportunities for Prevention. *MMWR Morbidity Mortal Wkly Rep.* 2016; 65:864–869.
5. Yende S, Austin S, Rhodes A, et al. Long-Term Quality of Life Among Survivors of Severe Sepsis: Analyses of Two International Trials. *Crit Care Med.* 2016; 44:1461–1467. [PubMed: 26992066]
6. Dellinger RP, Levy MM, Rhodes A, et al. Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock, 2012. *Intensive Care Med.* 2013; 39:165–228. [PubMed: 23361625]
7. Gao F, Melody T, Daniels DF, et al. The impact of compliance with 6-hour and 24-hour sepsis bundles on hospital mortality in patients with severe sepsis: a prospective observational study [Internet]. *Crit Care.* 2005; 9:R764–70. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16356225>. [PubMed: 16356225]
8. Surviving Sepsis Campaign Executive Committee. Updated Bundles in Response to New Evidence [Internet]. 2015. [cited 2017 Aug 25] Available from: http://www.survivingsepsis.org/SiteCollectionDocuments/SSC_Bundle.pdf
9. Angus DC, Barnato AE, Bell D, et al. A systematic review and meta-analysis of early goal-directed therapy for septic shock: the ARISE, ProCESS and ProMISe Investigators. *Intensive Care Med.* 2015; 41:1549–1560. [PubMed: 25952825]
10. Investigators A, Group ACT, Peake SL, et al. Goal-directed resuscitation for patients with early septic shock. *N Engl J Med.* 2014; 371:1496–1506. [PubMed: 25272316]
11. Umscheid CA, Betesh J, VanZandbergen C, et al. Development, implementation, and impact of an automated early warning and response system for sepsis. *J Hosp Med.* 2015; 10:26–31. [PubMed: 25263548]
12. Goodwin AJ, Rice DA, Simpson KN, et al. Frequency, cost, and risk factors of readmissions among severe sepsis survivors. *Crit Care Med.* 2015; 43:738–746. [PubMed: 25746745]
13. Castellanos-Ortega A, Suberviola B, Garcia-Astudillo LA, et al. Impact of the Surviving Sepsis Campaign protocols on hospital length of stay and mortality in septic shock patients: results of a three-year follow-up quasi-experimental study. *Crit Care Med.* 2010; 38:1036–1043. [PubMed: 20154597]
14. Damiani E, Donati A, Serafini G, et al. Effect of performance improvement programs on compliance with sepsis bundles and mortality: a systematic review and meta-analysis of observational studies. *PLoS One.* 2015; 10:e0125827. [PubMed: 25946168]
15. Bruce HR, Maiden J, Fedullo PF, et al. Impact of nurse-initiated ED sepsis protocol on compliance with sepsis bundles, time to initial antibiotic administration, and in-hospital mortality. *J Emerg Nurs.* 2015; 41:130–137. [PubMed: 25612516]
16. Semler MW, Weavind L, Hooper MH, et al. An Electronic Tool for the Evaluation and Treatment of Sepsis in the ICU. A Randomized Controlled Trial. *Crit Care Med.* 2015; 43:1595–1602. [PubMed: 25867906]
17. Fujishima S, Gando S, Saitoh D, et al. A multicenter, prospective evaluation of quality of care and mortality in Japan based on the Surviving Sepsis Campaign guidelines. *J Infect Chemother.* 2014; 20:115–120. [PubMed: 24462442]
18. Levy MM, Rhodes A, Phillips GS, et al. Surviving Sepsis Campaign: association between performance metrics and outcomes in a 7. 5-year study. *Crit Care Med.* 2015; 43:3–12. [PubMed: 25275252]
19. Hoo WE, Muehlberg K, Ferraro RG, et al. Successes and lessons learned implementing the sepsis bundle. *J Healthc Qual.* 2009; 31:5–9.
20. Xu JY, Chen QH, Liu SQ, et al. The Effect of Early Goal-Directed Therapy on Outcome in Adult Severe Sepsis and Septic Shock Patients: A Meta-Analysis of Randomized Clinical Trials. *Anesth Analg.* 2016
21. Rhodes A, Phillips G, Beale R, et al. The Surviving Sepsis Campaign bundles and outcome: results from the International Multicentre Prevalence Study on Sepsis (the IMPReSS study). *Intensive Care Med.* 2015; 41:1620–1628. [PubMed: 26109396]
22. Kadri SS, Rhee C, Strich JR, et al. Estimating Ten-Year Trends in Septic Shock Incidence and Mortality in United States Academic Medical Centers Using Clinical Data [Internet]. *Chest.* 2017;

- 151:278–285. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27452768>. [PubMed: 27452768]
23. Rhee C, Kadri S, Huang SS, et al. Objective Sepsis Surveillance Using Electronic Clinical Data [Internet]. *Infect Control Hosp Epidemiol*. 2016; 37:163–171. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26526737>. [PubMed: 26526737]
 24. Brandt BN, Gartner AB, Moncure M, et al. Identifying Severe Sepsis via Electronic Surveillance [Internet]. *Am J Med Qual*. 2015; 30:559–565. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24970280>. [PubMed: 24970280]
 25. Narayanan N, Gross AK, Pintens M, et al. Effect of an electronic medical record alert for severe sepsis among ED patients [Internet]. *Am J Emerg Med*. 2016; 34:185–188. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26573784>. [PubMed: 26573784]
 26. McRee L, Thanavaro JL, Moore K, et al. The impact of an electronic medical record surveillance program on outcomes for patients with sepsis [Internet]. *Hear Lung J Acute Crit Care*. 2014; 43:546–549. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24961633>.
 27. Westra BL, Landman S, Yadav P, et al. Secondary Analysis of an Electronic Surveillance System Combined with Multi-focal Interventions for Early Detection of Sepsis [Internet]. *Appl Clin Inform*. 2017; 8:47–66. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28097288>. [PubMed: 28097288]
 28. Mani S, Ozdas A, Aliferis C, et al. Medical decision support using machine learning for early detection of late-onset neonatal sepsis [Internet]. *J Am Med Informatics Assoc*. 2014; 21:326–336. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24043317>.
 29. Desautels T, Calvert J, Hoffman J, et al. Prediction of Sepsis in the Intensive Care Unit With Minimal Electronic Health Record Data: A Machine Learning Approach [Internet]. *JMIR Med informatics*. 2016; 4:e28. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27694098>.
 30. Taylor RA, Pare JR, Venkatesh AK, et al. Prediction of In-hospital Mortality in Emergency Department Patients With Sepsis: A Local Big Data-Driven, Machine Learning Approach [Internet]. *Acad Emerg Med*. 2016; 23:269–278. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26679719>. [PubMed: 26679719]
 31. Gultepe E, Green JP, Nguyen H, et al. From vital signs to clinical outcomes for patients with sepsis: a machine learning basis for a clinical decision support system [Internet]. *J Am Med Inform Assoc*. 2014; 21:315–25. [cited 2017 Sep 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23959843>. [PubMed: 23959843]
 32. Pruinelli L, Yadav P, Hangsleben A, et al. A Data Mining Approach to Determine Sepsis Guideline Impact on Inpatient Mortality and Complications [Internet]. *AMIA Jt Summits Transl Sci proceedings AMIA Jt Summits Transl Sci*. 2016; 2016:194–202. [cited 2017 Feb 6] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27570669>.
 33. Westra BL, Landman S, Yadav P, et al. Secondary Analysis of an Electronic Surveillance System Combined with Multi-focal Interventions for Early Detection of Sepsis [Internet]. *Appl Clin Inform*. 2017; 8:47–66. [cited 2017 Jul 25] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28097288>. [PubMed: 28097288]
 34. Austin PC. Comparing paired vs non-paired statistical methods of analyses when making inferences about absolute risk reductions in propensity-score matched samples [Internet]. *Stat Med*. 2011; 30 n/a-n/a[cited 2017 Aug 25] Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21337595>.
 35. Kirby JC, Speltz P, Rasmussen LV, et al. PheKB. a catalog and workflow for creating electronic phenotype algorithms for transportability [Internet]. *J Am Med Informatics Assoc*. 2016; 23:1046–1052. [cited 2017 Sep 6] Available from: <https://academic.oup.com/jamia/article-lookup/doi/10.1093/jamia/ocv202>.

Fig 1.

(a) Casual effect and (b) survival probabilities of the blood culture recommendation timing on in-hospital mortality between treatment and control groups.

Fig 2.

(a) Casual effect and (b) survival probabilities of the lactate recommendation timing on in-hospital mortality between treatment and control groups.

Fig 3.
(a) Casual effect and (b) survival probabilities of the antibiotics recommendation timing on in-hospital mortality between treatment and control groups.

Fig 4.

(a) Casual effect and (b) survival probabilities of the crystalloid recommendation timing on in-hospital mortality between case and control groups.

Table 1

Predictor variables for patients with severe sepsis and septic shock (n=5,072).

Variables	Median or Interquartile Ranges (IQR) or Counts (n) and Percentages	Variables	Median or Interquartile Ranges (IQR) or Counts (n) and Percentages
Age	66 [55 – 78]	Respiratory rate	20 [18 – 24]
Gender (Male)	2,588 (51.0)	Charlson Comorbidity Index	3.9 [1 – 6]
Race (Caucasian)	4,349 (85.7)	Comorbidities Severity Scores	
Ethnicity (Non-Hispanic)	4,822 (95.0)	Respiratory	2.2 [0 – 4]
Lactate	1.8 [1.8 – 3.2]	Cardiac	5.6 [1 – 9]
White blood cells	12.6 [7.9 – 17.4]	Liver	0.5 [0 – 0]
Median blood pressure	81.3 [67 – 96]	Cerebrovascular	0.3 [0 – 0]
Heart rate	97 [90 – 103]	Kidney	1.3 [0 – 2]
Temperature	98.5 [97 – 100]		