Published in final edited form as:

Neurotoxicology. 2018 March ; 65: 22–27. doi:10.1016/j.neuro.2018.01.005.

The Association between Early-life Relative Telomere Length and Childhood Neurodevelopment

Marina Oktapodas Feiler, MS^a, Deven Patel, PhD^b, Huiqi Li, PhD^c, Philip J. Meacham, MS^a, Gene E. Watson, DDS, PhD^a, Conrad Shamlaye, MBChB^d, Alison Yeates, PhD^e, Karin Broberg, PhD^f, and Edwin van Wijngaarden, PhD^{a,*}

^aUniversity of Rochester School of Medicine and Dentistry, Rochester, NY, USA

^bOccupational and Environmental Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, Rockville, MD

^cDivision of Occupational and Environmental Medicine, Lund University, Lund, Sweden

^dMinistry of Health, Republic of Seychelles, Seychelles

^eUlster University, Coleraine, UK

^fInstitute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

Abstract

Purpose—To examine the association between telomere length and neurodevelopment in children.

Methods—We examined the relationship between relative telomere length (rTL) and neurodevelopmental outcomes at 9 and 30 months, and 5 years of age in children enrolled in the Seychelles Child Development Study Nutrition Cohort 1 (NC1). Relative telomere length was measured in cord blood and in child blood at age five. Multivariable linear regression examined associations between neurodevelopmental outcomes and rTL adjusting for relevant covariates.

Results—Mean rTL was 1.18 at birth and 0.71 at age five. Increased cord blood rTL was associated with better scores on two neurodevelopmental tests, the psychomotor developmental index ($\beta = 4.01$; 95% confidence interval (CI)=0.17, 7.85) at age 30 months, and the Woodcock Johnson test of achievement letter-word score ($\beta = 2.88$; CI=1.21–4.56) at age five. The Woodcock Johnson test of achievement letter-word score remained statistically significant after two outliers were excluded ($\beta = 2.83$; CI=0.69, 4.97); the psychomotor developmental index did not ($\beta = 3.62$; CI=–1.28, 8.52). None of the neurodevelopmental outcomes at age five were associated with five-year rTL.

*Please address correspondence to: Edwin van Wijngaarden, PhD, Department of Public Health Sciences, University of Rochester Medical Center, 265 Crittenden Blvd, CU 420644, Rochester, NY 14642, edwin_van_wijngaarden@urmc.rochester.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusion—Although increased cord blood rTL was associated with better test scores for a few neurodevelopmental outcomes, this study found little consistent evidence of an association between rTL and neurodevelopment. Future studies with a larger sample size, longer follow-up, and other relevant biological markers (e.g. oxidative stress) are needed to clarify the role of rTL in neurodevelopment and its relevance as a potential surrogate measure for oxidative stress in the field of developmental neurotoxicity.

Keywords

children; epidemiology; cognition; language

Introduction

Telomeres are non-coding, nucleoprotein complexes at the ends of eukaryotic chromosomes whose function is to preserve genomic integrity. Telomeres naturally shorten due to incomplete DNA replication during cellular divisions, otherwise known as cellular aging (Blackburn, 2001; McEachern, Krauskopf, & Blackburn, 2000). Telomeres eventually reach a critical point in length where they lose their protective functions and the cells stop dividing and enter either apoptosis or cellular senescence (Gisselsson et al., 2001; Murnane, 2006; Verdun & Karlseder, 2007). Therefore, telomere length can be considered to be a measure of our “biological” age as opposed to our “chronological” age (Oeseburg, de Boer, van Gilst, & van der Harst, 2010). As a marker for cellular aging, telomere length (TL) is seen as a possible predictor of various age-related diseases, including cognitive decline. For example, telomere length has been associated with Alzheimer’s disease, (Roberts et al., 2014; Zhan et al., 2015) as well as poor cognitive function in adults (Kingma, de Jonge, van der Harst, Ormel, & Rosmalen, 2012; Valdes et al., 2010; Yaffe et al., 2011). Similarly, TL at young ages may be associated with cognition and neurodevelopment in early life or may be related to adult cognitive function in later life, likely as an indirect marker of other biological processes that may influence neurodevelopment such as oxidative stress (Epel et al., 2004; Houben, Moonen, van Schooten, & Hageman, 2008; Von Zglinicki, 2000, 2002). TL may be considered a surrogate measure for underlying oxidative stress and inflammation and potentially a good biomarker of neurotoxicity, as has been indicated in previous literature (Epel et al., 2004; Houben et al., 2008; Von Zglinicki, 2000, 2002).

The relationship between TL and neurodevelopmental outcomes in children has received little attention. Only four studies to date have examined the association with neurobehavioral outcomes in children (Costa Dde et al., 2015; Henje Blom et al., 2015; Z. Li et al., 2014; Wojcicki et al., 2015). Two of these studies have reported correlations between shorter TL and measures of depression and inattention in later childhood and adolescence (Costa Dde et al., 2015; Henje Blom et al., 2015). Other studies showed a shorter TL in young children with autism and defiant behavior (Z. Li et al., 2014; Wojcicki et al., 2015). Adding to this limited literature, our study examines the association between relative telomere length (rTL) and cognitive development, including motor, language, memory and problem-solving skills. These are more subtle aspects of development that have not been previously assessed in relation with TL.

Methods

Study population

The Seychelles Child Development Study (SCDS) is a series of longitudinal observational studies that evaluate the development of children in Seychelles, and examine if mercury exposure during pregnancy (due to a high fish diet) is associated with child development. The present analysis used data from the Nutrition Cohort 1 (NC1) (Davidson et al., 2008), a cohort of 300 mothers that were enrolled in 2001 during their first trimester of pregnancy. The inclusion criteria included mothers at least 16 years of age, native born of Seychelles, and residing on Mahé. Exclusion criteria included infants with major congenital anomalies and twins. Research protocols were reviewed and approved by the Institutional Review Boards of the University of Rochester, the Ministry of Health in Republic of Seychelles and the Regional Ethics Committee, Lund University. The procedures followed were in accordance with the Helsinki Declaration, and all participants gave informed consent. The present study examined child telomere length at birth and at the time of examination at approximately age 5 years in relation to developmental outcomes assessed in the children at 9 months, 30 months, and 5 years of age.

Blood collection

Cord blood samples were collected immediately after delivery into EDTA-containing tubes, from which whole blood was obtained and stored at -80°C until analysis. Similarly, children's venous non-fasting blood samples were collected in EDTA-containing tubes after completion of the 5-year developmental assessment, from which whole blood samples were obtained and stored at -80°C .

Telomere length assessment

DNA was isolated from peripheral blood using the Qiagen DNA blood Midi kit (Qiagen, Hilden, Germany). Relative telomere length (rTL) was measured using real-time PCR (7900HT, Applied Biosystems, Foster City, CA, USA), as described previously (Ameer et al., 2016; H. Li, Engstrom, Vahter, & Broberg, 2012). Briefly, master mixes were prepared, containing 0.5 U *Taq* Platina (Invitrogen, Carlsbad, CA, USA), 1×PCR Buffer, 0.8 mM dNTPs, 1.75 mM MgCl_2 , 0.3 mM SybrGreen (Invitrogen), 1×Rox (Invitrogen), and either telomere primers (0.45 μM of each primer), or hemoglobin beta chain (*HBB*) primers (0.45 μM for each primer). Five microliters of sample DNA (3 ng/ μl) was added to each reaction resulting in a final volume of 20 μl . A standard curve, a reference DNA, and a negative control were also included in each run, and all samples, standards, and controls were run in triplicate. The relative length of the telomeres was obtained through calculating the ratio (T/S) of the telomere repeat product to a single-copy gene product (S, here *HBB*) for each individual, by the formula $T/S = 2^{-\text{Ct}}$, where $\text{Ct} = \text{Ct}_{\text{telomere}} - \text{Ct}_{\text{HBB}}$. This ratio was then compared with the ratio of a reference DNA. The telomere length ratio is an arbitrary value. Relative telomere length was measured at birth from cord blood and again in blood collected at five years of age.

Neurodevelopmental assessment

We analyzed data from the Bayley Scales of Infant Development-II (BSID-II), a well-standardized measure of infant cognition and development that was administered at ages 9 and 30 months. The BSID-II yielded two endpoints: the mental developmental index (MDI) and psychomotor developmental index (PDI) (Davidson et al., 2008). We also examined the following developmental tests at five years of age: finger tapping (dominant and non-dominant hand), the Preschool Language Scale (total language score, verbal ability, and auditory comprehension), the Woodcock-Johnson Scholastic Achievement Test (letter-word recognition and applied problems), the Kaufman Brief Intelligence Test (verbal knowledge, matrices), and the Child Behavior Checklist (Strain et al., 2012).

Covariates

As in previous studies of this cohort, covariates were selected *a priori* based on their known association with developmental outcomes (Strain et al., 2008; Strain et al., 2012). Covariates included: child sex, birth weight, age of child at testing, Hollingshead socioeconomic status (SES) at birth, maternal IQ, maternal age at birth of child, family status (i.e. whether or not both parents resided with the child) at 9 months, and home environment at birth. Smoking was not included as a covariate because only eight mothers reported smoking during pregnancy.

Statistical analysis

Descriptive statistics for relevant sample characteristics were calculated, including the mean, median, and standard deviation for all continuous variables, the proportions for categorical variables, and the distribution of cord blood rTL, 5-year rTL, and all neurodevelopment outcomes. We examined correlation coefficients for the association between cord rTL, 5-year rTL, and change in rTL from birth to 5-years of age. Cord rTL and five-year rTL were poorly correlated (spearman $r=0.26$, $p=0.0007$; pearson $r=0.14$, $p=0.067$), whereas cord rTL and change in rTL were highly correlated (spearman $r=-0.90$, $p<0.0001$; pearson $r=-0.98$, $p<0.0001$). Therefore, the change in rTL was not further investigated as it did not contribute additional statistical information to cord blood rTL. Two outliers were identified in the cord blood rTL, defined as being at least three interquartile lengths above the third quartile (Tukey, 1977). Analyses of cord rTL were conducted with and without these observations. Appropriate assumptions were tested confirming regression models were appropriate for this analysis (Rosner, 2011).

Covariate-adjusted linear regression analyses of cord blood rTL were performed separately for 9-month, 30-month and 5-year developmental outcomes. Similarly, adjusted regression analyses modeled 5-year rTL against the outcomes at age five. We considered examining categories of shortened, maintained, and lengthened rTL using cut-points of 5% and 10% difference between five-year and cord blood rTL as proposed elsewhere (Wojcicki, Shiboski, et al., 2016). However, only a few children maintained or increased rTL using a 5% ($n=13$) or 10% ($n=15$) cut off; therefore, this was not further pursued.

In secondary analyses, we examined cord rTL in quartiles to assess the nature dose-response relationship between cord rTL and developmental outcomes that were statistically

significantly associated in continuous cord rTL analyses, in order to check consistency in our findings. P values <0.05 were considered statistically significant. All data management and analyses were performed using the SAS software system (SAS Institute Inc., Cary, NC, USA; version 9.4).

Results

Study population

Demographic, maternal and child characteristics are shown in Table 1. The sample size for cord blood rTL was n=184 and for the 5-year rTL was n=209. A similar number of males and females were included in the analysis, children had an average birth weight (SD) of 3247g (480), and had one parent or less residing in the household (53%). On average, mothers were 27 years of age, and had mean SES, PROCESS, and K-bit scores of 34, 152, and 86, respectively. Mean rTL was 1.18 at birth and decreased to 0.71 at age five. There was no considerable difference between males and females in mean TL in cord blood (1.18 and 1.19, respectively) or at age 5 (0.70 and 0.73, respectively). Other covariates were also not associated with either cord blood or 5-year rTL (data not shown).

Relative telomere length and neurodevelopmental outcomes at age 9 and 30 months, and age 5

Table 2 shows the adjusted results for the associations of cord blood and 5-year rTL with the various neurodevelopmental outcomes at the three different time points, including the full sample and after removing outliers (n=2). Overall, there was no clear association between rTL and neurodevelopmental outcomes in this study population. The majority of the outcomes show a slight positive association, however; the results are statistically imprecise and associations were not consistent across the different developmental outcomes and rTL measures. Cord blood rTL was associated with improved scores on the psychomotor development index ($\beta = 4.01$; 0.17, 7.85) at age 30 months and on the Woodcock Johnson test of achievement letter-word ($\beta = 2.88$; 95% CI: 1.21, 4.56) at age five (Table 2). When removing two outliers the Woodcock Johnson letter-word score remained statistically significant (Table 2), whereas the association with the psychomotor development index remained positive but lost precision and statistical significance (Table 2). Inference for the other outcomes was not affected after excluding the two outliers, despite the change in point estimate for some outcomes (e.g. MDI). In categorical cord rTL analysis we did not observe clear monotonic dose-response patterns for psychomotor development index (quartile 1: reference; quartile 2: $\beta = 5.42$; 95% CI: 0.03, 10.81; quartile 3: $\beta = 7.51$; 95% CI: 2.04, 12.98; quartile 4: $\beta = 4.71$; 95% CI: -0.75, 10.18) and the Woodcock Johnson Achievement letter-word scores (quartile 1: reference; quartile 2: $\beta = -0.27$; 95% CI: -2.69, 2.15; quartile 3: $\beta = 0.24$; 95% CI: -2.24, 2.72; quartile 4: ($\beta = 1.72$; 95% CI: -0.77, 4.20). Other neurodevelopmental outcomes were not statistically significantly associated with cord blood (Table 2), and none of the outcomes were associated with five-year rTL (Table 3).

Discussion

Our study examined rTL with an array of neurodevelopmental tests in young children. We did not find consistent associations between rTL and neurodevelopmental outcomes. Only a few of the neurodevelopmental outcomes showed positive associations with cord blood rTL, including the Woodcock Johnson Achievement letter-word test and psychomotor development index. We found no consistent dose-response relationship with these outcomes in categorical analyses however, and the psychomotor developmental index association was no longer significant after excluding two outliers; thus we should interpret our results cautiously. There were no significant differences in rTL between males and females, consistent with the lack of a sex difference in rTL in a case-control study of autism (Z. Li et al., 2014). Nevertheless, studies investigating the association with TL in children are limited (Costa Dde et al., 2015; Henje Blom et al., 2015; Z. Li et al., 2014; Wojcicki et al., 2015). No previous studies have investigated this association in infants and young children using the measures of neurodevelopment as we studied, but studies of other adverse cognitive or behavioral outcomes in children suggest an increased risk with decreases in TL (Costa Dde et al., 2015; Henje Blom et al., 2015; Z. Li et al., 2014; Wojcicki et al., 2015). In a case-control study of autism (n=239), children 4–6 years old with an autism diagnosis had a statistically significantly lower telomere length (0.88) compared with children without autism (1.01) (Z. Li et al., 2014). An additional study (n=108) was conducted investigating oppositional defiant behavior in 3–5 year olds and telomere length (Wojcicki et al., 2015). The results showed that oppositional defiant behavior was a predictor of shorter telomere length ($\beta = -359.25$ (95% CI: $-633.84, -84.66$), $p=0.01$) (Wojcicki et al., 2015). Lastly, a study of inattention and behavior assessed ADHD and telomere length in children aged 6–16 years of age (n=61) (Costa Dde et al., 2015). The hyperactive and impulsive dimensions of ADHD were found to be negatively correlated TL in children ($r = -0.34$, $p=0.0008$), while the inattention dimension was not found to be associated with TL ($p>0.05$) (Costa Dde et al., 2015). Sample sizes in these studies range from n=61 to n=239, and so the present study falls on the higher end of the spectrum with n=209. The methods for measuring telomere length varied across studies, and therefore results from previous studies cannot be directly compared to ours. Nevertheless, these studies still provide information on the consistency of evidence that telomere length plays some role on neurodevelopment and neurobehavioral outcomes in childhood and adolescence.

An important potential pathway to consider is that through oxidative stress and inflammation. Telomere length can be considered a biomarker of oxidative stress (Epel et al., 2004; Houben et al., 2008; Von Zglinicki, 2000, 2002). Life stress has also shown to accelerate telomere length shortening in healthy premenopausal women, suggesting that higher levels of life stress are associated with higher levels of oxidative stress (Epel et al., 2004). Chronic inflammation has shown to lead to persistent damage to telomeres and in return increase the rate of biological aging (Houben et al., 2008), and may influence neurodevelopment, (Andrews et al., 2008; Stolp & Dziegielewska, 2009; van der Burg et al., 2016). Thus, TL may be a surrogate marker for underlying oxidative stress and inflammation, which more directly impact neurodevelopmental outcomes, and as such, be a useful marker in longitudinal studies to reflect these processes. The use of TL as an

imperfect surrogate measure of underlying biological processes may have resulted in null findings in the present study.

More research has been done in adults to evaluate telomere length, aging and neuropsychological and cognitive outcomes (Kingma et al., 2012; Roberts et al., 2014; Valdes et al., 2010; Yaffe et al., 2011; Zhan et al., 2015). The PREVEND study in the Netherlands investigated intelligence and telomere length in adults (Kingma et al., 2012). Findings suggested increased general intelligence, measured by the General Aptitude-Test Battery (GATB), was associated with longer telomere length ($\beta = 0.163$, $p < .001$), after adjustment for additional covariates (Kingma et al., 2012). A study of healthy women used the Cambridge Neuropsychological Test Automated Battery (CANTAB) to assess neuropsychological aptitude and examined the association with telomere length (Valdes et al., 2010). Three dimensions of the CANTAB (delayed matching to sample, pattern recognition and space span) were all found to be positively correlated with longer TL ($p < .05$) (Valdes et al., 2010). The other three dimensions (paired associations learning, reaction time, spatial working memory) were found to be negatively correlated with TL (Valdes et al., 2010). Further analyses in children using sensitive cognitive test batteries such as CANTAB may shed further light on this association.

Several studies have assessed predictors of telomere length in children (Gilfillan et al., 2016; Wojcicki, Olveda, et al., 2016). One study in particular examining Latino children found female sex, higher maternal education, and child head circumference to be associated with longer cord blood telomere length (Wojcicki, Olveda, et al., 2016). Furthermore, shorter cord blood telomere length was associated with some level of oxidative stress in utero (preeclampsia, maternal hypertension, gestational diabetes), as well as low birth weight and preterm birth (Wojcicki, Olveda, et al., 2016). Additionally, a study assessing fetal telomere length with maternal and fetal glucose levels found inverse associations, $\beta = -0.563$, $p < 0.05$ and $\beta = -0.297$, $p < 0.05$, respectively (Gilfillan et al., 2016). These associations suggest possibly mechanisms by which prenatal factors may influence child telomere length and neurodevelopment. We did not find clear associations with our covariates other than family status.

While the sample size was relatively small resulting in imprecise estimates of association, the present study is the first to directly assess rTL and an array of tests assessing neurodevelopment in young children during the most critical period of brain development. This is also one of few studies to assess this association longitudinally rather than cross-sectionally (Costa Dde et al., 2015; Henje Blom et al., 2015; Z. Li et al., 2014; Wojcicki et al., 2015) using multiple rTL measurements and using a more comprehensive array of developmental outcomes than in previous studies. We were also able to account for important covariates.

Conclusion

In conclusion, our results do not strongly support an association between telomere length and child developmental outcomes at 5 years of age. Future studies with a larger sample

size, longer follow-up, and other relevant biological markers (e.g. oxidative stress) are needed to clarify the role of rTL in neurodevelopment.

Acknowledgments

Funding Sources: US National Institute of Environmental Health Sciences, National Institutes of Health (R01-ES010219, R01-ES015578, P30-ES01247); the European Union (Sixth Framework Programme; PHIME; FOOD-CT-2006-016253); the Swedish Research Council FORMAS; and the Government of Seychelles.

This research was supported by the US National Institute of Environmental Health Sciences, National Institutes of Health (R01-ES010219, R01-ES015578, P30-ES01247), the European Union (Sixth Framework Programme; PHIME; FOOD-CT-2006-016253); the Swedish Research Council FORMAS and by the Government of Seychelles. The contents reflect only the authors' views; the European Union is not liable for any use that may be made of the information.

Abbreviations

rTL	Relative telomere length
TL	Telomere length
SCDS	Seychelles Child Development Study
NC1	Nutritional Cohort 1
BSID-II	Bayley Scales of Infant Development-II
MDI	Mental Development Index
PDI	Psychomotor Development Index
SES	Socioeconomic status
GATB	General Aptitude-Test Battery
CANTAB	Cambridge Neuropsychological Test Automated Battery
HBB	Hemoglobin beta chain

References

- Ameer SS, Xu Y, Engstrom K, Li H, Tallving P, Nermell B, ... Broberg K. Exposure to Inorganic Arsenic Is Associated with Increased Mitochondrial DNA Copy Number and Longer Telomere Length in Peripheral Blood. *Front Cell Dev Biol.* 2016; 4:87.doi: 10.3389/fcell.2016.00087 [PubMed: 27597942]
- Andrews WW, Cliver SP, Biasini F, Peralta-Carcelen AM, Rector R, Alriksson-Schmidt AI, ... Hauth JC. Early preterm birth: association between in utero exposure to acute inflammation and severe neurodevelopmental disability at 6 years of age. *American Journal of Obstetrics and Gynecology.* 2008; 198(4):466e461–466.e411. doi:<http://dx.doi.org/10.1016/j.ajog.2007.12.031>. [PubMed: 18395043]
- Blackburn EH. Switching and signaling at the telomere. *Cell.* 2001; 106(6):661–673. [PubMed: 11572773]
- de Costa DS, Rosa DV, Barros AG, Romano-Silva MA, Malloy-Diniz LF, Mattos P, de Miranda DM. Telomere length is highly inherited and associated with hyperactivity-impulsivity in children with attention deficit/hyperactivity disorder. *Front Mol Neurosci.* 2015; 8:28.doi: 10.3389/fnmol.2015.00028 [PubMed: 26217174]

- Davidson PW, Strain JJ, Myers GJ, Thurston SW, Bonham MP, Shamlaye CF, ... Clarkson TW. Neurodevelopmental effects of maternal nutritional status and exposure to methylmercury from eating fish during pregnancy. *Neurotoxicology*. 2008; 29(5):767–775. DOI: 10.1016/j.neuro.2008.06.001 [PubMed: 18590763]
- Epel ES, Blackburn EH, Lin J, Dhabhar FS, Adler NE, Morrow JD, Cawthon RM. Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of Sciences of the United States of America*. 2004; 101(49):17312–17315. DOI: 10.1073/pnas.0407162101 [PubMed: 15574496]
- Gilfillan C, Naidu P, Gunawan F, Hassan F, Tian P, Elwood N. Leukocyte Telomere Length in the Neonatal Offspring of Mothers with Gestational and Pre-Gestational Diabetes. *PloS one*. 2016; 11(10):e0163824.doi: 10.1371/journal.pone.0163824 [PubMed: 27736899]
- Gisselsson D, Jonson T, Petersen A, Strombeck B, Dal Cin P, Hoglund M, ... Mandahl N. Telomere dysfunction triggers extensive DNA fragmentation and evolution of complex chromosome abnormalities in human malignant tumors. *Proc Natl Acad Sci U S A*. 2001; 98(22):12683–12688. DOI: 10.1073/pnas.211357798 [PubMed: 11675499]
- Henje Blom E, Han LK, Connolly CG, Ho TC, Lin J, LeWinn KZ, ... Yang TT. Peripheral telomere length and hippocampal volume in adolescents with major depressive disorder. *Transl Psychiatry*. 2015; 5:e676.doi: 10.1038/tp.2015.172 [PubMed: 26556285]
- Houben MJ, Moonen HJJ, van Schooten FJ, Hageman GJ. Telomere length assessment: Biomarker of chronic oxidative stress? *Free Radical Biology and Medicine*. 2008; 44(3):235–246. doi:<http://dx.doi.org/10.1016/j.freeradbiomed.2007.10.001>. [PubMed: 18021748]
- Kingma EM, de Jonge P, van der Harst P, Ormel J, Rosmalen JG. The association between intelligence and telomere length: a longitudinal population based study. *PloS one*. 2012; 7(11):e49356.doi: 10.1371/journal.pone.0049356 [PubMed: 23166646]
- Li H, Engstrom K, Vahter M, Broberg K. Arsenic exposure through drinking water is associated with longer telomeres in peripheral blood. *Chem Res Toxicol*. 2012; 25(11):2333–2339. DOI: 10.1021/tx300222t [PubMed: 22917110]
- Li Z, Tang J, Li H, Chen S, He Y, Liao Y, ... Chen X. Shorter telomere length in peripheral blood leukocytes is associated with childhood autism. *Sci Rep*. 2014; 4:7073.doi: 10.1038/srep07073 [PubMed: 25399515]
- McEachern MJ, Krauskopf A, Blackburn EH. Telomeres and their control. *Annu Rev Genet*. 2000; 34:331–358. DOI: 10.1146/annurev.genet.34.1.331 [PubMed: 11092831]
- Murnane JP. Telomeres and chromosome instability. *DNA Repair (Amst)*. 2006; 5(9–10):1082–1092. DOI: 10.1016/j.dnarep.2006.05.030 [PubMed: 16784900]
- Oeseburg H, de Boer RA, van Gilst WH, van der Harst P. Telomere biology in healthy aging and disease. *Pflugers Arch*. 2010; 459(2):259–268. DOI: 10.1007/s00424-009-0728-1 [PubMed: 19756717]
- Roberts RO, Boardman LA, Cha RH, Pankratz VS, Johnson RA, Druliner BR, ... Petersen RC. Short and long telomeres increase risk of amnesic mild cognitive impairment. *Mech Ageing Dev*. 2014; 141–142:64–69. DOI: 10.1016/j.mad.2014.10.002
- Rosner, B. *Fundamentals of Biostatistics*. 7. Brooks/Cole Cengage Learning; 2011. p. 448–452.
- Stolp HB, Dziegielewska KM. Review: Role of developmental inflammation and blood–brain barrier dysfunction in neurodevelopmental and neurodegenerative diseases. *Neuropathology and Applied Neurobiology*. 2009; 35(2):132–146. DOI: 10.1111/j.1365-2990.2008.01005.x [PubMed: 19077110]
- Strain J, Davidson PW, Bonham MP, Duffy EM, Stokes-Riner A, Thurston SW, ... Georger LA. Associations of maternal long-chain polyunsaturated fatty acids, methyl mercury, and infant development in the Seychelles Child Development Nutrition Study. *Neurotoxicology*. 2008; 29(5):776–782. [PubMed: 18590765]
- Strain J, Davidson PW, Thurston SW, Harrington D, Mulhern MS, McAfee AJ, ... Watson GE. Maternal PUFA status but not prenatal methylmercury exposure is associated with children's language functions at age five years in the Seychelles. *The Journal of nutrition*. 2012; 142(11):1943–1949. [PubMed: 23014496]
- Tukey, JW. *Exploratory data analysis*. 1977.

- Valdes AM, Deary IJ, Gardner J, Kimura M, Lu X, Spector TD, ... Cherkas LF. Leukocyte telomere length is associated with cognitive performance in healthy women. *Neurobiol Aging*. 2010; 31(6): 986–992. DOI: 10.1016/j.neurobiolaging.2008.07.012 [PubMed: 18718693]
- van der Burg JW, Sen S, Chomitz VR, Seidell JC, Leviton A, Dammann O. The role of systemic inflammation linking maternal body mass index to neurodevelopment in children: Inflammation and neurodevelopment. *Pediatric research*. 2016; 79(1–1):3–12. DOI: 10.1038/pr.2015.179 [PubMed: 26375474]
- Verdun RE, Karlseder J. Replication and protection of telomeres. *Nature*. 2007; 447(7147):924–931. DOI: 10.1038/nature05976 [PubMed: 17581575]
- Von Zglinicki T. Role of Oxidative Stress in Telomere Length Regulation and Replicative Senescence. *Annals of the New York Academy of Sciences*. 2000; 908(1):99–110. DOI: 10.1111/j.1749-6632.2000.tb06639.x [PubMed: 10911951]
- von Zglinicki T. Oxidative stress shortens telomeres. *Trends in Biochemical Sciences*. 2002; 27(7): 339–344. doi:[http://dx.doi.org/10.1016/S0968-0004\(02\)02110-2](http://dx.doi.org/10.1016/S0968-0004(02)02110-2). [PubMed: 12114022]
- Wojcicki JM, Heyman MB, Elwan D, Shiboski S, Lin J, Blackburn E, Epel E. Telomere length is associated with oppositional defiant behavior and maternal clinical depression in Latino preschool children. *Transl Psychiatry*. 2015; 5:e581.doi: 10.1038/tp.2015.71 [PubMed: 26080316]
- Wojcicki JM, Olveda R, Heyman MB, Elwan D, Lin J, Blackburn E, Epel E. Cord blood telomere length in Latino infants: relation with maternal education and infant sex. *J Perinatol*. 2016; 36(3): 235–241. DOI: 10.1038/jp.2015.178 [PubMed: 26633142]
- Wojcicki JM, Shiboski S, Heyman MB, Elwan D, Lin J, Blackburn E, Epel E. Telomere length change plateaus at 4 years of age in Latino children: associations with baseline length and maternal change. *Mol Genet Genomics*. 2016; 291(3):1379–1389. DOI: 10.1007/s00438-016-1191-2 [PubMed: 26965507]
- Yaffe K, Lindquist K, Kluse M, Cawthon R, Harris T, Hsueh WC, ... Cummings SR. Telomere length and cognitive function in community-dwelling elders: findings from the Health ABC Study. *Neurobiol Aging*. 2011; 32(11):2055–2060. DOI: 10.1016/j.neurobiolaging.2009.12.006 [PubMed: 20031273]
- Zhan L, Zhou J, Wang Y, Jin Y, Jahanshad N, Prasad G. ... For The Alzheimer's Disease Neuroimaging, I. Comparison of nine tractography algorithms for detecting abnormal structural brain networks in Alzheimer's disease. *Front Aging Neurosci*. 2015; 7:48.doi: 10.3389/fnagi.2015.00048 [PubMed: 25926791]

Highlights

- There was no clear association between cord blood relative telomere length, five-year relative telomere length and neurodevelopmental outcomes in this study population.
- The majority of the outcomes show a slight positive association with cord blood and five-year relative telomere length, although results are imprecise.
- Cord blood relative telomere length was statistically associated with improved scores on the psychomotor development index at age 30 months and on the Woodcock Johnson test of achievement letter-word.
- Other neurodevelopmental outcomes were not statistically significantly associated with either cord blood or five-year relative telomere length.

Table 1

Distribution and frequency of sociodemographic and telomere variables at each relative telomere length (rTL) measurement point: Seychelles Child Development Study Nutrition Cohort I.

Covariate	<u>Measurement Points of Relative Telomere Length</u>	
	Cord Blood (n=184)	5 years old (n=209)
	<i>N (%)</i>	
Child's Sex		
Male	87 (47)	103 (49)
Female	97 (53)	106 (51)
Family Status		
Both Parents	87 (47)	100 (48)
One Parent or Less	97 (53)	109 (52)
	<i>Mean (SD)</i>	
Relative Telomere Length	1.18 (0.50)	0.71 (0.10)
Child's Birth weight (g)	3247 (480)	3244 (470)
Maternal Age at Birth (yr)	27.35 (5.85)	27.89 (6.01)
Maternal K-BIT Score	86.34 (14.20)	86.29 (14.11)
PROCESS Score	152.58 (14.84)	152.43 (14.55)
Hollingshead SES at 9 months	33.76 (11.07)	34.36 (10.93)

Adjusted* associations between cord blood telomere length and cognitive outcomes in early childhood in the Seychelles Child Development Study Nutrition Cohort I.

Table 2

Model	Full Sample (n=184)				Outliers excluded (n=182)			
	N	β coefficient (95% CI)	P value	N	β coefficient (95% CI)	P value	N	β coefficient (95% CI)
9-month outcomes								
MDI	182	0.76 (-1.65, 3.16)	0.54	180	1.25 (-1.80, 4.31)	0.42		
PDI	180	0.83 (-2.24, 3.90)	0.60	178	3.19 (-0.67, 7.05)	0.10		
30-month outcomes								
MDI	183	1.20 (-1.50, 3.89)	0.38	181	-0.47 (-3.87, 2.94)	0.79		
PDI	180	4.01 (0.17, 7.85)	<0.001	178	3.62 (-1.28, 8.52)	0.15		
5-year outcomes								
Finger Tapping								
Dominant	180	-0.44 (-2.00, 1.12)	0.58	178	-0.35 (-2.34, 1.65)	0.73		
Non-dominant	180	0.71 (-0.72, 2.13)	0.33	178	1.11 (-0.69, 2.91)	0.23		
K-BIT								
Verbal Knowledge	180	0.77 (-0.06, 1.60)	0.07	178	0.72 (-0.34, 1.79)	0.18		
Matrices	180	-0.05 (-0.38, 0.28)	0.76	166	0.06 (-0.36, 0.47)	0.79		
Preschool Language Scale								
Auditory	180	0.32 (-0.47, 1.10)	0.42	178	0.15 (-0.84, 1.14)	0.77		
Verbal	180	0.41 (-0.52, 1.33)	0.39	178	0.27 (-0.92, 1.45)	0.66		
Total Language	180	0.72 (-0.81, 2.26)	0.35	178	0.41 (-1.54, 2.37)	0.68		
Woodcock-Johnson Test of Achievement								
Applied Problems	180	0.80 (-0.37, 1.96)	0.18	178	0.49 (-0.99, 1.98)	0.51		
Letter-Word	180	2.88 (1.21, 4.56)	0.001	178	2.83 (0.69, 4.97)	0.01		
Achenbach Child Behavior Checklist	180	-0.12 (-2.53, 2.30)	0.92	178	0.11 (-2.97, 3.19)	0.95		

* Adjusted for child sex, birth weight, Hollingshead SES, maternal KBIT, maternal age, family status, and PROCESS

Table 3

Adjusted* associations between telomere length at five years and cognitive outcomes at five years of age in the Seychelles Child Development Study Nutrition Cohort I.

Model	Full Sample (n=209)		
	N	β coefficient (95% CI)	P value
Finger Tapping			
Dominant	209	0.29 (−7.31, 7.89)	0.94
Non-dominant	209	3.01 (−3.84, 9.86)	0.37
K-BIT			
Verbal Knowledge	209	−1.36 (−5.19, 2.47)	0.48
Matrices	209	−0.99 (−2.61, 0.62)	0.23
Preschool Language Scale			
Auditory	209	−1.75 (−5.43, 1.93)	0.35
Verbal	209	−1.99 (−6.33, 2.35)	0.37
Total Language	209	−3.74 (−10.94, 3.46)	0.31
Woodcock-Johnson Test of Achievement			
Applied Problems	209	−0.54 (−6.16, 5.07)	0.85
Letter-Word	209	−1.55 (−9.63, 6.52)	0.70

* Adjusted for child sex, birth weight, Hollingshead SES, maternal KBIT, maternal age, family status, and PROCESS