Published in final edited form as:

Prev Med. 2017 November ; 104: 1–3. doi:10.1016/j.ypmed.2017.10.026.

Editorial for the special issue on behavior change, health, and health disparities 2017

Stephen T. Higgins

Vermont Center on Behavior and Health, Departments of Psychiatry and Psychological Science, University of Vermont, United States

Abstract

This Special Issue of *Preventive Medicine* (PM) is the 4th in a series on behavior change, health, and health disparities, a topic of critical importance to improving U.S. population health. The U.S. ranks near the bottom on measures of population health relative to other industrialized countries despite spending orders of magnitude more on health care than any other nation. Population health experts agree that the area of personal behavior, or lifestyle, such as substance abuse, physical inactivity/obesity, and non-adherence with medical regimens is the single largest contributor to this situation. These unhealthy behavior patterns disproportionately impact economically disadvantaged populations and other vulnerable populations and represent a major contributor to health disparities. Thus, behavior change represents an essential step in improving population health generally and curtailing health disparities more specifically. While perhaps more severe in the U.S., other industrialized countries are facing similar challenges with personal behavior patterns, adverse health impacts, and health disparities. Thus the topics discussed in this series have implications well beyond the U.S. In this 4th Special Issue we address (a) the potential health impacts of liberalizing laws on recreational marijuana use; (b) the ongoing challenge of tobacco use in vulnerable populations; and (b) the importance of weight management and physical activity in caring for vulnerable medical populations. Across each of these topics we include contributions from accomplished policymakers and scientists to acquaint readers with recent accomplishments and remaining knowledge gaps and challenges in these important topic areas.

Keywords

Behavior change; Population health; Chronic health conditions; Substance use disorders; Cannabis use: marijuana use; Marijuana legalization; Cigarette smoking; Tobacco use; *E*-cigarettes; Obesity; Weight management; Health disparities; Vulnerable populations

1. Introduction

This Special Issue of *Preventive Medicine* (PM) is the 4th in a series that focuses on behavior change, health, and health disparities. The first Special Issue appeared in November 2014 (<http://www.sciencedirect.com/science/journal/00917435/68/supp/C>), the 2nd in November 2015 (<http://www.sciencedirect.com/science/journal/00917435/80>) and the third in November 2016 (<http://www.sciencedirect.com/science/journal/00917435/92?sd=1>). Contributors to these Special Issues are selected from among participants in the annual conferences on Behavior Change, Health, and Health Disparities that are organized

by the Vermont Center on Behavior and Health (VCBH), a National Institutes of Health (NIH) and Food and Drug Administration (FDA) supported center of biomedical research excellence located at the University of Vermont (<http://www.uvm.edu/medicine/behaviorandhealth/>). This select subset of conferees was invited to contribute to this Special Issue because of their exemplary scholarship and outstanding conference presentations. Each contribution underwent thorough peer-review overseen by the Editor-in-Chief in coordination with the Guest Editor. Below I comment briefly on the rationale for organizing these annual conferences and associated publications as well the topics chosen for inclusion in this 4th Special Issue.

2. Unhealthy behavior patterns and risk for chronic disease and premature death

There is broad recognition that many individuals persist in behavior patterns (e.g., substance abuse, physical inactivity, unhealthy food choices, non-adherence with recommended medical regimens) that are strikingly harmful to their long-term health, and are a staggering financial burden on the health care systems of the U.S. and other industrialized countries (Higgins, 2015; Schroeder, 2016; Woolf and Aron, 2013). The current U.S. opioid addiction epidemic, which is having a devastating effect on U.S. population health, is an excellent example of this phenomenon (Compton et al., 2015; Sigmon et al., 2016; Volkow and Collins, 2017). Such unhealthy personal behavior patterns are estimated to be the largest contributor to premature deaths in the U.S. annually (~40%), with cigarette smoking alone being responsible for greater than 480,000 deaths/year in the U.S and 5 million globally (Higgins, 2015; Schroeder, 2016; Woolf and Aron, 2013). These behavior patterns increase risk for numerous chronic diseases, including cardiovascular disease, stroke, site-specific cancers, and type-2 diabetes. Importantly, because prevalence of these unhealthy behaviors is overrepresented among socioeconomically disadvantaged and other vulnerable populations, they are also important contributors to the unsettling problems of health disparities (Cutler and Lleras-Muney, 2010; Higgins and Chilcoat, 2009; Issacs and Schroeder, 2004; Kumanyika, 2012). Regarding costs to the health care system, direct medical costs of cigarette smoking in the U.S., for example, are estimated to exceed \$170 billion annually, and those associated with overweight and obesity in the U.S., as another example, approximately \$150 billion (Finkelstein et al., 2009; U.S. Department of Health and Human Services, 2014). These health-related behaviors are also major contributors to the dismal status of U.S. population health relative to other developed nations. For example, among the 34 developed nations with membership in the Organization for Economic Cooperation and Development, the U.S. ranks 25th in life expectancy, 29th in infant mortality, and 25th in maternal mortality (Higgins, 2015; Schroeder, 2016). These outcomes occur despite spending orders of magnitude more on healthcare than any other country. Importantly, when population health experts assess how to improve U.S. population health, the domain of personal behavior is identified as the area that offers the greatest potential for improvements (Higgins, 2014; Higgins, 2015; Schroeder, 2007; Schroeder, 2016). Hence, our overarching focuses on this topic area in the Vermont Center on Behavior and Health and in our annual conferences and associated Special Issues of *Preventive Medicine*.

3. Potential health impacts of legalizing recreational marijuana use

Legalization of marijuana use has gained considerable momentum in the U.S. with 28 states plus the District of Columbia (DC) legalizing medical marijuana use and 8 states plus DC legalizing recreational marijuana use, with similar liberalization of laws occurring in Canada and other countries (*NYTimes*, April 13, 2017). Such actions clearly have tremendous public health implications and it is important that those implications be considered using the best available scientific evidence. In this Special Issue we invited policy makers from Colorado (Ghosh et al., 2017, in this issue), the first U.S. state to legalize recreational marijuana use, Vermont (Chen and Searles, 2017, in this issue), a state currently considering legalization of recreational use, and the U.S.'s National Institute on Drug Abuse (Weiss and Wargo, 2017, in this issue) to provide a federal perspective on the health implications of legalizing recreational marijuana use. In addition to policy makers we invited contributions from scientific experts in the health impacts of marijuana use to address the implications of legalizing recreational marijuana use, including potential impacts on the epidemiology of marijuana use and risk perceptions among youth and adults (Carliner et al., 2017, in this issue), emergency medicine (Wang et al., 2017, in this issue), addiction risk (Budney and Borodovsky, 2017, in this issue), adolescent risks and potential interventions (Schuster et al., 2017; Walker, 2017, in this issue), and maternal and child health (Mark and Terplan, 2017, in this issue).

4. Tobacco use in vulnerable populations

While these recent political and legal developments regarding marijuana use demand careful scientific consideration in order to properly protect public health, we cannot to be distracted from other longstanding and more impactful behavioral health challenges, especially cigarette smoking. Cigarette smoking remains the largest contributor to premature death, being responsible for more than 480,000 premature deaths annually in the U.S. and 5,000,000 globally (U.S. Department of Health and Human Services, 2014). U.S. smoking prevalence dropped precipitously following the landmark U.S. Surgeon Generals report from approximately 42% in 1964 to 15% in 2015, a change estimated to have prevented 8 million premature deaths between 1964 and 2012 (Holford et al., 2014), and also netted tremendous health care cost savings (U.S. Department of Health and Human Services, 2014). Unfortunately those gains were disproportionately realized among the more affluent. Smoking prevalence decreased little or not at all among socioeconomically disadvantaged and other vulnerable populations (Higgins and Chilcoat, 2009; Chilcoat, 2009; Schroeder, 2016; U.S. Department of Health and Human Services, 2014). Thus, in this Special Issue we invited contributions on prevalence of cigarette smoking and use of other tobacco products among pregnant women (Higgins et al., 2017, in this issue; Kurti et al., 2017, in this issue), socioeconomically young adults (Villanti et al., 2017, in this issue), sexual minorities (Nayak et al., 2017, in this issue), and those residing in rural geographical areas (Doogan et al., 2017, in this issue). With the difficulties in promoting smoking cessation in these highly vulnerable populations, attention has appropriately turned to possible harm reduction strategies that might better protect these populations from the adverse health impacts of cigarette smoking, including substituting e-cigarettes for combusted tobacco cigarettes and reducing the maximal nicotine content of cigarettes to reduce the addiction potential of

cigarettes smoking. The latter is an especially timely topic in light of the recent announcement from U.S. Food and Drug Administration Commissioner Scott Gottlieb, MD, indicating that the agency is moving forward with consideration of a national policy that would reduce the standard for maximal nicotine content of cigarettes and other combusted tobacco products to very low levels in order to better protect the U.S. public health by decreasing addiction potential of cigarette smoking (U.S. Food and Drug Administration, 2017). We invited contributions assessing prevalence and risk perceptions of e-cigarettes in the U.S. adult population (Pericot-Valverde et al., 2017, in this issue), the relative dependence potential of e-cigarettes versus conventional tobacco cigarettes (Liu et al., 2017, in this issue), and the extent to which abruptly switching smokers of usual nicotine low nicotine content cigarettes to very low nicotine content cigarettes may disrupt behavioral and cognitive performance in regular smokers (Keith et al., 2017, in this issue).

5. Weight management and physical activity in vulnerable medical populations

Lastly, as was discussed above, physical inactivity, overweight, and obesity have a substantial adverse impact on individual and population health. There is growing recognition of the need to attend to these conditions as part of effective treatment for a wide variety of medical conditions. To address this important development, we invited contributions underscoring the importance of obesity as an unaddressed risk factor for coronary heart disease and the place of weight loss in effective rehabilitation from a cardiac event (Ades and Savage, 2017, in this issue), in the prevention and treatment of type-2 diabetes (Delahanty, 2017, in this issue), as a component of treating advanced cancer (Dittus et al., 2017, in this issue), and in efforts to prevent greater-than-recommended gestational weight gain and the serious adverse impacts of maternal obesity on perinatal outcomes (Phillips and Higgins, 2017, in this issue).

Acknowledgments

Funding

Preparation of this paper was supported in part by a Centers of Biomedical Research Excellence award P20GM103644 from the National Institute on General Medical Sciences, Tobacco Centers of Regulatory Science award P50DA036114-01 from the National Institute on Drug Abuse and Food and Drug Administration, and research awards R01HD075669 and R01HD078332 from the National Institute of Child Health and Human Development. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or Food and Drug Administration.

References

- Ades, PA., Savage, PD. Obesity in coronary heart disease: an unaddressed behavioral risk factor. *Prev Med.* 2017 Apr 13. <http://dx.doi.org/10.1016/j.ypmed.2017.04.013>. (Epub ahead of print)pii: S0091-7435(17)30133-0
- Budney, AJ., Borodovsky, JT. The potential impact of cannabis legalization on the development of cannabis use disorders. *Prev Med.* 2017 Jun 28. <http://dx.doi.org/10.1016/j.ypmed.2017.06.034>(Epub ahead of print)pii: S0091-7435(17)30238-4
- Carliner, H., Brown, QL., Sarvet, AL., Hasin, DS. Cannabis use, attitudes, and legal status in the U.S.: a review. *Prev Med.* 2017 Jul 11. <http://dx.doi.org/10.1016/j.ypmed.2017.07.008>. (Epub ahead of print)pii: S0091-7435(17)30255-4

- Chen, H., Searles, JS. Health considerations in regulating marijuana use in Vermont. *Prev Med.* 2017 Jun 8. <http://dx.doi.org/10.1016/j.ypmed.2017.06.004>. (Epub ahead of print)pii: S0091-7435(17)30208-6
- Chilcoat HD. An overview of the emergence of disparities in smoking prevalence, cessation, and adverse consequences among women. *Drug Alcohol Depend.* Oct; 2009 104(Suppl 1):S17–S23. Epub 2009 Jul 24. [PubMed: 19632070]
- Compton, WM., Boyle, M., Wargo, E. Prescription opioid abuse: problems and responses; *Prev Med.* 2015 Nov. p. 5-9.<http://dx.doi.org/10.1016/j.ypmed.2015.04.003>. (Epub 2015 Apr 11)
- Cutler DM, Lleras-Muney A. Understanding differences in health behaviors by education. *J Health Econ.* 2010 Jan; 29(1):1–28. [PubMed: 19963292]
- Delahanty, LM. Weight loss in the prevention and treatment of diabetes. *Prev Med.* 2017 Jul 28. <http://dx.doi.org/10.1016/j.ypmed.2017.07.022>. (Epub ahead of print)pii: S0091-7435(17)30269-4
- Dittus, KL., Grambling, RE., Ades, PA. Exercise interventions for individuals with advanced cancer: a systematic review. *Prev Med.* 2017 Jul 15. <http://dx.doi.org/10.1016/j.ypmed.2017.07.015>. (Epub ahead of print)pii: S0091-7435(17)30262-1
- Doogan, NJ., Roberts, ME., Wewers, ME., Stanton, CA., Keith, DR., Gaalema, DE., Kurti, AN., Redner, R., Cepeda-Benito, A., Bunn, JY., Lopez, AA., Higgins, ST. A growing geographic disparity: Rural and urban cigarette smoking trends in the United States. *Prev Med.* 2017 Mar 16. <http://dx.doi.org/10.1016/j.ypmed.2017.03.011>. (Epub ahead of print)pii: S0091-7435(17)30106-8
- Finkelstein, EA., Linnan, LA., Tate, DF., Leese, PJ. A longitudinal study on the relationship between weight loss, medical expenditures, and absenteeism among overweight employees in the WAY to health study; *J Occup Environ Med.* 512009 Dec. p. 1367-1373.<http://dx.doi.org/10.1097/JOM.0b013e3181c2bb56>
- Ghosh, TS., Vigil, DI., Maffey, A., Tolliver, R., Van Dyke, M., Kattari, L., Krug, H., Reed, JK., Wolk, L. Lessons learned after three years of legalized, recreational marijuana: the Colorado experience. *Prev Med.* 2017 Feb 21. <http://dx.doi.org/10.1016/j.ypmed.2017.02.021>. (Epub ahead of print)pii: S0091-7435(17)30076-2
- Higgins ST. Behavior change, health, and health disparities: an introduction. *Prev Med.* 2014; 68:1–4. [PubMed: 25456804]
- Higgins, ST. Editorial: 2nd special issue on behavior change, health, and health disparities; *Prev Med.* 2015 Nov. p. 1-4.<http://dx.doi.org/10.1016/j.ypmed.2015.07.020>
- Higgins ST, Chilcoat HD. Women and smoking: an interdisciplinary examination of socioeconomic influences. *Drug Alcohol Depend.* 2009 Oct 1.104(Suppl 1) <http://dx.doi.org/10.1016/j.jhealeco.2009.10.003>. (Epub 2009 Oct 31) S1-5. Epub 2009 Jul 8. doi: 10.1016/j.drugalcdep.2009.06.006
- Higgins, ST., Redner, R., Arger, CA., Kurti, AN., Priest, JS., Bunn, JY. Use of higher-nicotine/tar-yield (regular full-flavor) cigarettes is associated with nicotine dependence and smoking during pregnancy among U.S. women. *Prev Med.* 2017 Aug 5. <http://dx.doi.org/10.1016/j.ypmed.2017.07.029>. (Epub ahead of print)pii: S0091-7435(17)30276-1
- Holford TR, Meza R, Warner KE, Meernik C, Jeon J, Moolgavkar SH, Levy DT. Tobacco control and the reduction in smoking-related premature deaths in the United States, 1964–2012. *JAMA.* 2014; 311:164–171. <http://dx.doi.org/10.1001/jama.2013.285112>. [PubMed: 24399555]
- Issacs SL, Schroeder SA. Class—the ignored determinant of the nation’s health. *N Engl J Med.* 2004 Sep 9; 351(11):1137–1142. <http://dx.doi.org/10.1056/NEJMs040329>. [PubMed: 15356313]
- Keith, DR., Kurti, AN., Davis, DR., Zvorsky, IA., Higgins, ST. A review of the effects of very low nicotine content cigarettes on behavioral and cognitive performance. *Prev Med.* 2017 Jun 21. <http://dx.doi.org/10.1016/j.ypmed.2017.06.016>. (Epub ahead of print)pii: S0091-7435(17)30221-9
- Kumanyika, S. Health disparities research in global perspective: new insights and new directions; *Annu Rev Public Health.* 2012 Apr. p. 1-5.<http://dx.doi.org/10.1146/annurev-publhealth-112911-155504>. (Epub 2012 Jan 3)
- Kurti, AN., Redner, R., Lopez, AA., Keith, DR., Villanti, AC., Stanton, CA., Gaalema, DE., Bunn, JY., Doogan, NJ., Cepeda-Benito, A., Roberts, ME., Phillips, J., Higgins, ST. Tobacco and nicotine delivery product use in a national sample of pregnant women. *Prev Med.* 2017 Aug 5. <http://dx.doi.org/10.1016/j.ypmed.2017.07.030>. (Epub ahead of print)pii: S0091-7435(17)30277-3

- Liu, G., Wasserman, E., Kong, L., Foulds, J. A comparison of nicotine dependence among exclusive E-cigarette and cigarette users in the PATH study. *Prev Med.* 2017 Apr 4. <http://dx.doi.org/10.1016/j.ypmed.2017.04.001>. (Epub ahead of print)pii: S0091-7435(17)30122-6
- Mark, K., Terplan, M. Cannabis and pregnancy: maternal child health implications during a period of policy liberalization. *Prev Med.* 2017 May 18. <http://dx.doi.org/10.1016/j.ypmed.2017.05.012>. (Epub ahead of print)pii: S0091-7435(17)30175-5
- Nayak, P., Salazar, LF., Kota, KK., Pechacek, TF. Prevalence of use and perceptions of risk of novel and other alternative tobacco products among sexual minority adults: results from an online national survey, 2014–2015. *Prev Med.* 2017 Jun 1. <http://dx.doi.org/10.1016/j.ypmed.2017.05.024>. (Epub ahead of print)pii: S0091-7435(17)30189-5
- NY Times. [Accessed date: 16 October 2017 April 13] Trudeau unveils bill legalizing recreational marijuana in Canada. 2017. https://www.nytimes.com/2017/04/13/world/canada/trudeau-marijuana.html?_r=0
- Pericot-Valverde, I., Gaalema, DE., Priest, JS., Higgins, ST. E-cigarette awareness, perceived harmfulness, and ever use among U.S. adults. *Prev Med.* 2017 Jul 17. <http://dx.doi.org/10.1016/j.ypmed.2017.07.014>. (Epub ahead of print)pii: S0091-7435(17)30261-X
- Phillips, JK., Higgins, ST. Applying behavior change techniques to weight management during pregnancy: Impact on perinatal outcomes. *Prev Med.* 2017 Jul 28. <http://dx.doi.org/10.1016/j.ypmed.2017.07.023>. (Epub ahead of print)pii: S0091-7435(17)30270-0
- Schroeder SA. Shattuck lecture: we can do better—improving the health of the American people. *N Engl J Med.* 2007; 357(12):1221–1228. [PubMed: 17881753]
- Schroeder SA. American health improvement depends upon addressing class disparities. *Prev Med.* 2016 Nov.(92):6–15.
- Schuster, RM., Fontaine, M., Nip, E., Zhang, H., Hanly, A., Evins, EA. Prolonged cannabis withdrawal in young adults with lifetime psychiatric illness. *Prev Med.* 2017 Feb 27. <http://dx.doi.org/10.1016/j.ypmed.2017.02.019>. (Epub ahead of print)pii: S0091-7435(17)30075-0
- Sigmon SC, Ochalek TA, Meyer AC, Hruska B, Heil SH, Badger GJ, Rose G, Brooklyn JR, Schwartz RP, Moore BA, Higgins ST. Interim buprenorphine vs. waiting list for opioid dependence. *N Engl J Med.* 2016; 375(25):2504–2505. [PubMed: 28002704]
- U.S. Department of Health and Human Services. The Health Consequences of Smoking—50 Years of Progress: a Report of the Surgeon General. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; Atlanta, GA: 2014.
- U.S. Food and Drug Administration. [Accessed date: 1 October 2017] FDA news release: FDA announces comprehensive regulatory plan to shift trajectory of tobacco-related disease, death. 2017. <https://www.fda.gov/newsevents/newsroom/pressannouncements/ucm568923.htm>
- Villanti, AC., Johnson, AL., Rath, JM. Beyond education and income: Identifying novel socioeconomic correlates of cigarette use in U.S. young adults. *Prev Med.* 2017 Jun 21. <http://dx.doi.org/10.1016/j.ypmed.2017.06.019>. (Epub ahead of print)pii: S0091-7435(17)30223-2
- Volkow ND, Collins FS. The role of science in addressing the opioid crisis. *N Engl J Med.* 2017; 377:391–394. <http://dx.doi.org/10.1056/NEJMSr1706626>. (Epub 2017 May 31). [PubMed: 28564549]
- Walker, DD. Legalization of cannabis: Considerations for intervening with adolescent consumers. *Prev Med.* 2017 Jul 17. <http://dx.doi.org/10.1016/j.ypmed.2017.07.010>. (Epub ahead of print)pii: S0091-7435(17)30257-8
- Wang, GS., Hall, K., Vigil, D., Banerji, S., Monte, A., VanDyke, M. Marijuana and acute health care contacts in Colorado. *Prev Med.* 2017 Mar 30. <http://dx.doi.org/10.1016/j.ypmed.2017.03.022>. (Epub ahead of print)pii: S0091-7435(17)30120-2
- Weiss, SRB., Wargo, EM. Commentary: navigating the complexities of marijuana. *Prev Med.* 2017 Jun 23. <http://dx.doi.org/10.1016/j.ypmed.2017.06.026>. (Epub ahead of print)pii: S0091-7435(17)30230-X
- Woolf, SH., Aron, L. US Health in International Perspective: Shorter Lives, Poorer Health. National Academies Press; 2013.