Published in final edited form as:

Synthesis (Stuttg). 2018 ; 50(3): 539–547. doi:10.1055/s-0036-1591500.

A general protocol for radical anion [3 + 2] cycloaddition enabled by tandem Lewis acid photoredox catalysis

Adrian G Amador^a, Evan M. Sherbrook^a, Zhan Lu^b, and Tehshik P. Yoon^{*,a}

^aDepartment of Chemistry, University of Wisconsin-Madison, 1101 University Avenue, Madison, WI 53706 USA

^bDepartment of Chemistry, Zhejiang University, Hangzhou 310058, China

Abstract

We report herein a method for intermolecular [3 + 2] cycloaddition between aryl cyclopropyl ketones and alkenes involving the combination of Lewis acid and photoredox catalysis. In contrast to other more common methods for [3 + 2] cycloaddition, these conditions operate using a broad range of both electron-rich and electron-deficient reaction partners. The critical factors predicting the success of these reactions is the redox potential of the cyclopropyl ketone and the ability of the alkene to stabilize a key radical intermediate.

Graphical Abstract

Keywords

Cycloaddition; photoredox; photocatalysis; cyclopropane; cyclopentane

Five-membered carbocycles feature prominently in a variety of complex natural products and many other small-molecule compounds of biological interest. As a result, significant research efforts have been directed towards the discovery of selective and flexible methods to synthesize cyclopentanes.¹ While numerous successful strategies have been developed (e.g. cyclizations,² ring expansions,³ etc), [3 + 2] cycloaddition reactions are particularly appealing because they provide a direct means to access a diverse array of cyclopentanoid frameworks from relatively simple building blocks. The most common strategies for [3 + 2]

* indicates the main/corresponding author. tyoon@chem.wisc.edu.

Supporting Information

Supporting Information Supporting information for this article is available online at <https://doi.org/10.1055/s-0036-1591500>.

cycloadditions involve 1,3-dipolar and zwitterionic reagents,⁴ vinyl carbenoids,⁵ or reductive cycloadditions.⁶ While the existence of these various methods offers flexibility in the synthesis of complex cyclopentane-containing compounds, they are generally sensitive to the polarity of the coupling partners and are often incompatible with polar functional groups.

Organoradical intermediates can be synthetically attractive because they often react with both electron-rich and electron-poor partners and are generally tolerant of a range of common functional groups. Our group has been interested in the development of cycloaddition methods that take advantage of radical anion intermediates generated using visible light photocatalysis.⁷ We recently reported two photocatalytic methods that result in the formal [3+2] cycloaddition of aryl cyclopropyl ketones with alkenes (Scheme 1).^{8,9,10} In these reactions, a Lewis acid co-catalyst is used to activate the cyclopropyl ketone towards one-electron reduction by a photogenerated Ru^{I} catalyst; the resulting ketyl radical reacts with an alkene coupling partner to afford a new five-membered carbocycle. The correct choice of Lewis acid and careful tuning of the coordination sphere of the Lewis acid proved to be critical for the success of these reactions. Unfortunately, the scope of these reactions has proven to be somewhat limited: the optimal conditions for intramolecular [3 + 2] cycloadditions did not translate smoothly to intermolecular reactions, and the optimal conditions for enantioselective intermolecular cycloadditions unfortunately required the use of relatively reactive cyclopropanes and electron-rich alkene partners to achieve reasonable yields. In order to perform a more comprehensive survey of the synthetic capabilities and limitations of this unique strategy for [3 + 2] cycloaddition, we became interested in developing robust conditions that allow for efficient racemic [3 + 2] cycloadditions with a relatively broad substrate scope. The results of these investigations are described herein.

Our initial optimization studies focused on the reaction of simple unsubstituted cyclopropyl phenyl ketone **1** with a relatively unpolarized styrene partner. We first assessed this reaction under conditions previously reported for the intramolecular version of this reaction, employing 1 equiv. of the Lewis acid $\text{La}(\text{OTf})_3$, 2.5 mol % of the photocatalyst $\text{Ru}(\text{bpy})_3\text{Cl}_2$, and 5 equiv. of TMEDA as reductive quencher. An initial experiment provided the desired cycloadduct in 44% yield after 24 h (Table 1, entry 1). Alternative solvents showed no improvements in the overall efficiency of the reaction (entries 2–4). The identity of the Lewis acid co-catalyst had a more positive effect. A screen of Lewis acids revealed that lanthanide triflates are uniquely competent in this transformation and that $\text{Gd}(\text{OTf})_3$ offers the greatest yield among those screened (entries 5–7). Alternate tertiary amine additives proved to be less effective than TMEDA, consistent with its proposed dual role as a chelating ligand and a reductive quencher in this reaction (entries 8 and 9). The loading of $\text{Gd}(\text{OTf})_3$ could be decreased to 50 mol% with a modest decrease in yield (entry 10), which could be restored by lowering the concentration of TMEDA (entry 11). Lowering the loading of the photocatalyst to 1 mol% did not appear to have an appreciable influence on the effectiveness of the reaction. These optimal conditions provided the desired cycloadduct in 71% yield after 20 h (entry 12). Control reactions verified that the photocatalyst, Lewis acid, quencher, and light source are all required for any product formation, validating the photocatalytic nature of this reaction.

Having identified optimized reaction conditions, we next explored the generality of this transformation. The scope with respect to the cyclopropane substrate proved to be relatively broad (Figure 1). A heteroaryl cyclopropyl ketone was well tolerated (**3**).¹¹ In general, aryl and alkyl substituents on the cyclopropane provided superior reactivity compared to the unsubstituted cyclopropyl phenyl ketone, presumably due to the increased stability of the radical resulting from homolytic opening of the cyclopropane ring (**4** and **5**). Multiple alkyl substitutions were also well tolerated, allowing for the construction of quaternary centers (**6** and **7**). Interestingly this cycloaddition also tolerated electron-withdrawing substituents (e.g., ester and ketone, **8** and **9**). This is in stark contrast to the majority of methodologies available for [3 + 2] cycloadditions which tend to be very sensitive to the polarity of each coupling partner. Cyclopropanes substituted on all three carbons react smoothly (**10**). When ring-opening of an unsymmetrically substituted cyclopropane can result in the formation of two different radical intermediates, the major product is formed with excellent regioselectivity consistent with reaction through the more stable radical (**11**).

The scope with respect to the alkene reaction partner is summarized in Figure 2. A variety of alkenes participate in this reaction, with the limitation that the olefin must bear a radical-stabilizing substituent. Thus, styrenes are good coupling partners for this transformation: both electron-rich and electron-poor styrenes react smoothly (**12** and **13**). Neither reactive halides nor the introduction of steric bulk on the aryl ring of the styrene adversely impact the transformation (**14** and **15**). Substitutions at the α -position of styrenes were also well tolerated (**16–18**). In line with our design plan, a range of alkene reaction partner with quite varied polarities are readily tolerated. Very electron-rich heteroatom-substituted coupling partners such as a vinylcarbazole (**19**) and a silyl enol ether (**20**) readily undergo [3 + 2] cycloaddition. Electron-deficient alkenes such as acrylates (**21**) also worked well under these conditions. Interestingly 1,3-butadiene was also an efficient coupling partners (**22**). The broad scope of the olefin coupling partner thus represents a complementary feature to many of the existing [3 + 2] cycloaddition methodologies. However, 1,2-disubstituted alkenes such as β -methylstyrene failed to produce identifiable quantities of cycloadduct and represent a limitation of this method.

Figure 3 outlines a mechanism for this intermolecular [3 + 2] cycloaddition analogous to that previously proposed for the intramolecular method. Photoexcitation of $\text{Ru}(\text{bpy})_3^{2+}$ with visible light and subsequent reductive quenching by TMEDA results in the generation of $\text{Ru}(\text{bpy})_3^{1+}$, which can then reduce the [Gd]–cyclopropane complex ([Gd]–**23**). The ketyl radical anion then undergoes reversible ring opening to a distonic radical anion. This reactive intermediate then undergoes stepwise [3 + 2] cycloaddition with an olefin reaction partner to produce the ketyl radical of the product. Oxidation to produce the neutral cycloadduct occurs either through reduction of the radical cation of TMEDA or through a radical chain process¹² to produce another equivalent of the radical anion of [Gd]–**23**. The rapid reversibility of the ring-opening of ketyl radical anions has been studied extensively by Tanko and coworkers.¹³ To support the relevance of this process in this transformation, diastereomerically pure *cis*-**23** was observed to undergo isomerization to *trans*-**23** within 2 h upon exposure to the reaction conditions (Scheme 2). This isomerization only occurs in the presence of photocatalyst, Lewis acid, and TMEDA, which suggests that the isomerization is

occurring through an electron-transfer process. Additionally, we observed the formation of 10% yield of reductive ring cleavage product **24**, which constitutes a significant byproduct in many of the lower-yielding cycloadditions.

In conclusion, we have identified conditions for an efficient intermolecular [3 + 2] cycloaddition between a range of simple aryl cyclopropyl ketones and olefins. The choice of Lewis acid and loading of the reductive quencher proved critical in attaining efficient reactivity. This [3 + 2] cycloaddition provides a valuable complement to the existing methodologies in the literature for the construction of five-membered carbocycles. Particularly notable is the relative insensitivity of this cycloaddition to the polarity of the substrates. The critical constraints are the reduction potential of the cyclopropane and the ability of the olefin coupling partner to stabilize a key radical intermediate in the step-wise cycloaddition. This method should thus prove to be a valuable tool for the synthesis of a variety of structurally complex five-membered ring compounds.

The experimental section has no title; please leave this line here.

For details of equipment, sources of chemicals etc., see the Supporting Information.

General Procedure

An oven-dried Schlenk tube equipped with a magnetic stir-bar was charged with Gd(OTf)₃ (0.5 equivalents). A flame-dried vial was charged with cyclopropane (1 equiv.), alkene (5 equiv.), Ru(bpy)₃(PF₆)₂ (1 mol %), TMEDA (1.5 equiv.) and MeCN (0.2 M concentration). The contents of the vial were then transferred to the Schlenk tube. The reaction mixture was then thoroughly degassed through three freeze-pump-thaw cycles, then backfilled with N₂. The reaction flask was then placed in front of a 23 W (1380 lumen) compact fluorescent lamp and stirred at room temperature. Upon consumption of starting material, the reaction was diluted with 1:1 Et₂O/pentanes and passed through a short plug of silica. The filtrate was concentrated and the residue purified by column chromatography.

Phenyl(*trans* 2-phenylcyclopentyl)methanone (**2**)

Yield: 71.1 mg (71%, 8:1 d.r.); *R*_f = 0.3 (Et₂O/pentanes, 1:9). ¹H and ¹³C NMR were consistent with the literature¹⁴

(1-methyl-1H-imidazol-2-yl)(2-phenylcyclopentyl)methanone (**3**)

Colorless oil; Yield: 66.1 mg (65%, 6:1 d.r.); *R*_f = 0.3 (Acetone/pentanes, 1:3).

IR (ATR): 2983, 1705, 1274, 1080 cm⁻¹.

¹H NMR (CDCl₃, 500 MHz) δ 7.31 – 7.27 (m, 2H), 7.23 (dd, *J* = 8.5, 6.9 Hz, 2H), 7.16 – 7.09 (m, 2H), 7.09 (d, *J* = 0.9 Hz, 1H), 6.96 (d, *J* = 0.9 Hz, 1H), 4.32 (td, *J* = 9.6, 8.0 Hz, 1H), 3.95 (s, 3H), 3.61 (td, *J* = 9.7, 7.8 Hz, 1H), 2.45 – 2.33 (m, 1H), 2.30 – 2.18 (m, 1H), 2.01 – 1.88 (m, 1H), 1.91 – 1.77 (m, 3H).

¹³C NMR (CDCl₃, 101 MHz): δ = 194.72, 144.32, 143.19, 129.05, 128.29, 127.48, 126.89, 126.01, 54.54, 47.78, 36.19, 35.55, 32.66, 25.21.

HRMS (ESI): m/z $[M+H]^+$ calcd for $[C_{16}H_{19}N_2O]^+$: 255.1492; found: 255.1492.

2,4-diphenylcyclopentyl(phenyl)methanone (4)

Colorless oil; Yield: 111.1 mg (74%, 5:1 d.r.); R_f = 0.3 (Et₂O/pentanes, 1:9).

IR (ATR): 2978, 1718, 1305, 1272, 1150 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.84 (d, J = 7.3 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.42 – 7.13 (m, 12H), 4.16 – 3.82 (m, 2H), 3.58 (dt, J = 17.9, 9.0 Hz, 1H), 2.69 – 2.52 (m, 1H), 2.48 – 2.28 (m, 2H), 2.26 – 2.04 (m, 1H).

¹³C NMR (CDCl₃, 101 MHz): δ = 200.51, 143.26, 136.96, 136.34, 132.98, 132.90, 128.63, 128.54, 128.47, 128.36, 127.45, 127.33, 126.49, 54.72, 47.44, 45.57, 37.52, 34.81.

HRMS (ESI): m/z $[M+H]^+$ calcd for $[C_{24}H_{22}O]^+$: 326.1671; found: 326.1670.

1-(3-Benzoyl-4-phenylcyclopentyl)ethanone (9)

Colorless oil; Yield: 21.1 mg (36%, 5:1 d.r.); R_f = 0.3 (Et₂O/pentanes, 1:15).

IR (ATR): 2966, 1708, 1683, 1262, 1114 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.72 (d, J = 8.6 Hz, 2H), 7.45 – 7.38 (m, 1H), 7.30 (dd, J = 8.4, 7.1 Hz, 2H), 7.23 – 7.04 (m, 4H), 6.97 – 6.84 (m, 1H), 3.85 (q, J = 8.7 Hz, 1H), 3.60 (q, J = 8.7 Hz, 1H), 3.19 (qd, J = 8.5, 5.9 Hz, 1H), 2.55 – 2.32 (m, 2H), 2.23 – 2.18 (m, 1H), 2.15 (s, 3H), 2.04 (dt, J = 13.3, 9.4 Hz, 1H).

¹³C NMR (CDCl₃, 101 MHz): δ = 208.91, 200.61, 143.46, 136.73, 133.00, 128.59, 128.49, 128.40, 127.27, 126.55, 54.47, 51.45, 47.51, 36.23, 33.80, 28.41.

HRMS (ESI): m/z $[M+H]^+$ calcd for $[C_{20}H_{21}O_2]^+$: 293.1536; found: 293.1535.

(4-Methyl-2-phenylcyclopentyl)(phenyl)methanone (5)

White solid (m.p. = 44.9–51 °C); Yield: 49.1 mg (93%, 11:1 d.r.); R_f = 0.3 (Et₂O/pentanes, 1:10).

IR (ATR): 2957, 2868, 1681, 1602, 1451, 1266, 1012 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.74 (d, J = 7.4 Hz, 2H), 7.42 (t, J = 7.4 Hz, 1H), 7.30 (t, J = 7.7 Hz, 2H), 7.24 – 7.11 (m, 4H), 7.13 – 7.00 (m, 1H), 3.90 – 3.69 (m, 2H), 2.48 – 2.22 (m, 2H), 2.02 (dt, J = 15.2, 7.8 Hz, 1H), 1.82 (ddd, J = 13.0, 8.8, 7.2 Hz, 1H), 1.55 – 1.43 (m, 1H), 1.03 (d, J = 6.5 Hz, 3H).

¹³C NMR (CDCl₃, 101 MHz): δ = 201.90, 145.50, 132.80, 128.55, 128.44, 128.41, 127.37, 127.30, 126.07, 55.96, 46.89, 42.50, 41.01, 34.44, 20.78.

HRMS (ESI): m/z $[M+NH_4]^+$ calcd for $[C_{19}H_{24}ON]^+$: 282.1852; found: 282.1853.

(4,4-Dimethyl-2-phenylcyclopentyl)(phenyl)methanone (6)

Yield: 52.3 mg (94%, 2:1 d.r.). R_f = 0.3 (Et₂O/pentanes, 1:10)

Major diastereomer—White solid (m.p. = 105.3–107.8 °C);

IR (ATR): 2941, 2863, 1678, 1448, 1283, 1015 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.82 (d, J = 8.4 Hz, 2H), 7.48 (t, J = 7.4 Hz, 1H), 7.37 (t, J = 8.0 Hz, 2H), 7.29 – 7.18 (m, 4H), 7.12 (t, J = 7.0 Hz, 1H), 4.04 – 3.84 (m, 2H), 2.12 (dd, J = 13.0, 9.2 Hz, 1H), 2.04 (dd, J = 12.7, 6.8 Hz, 1H), 1.86 (t, J = 11.9 Hz, 1H), 1.73 (dd, J = 13.1, 7.8 Hz, 1H), 1.21 (s, 3H), 1.13 (s, 3H).

¹³C NMR (CDCl₃, 101 MHz): δ = 201.72, 144.08, 137.00, 132.80, 128.43, 128.43, 128.38, 127.39, 126.15, 54.88, 49.42, 46.79, 46.76, 39.02, 30.59, 29.55

HRMS (ESI): m/z [M+H]⁺ calcd for [C₂₀H₂₃O]⁺: 279.1743; found: 279.1743.

Minor diastereomer—White solid (m.p. = 71.6–73.4 °C);

IR (ATR): 2953, 1927, 2866, 1679, 1463, 1448, 1367, 1221 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.88 (d, J = 8.2 Hz, 2H), 7.62 (d, J = 8.3 Hz, 2H), 7.33 – 7.06 (m, 5H), 3.98 (q, J = 9.3 Hz, 1H), 3.62 (ddd, J = 11.3, 9.4, 7.6 Hz, 1H), 3.02 (ddd, J = 16.5, 9.2, 7.3 Hz, 1H), 2.63 – 2.38 (m, 2H), 2.25 – 2.00 (m, 2H), 1.49 (s, 6H).

¹³C NMR (CDCl₃, 101 MHz): δ = 202.23, 142.11, 136.99, 132.03, 128.46, 127.96, 127.93, 127.71, 125.98, 50.97, 48.93, 48.45, 43.26, 38.47, 29.40, 28.53

HRMS (ESI): m/z [M+H]⁺ calcd for [C₂₀H₂₃O]⁺: 279.1743; found: 279.1741.

Phenyl(3-phenylspiro[4.4]nonan-2-yl)methanone (7)

Yield: 51.1 mg (84%, 1:1 d.r.). R_f = 0.3 (Et₂O/pentanes, 1:30)

Major diastereomer—Colorless oil.

IR (ATR): 2945, 2858, 1680, 1448, 1216 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.83 (d, J = 7.1 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.38 (t, J = 7.6 Hz, 2H), 7.32 – 7.18 (m, 4H), 7.13 (t, J = 6.9 Hz, 1H), 3.97 – 3.82 (m, 2H), 2.21 (dd, J = 12.9, 9.6 Hz, 1H), 2.12 (dd, J = 12.6, 6.5 Hz, 1H), 2.02 – 1.88 (m, 1H), 1.83 (dd, J = 12.9, 6.9 Hz, 1H), 1.77 – 1.49 (m, 8H).

¹³C NMR (CDCl₃, 101 MHz): δ = 201.76, 144.22, 136.98, 132.77, 128.44, 128.42, 128.36, 127.38, 126.13, 54.64, 50.57, 47.52, 46.91, 44.98, 40.28, 39.78, 24.40, 24.34.

HRMS (ESI): m/z [M+H]⁺ calcd for [C₂₂H₂₅O]⁺: 305.1900; found: 305.1899.

Minor diastereomer—Colorless oil.

IR (ATR): 2950, 2863, 1679, 1448, 1220, 1023 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.56 (d, J = 7.2 Hz, 2H), 7.36 (t, J = 7.3 Hz, 1H), 7.29 – 7.18 (m, 2H), 6.97 (7.02 – 6.90 (m, 5H), 4.34 – 4.21 (m, 1H), 3.71 (td, J = 10.4, 7.6 Hz, 1H), 2.39 (dd, J = 13.0, 9.0 Hz, 1H), 2.16 – 1.97 (m, 2H), 1.90 – 1.47 (m, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 202.33, 142.17, 138.18, 132.06, 128.39, 127.97, 127.96, 127.72, 125.97, 50.89, 49.81, 48.59, 46.82, 41.52, 39.26, 38.89, 24.81, 24.66.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{22}\text{H}_{25}\text{O}]^+$: 305.1900; found: 305.1900.

tert-butyl 3-benzoyl-4-phenylcyclopentanecarboxylate (8)

Yield: 131.8 mg (94%, 3:1 d.r.). R_f = 0.3 (Et_2O /pentanes, 1:5)

Major diastereomer—White solid (m.p. = 97.1–98.5 $^{\circ}\text{C}$).

IR (ATR): 2980, 1722, 1682, 1368, 1265, 1151 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.81 (d, J = 7.9 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.37 (t, J = 7.7 Hz, 2H), 7.28–7.22 (m, 4H), 7.16 (dq, J = 9.1, 4.3 Hz, 1H), 3.94 – 3.73 (m, 2H), 3.17 – 3.00 (m, 1H), 2.58 – 2.44 (m, 2H), 2.22 (dt, J = 13.3, 9.0 Hz, 1H), 2.13 (dt, J = 13.1, 9.0 Hz, 1H), 1.46 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.56, 174.21, 143.74, 136.85, 132.92, 128.52, 128.47, 128.39, 127.31, 126.42, 80.52, 54.75, 47.14, 44.46, 37.31, 35.36, 28.08.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{23}\text{H}_{27}\text{O}_3]^+$: 351.1955; found: 351.1955.

Minor diastereomer—White solid (m.p. = 111.5–113.0 $^{\circ}\text{C}$).

IR (ATR): 2981, 1719, 1684, 1365, 1262, 1150 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.82 (d, J = 7.6 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.37 (t, J = 7.7 Hz, 2H), 7.28–7.22 (m, 4H), 7.15 (t, J = 6.8 Hz, 1H), 3.99 (q, J = 9.3 Hz, 1H), 3.68 (dt, J = 11.0, 8.4 Hz, 1H), 3.00 (ddd, J = 16.6, 9.2, 7.5 Hz, 1H), 2.62 – 2.43 (m, 2H), 2.14 (dt, J = 13.2, 8.9 Hz, 2H), 1.48 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.32, 174.92, 142.96, 136.62, 133.04, 128.51, 128.50, 128.50, 127.42, 126.53, 80.49, 53.70, 48.34, 44.20, 38.84, 34.83, 28.13.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{23}\text{H}_{30}\text{O}_3\text{N}]^+$: 368.2220; found: 368.2220.

Di-tert-butyl 3-benzoyl-4-phenylcyclopentane-1,2-dicarboxylate (10)

Yield: 84.7 mg (94%, 2:1 d.r.). R_f = 0.3 (Et_2O /pentanes, 1:5)

Major diastereomer—White solid (m.p. = 96–97 $^{\circ}\text{C}$).

IR (ATR): 2977, 1724, 1681, 1393, 1368, 1257, 1158 cm^{-1} .

¹H NMR (CDCl₃, 400 MHz): δ = 7.70 (d, J = 7.7 Hz, 2H), 7.43 (t, J = 7.3 Hz, 1H), 7.29 (t, J = 7.7 Hz, 2H), 7.24–7.2 (m, 4H), 7.14 (dq, J = 8.2, 5.5, 4.8 Hz, 1H), 4.17 (t, J = 9.5 Hz, 1H), 3.64 (dt, J = 25.1, 8.8 Hz, 2H), 3.36 (dq, J = 9.3, 5.5 Hz, 1H), 2.51 (ddd, J = 13.4, 8.1, 5.5 Hz, 1H), 2.27 (dt, J = 13.2, 9.8 Hz, 1H), 1.49 (s, 9H), 1.25 (s, 9H).

¹³C NMR (CDCl₃, 101 MHz): δ = 200.49, 173.03, 172.54, 141.91, 137.22, 132.93, 128.59, 128.40, 128.29, 127.20, 126.81, 81.21, 80.89, 57.00, 53.04, 49.76, 47.34, 37.16, 28.05, 27.78.

HRMS (ESI): m/z [M+H]⁺ calcd for [C₂₈H₃₅O₅]⁺: 451.2479; found: 451.2479.

Minor diastereomer—White solid (m.p. = 83.7–85.1 °C).

IR (ATR): 2978, 2936, 1725, 1680, 1367, 1257, 1154 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.57 (d, J = 7.0 Hz, 2H), 7.46–7.33 (m, 1H), 7.27–7.23 (m, 2H), 7.07–6.91 (m, 5H), 4.50 (dd, J = 10.0, 7.7 Hz, 1H), 3.93 (dd, J = 10.3, 7.7 Hz, 1H), 3.70 (td, J = 10.0, 7.3 Hz, 1H), 3.10 (td, J = 10.5, 7.9 Hz, 1H), 2.57–2.36 (m, 2H), 1.53 (s, 9H), 1.43 (s, 9H).

¹³C NMR (CDCl₃, 101 MHz): δ = 172.61, 132.42, 128.32, 128.07, 128.05, 127.94, 126.59, 53.83, 48.52, 48.27, 36.85, 28.15, 28.08.

HRMS (ESI): m/z [M+H]⁺ calcd for [C₂₈H₃₅O₅]⁺: 451.2479; found: 451.2478.

tert-Butyl 5-benzoyl-2,2-dimethyl-4-phenylcyclopentanecarboxylate (11)

White solid (m.p. = 134.7–138.5 °C); Yield: 74.2 mg (97%, 20:1 d.r.); R_f = 0.3 (Et₂O/pentanes, 1:10).

IR (ATR): 2960, 1722, 1667, 1369, 1153 cm⁻¹.

¹H NMR (CDCl₃, 400 MHz): δ = 7.70 (d, J = 7.1 Hz, 2H), 7.42 (t, J = 7.4 Hz, 1H), 7.28 (t, J = 8.1 Hz, 4H), 7.20 (t, J = 7.6 Hz, 2H), 7.11 (t, J = 7.3 Hz, 1H), 4.37 (dd, J = 10.2, 8.1 Hz, 1H), 3.68 (td, J = 10.5, 8.4 Hz, 1H), 2.86 (d, J = 8.1 Hz, 1H), 2.15–1.94 (m, 2H), 1.40 (s, 9H), 1.28 (s, 3H), 1.18 (s, 3H).

¹³C NMR (CDCl₃, 101 MHz): δ = 201.86, 173.23, 142.94, 137.18, 132.79, 128.47 (d, J = 6.8 Hz), 128.22, 127.58, 126.46, 80.96, 59.98, 57.17, 49.51, 47.24, 42.31, 30.24, 28.09, 26.41.

HRMS (ESI): m/z [M+H]⁺ calcd for [C₂₅H₃₁O₃]⁺: 379.2268; found: 379.2268.

tert-Butyl 3-benzoyl-4-(4-methoxyphenyl)cyclopentanecarboxylate (12)

Yield: 64.2 mg (84%, 3:1 d.r.). R_f = 0.3 (Acetone/pentanes, 1:29)

Major diastereomer—White solid (m.p. = 95.5–98.5 °C).

IR (ATR): 2975, 1721, 1679, 1612, 1513, 1366, 1246, 1012, cm⁻¹.

^1H NMR (CDCl_3 , 400 MHz): δ = 7.81 (d, J = 6.9 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.37 (t, J = 7.8 Hz, 2H), 7.15 (d, J = 8.7 Hz, 2H), 6.78 (d, J = 8.6 Hz, 2H), 3.82 (td, J = 9.5, 8.0 Hz, 1H), 3.77 – 3.71 (m, 1H), 3.75 (s, 3H), 3.07 (qd, J = 8.4, 5.7 Hz, 1H), 2.54 – 2.43 (m, 2H), 2.21 (ddd, J = 13.2, 9.8, 8.3 Hz, 1H), 2.08 (dt, J = 13.3, 9.1 Hz, 1H), 1.46 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.72, 174.30, 158.12, 136.92, 135.69, 132.90, 128.47, 128.39, 128.23, 113.90, 80.46, 55.24, 54.88, 46.54, 44.30, 37.40, 35.27, 28.08.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{24}\text{H}_{29}\text{O}_4]^+$: 381.2060; found: 381.2053.

Minor diastereomer—White solid (m.p. = 110.5–113.5 °C).

IR (ATR): 2975, 1723, 1680, 1514, 1448, 1367, 1249, 1220, 1152, 1036, 830, 701 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.82 (d, J = 7.2 Hz, 2H), 7.50 (t, J = 7.4 Hz, 1H), 7.37 (t, J = 7.7 Hz, 2H), 7.18 (d, J = 8.6 Hz, 2H), 6.78 (d, J = 8.7 Hz, 2H), 3.93 (q, J = 9.4 Hz, 1H), 3.74 (s, 3H), 3.69 – 3.54 (m, 1H), 2.99 (ddd, J = 16.6, 9.3, 7.3 Hz, 1H), 2.54 – 2.41 (m, 2H), 2.18 – 2.01 (m, 2H), 1.48 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.51, 175.01, 158.20, 136.69, 134.93, 133.01, 128.49, 128.49, 128.34, 113.89, 80.46, 55.24, 53.85, 47.74, 44.11, 38.92, 34.76, 28.14.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{24}\text{H}_{32}\text{NO}_4]^+$: 398.2326; found: 398.2322.

***tert*-Butyl 3-benzoyl-4-(4-trifluoromethylphenyl)cyclopentanecarboxylate (13)**

Yield: 57.6 mg (68%, 2:1 d.r.). R_f = 0.3 (Acetone/pentanes, 1:49)

Major diastereomer—White solid (m.p. = 106.9–108.4 °C).

IR (ATR): 2978, 1724, 1682, 1619, 1449, 1368, 1326, 1258, 1156, 1124, 1069, 1017, 842, 701 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.82 (d, J = 7.1 Hz, 2H), 7.55 – 7.47 (m, 3H), 7.40 (t, J = 7.8 Hz, 2H), 7.36 (d, J = 8.1 Hz, 2H), 3.93 – 3.81 (m, 2H), 3.15 – 3.07 (m, 1H), 2.55 (qd, J = 8.2, 4.1 Hz, 2H), 2.19 (ddd, J = 13.1, 9.5, 8.0 Hz, 1H), 2.12 (dt, J = 13.5, 9.2 Hz, 1H), 1.46 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 199.89, 173.95, 147.72, 136.59, 133.17, 128.75 (q, J = 32.5 Hz), 128.59, 128.35, 127.70, 125.45 (q, J = 3.9 Hz), 124.47 (q, J = 272.9 Hz), 80.73, 54.76, 46.57, 44.28, 37.02, 35.45, 28.05

^{19}F NMR (CDCl_3 , 377 MHz): δ = –62.46.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{24}\text{H}_{26}\text{F}_3\text{O}_3]^+$: 419.1829; found: 419.1826.

Minor diastereomer—White solid (m.p. = 122.8–125.9 °C).

IR (ATR): 2974, 1722, 1677, 1369, 1327, 1281, 1247, 1225, 1164, 1138, 1170, 1018, 842, 699 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.84 (d, J = 7.1 Hz, 2H), 7.57 – 7.46 (m, 3H), 7.40 (t, J = 7.5 Hz, 4H), 3.98 (q, J = 9.4 Hz, 1H), 3.83 – 3.71 (m, 1H), 3.01 (ddd, J = 16.4, 9.2, 7.3 Hz, 1H), 2.60 – 2.48 (m, 2H), 2.20 – 2.06 (m, 2H), 1.49 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.62, 174.73, 147.10, 136.36, 133.29, 128.80 (q, J = 31.1 Hz), 128.62, 128.45, 127.82, 125.44 (q, J = 3.7 Hz), 124.16 (q, J = 271.3), 80.72, 53.75, 47.67, 44.03, 38.34, 34.99, 28.11.

^{19}F NMR (CDCl_3 , 377 MHz): δ = -62.47.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{24}\text{H}_{29}\text{F}_3\text{NO}_3]^+$: 436.2095; found: 436.2093.

***tert*-Butyl 3-benzoyl-4-(4-bromophenyl)cyclopentanecarboxylate (14)**

Yield: 55.3 mg (64%, 2:1 d.r.). R_f = 0.3 (Acetone/pentanes, 1:29)

Major diastereomer—White solid (m.p. = 103.1–104.4 °C).

IR (ATR): 2976, 1722, 1681, 1490, 1448, 1366, 1257, 1150, 1074, 1010, 848, 821, 701 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.82 (d, J = 7.3 Hz, 2H), 7.52 (t, J = 7.4 Hz, 1H), 7.42 – 7.34 (m, 4H), 7.12 (d, J = 8.4 Hz, 2H), 3.83 – 3.74 (m, 2H), 3.08 (qd, J = 8.2, 5.7 Hz, 1H), 2.57 – 2.45 (m, 2H), 2.23 – 2.13 (m, 1H), 2.07 (dt, J = 13.4, 9.2 Hz, 1H), 1.45 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.16, 174.05, 142.62, 136.69, 133.11, 131.56, 129.10, 128.57, 128.35, 120.13, 80.65, 54.77, 46.40, 44.26, 37.09, 35.41, 28.06.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{23}\text{H}_{26}\text{BrO}_3]^+$: 429.1060; found: 429.1074.

Minor diastereomer—White solid (m.p. = 129.1–132.8 °C).

IR (ATR): 2975, 1722, 1680, 1490, 1448, 1367, 1284, 1246, 1220, 1152, 1074, 1011, 823, 700 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.83 (d, J = 7.3 Hz, 2H), 7.52 (t, J = 7.4 Hz, 1H), 7.44 – 7.32 (m, 4H), 7.15 (d, J = 8.4 Hz, 2H), 3.92 (q, J = 9.4 Hz, 1H), 3.65 (ddd, J = 11.3, 9.5, 7.6 Hz, 1H), 2.99 (ddd, J = 16.5, 9.3, 7.3 Hz, 1H), 2.50 (ddd, J = 13.3, 9.8, 6.8 Hz, 2H), 2.17 – 2.02 (m, 2H), 1.48 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.92, 174.82, 141.95, 136.47, 133.23, 131.56, 129.20, 128.60, 128.46, 120.25, 80.64, 53.74, 47.56, 44.03, 38.46, 34.93, 28.12.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{23}\text{H}_{29}\text{BrNO}_3]^+$: 446.1326; found: 446.1313.

***tert*-Butyl 3-benzoyl-4-(2-methylphenyl)cyclopentanecarboxylate (15)**

Yield: 60.5 mg (83%, 2:1 d.r.). R_f = 0.3 (Et_2O /pentanes, 1:49)

Major diastereomer—White solid (m.p. = 70.7–74.1 °C).

IR (ATR): 2975, 1723, 1681, 1448, 1367, 1218, 1149, 1101, 848, 753, 700 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.81 (d, J = 7.3 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.37 (t, J = 7.7 Hz, 2H), 7.26 (d, J = 7.7 Hz, 1H), 7.16 (dt, J = 7.9, 4.3 Hz, 1H), 7.10 – 7.01 (m, 2H), 4.08 – 3.92 (m, 2H), 3.09 (td, J = 8.5, 6.5 Hz, 1H), 2.59 – 2.45 (m, 2H), 2.31 (s, 3H), 2.24 (dt, J = 13.1, 8.8 Hz, 1H), 1.96 (dt, J = 13.3, 8.5 Hz, 1H), 1.46 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.83, 174.40, 142.30, 136.88, 136.65, 133.05, 130.57, 128.60, 128.49, 126.38, 126.28, 125.49, 80.60, 54.16, 44.80, 43.00, 37.70, 35.29, 28.21, 20.00.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{24}\text{H}_{29}\text{O}_3]^+$: 365.2111; found: 365.2105.

Minor diastereomer—Colorless oil.

IR (ATR): 2976, 1723, 1680, 1448, 1367, 1290, 1216, 1151, 848, 575, 700 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.81 (d, J = 7.4 Hz, 2H), 7.49 (t, J = 7.4 Hz, 1H), 7.41 – 7.31 (m, 3H), 7.23 – 7.12 (m, 1H), 7.10 – 7.01 (m, 2H), 4.09 (q, J = 9.0 Hz, 1H), 3.88 (dt, J = 11.0, 8.3 Hz, 1H), 3.01 (p, J = 8.9 Hz, 1H), 2.51 (dq, J = 13.5, 7.3 Hz, 2H), 2.28 (s, 3H), 2.23 – 2.12 (m, 1H), 2.05 – 1.91 (m, 1H), 1.48 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.48, 174.93, 141.45, 136.53, 136.33, 133.01, 130.31, 128.46, 128.46, 126.42, 126.20, 125.81, 80.45, 53.47, 44.43, 43.81, 39.39, 34.61, 28.13, 19.83.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{24}\text{H}_{32}\text{NO}_3]^+$: 382.2377; found: 382.2372.

tert-Butyl 3-benzoyl-4-(9H-carbazol-9-yl)cyclopentanecarboxylate (19)

Yield: 63.5 mg (71%, 2:1 d.r.). R_f = 0.3 (Acetone/pentanes, 1:29)

Major diastereomer—Colorless oil.

IR (ATR): 2976, 1724, 1682, 1597, 1484, 1453, 1367, 1337, 1227, 1154, 847, 750, 724, 698 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 8.04 (d, J = 7.7 Hz, 2H), 7.62 (d, J = 7.2 Hz, 2H), 7.57 (d, J = 8.3 Hz, 2H), 7.44 (t, J = 7.8 Hz, 2H), 7.35 (t, J = 7.5 Hz, 1H), 7.27 – 7.14 (m, 4H), 5.94 (q, J = 9.1 Hz, 1H), 4.66 (q, J = 9.1 Hz, 1H), 3.40 (p, J = 8.1 Hz, 1H), 2.85 – 2.57 (m, 3H), 2.38 (dt, J = 13.3, 9.1 Hz, 1H), 1.50 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 199.53, 173.42, 139.31, 136.18, 133.14, 128.39, 128.26, 125.66, 123.41, 120.39, 119.08, 109.76, 81.12, 56.17, 49.49, 44.09, 34.36, 31.36, 28.10.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{29}\text{H}_{30}\text{NO}_3]^+$: 440.2220; found: 440.2217.

Minor diastereomer—Colorless oil.

IR (ATR): 2977, 1722, 1681, 1597, 1484, 1453, 1367, 1337, 1221, 1151, 750, 724, 699 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 8.05 (d, J = 7.7 Hz, 2H), 7.71 (d, J = 7.2 Hz, 2H), 7.64 (d, J = 7.0 Hz, 2 Hz), 7.49 – 7.35 (m, 3H), 7.28 – 7.17 (m, 5H), 5.81 (dt, J = 11.0, 8.5 Hz, 1H), 4.85 – 4.74 (m, 1H), 3.09 (dt, J = 15.9, 7.8 Hz, 1H), 2.97 – 2.84 (m, 1H), 2.76 (ddd, J = 13.5, 9.9, 7.7 Hz, 1H), 2.51 (dt, J = 13.3, 8.1 Hz, 1H), 2.30 (dt, J = 14.0, 7.6 Hz, 1H), 1.52 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.20, 174.24, 139.40, 135.75, 133.31, 128.49, 128.47, 125.71, 123.40, 120.34, 119.11, 109.90, 81.07, 56.75, 47.76, 42.78, 33.31, 32.53, 28.15.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{29}\text{H}_{30}\text{NO}_3]^+$: 440.2220; found: 440.2217.

***tert*-Butyl 3-benzoyl-4-(4-methylphenyl)cyclopentanecarboxylate (16)**

Yield: 63.0 mg (85%, 2.0:1 d.r.). R_f = 0.3 (Acetone/toluene, 0:100 to 10:90)

Major diastereomer—White solid (m.p. = 74.2–79.5 $^{\circ}\text{C}$).

IR (ATR): 2977, 1675, 1597, 1447, 1367, 1285, 1253, 848, 732, 700 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.52 (d, J = 7.0 Hz, 2H), 7.41 (t, J = 7.4 Hz, 1H), 7.31 – 7.16 (m, 7H), 4.09 (dd, J = 9.8, 6.7 Hz, 1H), 3.08 (dt, J = 18.3, 9.3 Hz, 1H), 2.63 (dt, J = 13.3, 10.0 Hz, 1H), 2.53 (dd, J = 13.7, 9.2 Hz, 1H), 2.29 – 2.14 (m, 2H), 1.49 (s, 9H), 1.26 (s, 3H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.35, 174.34, 148.87, 137.78, 132.57, 128.39, 128.38, 128.15, 126.15, 125.88, 80.38, 57.65, 49.53, 45.42, 43.63, 32.71, 28.14, 24.30.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{24}\text{H}_{29}\text{O}_3]^+$: 365.2112; found: 365.2115.

Minor diastereomer—Colorless oil.

IR (ATR): 2977, 1724, 1676, 1449, 1369, 1258, 1222, 1154, 1086, 1026, 803, 700 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.49 (d, J = 7.1 Hz, 2H), 7.37 (t, J = 7.4 Hz, 1H), 7.31 (d, J = 8.1 Hz, 2H), 7.25 – 7.09 (m, 5H), 4.17 (t, J = 8.7 Hz, 1H), 3.17 (qd, J = 9.0, 5.6 Hz, 1H), 2.68 – 2.57 (m, 2H), 2.28 (ddd, J = 13.9, 8.6, 5.6 Hz, 1H), 2.17 (dd, J = 13.1, 8.2 Hz, 1H), 1.49 (s, 9H), 1.30 (s, 3H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.56, 175.76, 147.07, 137.78, 132.47, 128.29, 128.23, 128.05, 126.21, 126.19, 80.42, 56.74, 50.65, 46.99, 42.29, 32.17, 28.15, 22.14.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{24}\text{H}_{32}\text{NO}_3]^+$: 382.2377; found: 382.2376.

***tert*-Butyl 2-benzoyl-2',3'-dihydrospiro[cyclopentane-1,1'-indene]-4-carboxylate (17)**

Yield: 61.7 mg (83%, 2:1 d.r.). R_f = 0.3 (Acetone/pentanes, 1:49)

Major diastereomer—Colorless oil.

IR (ATR): 2975, 2935, 1724, 1672, 1478, 1448, 1392, 1367, 1225, 1151, 1002, 849, 756, 732, 691 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.49 (d, J = 7.0 Hz, 2H), 7.40 – 7.32 (m, 2H), 7.26 (m, 1H), 7.17 (t, J = 7.8 Hz, 2H), 7.09 (t, J = 7.3 Hz, 1H), 6.93 (d, J = 6.9 Hz, 1H), 3.91 (dd, J = 10.4, 6.9 Hz, 1H), 3.06 (dt, J = 17.3, 8.3 Hz, 1H), 2.73 – 2.60 (m, 2H), 2.44 (dd, J = 13.3, 9.1 Hz, 1H), 2.38 – 2.22 (m, 4H), 1.93 – 1.83 (m, 1H), 1.48 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.23, 174.36, 149.28, 143.69, 137.38, 132.57, 128.16, 128.02, 127.08, 126.62, 124.69, 122.27, 80.35, 59.06, 56.04, 44.49, 43.23, 36.75, 33.47, 30.97, 28.12.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{25}\text{H}_{29}\text{O}_3]^+$: 377.2111; found: 377.2107.

Minor diastereomer—Colorless oil.

IR (ATR): 2976, 2932, 1720, 1671, 1448, 1367, 1220, 1150, 908, 846, 729, 690 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.49 – 7.41 (m, 3H), 7.31 (t, J = 7.4 Hz, 1H), 7.26 – 7.18 (m, 1H), 7.11 (t, J = 7.8 Hz, 2H), 7.03 (t, J = 7.6 Hz, 1H), 6.84 (d, J = 7.5 Hz, 1H), 4.05 (t, J = 8.7 Hz, 1H), 3.11 (tdd, J = 9.9, 7.6, 6.1 Hz, 1H), 2.71 (dt, J = 13.6, 9.7 Hz, 1H), 2.67 – 2.57 (m, 1H), 2.54 (dd, J = 12.8, 9.8 Hz, 1H), 2.47 – 2.25 (m, 2H), 2.27 – 2.09 (m, 2H), 1.76 (dt, J = 12.4, 8.9 Hz, 1H), 1.50 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.42, 175.61, 147.06, 143.90, 137.16, 132.45, 128.14, 127.82, 127.19, 126.54, 124.57, 122.60, 80.39, 60.28, 55.35, 46.18, 42.58, 34.90, 31.96, 30.78, 28.17.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{25}\text{H}_{32}\text{NO}_3]^+$: 394.2377; found: 394.2374.

tert-Butyl 2-benzoyl-3',4'-dihydro-2'H-spiro[cyclopentane-1,1'-naphthalene]-4-carboxylate (18)

Yield: 66.6 mg (84%, 2:1 d.r.). R_f = 0.3 (Acetone/pentanes, 1:49)

Major diastereomer—White solid (m.p = 104.0–106.8 °C).

IR (ATR): 2974, 2936, 1724, 1673, 1447, 1366, 1254, 1230, 1152, 754, 720, 691 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.52 (d, J = 7.9 Hz, 1H), 7.44 (d, J = 8.4 Hz, 2H), 7.35 (t, J = 7.4 Hz, 1H), 7.27 (d, J = 6.2 Hz, 1H), 7.14 (t, J = 7.8 Hz, 2H), 7.10 (t, J = 7.7 Hz, 1H), 6.90 (d, J = 7.6 Hz, 1H), 4.16 (dd, J = 11.6, 6.4 Hz, 1H), 3.10 (dtd, J = 11.6, 9.3, 7.2 Hz, 1H), 2.72 (q, J = 11.8 Hz, 1H), 2.57 (dt, J = 16.5, 5.5 Hz, 1H), 2.36 (qd, J = 11.7, 10.4, 4.5 Hz, 2H), 2.23 (ddd, J = 13.5, 7.9, 5.5 Hz, 2H), 1.81 – 1.64 (m, 2H), 1.66 – 1.50 (m, 2H), 1.47 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 201.38, 174.27, 143.94, 137.78, 137.47, 132.45, 129.28, 128.34, 128.08, 126.70, 126.46, 125.88, 80.39, 58.96, 49.24, 48.71, 43.93, 34.36, 33.26, 29.98, 28.13, 20.23.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{26}\text{H}_{31}\text{O}_3]^+$: 391.2273; found: 391.2263.

Minor diastereomer—Colorless oil.

IR (ATR): 2976, 2937, 1723, 1674, 1447, 1366, 1218, 1151, 733 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.67 (d, J = 8.0 Hz, 1H), 7.46 (d, J = 8.2 Hz, 2H), 7.32 (t, J = 7.4 Hz, 1H), 7.30 – 7.21 (m, 1H), 7.13 (t, J = 7.8 Hz, 2H), 7.04 (t, J = 7.3 Hz, 1H), 6.85 (d, J = 7.6 Hz, 1H), 4.32 (t, J = 8.3 Hz, 1H), 3.17 – 3.04 (m, 1H), 2.73 (dt, J = 13.7, 8.7 Hz, 1H), 2.52 (dt, J = 16.1, 5.0 Hz, 1H), 2.40 – 2.22 (m, 4H), 1.73 (ddd, J = 13.1, 9.4, 3.4 Hz, 1H), 1.67 – 1.49 (m, 3H), 1.49 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 202.08, 175.59, 141.75, 138.18, 137.82, 132.34, 129.00, 128.31, 127.96, 127.36, 126.41, 125.88, 80.36, 58.12, 50.62, 49.53, 42.83, 32.01, 31.97, 30.32, 28.16, 20.50.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{26}\text{H}_{34}\text{NO}_3]^+$: 408.2533; found: 408.2529.

tert-Butyl 3-benzoyl-4-vinylcyclopentanecarboxylate (22)

Yield: 124.7 mg (67%, 3:1 d.r.). R_f = 0.3 (Et_2O /pentanes, 1:9)

Major diastereomer—Colorless oil.

IR (ATR): 2977, 2933, 1725, 1681, 1449, 1367, 1258, 1152 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.87 (d, J = 7.3 Hz, 2H), 7.48 (t, J = 7.4 Hz, 1H), 7.39 (t, J = 7.6 Hz, 2H), 5.71 (ddd, J = 17.4, 10.2, 7.4 Hz, 1H), 4.95 (d, J = 17.1 Hz, 1H), 4.87 (d, J = 10.3 Hz, 1H), 3.49 (q, J = 8.3 Hz, 1H), 3.10 (p, J = 7.9 Hz, 1H), 2.95 – 2.73 (m, 1H), 2.28 (dt, J = 13.1, 8.2 Hz, 1H), 2.17 (ddd, J = 14.0, 8.0, 6.4 Hz, 1H), 2.07 (dt, J = 13.1, 9.1 Hz, 1H), 1.76 (dt, J = 13.1, 8.6 Hz, 1H), 1.37 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): δ = 200.70, 174.32, 140.11, 137.02, 132.99, 128.56, 128.46, 114.72, 80.39, 52.21, 45.98, 43.86, 35.09, 34.47, 28.06.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{19}\text{H}_{25}\text{O}_3]^+$: 301.1804; found: 301.1803.

Minor diastereomer—Colorless oil.

IR (ATR): 2981, 2934, 1719, 1678, 1368, 1155 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): δ = 7.90 (d, J = 7.3 Hz, 2H), 7.54 (t, J = 7.4 Hz, 1H), 7.44 (t, J = 7.6 Hz, 2H), 5.61 (dt, J = 16.9, 9.7 Hz, 1H), 4.81 – 4.71 (m, 2H), 3.92 (q, J = 9.0 Hz, 1H), 3.05 (p, J = 8.0 Hz, 1H), 2.83 (dt, J = 17.3, 8.7 Hz, 1H), 2.48 (dt, J = 13.2, 10.0 Hz, 1H),

2.22 (dt, $J = 13.4, 8.0$ Hz, 1H), 2.12 (dt, $J = 13.9, 7.4$ Hz, 1H), 1.99 (ddd, $J = 13.3, 8.9, 7.4$ Hz, 1H), 1.47 (s, 9H).

^{13}C NMR (CDCl_3 , 101 MHz): $\delta = 200.46, 174.29, 138.13, 137.62, 132.81, 128.48, 128.31, 115.40, 80.30, 50.58, 47.40, 44.08, 35.63, 31.60, 28.10$.

HRMS (ESI): m/z $[\text{M}+\text{NH}_4^+]^+$ calcd for $[\text{C}_{19}\text{H}_{28}\text{NO}_3]^+$: 318.2069; found: 318.2067.

ethyl 4-benzoyl-3-((tert-butyldimethylsilyl)oxy)-3-phenylcyclopentane carboxylate (20)

Colorless oil; Yield: 80.8 mg (94%, 11:1 d.r.); $R_f = 0.3$ (Et_2O /pentanes, 1:5).

IR (ATR): 2960, 1722, 1667, 1369, 1153 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): $\delta = 7.76$ (d, $J = 7.76$ Hz, 2H), 7.5 (t, $J = 7.8$ Hz, 1H), 7.4 (t, $J = 7.8$ Hz, 2H), 7.35–7.3 (m, 2H), 7.2–7.1 (m, 3H), 4.4 (ddd, $J = 8.6, 3.6, 1.3$ Hz, 1H), 4.25 (q, $J = 7.1$ Hz, 2H), 3.43 (tt, $J = 10.8, 7.4$ Hz, 1H), 3.15 (dd, $J = 13.1, 10.9$ Hz, 1H), 2.7 (dt, $J = 13.1, 9.8$ Hz, 1H), 2.57 (dd, $J = 13.1, 7.4$ Hz, 1H), 2.41 (ddd, $J = 13.1, 6.7, 4.1$ Hz, 1H), 1.34 (t, $J = 7.1$ Hz, 3H), 0.97 (s, 9H), -0.07 (s, 3H), -0.45 (s, 3H).

^{13}C NMR (CDCl_3 , 101 MHz): $\delta = 201.33, 175.22, 142.04, 137.67, 132.49, 128.21, 127.65, 127.43, 127.05, 88.15, 60.63, 58.64, 41.92, 40.43, 31.81, 25.96, 18.30, 14.28, -2.76, -3.47$.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{27}\text{H}_{37}\text{O}_4\text{Si}]^+$: 453.2456; found: 453.2456.

methyl 2-benzoyl-4,4-dimethylcyclopentanecarboxylate (21)

Colorless oil; Yield: 44.3 mg (85%, 4:1 d.r.); $R_f = 0.3$ (Et_2O /pentanes, 1:19).

IR (ATR): 2960, 1722, 1667, 1369, 1153 cm^{-1} .

^1H NMR (CDCl_3 , 400 MHz): $\delta = 8.03 - 7.92$ (m, 2H), 7.60 – 7.52 (m, 1H), 7.47 (t, $J = 7.5$ Hz, 2H), 4.24 (dt, $J = 10.0, 8.6$ Hz, 1H), 3.71 (q, $J = 9.0$ Hz, 1H), 3.63 (s, 3H), 2.04 (dd, $J = 12.8, 10.0$ Hz, 1H), 1.97 (dd, $J = 12.7, 8.9$ Hz, 1H), 1.82 (dd, $J = 12.7, 9.4$ Hz, 1H), 1.59 (ddd, $J = 12.9, 8.6, 0.8$ Hz, 1H), 1.11 (s, 3H), 1.03 (s, 3H).

^{13}C NMR (CDCl_3 , 101 MHz): $\delta = 200.88, 175.88, 136.51, 133.06, 128.62, 128.57, 51.86, 49.43, 46.00, 44.94, 44.52, 39.99, 29.15, 28.71$.

HRMS (ESI): m/z $[\text{M}+\text{H}^+]^+$ calcd for $[\text{C}_{16}\text{H}_{21}\text{O}_3]^+$: 260.14; found: 260.15.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Information

This work was conducted using funding from the NIH (GM095666). The mass spectroscopy facilities at UW-Madison are funded in part by the NIH (S10OD020022), as are the NMR facilities (S10 OD012245).

References

1. For recent reviews: Hashimoto T, Maruoka K. Chem Rev. 2015; 115:5366. [PubMed: 25961125] Heasley B. Eur J Org Chem. 2009:1477–1489. Hartley RC, Caldwell ST. J Chem Soc Perkin Trans 1. 2000:477.
2. For examples of cyclizations: Parr BT, Li Z, Davies HML. Chem Sci. 2011; 2:2378. [PubMed: 22708053] Raimondi W, Lettieri G, Dulcère JP, Bonne D, Rodriguez J. Chem Commun. 2010; 46:7247. Unger R, Weisser F, Chinkov N, Stanger A, Cohen T, Marek I. Org Lett. 2009; 11:1853. [PubMed: 19290654] Marcé P, Díaz Y, Matheu MI, Castilló S. Org Lett. 2008; 10:4735. [PubMed: 18816138] Crimmins MT, King BW. J Org Chem. 1996; 61:4192. [PubMed: 11667311]
3. For examples of ring expansions: Yin Q, You SL. Org Lett. 2014; 16:1810. [PubMed: 24617607] Zhu DY, Hang FM, Wang SH. Chem Eur J. 2015; 21:15502. [PubMed: 26387785]
4. For recent examples of zwitterionic intermediates in [3 + 2] cycloadditions: Trost BM, Ehmke V. Org Lett. 2014; 16:2708. [PubMed: 24787546] Grover HK, Emmet MR, Kerr MA. Org Biomol Chem. 2015; 13:655. [PubMed: 25425071] Trost BM, Morris PJ, Sprague SJ. J Am Chem Soc. 2012; 134:17823. [PubMed: 23030714] Zhou W, Wang H, Tao M, Zhu CZ, Lin TY, Zhang J. Chem Sci. 2017; 8:4660. [PubMed: 28936335]
5. Recent example with vinyl carbenoids: Briones JF, Davies HML. J Am Chem Soc. 2013; 135:13314. [PubMed: 23978108]
6. Recent example of reductive cycloaddition: Ahlin JSE, Donets PA, Cramer N. Angew Chem Int Ed. 2014; 53:13229.
7. Yoon TP. ACS Catal. 2013; 3:895. [PubMed: 23691491]
8. (a) Lu Z, Shen M, Yoon TP. J Am Chem Soc. 2011; 133:1162. [PubMed: 21214249] (b) Amador AG, Sherbrook EM, Yoon TP. J Am Chem Soc. 2016; 138:4722. [PubMed: 27015009]
9. For a conceptually similar photocatalytic radical anion cycloaddition of nitrocyclopropanes by one of us, see: Wang C, Ren X, Xie H, Lu Z. Chem Eur J. 2015; 21:9676. [PubMed: 26037066]
10. For photocatalytic radical cation cycloadditions of aminocyclopropanes, see: Maity S, Zhu M, Shinabery RS, Zheng N. Angew Chem Int Ed. 2012; 51:222. Nguyen TH, Morris SA, Zheng N. Adv Synth Catal. 2014; 356:2831. [PubMed: 26120292]
11. For leading references on the use of C-acylimidazole auxiliaries in photoredox catalysis, see: Tyson EL, Farney EP, Yoon TP. Org Lett. 2012; 14:1110. [PubMed: 22320352] Huo H, Shen X, Wang C, Zhang L, Röse P, Chen LA, Harms K, Marsch M, Hilt G, Meggers E. Nature. 2014; 515:100. [PubMed: 25373679] Huang X, Webster RD, Harms K, Meggers E. J Am Chem Soc. 2016; 138:12636. [PubMed: 27577929]
12. Cismesia MA, Yoon TP. Chem Sci. 2015; 6:5426. [PubMed: 26668708]
13. (a) Tanko JM, Drumright RE. J Am Chem Soc. 1992; 114:1844. (b) Chahma M, Li X, Phillips JP, Schwartz P, Brammer LE, Wang Y, Tanko JM. J Phys Chem A. 2005; 109:3372. [PubMed: 16833672] (c) Tanko JM, Drumright RE. J Am Chem Soc. 1990; 112:5362.
14. Zimmerman HE, Wang P. J Org Chem. 2003; 68:9226. [PubMed: 14629140]

Figure 1. Reagents and conditions

Cyclopropane **1** (1 equiv), alkene partner (5 equiv), $\text{Ru}(\text{bpy})_3(\text{PF}_6)_2$ (1 mol%), $\text{Gd}(\text{OTf})_3$ (0.5 equiv), and TMEDA (1.5 equiv) in MeCN (0.2 M). Reactions were irradiated with a 23 W compact fluorescent bulb. Yields are of isolated cyclopentane. Diastereomer ratios were determined by ^1H NMR analysis of the unpurified products.

Figure 2. Reagents and conditions

Cyclopropane **1** (1 equiv), alkene partner (5 equiv), Ru(bpy)₃(PF₆)₂ (1 mol%), Gd(OTf)₃ (0.5 equiv), and TMEDA (1.5 equiv) in MeCN (0.2 M). Reactions were irradiated with a 23 W compact fluorescent bulb. Yields are of isolated cyclopentane. Diastereomer ratios were determined by ¹H NMR analysis of the unpurified products.

Figure 3.
Proposed mechanism for intermolecular [3 + 2] cycloaddition

Scheme 1.

Top: Racemic photocatalytic intramolecular [3 + 2] cycloadditions. Bottom:
Enantioselective photocatalytic intermolecular [3 + 2] cycloadditions.

Scheme 2. Isomerization of *cis*-cyclopropane under photocatalytic conditions

Table 1Optimization of reaction conditions^a

Entry	Solvent	Lewis acid	Base	Yield ^b
1	MeCN	La(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	44%
2	EtOAc	La(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	32%
3	DMF	La(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	0%
4	CH ₂ Cl ₂	La(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	0%
5	MeCN	Zn(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	0%
6	MeCN	Er(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	55%
7	MeCN	Gd(OTf) ₃ (1 equiv)	TMEDA (5 equiv)	69%
8	MeCN	Gd(OTf) ₃ (1 equiv)	<i>i</i> Pr ₂ NEt (5 equiv)	9%
9	MeCN	Gd(OTf) ₃ (1 equiv)	PMP (5 equiv)	7%
10	MeCN	Gd(OTf) ₃ (0.5 equiv)	TMEDA (5 equiv)	49%
11	MeCN	Gd(OTf) ₃ (0.5 equiv)	TMEDA (1.5 equiv)	69%
12 ^c	MeCN	Gd(OTf) ₃ (0.5 equiv)	TMEDA (1.5 equiv)	71%

^a[0.2 M] in indicated solvent, 0.04 mmol cyclopropane, 2.5 % [Ru].^bYields determined by ¹H NMR using phenanthrene as an internal standard.^cUsing 1 % Ru(bpy)₃(PF₆)₂