Published in final edited form as:

Hepatology. 2018 April ; 67(4): 1441–1457. doi:10.1002/hep.29540.

CHOP-induced loss of intestinal epithelial stemness contributes to bile duct ligation-induced cholestatic liver injury

Runping Liu^{1,2}, Xiaojiaoyang Li^{1,2}, Zhiming Huang³, Derrick Zhao¹, Bhagyalaxmi Sukka Ganesh¹, Guanhua Lai⁴, William M. Pandak^{2,5}, Phillip B Hylemon^{1,2,5}, Jasmohan S Bajaj^{2,5}, Arun J Sanyal⁵, and Huiping Zhou^{1,2,3,5}

¹Department of Microbiology and Immunology, Medical College of Virginia Campus, Richmond, Virginia, 23298

²McGuire Veterans Affairs Medical Center, Virginia Commonwealth University, Richmond, Virginia, 23298

³Department of Gastroenterology, the First Affiliated Hospital, Wenzhou Medical University, Wenzhou, Zhejiang, China

⁴Department of Pathology, Medical College of Virginia Campus, Virginia Commonwealth University, Richmond, Virginia, 23298

⁵Department of Internal Medicine/GI Division, Virginia Commonwealth University, Richmond, Virginia, 23298

Abstract

Impaired intestinal barrier function promotes the progression of various liver diseases including cholestatic liver disease. The close association of primary sclerosing cholangitis (PSC) with inflammatory bowel disease highlights the importance of the gut-liver axis. It has been reported that bile duct ligation (BDL)-induced liver fibrosis is significantly reduced in C/EBP homologous protein knock out (CHOP^{-/-}) mice. However, the underlying mechanisms remain unclear. In the current study, we demonstrate that BDL induces striking and acute hepatic ER stress responses after 1 day, which return to normal after 3 days. No significant hepatocyte apoptosis is detected 7 to 14 days following BDL. However, the inflammatory response is significantly increased after 7 days, which is similar to what we found in human PSC liver samples. BDL-induced loss of stemness in intestinal stem cells (ISCs), disruption of intestinal barrier function, bacterial translocation, activation of hepatic inflammation and M2 macrophage polarization and liver fibrosis are significantly reduced in CHOP^{-/-} mice. In addition, intestinal organoids derived from CHOP^{-/-} mice contain more and longer crypt structures than those from WT mice, which is consistent with the upregulation of stem cell markers (Lgr5, Olfm4 and Sox9) and *in vivo* findings

Address correspondence to: Huiping Zhou, Ph.D, Department of Microbiology & Immunology, Virginia Commonwealth University, McGuire Veterans Affairs Medical Center, P.O. Box 980678, Richmond, VA 23298-0678, Tel: 804-828-6817; Fax: 804-828-0676, huiping.zhou@vcuhealth.org.

Contributions: RL and HZ conceived the original ideas, designed the study, analyzed the data and wrote the manuscript; RL, XL, DZ, BSG carried out the experiments and data analysis. GL helped by doing histology analysis. JSB was responsible for human sample collection; PBH, WMP, JSB, AJS helped data with data analysis and reviewed manuscript.

Competing interests
None declared.

that CHOP^{-/-} mice have longer villi and crypts as compared to WT mice. Similarly, the mRNA levels of CD14, IL-1 β , TNF- α and MCP-1 are increased and stem cell proliferation is suppressed in the duodenum of cirrhotic patients.

Conclusion—Activation of ER stress and subsequent loss of stemness of ISCs plays a critical role in BDL-induced systemic inflammation and cholestatic liver injury. Modulation of the ER stress response represents a potential therapeutic strategy for cholestatic liver diseases as well as other inflammatory diseases

Keywords

ER stress; Bile duct ligation; loss of stemness; cholestasis; inflammation

Introduction

Bile duct obstruction is a major driving force that causes chronic cholestasis and eventually progressive liver fibrosis and cirrhosis regardless of different etiologies (1). The accumulation of hydrophobic bile acids is believed to induce chronic inflammation and hepatocyte apoptosis (2). Recent clinical studies indicate that disruption of intestinal barrier function and subsequent bacterial translocation also contribute to cholestatic liver disease progression, such as primary sclerosing cholangitis (PSC), primary biliary cholangitis (PBC) and non-cholestatic liver diseases by inducing hepatic inflammation (3). It has been reported that PSC represents one of the most common extra-intestinal manifestations of inflammatory bowel disease (IBD) (3). The close association between PSC and IBD highlights the importance of the gut-liver axis (4). Furthermore, disruption of the intestinal barrier was not only observed in obstructive jaundice patients, but also in bile duct ligation (BDL) rodent models of cholestatic liver diseases (5, 6). However, the underlying mechanisms remain largely unknown.

The intestinal epithelium represents the largest epithelial surface in our body (7). The maintenance of intestinal epithelial integrity is not only important for nutrient absorption, but also critical to prevent invasion of the pathogenic microorganisms from the luminal contents. During the last decade, increasing evidence indicates that activation of the endoplasmic reticulum (ER) stress-mediated unfolding protein response (UPR) plays a central role in the maintenance of gastrointestinal mucosal homeostasis (8). Extended activation of ER stress and dysregulated UPR induces the expression of the CCAAT/enhancer-binding protein homologous protein (CHOP), an apoptosis-promoting transcription factor, which has been identified as a major contributor to IBD because it induces epithelial cell death, impairs mucosal barrier function and activates the inflammatory response (8–10). Previous studies reported that ablation of CHOP alleviated dextran sulfate sodium- and trinitrobenzene sulfonic acid-induced experimental colitis (11–13). We also reported that CHOP^{-/-} mice were protected from HIV protease inhibitor (strong ER stress inducer)-induced disruption of intestinal barrier function and hepatic liver injury (14, 15). The intestinal epithelium is highly proliferative and is organized into crypts and villi structures. All types of intestinal epithelia cells are differentiated from a common and constantly renewing leucine-rich repeat-containing G-protein coupled receptor 5 positive (Lgr5⁺) intestinal stem cells (ISCs) (8, 16). Maintenance of normal physiological function of the ER

is critical for cellular differentiation as a large number of specialized proteins require processing in the ER. ER stress has been linked to a loss of intestinal epithelial stemness (17).

It has been reported that BDL-induced cholestatic liver injury was attenuated in CHOP^{-/-} mice through inhibition of hepatocyte apoptosis (18). Although BDL-induced hepatocyte apoptosis was significantly reduced in CHOP^{-/-} mice in 2 days, no significant difference between wild type (WT) and CHOP^{-/-} mice was observed after two weeks (18). These results suggest the other unidentified mechanisms may play more important roles in BDL-induced cholestatic liver injury than CHOP-mediated hepatocyte apoptosis. Based on our previous studies and studies from others, we hypothesize that ER stress-induced disruption of intestinal barrier function plays a critical role in cholestatic liver injury. Here, we used a combination of an *in vivo* 14-day BDL mouse model and *in vitro* cellular techniques to define the mechanisms underlying the CHOP deficiency-mediated protective effects against BDL-induced cholestatic liver injury. Our data reveal a critical role of CHOP in regulating the proliferation and differentiation of ISCs and intestinal epithelia homeostasis. ER stress-mediated rapid loss of intestinal epithelial stemness represents an important mechanism underlying BDL-induced disruption of gut-liver axis and cholestatic liver injury.

Materials and Methods

Animal Studies

C57BL/6J WT mice and CHOP^{-/-} mice (both male and female, 8–10 weeks old) were purchased from the Jackson Laboratory (Bar Harbor, ME) and housed under a 12h light/12h dark cycle with free access to water and normal chow. BDL or sham operation was performed on WT and CHOP^{-/-} mice as described previously for 1, 3, 5, 7 or 14 days (19, 20). At the end of the experiment, the mice were sacrificed for further studies. All animal study protocols were approved by the Institutional Animal Care and Use Committee of Virginia Commonwealth University.

Human liver and intestine samples

De-identified frozen PSC patient liver tissues and normal human liver tissues were obtained through the Liver Tissue Cell Distribution System (LTCDS) (Minneapolis, Minnesota), which was funded by NIH Contract# HSN276201200017C. After written informed consent, outpatients with compensated cirrhosis and age- and sex-matched healthy controls were prospectively recruited. Cirrhosis was diagnosed using fibroscan, liver biopsy or by endoscopic or radiological evidence of portal hypertension in patients with chronic liver disease. None of the patients were abusing alcohol or illicit drugs or were on antibiotics or probiotics. Pinch biopsies from the second part of the duodenum were obtained during an upper endoscopy and were flash frozen in the endoscopy suite and stored at -80°C until further analysis. This protocol was approved by the Institutional Review Board of the McGuire VA Medical Center in Richmond.

Bacterial translocation

Blood and mesenteric lymph nodes (MLNs) were harvested in sterile conditions. Blood and homogenized MLNs were diluted in series and plated on Blood Agar Plates. After 72h incubation at 37°C in aerobic conditions, colony-forming units (CFUs) were counted and calculated as described in a previous study (21).

Isolation of intestinal crypts and intestinal organoid culture

The intestinal crypts and intestinal organoids were isolated and cultured as described previously (22, 23).

Statistical analysis

The results are presented as the mean $\pm$ SE and are from at least three independent experiments. One-way ANOVA and post-test was performed to analyze the differences between multiple groups by GraphPad Prism 5 (GraphPad, San Diego, CA). P values of 0.05 were considered statistically significant.

Please see Supporting Materials and Methods for more-detailed information for this section.

Results

BDL-induced cholestatic liver injury was reduced in CHOP^{-/-} mice

To investigate whether CHOP deficiency had a protective effect against BDL-induced cholestatic liver injury, both age- and sex-matched WT and CHOP^{-/-} mice were subjected to BDL or a sham operation for 14 days. The overall survival rate with BDL surgery in CHOP^{-/-} mice was higher than that in WT mice (Supplementary Fig. 1A). BDL-induced increase of serum total bile acid (TBA), ALP, AST and ALT was significantly reduced in CHOP^{-/-} mice (Fig. 1A). Histological analysis of liver sections indicated that BDL-induced cholestatic liver injury characterized by inflammatory cell infiltration, bile duct over-proliferation, collagen deposition and fibrosis was also significantly reduced in the absence of CHOP (Fig. 1B–C). Consistent with immunohistochemical staining for cytokeratin-19 (CK-19), real-time PCR results also showed that BDL-induced upregulation of CK-19 was markedly reduced in CHOP^{-/-} mice (Supplementary Fig. 1B–D), which was accompanied by a reduction of hepatic stellate cells activation genes, including collagen I, alpha smooth muscle actin (α -SMA), lysyl oxidase like 2 (LOXL2) and Fibronectin (Fig. 1D). In addition, BDL-induced increase of hepatic TBA and hydroxyproline levels in WT mice was also reduced in CHOP^{-/-} mice (Fig. 1E and F).

BDL-induced hepatic inflammatory responses were inhibited in CHOP^{-/-} mice

Profound inflammation is often associated with severe cholestatic liver injury. Consistent with H&E staining indicating massive inflammatory cell infiltration (Fig. 1B), immunohistochemical staining of cluster of differentiation (CD)11b and F4/80 further showed that BDL significantly increased recruitment of CD11b⁺ inflammatory cells into the periportal area accompanied by an increased number of monocyte-derived macrophages (CD11b⁺/F4/80⁺) in WT mice, which was reduced in CHOP^{-/-} mice (Fig. 2A). Real-time PCR results further showed that the mRNA levels of CD68 and F4/80, macrophage markers,

were upregulated by 10- and 8-fold, respectively, in WT BDL mice, but only 3- and 2-fold, respectively, in CHOP^{-/-} BDL mice (Fig. 2B). In addition, CD11b, a marker for recruited inflammatory cells, was also significantly elevated in WT BDL mice, but much less so in CHOP^{-/-} BDL mice (Fig. 2B). As expected, the pro-inflammatory chemokines and cytokines, including monocyte chemoattractant protein-1 (MCP-1), C-X-C motif chemokine 10 (CXCL10), tumor necrosis factor- α (TNF- α) and interleukin 6 (IL-6), were markedly induced by BDL in WT mice, but not in CHOP^{-/-} mice. These results indicate that BDL-induced hepatic inflammatory response was significantly reduced in the absence of CHOP (Fig. 2C–D).

Macrophages have a remarkable plasticity and can differentiate into functionally diverse subtypes, e.g. “classically activated” M1 and “alternatively activated” M2 macrophages (24). Persistent activation of M2 macrophages promotes fibrosis progression. To further define the role of CHOP in promoting hepatic liver fibrosis, we measured the mRNA levels of IL-10, an anti-inflammatory cytokine mainly secreted by M2 macrophages, and several pro-fibrotic cytokines and chemokines, such as tumor growth factor-beta (TGF- β), Resistin-like molecular alpha (Fizz-1), C-C chemokine ligand (CCL)17, CCL22 and CCL24. As shown in Fig. 2E–F, BDL-induced upregulation of IL-10, TGF- β , FIZZ-1, CCL17, CCL22 and CCL24 in WT mice was significantly reduced in CHOP^{-/-} mice.

ER stress-induced hepatocyte apoptosis is not the major trigger to activate hepatic innate immune response in 14-day BDL mouse model

Cholestatic liver injury is associated with dysregulation of the immune response and aberrant expression of inflammatory mediators induced by leak-gut-derived bacteria and endogenous sterile danger signals released from apoptotic hepatocytes (25). As shown in Fig. 3A, BDL induced rapid activation of the UPR in the liver as indicated by a transient upregulation of CHOP, activating transcription factor 4 (ATF4) and 78kDa glucose regulated protein (GRP78) after 1 day that returned to base level after 3 days. In 14-day BDL mice, there was only a slight upregulation of ATF4. Consistent with these findings and previous reports, no significant difference in hepatocyte apoptosis was observed between WT and CHOP^{-/-} mice 14 days following BDL by TdT-Mediated dUTP Nick-End Labeling (TUNEL) assay (Supplementary Fig. 2A) (18). Western blot analysis and immunofluorescence staining of cleaved caspase-3 further confirmed our observations (Fig. 3B–C and Supplementary Fig. 2B). However, the mRNA levels of toll-like receptor (TLR) 2, TLR4, CD14, IL-18 and IL-1 β , key players involved in the bacterial infection-induced innate immunity activation, were not changed during the first few days, but markedly elevated 3–5 days after BDL, reaching peak levels after 7 days and persisting until 14 days (Fig. 3D and F). Similarly, mRNA levels of TLR2, TLR4, CD14, IL-18 and IL-1 β were also significantly elevated in the livers of PSC patients compared to normal controls with most of them being associated with IBD and massive bacterial translocation (Fig 3E). No significant change of the UPR genes was observed (Supplementary Fig. 3). These findings suggest that BDL-induced activation of innate immunity and hepatic inflammation observed 14 days post-BDL is independent of ER stress-induced hepatocyte apoptosis. However, CHOP deficiency significantly ameliorated BDL-induced upregulation of TLR2, TLR4,

CD14, IL-18 and IL-1 β in the liver (Fig. 3F), which indicates the involvement of gut bacterial translocation.

To further identify the specific hepatic cells involved in BDL-induced activation of the UPR and innate immunity, we isolated primary hepatocytes, Kupffer cells, cholangiocytes and HSC from WT BDL and sham control mice and the purity of the isolated cells was checked by real-time RT-PCR with specific hepatic cell markers, CK-19 for MPC, CYP1A2 for primary hepatocytes, f4/80 for Kupffer cells, Desmin for HSC (Supplementary Fig. 4) (26, 27). As shown in Supplementary Fig. 5A, BDL induced CHOP expression in hepatocytes, not in other hepatic cells. However, BDL significantly induced TLR2, TLR4, CD14, IL-18 and IL-1 β in Kupffer cells and hepatocytes (Supplementary Fig. 5B–F).

CHOP deficiency protects against BDL-induced disruption of intestinal barrier function and bacterial translocation

Disruption of internal barrier integrity induces bacterial translocation into other organs and circulation and results in systemic inflammation. Our previous study showed that CHOP deficiency reduced HIV protease inhibitor-induced increase of intestinal permeability (15). Similarly, the BDL-induced increase of gut permeability was also significantly reduced in CHOP^{-/-} mice (Fig. 4A). We further detected the bacteria in the MLNs, the first stop on the route of bacterial translocation, and in the blood. Significant bacterial translocation to the MLNs was first observed at 5 days and increased continuously until 14 days (Supplementary Fig. 6A and Fig. 4B), which was consistent with the time course of innate immunity activation in the liver (Fig. 3C) and intestine (Supplementary Fig. 6B). As shown in Fig. 4B, BDL-induced massive bacterial translocation in MLNs was significantly reduced in CHOP^{-/-} mice. In addition, the colony counts of the bacteria from the MLNs of WT BDL mice was significantly higher than that of bacteria level from the MLNs of CHOP^{-/-} mice. Consistently, more bacteria were detected in the blood of WT BDL mice when compared to that of CHOP^{-/-} mice (Fig. 4C). Furthermore, H&E staining of intestines identified more inflammatory cell infiltration and lymphatic vessel dilation in WT mice than in CHOP^{-/-} mice (Fig. 4D). No obvious inflammatory cell infiltration was detected in the colon (Supplementary Fig. 7A). However, BDL significantly upregulated IL-1 β , TNF- α and MCP-1 in both the intestine and colon, which was alleviated in CHOP^{-/-} mice (Fig. 4E and Supplementary Fig. 7B).

CHOP deficiency reduced BDL-induced disruption of tight junction

Maintenance of the intestinal barrier integrity depends on a variety of mucosal structural components, which are tightly regulated in homeostasis (28). The intercellular junctions are composed of the tight junction, adherens junction and desmosome. The tight junction is the primary determinant of paracellular permeability, and the adherens junction and desmosomes provide the adhesive forces necessary for maintenance of cell-cell interaction (28). To determine whether BDL-induced increase of intestinal permeability is caused by disruption of intercellular junctions, we first examined the mRNA and protein expression levels of tight junction proteins, zonula occludens-1 (ZO-1) and Occludin (OCLN) and adherens junction protein, E-cadherin (E-cad). As shown in Fig. 5A and B, BDL did not affect the mRNA and protein expression of ZO-1, OCLN and E-cad. However, impaired

tight junction structure in the small intestinal epithelium was observed in WT BDL mice. As shown in Fig. 5C, BDL induced a mislocation of OCLN in WT mice, which was significantly improved in CHOP^{-/-} BDL mice. In the colon, BDL significantly suppressed both mRNA and protein expression of OCLN, ZO-1 and E-cad accompanied with the loss of tight junction structure in WT mice, but not in CHOP^{-/-} mice (Supplementary Fig. 8A–C).

CHOP deficiency inhibits BDL-induced intestinal epithelial cell apoptosis

Our previous study demonstrated that ER stress-induced intestinal epithelial cell apoptosis was reduced in CHOP^{-/-} mice (15). To determine whether the CHOP deficiency-mediated protective effect against BDL-induced disruption of intestinal barrier function was due to an inhibition of cell apoptosis in the intestine, we did TUNEL assay in small intestinal sections. As shown in Fig. 5D, following BDL, intestinal epithelial cell apoptosis was significantly increased in WT mice. Apoptotic cells located in the intestinal crypts were also observed, as indicated by black arrows. As expected, intestinal epithelial cell apoptosis was markedly inhibited in CHOP^{-/-} mice. Similar results were also found in the colon (Supplementary Fig. 8D). Consistent with these findings, the pro-apoptotic mediator, CHOP, was significantly induced in WT BDL mice both in the small intestine and colon (Fig. 5E and Supplementary Fig. 8E). BDL induced an upregulation of spliced X-box binding protein-1 (XBP-1s), ATF4 and GRP78 in the intestine of both WT and CHOP^{-/-} mice 14 days following BDL (Fig. 5F), but not in the colon (Supplementary Fig. 8F). The overlapped time courses of activation of hepatic inflammation with ER stress-induced apoptosis in intestinal epithelium indicates that gut-derived bacteria may play a critical role in activation of innate immune response in the liver (Supplementary Fig. 6B–C).

CHOP deficiency relieves BDL-induced suppression of ISCs

The adult intestinal epithelium undergoes rapid renewal to replace the lost epithelial cells caused by toxic contents in the intestinal lumen. ISCs, a small population of cells that reside in the bottom of intestinal crypts, are responsible for daily self-renewal of intestinal epithelium. ISC-derived highly proliferative progenitor cells, transit-amplifying cells, differentiate into the absorptive or secretory epithelial lineage and further migrate to intestinal villi to maintain intestinal epithelium integrity (29). Interestingly, the length of small intestine of CHOP^{-/-} mice was longer than that of WT mice (Supplementary Fig. 9). Histological analysis indicated that BDL significantly reduced the length of both villi and crypts in WT mice, but not in CHOP^{-/-} mice (Figs. 4D and 6A). These results suggest that BDL-induced CHOP expression may inhibit intestinal epithelium renewal by suppressing the proliferation and differentiation of ISCs. To further delineate the underlying mechanisms, the small intestinal crypts were isolated. The real-time PCR results indicate that BDL significantly activated the expression of Lgr5 and Olfm4, the active ISC markers, 3 days after BDL. However, the expression levels of Lgr5 and olfactomedin (Olfm) 4 were significantly inhibited 5–14 days after BDL in WT mice (Fig. 6B). Interestingly, BDL had no significant effect on the expression of both Lgr5 and Olfm4 in CHOP^{-/-} mice. Similarly, BDL-induced inhibition of Lgr5 in the colon of WT mice was also reversed in CHOP^{-/-} mice (Supplementary Fig. 10A).

ISC self-renewal and maintenance of stemness highly depends on several signaling pathways in the stem cell niche, especially the Wnt/ β -catenin pathway. c-Myc, a downstream signaling molecule in the Wnt/ β -catenin signaling pathway, is critical for the expression of a core set of Wnt-target genes. As shown in Fig. 6C, the mRNA level of c-Myc was downregulated 5 days after BDL and persisted until 14 days in WT mice, but remained unchanged in CHOP^{-/-} mice. IHC results further demonstrated that c-Myc was markedly downregulated in both crypt base stem cells and crypt TA cells in BDL WT mice, but not in BDL CHOP^{-/-} mice 14 days after BDL. In support of this finding, Ki67 was also examined. As shown in Fig. 6C–D, its expression was significantly inhibited in 7–14 days after BDL in WT mice but not in CHOP^{-/-} mice. Similar results were also found in the colon crypts (Supplementary Fig. 10B and C). To further examine the effect of BDL on intestinal epithelial cell turnover, BrdU staining was performed. As shown in Fig. 6E, BDL induced a significant suppression of intestinal epithelium turnover in WT mice in 7–14 days after BDL, but not in CHOP^{-/-} mice. These results indicate that CHOP deficiency significantly relieved BDL-induced suppression of ISCs and intestinal epithelium turnover.

It has been reported that activation of the UPR resulted in rapid loss of stemness (17). Here, we found that BDL gradually induced CHOP, ATF4 and GRP78 activation starting from 5 days after BDL in intestinal crypts, which shared a similar time course of bacterial translocation and loss of stemness in intestinal crypt cells (Supplementary Fig. 6A, Fig. 6B and F). Although no significant difference in GRP78 expression between WT and CHOP^{-/-} mice was observed, activation of ATF4 in intestinal crypts was inhibited in CHOP^{-/-} mice (Fig. 6F). These results suggest that CHOP is a major player in ER stress-induced suppression of ISC differentiation.

CHOP deficiency protects lipopolysaccharide (LPS)- and ER stress-induced loss of stemness in cultured intestinal epithelial organoids

To delineate the role of CHOP on regulating ISC proliferation and differentiation, the organoids of primary intestinal epithelium from WT and CHOP^{-/-} mice were cultured. As shown in Fig. 7A, intestinal organoids were formed in 2 days and markedly expanded in 10 days. The overall size of CHOP^{-/-} intestinal organoids was significantly larger than that of WT organoids. In addition, intestinal organoids derived from CHOP^{-/-} mice consisted of more crypts that were longer in structure than those from WT mice, which was consistent with the *in vivo* findings that CHOP^{-/-} mice had a longer intestine with more intestinal villi and crypts when compared to WT mice (Supplementary Fig. 9 and Fig. 6A).

To further determine the direct effect of bacterial toxin and ER stress activation on ISC differentiation, LPS and thapsigargin (TG), a known ER stress inducer, were added to the organoid culture medium at day 4 until day 10. As shown in Fig. 7B, no significant impact on both WT and CHOP^{-/-} intestinal organoids was found for the first 2 days of treatment, but LPS induced a more differentiated epithelial cell shedding in the lumen-like structure of WT organoids (indicated by white arrows) and led to the destruction of WT organoid crypts (indicated by black arrows) after 4 days. Strikingly, much less epithelial cell shedding and no destruction of crypts was observed in CHOP^{-/-} organoids. Furthermore, TG rapidly inhibited intestinal organoid formation as indicated by the smaller size and fewer crypts, 2

days after treatment in WT organoids, but had no significant effect on CHOP^{-/-} organoids, even after 6-day treatment. In line with these findings, the mRNA level of the active stem cell marker Lgr5 was significantly higher in CHOP^{-/-} intestinal organoids (Fig 7C). In addition, both LPS and TG significantly inhibited Ki67 expression in WT intestinal organoids indicating reduced stem cell proliferation, but had no effect on CHOP^{-/-} organoids (Fig. 7D). In support of the real-time PCR results, immunofluorescence staining also showed decreased numbers of proliferative crypt cells, indicated by nuclear staining of proliferating cell nuclear antigen (PCNA) (White *), in LPS- and TG-treated WT organoids, but not in CHOP^{-/-} organoids (Supplementary Fig. 11). Furthermore, both LPS and TG significantly induced CHOP expression in WT intestinal organoids, indicating the critical role of ER stress-induced CHOP expression in inhibiting ISC proliferation and differentiation (Fig. 7E).

Activation of ER stress and inflammation in the duodenum of cirrhotic patients

To further determine the clinical relevance of activation of intestinal ER stress and inflammatory response in human patients with cirrhosis, we examined the mRNA levels of CHOP, ATF4, GRP78 and Xbp1s in duodenal biopsy samples from thirteen cirrhotic patients and a similar number of age-/sex-matched healthy controls (cirrhosis 56±5 years old, controls 54±6 years old, 8 men in both groups). All cirrhotic patients had active hepatitis C as the etiology of cirrhosis. Their mean model for end-stage liver disease (MELD) score of 9±4 and all were Child class A. As shown in Supplementary Fig. 12A, mRNA levels of CHOP, ATF4 and GRP78 were increased in the intestine of cirrhotic patients. Similar to our findings in BDL mice, intestinal mRNA levels of CD14, IL-1β, TNF-α and MCP-1 were also increased in cirrhotic patients (Supplementary Fig. 12B). In addition, the mRNA levels of LGR5, SOX-9, c-Myc and Ki67 were reduced indicating the suppression of stem cell proliferation (Supplementary Fig.12C).

Discussion

Bile duct obstruction plays a critical role in cholestatic liver disease progression, but the pathological mechanisms remain incompletely understood. The current study identifies a novel mechanism underlying BDL-induced cholestatic liver injury. Our results indicate that CHOP-mediated rapid loss of intestinal epithelial stemness and subsequent disruption of the gut-liver axis plays a crucial role in cholestatic liver disease progression.

CHOP is a key transcription factor responsible for ER stress-mediated cell injury and has been implicated in various pathological processes including gastrointestinal disorders and cholestatic liver injury (7, 10, 18, 30–33). A previous study reported that CHOP^{-/-} mice were protected from BDL-induced cholestatic liver injury without significant effect on hepatocyte apoptosis (18). Another study with the CCl₄-induced acute liver injury mouse model reported that CHOP was rapidly upregulated after CCl₄ treatment. However, CHOP deficiency had no protective effect against CCl₄-induced liver injury (34). These studies suggest that the role of CHOP in BDL-induced cholestatic liver injury and acute liver injury may be different. In addition, CHOP was rapidly activated in the liver 1 day after BDL, but returned to basal levels after 3 days. However, the innate immune response was significantly

activated 3 days after BDL. Furthermore, no significant difference of hepatocyte apoptosis were observed 14 days after BDL between WT and CHOP^{-/-} mice. All of these results suggest that CHOP-mediated hepatocyte apoptosis may only play a minimal role in BDL-cholestatic liver injury.

Inflammation is a remarkable feature of the cholangiopathies (25). To further identify the key mechanisms underlying CHOP deficiency-mediated protective effect against BDL-cholestatic injury, we examined the effect of CHOP on the gut-liver axis under cholestatic conditions. A fundamental function of the intestinal epithelium is to prevent the underlying immune system and the remainder of the body from harmful substances and pathogens (28). Compromised intestinal barrier function has been associated with various human diseases including liver diseases, such as PSC (35). The intestinal epithelium is a polarized single cell-layer organized in crypts and villi, which is constantly exposed to various microbial toxins and ER stress inducers and has an extremely high turnover rate (7). Excessive loss or reduced renewal of epithelial cells results in the disruption of epithelial layer integrity and an increase of intestinal permeability. In the BDL mouse model, the time courses of BDL-induced bacterial translocation, activation of intestinal ER stress, induction of intestinal epithelium apoptosis, suppression of intestinal turnover and activation of hepatic inflammation, but not hepatocyte apoptosis notably overlapped. BDL-induced liver injury and disruption of intestinal barrier function were all significantly reduced in CHOP^{-/-} mice. (Figs. 1, 2, 4, 5, and Supplementary Fig. 1–2). By using both the *in vivo* BDL mouse model and the *in vitro* intestinal organoid culture model, we identified the important mechanisms of BDL-induced cholestatic liver injury. Furthermore, we also confirmed that our findings in BDL mice are clinically relevant to human patients with cirrhosis, even without cholestatic injury as the etiology (Supplementary Figs. 3&11). Our data strongly support our hypothesis that ER stress-mediated disruption of intestinal barrier function plays a critical role in cholestatic liver injury.

It has been reported that ER stress causes rapid loss of intestinal epithelial stemness through activation of the UPR (17). Maintenance of self-renewal and proliferation of ISCs is critical for physiological function of the intestines, especially for post-injury intestinal repair (29, 36). Our study demonstrates for the first time that BDL-induced activation of ER stress and the UPR, especially CHOP activation, is associated with a loss of intestinal epithelial stemness and further disruption of intestinal epithelium turnover. BDL had no direct effect on the expression of intestinal epithelial junction proteins, but increased intestinal epithelial cell apoptosis and suppressed epithelium turnover, which subsequently induced dislocation of OCLN and E-Cad, resulting in the disruption of intestinal barrier integrity. Most importantly, CHOP deficiency significantly increased proliferation of ISCs as indicated by an increase of size and number of intestinal crypts in *in vitro* organoid culture (Fig. 7). This was consistent with the *in vivo* findings that CHOP^{-/-} mice had longer intestines and colons as well as more crypts and villi in the intestine (Figs. 4 and 6). Furthermore, both LPS- and TG-induced loss of stemness was attenuated with CHOP deficiency (Fig. 7).

In addition to the recently established correlation between PSC and IBD, intestinal barrier dysfunction-induced bacterial translocation was suggested as a critical cause of end-stage liver diseases (5). It has been reported that obstruction of bile flow causes overgrowth of

intestinal bacteria and mucosal injury, which can lead to bacterial translocation and systemic infection (37). It also has been reported that enterohepatic bacterial infections suppress the intestinal expression of FGF15. The decrease in ileal FGF15 expression was associated with liver bacterial colonization and hepatobiliary pathophysiology (38). We also found that BDL induced down-regulation of intestinal FGF15 and SHP expression, which was reversed in CHOP^{-/-} mice (Supplementary Fig. 13). In addition, the basal levels of intestinal and hepatic SHP mRNAs were higher in CHOP^{-/-} mice than those in WT mice. Consistent with previously published studies, we also found that the inflammatory mediators were significantly upregulated in human PSC patient livers (Fig. 3E) (39). Our results indicate that BDL induced bacterial translocation and massive infiltration of inflammatory cells, which was markedly reduced in the absence of CHOP.

Activation of Kupffer cells (KCs) plays a pivotal role in the pathogenesis of liver fibrosis (40). Multiple lines of evidence from previous reports and our current study suggest that persistent inflammation further triggers macrophage phenotype transition from pro-inflammatory M1 (classically activated) macrophages to M2 (alternatively activated) macrophages (41–43). M2 macrophages promote wound healing and damaged tissue regeneration through the production of a set of pro-fibrotic cytokines and growth factors, including TGF- β , CCL-17, CCL-22, CCL-24 and Fizz-1 (24, 44). Macrophage depletion attenuates extracellular matrix deposition and ductular reaction in a mouse model of chronic cholangiopathies (45). In addition, ER stress has been identified as a key regulator of macrophage differentiation (46). Our study suggests that the protective effect of CHOP deficiency against BDL-induced hepatic inflammation and cholestatic injury was mainly due to the improvement of intestinal barrier function and inhibition of bacterial translocation and inflammation.

In summary, the current study demonstrates that ER stress-mediated intestinal barrier dysfunction represents a key event in BDL-induced cholestatic liver injury. As illustrated in Fig. 8, ER stress-induced activation of CHOP suppresses ISC proliferation and intestinal epithelial cell renewal, which leads to disruption of intestinal barrier function, massive bacterial translocation, activation of KCs and periportal inflammation and eventually results in fibrosis in the liver. Here, we have provided strong evidence that targeting CHOP represents a novel therapeutic strategy for cholestatic liver diseases as well as other inflammatory liver diseases.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding support: This work was supported by National Institutes of Health Grant R01 DK104893 (to HZ and PBH), R01DK-057543 (to PBH and HZ), VA Merit Review I01BX001390 (to HZ) and I0CX001076 (to JSB), National Natural Science Foundation of China Grants 81070245 and 81270489 (to H.Z.); Massey Cancer Center pilot grant (to HZ and PBH). Microscopy was performed at the VCU Microscopy Facility, supported in part by funding from NIH-NCI Cancer Center Grant P30 CA016059.

We would like to thank Elaine Kennedy and Yunzhou Li from Forte Editing Group for proofreading and editing. We would like to thank Dr. Shannon Glaser (Texas A&M HSC, Temple, TX) for helping with isolation of primary hepatocytes, Kupffer cells, cholangiocytes and HSCs from BDL mice.

Abbreviation

PSC	primary sclerosing cholangitis
PBC	primary biliary cholangitis
IBD	inflammatory bowel disease
BDL	bile duct ligation
ER	endoplasmic reticulum
UPR	unfolding protein response
CHOP	C/EBP homologous protein
Lgr5	leucine rich repeat containing G protein-coupled receptor 5
ISCs	intestinal stem cells
WT	wild type
MLN	mesenteric lymph node
CFUs	colony-forming units
TBA	total bile acid
CK-19	cytokeratin-19
HSC	hepatic stellate cells
α-SMA	alpha smooth muscle actin
LOXL2	lysyl oxidase like 2
CD	cluster of differentiation
MCP-1	monocyte chemotactic protein-1
CXCL10	C-X-C motif chemokine 10
TNF-α	tumor necrosis factor-alpha
IL	interleukin
TGF-β	tumor growth factor-beta
FIZZ1	Resistin-like molecular alpha
CCL	C-C chemokine ligand
ATF4	activating transcription factor 4

GRP78	78kDa glucose regulated protein
TUNEL	TdT-Mediated dUTP Nick-End Labeling
TLR	toll like receptor
ZO-1	zonula occluden-1
OCLN	Occludin
E-cad	E-cadherin
XBP-1s	X-box binding protein 1 spliced
Olfm4	olfactomedin 4
LPS	liposaccharide
TG	thapsigargin
PCNA	Proliferating cell nuclear antigen
KCs	Kupffer cells

References

1. Liu R, Zhao R, Zhou X, Liang X, Campbell DJ, Zhang X, Zhang L, et al. Conjugated bile acids promote cholangiocarcinoma cell invasive growth through activation of sphingosine 1-phosphate receptor 2. *Hepatology*. 2014; 60:908–918. [PubMed: 24700501]
2. Hylemon PB, Zhou H, Pandak WM, Ren S, Gil G, Dent P. Bile acids as regulatory molecules. *J Lipid Res*. 2009; 50:1509–1520. [PubMed: 19346331]
3. Chazouilleres O. Novel Aspects in the Management of Cholestatic Liver Diseases. *Dig Dis*. 2016; 34:340–346. [PubMed: 27170387]
4. Eksteen B. The Gut-Liver Axis in Primary Sclerosing Cholangitis. *Clin Liver Dis*. 2016; 20:1–14. [PubMed: 26593287]
5. Wiest R, Lawson M, Geuking M. Pathological bacterial translocation in liver cirrhosis. *J Hepatol*. 2014; 60:197–209. [PubMed: 23993913]
6. Louvet A, Mathurin P. Alcoholic liver disease: mechanisms of injury and targeted treatment. *Nat Rev Gastroenterol Hepatol*. 2015; 12:231–242. [PubMed: 25782093]
7. Delgado ME, Grabinger T, Brunner T. Cell death at the intestinal epithelial front line. *FEBS J*. 2016; 283:2701–2719. [PubMed: 26499289]
8. Luo K, Cao SS. Endoplasmic reticulum stress in intestinal epithelial cell function and inflammatory bowel disease. *Gastroenterol Res Pract*. 2015; 2015:328791. [PubMed: 25755668]
9. Zhou H, Liu R. ER stress and hepatic lipid metabolism. *Front Genet*. 2014; 5:112. [PubMed: 24847353]
10. Cao SS. Endoplasmic reticulum stress and unfolded protein response in inflammatory bowel disease. *Inflamm Bowel Dis*. 2015; 21:636–644. [PubMed: 25581827]
11. Kaser A, Adolph TE, Blumberg RS. The unfolded protein response and gastrointestinal disease. *Semin Immunopathol*. 2013; 35:307–319. [PubMed: 23588234]
12. Sabino J, Vieira-Silva S, Machiels K, Joossens M, Falony G, Ballet V, Ferrante M, et al. Primary sclerosing cholangitis is characterised by intestinal dysbiosis independent from IBD. *Gut*. 2016; 65:1681–1689. [PubMed: 27207975]
13. Hue S, Ahern P, Buonocore S, Kullberg MC, Cua DJ, McKenzie BS, Powrie F, et al. Interleukin-23 drives innate and T cell-mediated intestinal inflammation. *J Exp Med*. 2006; 203:2473–2483. [PubMed: 17030949]

14. Wang Y, Zhang L, Wu X, Gurley EC, Kennedy E, Hylemon PB, Pandak WM, et al. The role of CCAAT enhancer-binding protein homologous protein in human immunodeficiency virus protease-inhibitor-induced hepatic lipotoxicity in mice. *Hepatology*. 2013; 57:1005–1016. [PubMed: 23080229]
15. Wu X, Sun L, Zha W, Studer E, Gurley E, Chen L, Wang X, et al. HIV protease inhibitors induce endoplasmic reticulum stress and disrupt barrier integrity in intestinal epithelial cells. *Gastroenterology*. 2010; 138:197–209. [PubMed: 19732776]
16. Tan S, Barker N. Epithelial stem cells and intestinal cancer. *Semin Cancer Biol*. 2015; 32:40–53. [PubMed: 24560652]
17. Heijmans J, van Lidth de Jeude JF, Koo BK, Rosekrans SL, Wielenga MC, van de Wetering M, Ferrante M, et al. ER stress causes rapid loss of intestinal epithelial stemness through activation of the unfolded protein response. *Cell Rep*. 2013; 3:1128–1139. [PubMed: 23545496]
18. Tamaki N, Hatano E, Taura K, Tada M, Kodama Y, Nitta T, Iwaisako K, et al. CHOP deficiency attenuates cholestasis-induced liver fibrosis by reduction of hepatocyte injury. *Am J Physiol Gastrointest Liver Physiol*. 2008; 294:G498–505. [PubMed: 18174271]
19. Wang Y, Aoki H, Yang J, Peng K, Liu R, Li X, Qiang X, Sun L, Gurley EC, Lai G, Zhang L, Laing G, Nagahashi M, Takebe K, Pandak WM, Hylemon PB, Zhou H. The role of S1PR2 in bile acid-induced cholangiocyte proliferation and cholestasis-induced liver injury in mice. *Hepatology*. 2017 In press.
20. Aoki H, Aoki M, Yang J, Katsuta E, Mukhopadhyay P, Ramanathan R, Woelfel IA, et al. Murine model of long-term obstructive jaundice. *J Surg Res*. 2016; 206:118–125. [PubMed: 27916350]
21. Fouts DE, Torralba M, Nelson KE, Brenner DA, Schnabl B. Bacterial translocation and changes in the intestinal microbiome in mouse models of liver disease. *J Hepatol*. 2012; 56:1283–1292. [PubMed: 22326468]
22. Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen M, Haegbarth A, et al. Identification of stem cells in small intestine and colon by marker gene *Lgr5*. *Nature*. 2007; 449:1003–1007. [PubMed: 17934449]
23. Sato T, Stange DE, Ferrante M, Vries RG, Van Es JH, Van den Brink S, Van Houdt WJ, et al. Long-term expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and Barrett's epithelium. *Gastroenterology*. 2011; 141:1762–1772. [PubMed: 21889923]
24. Heymann F, Trautwein C, Tacke F. Monocytes and macrophages as cellular targets in liver fibrosis. *Inflamm Allergy Drug Targets*. 2009; 8:307–318. [PubMed: 19534673]
25. Hirschfield GM, Heathcote EJ, Gershwin ME. Pathogenesis of cholestatic liver disease and therapeutic approaches. *Gastroenterology*. 2010; 139:1481–1496. [PubMed: 20849855]
26. Assani K, Shrestha CL, Robledo-Avila F, Rajaram MV, Partida-Sanchez S, Schlesinger LS, Kopp BT. Human Cystic Fibrosis Macrophages Have Defective Calcium-Dependent Protein Kinase C Activation of the NADPH Oxidase, an Effect Augmented by *Burkholderia cenocepacia*. *J Immunol*. 2017
27. Yu Y, Sitaraman S, Gewirtz AT. Intestinal epithelial cell regulation of mucosal inflammation. *Immunol Res*. 2004; 29:55–68. [PubMed: 15181270]
28. Odenwald MA, Turner JR. The intestinal epithelial barrier: a therapeutic target? *Nat Rev Gastroenterol Hepatol*. 2017; 14:9–21. [PubMed: 27848962]
29. Barker N. Adult intestinal stem cells: critical drivers of epithelial homeostasis and regeneration. *Nat Rev Mol Cell Biol*. 2014; 15:19–33. [PubMed: 24326621]
30. Cao J, Dai DL, Yao L, Yu HH, Ning B, Zhang Q, Chen J, et al. Saturated fatty acid induction of endoplasmic reticulum stress and apoptosis in human liver cells via the PERK/ATF4/CHOP signaling pathway. *Mol Cell Biochem*. 2012; 364:115–129. [PubMed: 22246806]
31. Chikka MR, McCabe DD, Tyra HM, Rutkowski DT. C/EBP homologous protein (CHOP) contributes to suppression of metabolic genes during endoplasmic reticulum stress in the liver. *J Biol Chem*. 2013; 288:4405–4415. [PubMed: 23281479]
32. Rao J, Zhang C, Wang P, Lu L, Qian X, Qin J, Pan X, et al. C/EBP homologous protein (CHOP) contributes to hepatocyte death via the promotion of ERO1 α signalling in acute liver failure. *Biochem J*. 2015; 466:369–378. [PubMed: 25387528]

33. Cao SS. Epithelial ER Stress in Crohn's Disease and Ulcerative Colitis. *Inflamm Bowel Dis*. 2016; 22:984–993. [PubMed: 26950312]
34. Campos G, Schmidt-Heck W, Ghallab A, Rochlitz K, Putter L, Medinas DB, Hetz C, et al. The transcription factor CHOP, a central component of the transcriptional regulatory network induced upon CCl4 intoxication in mouse liver, is not a critical mediator of hepatotoxicity. *Arch Toxicol*. 2014; 88:1267–1280. [PubMed: 24748426]
35. Konig J, Wells J, Cani PD, Garcia-Rodenas CL, MacDonald T, Mercenier A, Whyte J, et al. Human Intestinal Barrier Function in Health and Disease. *Clin Transl Gastroenterol*. 2016; 7:e196. [PubMed: 27763627]
36. Peterson LW, Artis D. Intestinal epithelial cells: regulators of barrier function and immune homeostasis. *Nat Rev Immunol*. 2014; 14:141–153. [PubMed: 24566914]
37. Inagaki T, Moschetta A, Lee YK, Peng L, Zhao G, Downes M, Yu RT, et al. Regulation of antibacterial defense in the small intestine by the nuclear bile acid receptor. *Proc Natl Acad Sci U S A*. 2006; 103:3920–3925. [PubMed: 16473946]
38. Romain G, Tremblay S, Arena ET, Antunes LC, Covey S, Chow MT, Finlay BB, et al. Enterohepatic bacterial infections dysregulate the FGF15-FGFR4 endocrine axis. *BMC Microbiol*. 2013; 13:238. [PubMed: 24165751]
39. Pichler C, Boettler T, Thimme R. Cholestatic liver disease. *Dtsch Med Wochenschr*. 2016; 141:1683–1687. [PubMed: 27855457]
40. Cheng Y, Tian Y, Xia J, Wu X, Yang Y, Li X, Huang C, et al. The role of PTEN in regulation of hepatic macrophages activation and function in progression and reversal of liver fibrosis. *Toxicol Appl Pharmacol*. 2017; 317:51–62. [PubMed: 28095306]
41. Wynn TA. Fibrotic disease and the T(H)1/T(H)2 paradigm. *Nat Rev Immunol*. 2004; 4:583–594. [PubMed: 15286725]
42. Tacke F, Zimmermann HW. Macrophage heterogeneity in liver injury and fibrosis. *J Hepatol*. 2014; 60:1090–1096. [PubMed: 24412603]
43. Braga TT, Agudelo JS, Camara NO. Macrophages During the Fibrotic Process: M2 as Friend and Foe. *Front Immunol*. 2015; 6:602. [PubMed: 26635814]
44. Zimmermann HW, Tacke F. Modification of chemokine pathways and immune cell infiltration as a novel therapeutic approach in liver inflammation and fibrosis. *Inflamm Allergy Drug Targets*. 2011; 10:509–536. [PubMed: 22150762]
45. Best J, Verhulst S, Syn WK, Lagaisse K, van Hul N, Heindryckx F, Sowa JP, et al. Macrophage Depletion Attenuates Extracellular Matrix Deposition and Ductular Reaction in a Mouse Model of Chronic Cholangiopathies. *PLoS One*. 2016; 11:e0162286. [PubMed: 27618307]
46. Oh J, Riek AE, Weng S, Petty M, Kim D, Colonna M, Cella M, et al. Endoplasmic reticulum stress controls M2 macrophage differentiation and foam cell formation. *J Biol Chem*. 2012; 287:11629–11641. [PubMed: 22356914]

Fig. 1. CHOP deficiency protected BDL-induced liver injury

WT and CHOP^{-/-} mice were subjected to BDL or sham operation for 14 days. **(A)** Total bile acid (TBA) levels, ALP, AST and ALT activity levels in serum were determined. **(B)** Representative images of H&E, Masson's Trichrome staining of liver paraffin sections. Scale bar 100μm. **(C)** Metavir fibrosis score based on H&E and Masson's Trichrome images. **(D)** The relative mRNA levels of α-SMA, Collagen-1, LOXL2 and Fibronectin in the liver were measured by qRT-PCR and normalized using HPRT1 as an internal control. **(E)** Hepatic TBA levels. **(F)** Hepatic hydroxyproline levels. Results are represented as Mean

± S.E. (n=6–9). Statistical significance relative to the WT sham group, *p<0.05, **p<0.01, ***p<0.001; relative to the WT BDL group, #p<0.05, ##p<0.01, ###p<0.001.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 2. CHOP deficiency attenuated BDL-induced hepatic inflammatory responses

(A) Representative images of IHC against CD11b, F4/80, merged images with counterstaining of DAPI, and magnified views of portal area from liver sections of 14-day BDL or sham WT and CHOP^{-/-} mice. Scale bar 50 μm. (B–F) The relative mRNA levels of target genes in the liver of 14-day BDL or sham WT and CHOP^{-/-} mice were measured by qRT-PCR and normalized using HPRT1. Results are represented as Mean ± S.E. (n=6–9). Statistical significance relative to the WT sham group, *p<0.05, **p<0.01, ***p<0.001; relative to the WT BDL group, #p<0.05, ##p<0.01.

Fig. 3. The effect of CHOP deficiency on BDL-induced hepatocyte apoptosis and innate immunity

(A) The relative mRNA levels of CHOP, ATF4 and GRP78 in the livers of WT mice 1, 3, 5, 7 and 14 days after BDL. (B) Representative immunoblot images of cleaved-caspase-3 in liver samples of 14-day BDL or sham WT and CHOP^{-/-} mice. The relative protein levels of cleaved-caspase-3 were analyzed using β-Actin as a loading control. (C) WT and CHOP^{-/-} mice were subjected to BDL or sham operation for 14 days. Representative images of immunofluorescent staining against cleaved-caspase-3 and merged images with DAPI counterstaining are shown. The number of cleaved-caspase-3 positive cells was counted. Scale bar 50 μm. (D) The relative mRNA levels of TLR2, TLR4, CD14, IL-18 and IL-1β, in

GRP78 in the liver of WT mice 1, 3, 5, 7 and 14 days after BDL. **(E)** The relative mRNA levels of TLR2, TLR4, CD14, IL-18 and IL-1 β in the livers of normal human and PSC patient. **(F)** The relative mRNA levels of TLR2, TLR4, CD14, IL-18 and IL-1 β in the liver of WT and CHOP^{-/-} mice with 14-day BDL. Results are represented as mean $\pm$ S.E. from each group (n=5–9 for mice samples, n=10 for human samples). Statistical significance relative to the WT sham group or BDL 0 day group, *p<0.05, **p<0.01, ***p<0.001; relative to the WT BDL group, #p<0.05, ##p<0.01, ###p<0.001.

Fig. 4. CHOP^{-/-} mice were protected from BDL-induced disruption of intestinal barrier function and bacterial translocation

WT and CHOP^{-/-} mice were subjected to BDL or sham operation for 14 days. (A) FITC-Dextran levels in the serum. Colony-forming units (CFUs) of bacteria from (B) MLNs and (C) blood. (D) Representative images of H&E staining of small intestine sections and magnified views. Yellow arrows indicate infiltration of inflammatory cells into intestinal epithelium. Red arrows indicate dilated lymph vessel. Scale bar 200 μm . (E) The relative mRNA levels of IL-1 β , TNF- α and MCP-1 in the small intestine were measured by qRT-PCR and normalized using HPRT1. Results are represented as Mean $\pm$ S.E. from each group (n=5–8). Statistical significance relative to the WT sham group, *p<0.05, **p<0.01, ***p<0.001; relative to the WT BDL group, #p<0.05, ##p<0.01.

Fig. 5. CHOP deficiency attenuated BDL-induced loss of tight junctions and apoptosis in small intestine

WT and CHOP^{-/-} mice were subjected to BDL or sham operation for 14 days. **(A)** The relative mRNA levels of OCLN, ZO-1 and E-Cad in the intestine, normalized using HPRT1. **(B)** The protein levels of OCLN, E-Cad and β -Actin in the small intestine villi were determined by Western blot analysis. Representative images are shown. **(C)** Representative images of immunofluorescent staining of small intestine sections against E-Cadherin, OCLN, and merged images with counterstaining of DAPI are shown. Scale bar 50 μ m. **(D)** TUNEL assay of small intestine sections. The number of TUNEL positive cells in intestinal epithelium is shown. Scale bar 50 μ m. **(E-F)** The relative mRNA levels of CHOP, XBP-1s,

ATF4 and GRP78 in intestine were measured by qRT-PCR and normalized using HPRT1. Results are represented as Mean $\pm$ S.E. from each group (n=5–8). Statistical significance relative to the WT sham group, *p<0.05, ***p<0.001; relative to the WT BDL group, #p<0.05.

Fig. 6. CHOP^{-/-} mice were protected from BDL-induced suppression of ISCs
 (A) WT and CHOP^{-/-} mice were subjected to BDL or sham operation for 14 days. The length of intestinal villi and intestinal crypts was measured based on the H&E staining images. (B–C) WT and CHOP^{-/-} mice were subjected to BDL or sham operation for 3, 5, 7 or 14 days. The relative mRNA levels of Lgr5, Olfm4, c-Myc and Ki67 in isolated intestinal crypts were measured by qRT-PCR and normalized using HPRT1. (D) IHC staining against c-Myc, Ki67 in the small intestine, and magnified Ki67 staining in the intestinal crypt base from WT and CHOP^{-/-} mice with 14-day BDL. Scale bar 50 μ m. (E) The representative images of BrdU staining in the small intestine from WT and CHOP^{-/-} mice subjected to BDL for 3, 7, and 14 days. Scale bar 200 μ m. (F) The relative mRNA levels of CHOP, ATF4

and GRP78 in the small intestine from WT and CHOP^{-/-} mice 3, 7, and 14 days after BDL were measured by qRT-PCR and normalized using HPRT1. Results are represented as Mean $\pm$ S.E. from each group (n=6–10). Statistical significance relative to the WT sham group or BDL 0-day group, *p<0.05, **p<0.01; relative to the WT BDL group, #p<0.05, ##p<0.01.

Fig. 7. CHOP deficiency prevented LPS- and TG-induced suppression of intestinal organoid growth

(A) Representative images of WT and CHOP^{-/-} intestinal organoids cultured for 2, 4, 6, 8 and 10 days. (B) WT and CHOP^{-/-} intestinal organoids were cultured for 4 days, then treated with LPS (500 ng/ml) or TG (200 nM) for 6 days. Representative images at 2, 4, 6 days are shown. (C–E) The relative mRNA levels of Lgr5, Ki67 and CHOP in crypt structures of intestinal organoids after 6-day treatment were measured by qRT-PCR and normalized using HPRT1. Results are represented as Mean ± S.E. from each group (n=3). Statistical significance relative to the WT organoids vehicle control group, *p<0.05, **p<0.01; relative to the WT organoids counterparts, #p<0.05, ###p<0.001.

Fig. 8.
Schematic diagram of the proposed mechanism underlying the protective effects of CHOP deficiency on BDL-induced intestine dysfunction and cholestatic liver injury.