Published in final edited form as:

Sci Signal. ; 11(520): . doi:10.1126/scisignal.aan6480.

A role for corticotrophin releasing factor signaling in the lateral habenula and its modulation by early life stress*

Michael E. Authement^{1,*}, Ludovic D. Langlois^{1,*}, Ryan D. Shepard^{1,*}, Caroline A. Browne¹, Irwin Lucki¹, Haifa Kassis¹, and Fereshteh S. Nugent^{1,†}

¹Uniformed Services University of the Health Sciences, Department of Pharmacology, Bethesda, Maryland 20814, USA

Abstract

Centrally released corticotrophin releasing factor or hormone (extrahypothalamic CRF or CRH) in the brain is involved in behavioral and emotional responses to stress. The lateral habenula (LHb) is an epithalamic brain region involved in value-based decision making and stress evasion. Through its inhibition of dopamine-mediated reward circuitry, increased activity of the LHb is associated with addiction, depression, schizophrenia and behavior disorders. Here, we found that extrahypothalamic CRF neurotransmission increased neuronal excitability in the LHb. Through its receptor CRFR1 and subsequently protein kinase A (PKA), CRF application increased the intrinsic excitability of LHb neurons by affecting changes in small- and large-conductance SK- and BK-type K⁺ channels. CRF also reduced inhibitory GABAergic synaptic transmission onto LHb neurons through endocannabinoid (eCB)-mediated retrograde signaling. Maternal deprivation is a severe early life stress that alters CRF neural circuitry and is likewise associated with abnormal mental health later in life. LHb neurons from pups deprived of maternal care exhibited increased intrinsic excitability, reduced GABAergic transmission, decreased abundance of SK2 channel protein, and increased activity of PKA, without any significant changes in *Crh* or *Crhr1* expression. Notably, maternal deprivation blunted the response of LHb neurons to subsequent, acute CRF exposure. Activating SK channels or inhibiting postsynaptic PKA activity prevented the effects of both CRF and maternal deprivation on LHb intrinsic excitability, thus identifying potential pharmacological targets to reverse central CRF circuit dysregulation in patients with associated disorders.

*"This manuscript has been accepted for publication in Science Signaling. This version has not undergone final editing. Please refer to the complete version of record at <http://www.sciencesignaling.org/>. The manuscript may not be reproduced or used in any manner that does not fall within the fair use provisions of the Copyright Act without the prior, written permission of AAAS."

†Corresponding author. fereshteh.nugent@usuhs.edu.

*These authors contributed equally to this work.

Author contributions: F.S.N designed the research; M.E.A, L.D.L, R.D.S, H.K, and F.S.N performed the experiments; C.A.B and I.L helped to design, perform, analyze and interpret qPCR experiments. M.E.A, L.D.L, R.D.S, and F.S.N analyzed the data and prepared the figures; M.E.A., L.D.L, R.D.S and F.S.N wrote the paper.

Competing interests: The authors declare that they have no conflicts of interest.

Introduction

Corticotropin releasing factor [CRF; also known as corticotropin releasing hormone (CRH)] mediates behavioral responses to stress and is implicated in a multitude of stress and mood related disorders. Although widely known for its participation in stress-induced activation of the hypothalamic-pituitary-adrenal (HPA) axis, CRF also acts within other regions of the brain where it directly contributes to stress responsive behavior. This extrahypothalamic CRF system includes brain regions whose activities are implicated in mental illness (1–4). Many of these brain regions, including the ventral tegmental area (VTA), extended amygdala, raphe nuclei, the bed nucleus of the stria terminalis (BNST) and medial prefrontal cortex (mPFC) are under intense investigation with regards to their role in the extrahypothalamic CRF system (5–8). To date, these studies have demonstrated that the action of CRF is not only region-specific, owing to circuit, cell-type, and pathway specificity, but that CRF's action is also dependent on the organism's previous exposures to stress (9–11).

To extend our knowledge of the extrahypothalamic CRF system to other brain regions, we investigated the role of CRF in the lateral habenula (LHb). The LHb is a highly stress-responsive brain region due to its ability to convey negative reward information to midbrain monoaminergic systems (12). Dysfunction of the LHb is implicated in stress-related disorders (13–16). This area is activated by reward omission and aversive stimuli but inactivated by unexpected reward. The LHb influences behavior through its inhibitory control of midbrain dopamine (DA) neurons. Indeed, optogenetic activation of LHb neurons projecting to rostromedial tegmental nucleus (RMTg) GABAergic neurons is sufficient to elicit behavioral avoidance (17). Other studies have demonstrated the role of fast synaptic transmission mediated by GABA and glutamate, as well as neuromodulatory effects of serotonin, norepinephrine and DA in the LHb (18); however, no study to date has investigated CRF neurotransmission in the LHb. The CRF receptors CRFR1 and CRFR2 (also known as CRHR1 and CRHR2) are G-protein coupled receptors (GPCRs). Activation of these receptors has a diverse range of neuronal effects depending on the brain region in which they are expressed. In general, activation of CRFR1 is anxiogenic whereas CRFR2 activation is anxiolytic (4). Although, the LHb shows immunoreactivity and mRNA expression for both CRF and CRFR1 (19–21), and receives inputs from the structures that comprise the extrahypothalamic CRF system, its involvement in this system is unexplored.

In this study, we investigated the role of CRF signaling on synaptic transmission and neuronal excitability in the rat LHb. We also examined the experience-specific role of CRF signaling in the LHb using an established model of severe early life stress, maternal deprivation (MD), that is associated with persistent hyperactivity of the HPA axis, CRF signaling and DA dysfunction (22–24). In fact, MD impairs GABAergic control of VTA DA neurons through epigenetic impairment of GABAergic plasticity which in part contributes to MD-induced dysregulation of DA neurotransmission (25). CRF can modulate DA signaling and alter mesolimbic reward function (26, 27). Therefore, an altered CRF signaling within the LHb as an upstream input to the VTA could play a crucial part in DA dysfunction associated with MD stress model. Here, we found that CRF-CRFR1-protein kinase A (PKA) signaling in the LHb increased intrinsic neuronal excitability through a G-protein dependent

activation of Ca^{2+} signaling and changes in afterhyperpolarizations (AHPs) mediated by the Ca^{2+} -activated K^{+} channels, BK and SK channels, in a PKA-dependent manner. Additionally, we found that changes in the abundance of SK2 channels and PKA signaling within the LHb mediated the increases in LHb intrinsic neuronal excitability by an acute episode of MD, which blunted the excitatory action of CRF within the LHb after MD. Although CRF suppressed GABAergic synaptic transmission onto LHb neurons through an eCB retrograde signaling, this effect remained intact after MD. Our study provides evidence for the existence of CRF-CRFR1-PKA signaling within the LHb, which may mediate part of stress-relevant effects of CRF through changes in synaptic transmission and intrinsic excitability of LHb neurons. The recruitment of the CRF stress systems within the LHb may play a key role in the development of stress-, addiction- and depressive-related behaviors.

Results

CRF increased the excitability of LHb neurons

Extrahypothalamic CRF mediates behavioral responses to stress. The region-specific actions of CRF have been defined in brain regions that are known to be involved in stress-related behaviors to date, no study has investigated the role of CRF signaling in the LHb. Therefore, we first tested the action of CRF on the excitability of LHb neurons. Neuronal excitability is influenced by both synaptic transmission and neuronal intrinsic properties. Action potentials (APs) generated by depolarizing steps were first measured in LHb neurons with intact fast synaptic transmission. We found that CRF significantly increased the firing of neurons. This may involve changes in fast synaptic transmission as well as intrinsic excitability by CRF (Figure 1, A and B). To isolate the action of CRF on the intrinsic excitability of the cell, we repeated the same experiment in the absence of fast synaptic transmission [by blocking GABA_A (gamma-aminobutyric acid) receptors (GABA_ARs), AMPA (α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) receptors (AMPA), and NMDA (N-methyl-D-aspartate) receptors (NMDARs) in the perfusate] and found that CRF still significantly increased neuronal excitability (Figure 1, C and D). The excitatory effect of CRF on neuronal excitability of LHb neurons was maximally achieved by 20–30 min of bath application, remained stable over time and persisted even after washout of CRF (figure S1). With fast synaptic transmission blocked, we then analyzed CRF's effects on different characteristics of APs and intrinsic neuronal excitability including fast AHPs (fAHPs), medium AHPs (mAHPs), threshold potential, half width and amplitude of AP and input resistance (R_{in}). Of these, only fAHP, mAHP, R_{in} and AP threshold measurements were significantly different after CRF. AHPs are mainly mediated by multiple Ca^{2+} -dependent K^{+} channels that tend to be inhibitory (reduce neuronal excitability) (28). Although fAHPs were increased after CRF application, CRF was associated with decreased mAHPs and increased R_{in} . CRF also lowered AP threshold (Figure 1, E and F; figure S1; and table S1). The caveat for the measurements of AHPs in response to sustained membrane depolarization is the inability to produce similar numbers of APs in different conditions because these various AHPs are to varying degrees dependent on intracellular calcium concentration.

CRF did not affect glutamatergic synaptic transmission but decreased GABAergic synaptic transmission through retrograde eCB signaling

Although CRF increased neuronal excitability in the absence of fast synaptic transmission, it was likely that CRF also altered synaptic transmission to promote LHb hyperexcitability. To test this, we examined the effects of CRF on the frequency and amplitude of AMPAR-mediated miniature excitatory postsynaptic currents (mEPSCs) and GABA_AR-mediated miniature inhibitory postsynaptic currents (mIPSCs) recorded from LHb neurons. CRF did not have a significant effect on either the frequency or amplitude of mEPSCs, suggesting that CRF did not affect the presynaptic release of glutamate, nor the function of postsynaptic AMPARs (Figure 2A). However, we found that mIPSCs were significantly reduced in the frequency and amplitude; whereas the mean amplitude of mIPSC was not significantly different after CRF, we detected a significant shift in the cumulative probability plot for amplitude by CRF (Figure 2B), suggesting that CRF reduced GABAergic inhibition in LHb neurons through significant decreases in the release of GABA from presynaptic terminals and perhaps reduction in postsynaptic GABA_AR number or function. The suppressing effect of CRF on GABAergic transmission persisted even after washout of CRF (figure S2). As CRF depressed GABAergic transmission without altering glutamatergic transmission in control rats that had been deprived of maternal care (non-MD rats), we next recorded evoked EPSCs and IPSCs in LHb neurons by stimulating afferents in the stria medullaris before and after application of CRF to quantify synaptic drive (Excitation and inhibition balance; EPSC/IPSC or E/I, ratio) onto LHb neurons and examine whether CRF shifts the excitatory-inhibitory balance toward excitation. However, E/I ratio measurements did not change after CRF application (figure S3A) suggesting that the balance of excitation to inhibition remains unchanged after CRF.

Given the well-known ability of eCB signaling to suppress GABA release from presynaptic GABAergic terminals (29), we investigated whether CRF induced an eCB-mediated GABAergic depression in the LHb. First, we blocked postsynaptic G-protein signaling in the neuron by intracellular administration of GDP β S, a non-hydrolyzable GDP analog, through the patch pipette and tested the effects of CRF on mIPSCs in the same neurons. Intra-pipette GDP β S completely prevented CRF-induced decreases in the amplitude and frequency of mIPSCs (Figure 3A). Consistently, we found that activation of CB1 cannabinoid receptors by WIN55,212-2, a CB1 receptor agonist, depressed GABAergic synaptic transmission (Figure 3B). AM251, a CB1 receptor antagonist, not only blocked the suppressing effect of CRF on presynaptic GABA release but also led to a left-ward shift in the cumulative probability curve of inter-event intervals indicative of an increase in the probability of presynaptic GABA release (Figure 3C). Notably, we also detected significant reductions in the mean amplitude of mIPSCs and a left-ward shift in the cumulative amplitude probability curve of mIPSC amplitude by WIN55,212-2 (Figure 3B), similar to the effect of CRF (Figure 2B). Although we assume that significant changes in the probability of GABA release by CRF and WIN55,212-2 affected not only the frequency but also the amplitude distribution of mIPSCs in these recordings (30), it is also possible that CRF affected postsynaptic GABA_A-receptor mediated inhibition independent of eCB-CB1 receptor signaling as the postsynaptic effect of CRF on mIPSC amplitude which was not blocked by AM251 (Figure 3C).

CRF activated CRFR1 and acted through postsynaptic Ca^{2+} signaling

Given that the LHb is involved in the processing of aversive stimuli and has also been shown to be hyperactive in depressed and stress-related behavioral states, we hypothesized that the ability of CRF to increase LHb neuronal excitability is achieved through binding to postsynaptic CRFR1 rather than CRFR2. In all the following experiments characterizing the mechanisms underlying CRF-induced increases in LHb intrinsic excitability, AP recordings were performed in the presence of synaptic blockers similar to Figure 1C (AMPA, NMDA, and GABA_A receptors were blocked). Indeed, in the presence of antalarmin, a selective CRFR1 antagonist, we found that CRF's excitatory action on intrinsic excitability was completely blocked (Figure 4A). On the other hand, CRF significantly increased the intrinsic excitability of LHb neurons in the presence of antisauvagine as a CRFR2-selective antagonist (Figure 4B). To further confirm CRF's action through postsynaptic CRFR1, a G-coupled protein receptor, we blocked postsynaptic G-protein signaling in the neuron by intracellular administration of GDPβS through the patch pipette. Intra-pipette GDPβS prevented CRF-induced increase in LHb intrinsic excitability (Figure 4C). Both intracellular release of Ca^{2+} from internal stores and extracellular Ca^{2+} influx have previously been shown to be involved in CRFR1 signaling in other brain areas (31, 32). To further investigate whether an increase in postsynaptic Ca^{2+} signaling is involved downstream of CRFR1 activation, BAPTA (a fast Ca^{2+} chelator) was used intracellularly through the patch pipette. As Ca^{2+} signaling is involved in activation of some of cationic currents including Na^+ channels in LHb neurons (33), not surprisingly we detected that inclusion of BAPTA in the patch pipette blunted neuronal firing prior to CRF treatment. Nevertheless, intra-pipette application of BAPTA blocked CRF's ability to increase intrinsic excitability of LHb neurons (Figure 4D). We next sought to determine how exactly Ca^{2+} signaling through CRFR1 activation could lead to increased intrinsic excitability of LHb neurons.

CRF affected SK and BK channel function in LHb neurons in a PKA-dependent manner

Changes in intrinsic conductances through K^+ channels, such as SK and BK channels, influence the firing of neurons (34). In our earlier experiments evaluating the effects of CRF on intrinsic excitability of LHb neurons, we observed that CRF altered AHPs coincident with higher input resistance and enhanced excitability while significantly lowered AP threshold in LHb neurons (Figure 1, E and F; and table S1). The fAHP amplitude was enhanced by CRF suggesting an increase in BK channel function by CRF. However, the mAHPs measured in response to depolarization was decreased after bath application of CRF, suggesting a decreased function of SK channels. We also detected an increase in R_{in} by CRF which could support CRF-induced increase in LHb intrinsic excitability. First, we tested whether a SK channel blocker, apamin, could mimic and occlude CRF-induced increase in LHb intrinsic excitability through decrease in mAHP. We found that apamin significantly increased LHb intrinsic excitability and furthermore occluded CRF's action on LHb neuronal excitability (Figure 5A). The effect of apamin on intrinsic excitability was also accompanied by an increase in the amplitude of fAHP, a decrease in the amplitude of mAHP (an apamin-sensitive component of mAHP) with a parallel increase in R_{in} and a lower AP threshold (Figure 5A; table S2). To further confirm that increasing SK channel function could prevent CRF-induced increase in intrinsic excitability, we used a positive modulator of SK channels (Cyclohexyl-[2-(3,5-dimethyl-pyrazol-1-yl)-6-methyl-pyrimidin-4-yl]-amine,

CYPPA). CYPPA did not affect intrinsic neuronal excitability by itself, but blocked the excitatory effect of CRF on LHb intrinsic excitability (Figure 5B). CYPPA did not significantly affect fAHP, mAHP, Rin or AP threshold measurements (Figure 5B, table S3). Although a decrease in BK-mediated fAHP seems to mediate an increase in neuronal excitability, an increase in BK channel activity can also facilitate high frequency firing of neurons through inactivation of other K^+ channels (35). So we tested the possibility that CRF may first involve BK channels which could subsequently inactivate SK channels to promote neuronal intrinsic excitability. BK channel blocker, iberiotoxin, slightly increased LHb intrinsic excitability but significantly attenuated CRF's excitatory effects while blocking both CRF-induced changes in AHPs and Rin (Fig. 5C). No significant changes were detected in AHPs and Rin after CRF application when iberiotoxin was present in the bath application (Figure 5C). Activation of CRF-CRFR1 signaling initiates PKA activity so we next examined whether activation of PKA in LHb neurons affected BK and SK channels to increase LHb neuronal excitability. PKA-mediated phosphorylation of the SK2 channel subunit (which contributes to the apamin-sensitive component of mAHP) has been shown to dramatically decrease surface localization of SK2 channels thereby promoting neuronal excitability and the induction of synaptic plasticity at Schaffer collateral synapses and hippocampal-dependent learning (36–38). On the other hand, BK channel activity can be enhanced through PKA-mediated phosphorylation of BK α subunit (the pore forming subunit of BK channels) (39). Intracellular inclusion of PKI(6–22), a membrane impermeant PKA inhibitor, significantly blocked CRF-induced enhancement of intrinsic excitability and CRF-induced changes in AHPs and Rin (Figure 5D; table S4). Although a slight reduction in LHb intrinsic excitability was observed within 15–30 min after initiation of the whole cell recording with PKI(6–22)-filled pipettes, this change in intrinsic excitability was not significantly different from the excitability of LHb neurons recorded within the same time frame from non-MD rats with patch pipettes filled with normal drug-free internal solutions (figure S4, and table S4).

Early-life stress increased the spontaneous firing and excitability of LHb neurons by decreasing the function and abundance of SK channels

Given that early life stresses including maternal deprivation (MD) induce long-lasting alterations in CRF-CRFR1 signaling in the brain (24), it was feasible to assume that an adverse early-life experience like MD would affect the function of LHb neurons and alter their responsiveness to acute CRF. In fact, MD modifies DA signaling from the VTA through epigenetic modifications of critical signaling pathways involved in trafficking of GABA_A receptors onto VTA DA neurons (25). Considering the potent inhibitory control of VTA DA neurons by the LHb, we chose to use the same model of severe early life stress in this work to determine whether the response of LHb neurons to extrahypothalamic CRF is affected by exposure to early life stress. The protocol used was of an acute MD, in which rats were isolated at P9 for 24 hours, then returned to the litter. First, we hypothesized that this period of MD would increase LHb neuronal excitability. To test this hypothesis, we recorded LHb neuronal spontaneous activity using cell-attached recordings in voltage-clamp mode to determine the percentage of neurons which were spontaneously active and of those, the average spike frequency. In this set of experiments, synaptic transmission was intact. We observed that the portion of spontaneously active LHb neurons was significantly larger in

maternally deprived rats (MD rats) compared to control non-MD rats. Moreover, LHb neurons displaying spontaneous activity had remarkably higher spike frequencies in MD rats compared to controls (Figure 6A).

In addition to cell-attached recordings of basal neuronal firing, we also tested whether LHb neurons from MD rats exhibit enhanced firing in response to depolarization compared to control non-MD rats in the absence or presence of fast synaptic transmission. Although we did not detect a significant difference in LHb neuronal excitability between non-MD and MD rats in intact synaptic transmission in response to 10–100pA depolarizing current steps, MD significantly increased LHb intrinsic excitability in response to depolarization in this range of current injections when fast synaptic transmission was blocked (Figure 6, B and C). Furthermore, the MD-induced increase in intrinsic excitability was associated with a significant reduction in mAHP amplitude and a significantly higher input resistance without any change in the amplitude of fAHP, suggesting that this increase may be due to a decreased function and/or abundance of SK channels (Figure 6D). Except mAHP and Rin, all the other measured passive and active properties of LHb neurons including RMP were unaffected following MD (table S5). Consistently, Western blot analysis of LHb tissues from non-MD and MD rats demonstrated that SK2 but not BK channel α subunit protein abundance was significantly decreased in MD rats compared to controls (Figure 6E; and figure S6). In contrast, we did not observe any significant changes in protein abundance of SK1 and SK3 (figure S5).

MD potentiated both glutamatergic and GABAergic synaptic transmission while increased E/I ratio

We found that MD-induced changes in glutamatergic synapses supported increases in LHb neuronal excitability, given that LHb neurons from MD rats exhibited significantly higher AMPA/NMDA ratios compared to control non-MD rats (figure S7A). In support of this postsynaptic glutamatergic potentiation, there was a significant increase in the cumulative probability of AMPAR-mediated mEPSC amplitudes after MD, although the average amplitudes were not significantly different between MD and non-MD rats. Moreover, we detected a marked increase in the average and cumulative probability of frequencies of mEPSCs suggesting an increase in presynaptic glutamate release by MD (Figure 7A). We also recorded evoked AMPAR-mediated EPSCs at different holding potentials to measure AMPAR rectification and determine whether MD induced any changes in synaptic AMPAR subunit composition. AMPA EPSC rectification and EPSC IV plots were unchanged after MD (figure S7B).

Although the average amplitudes of GABA_A receptor-mediated mIPSCs were not significantly different between MD and non-MD rats, we found that the cumulative probability of mIPSC amplitudes was significantly increased after MD. Moreover, the average and cumulative probability of frequency of mIPSCs were significantly increased in MD compared to non-MD rats. This suggests that MD potentiated GABAergic synaptic transmission pre- and postsynaptically (Figure 7B). While both excitatory and inhibitory synapses onto LHb seemed to be potentiated by MD, we detected an increase in E/I ratio

following MD suggesting a shift in the balance between excitation and inhibition in synaptic inputs towards excitation following MD (Figure 7C).

The effects of CRF and apamin on LHb intrinsic excitability were blunted following MD due to increased PKA signaling

Depending of the timing and severity of the stress, CRF release can have long lasting regional effects and change subsequent responses to CRF exposure. Here, we tested whether the responsiveness of LHb neurons to acute exposure to CRF was altered by MD, with the hypothesis being that CRF is excessively released during the episode of MD, leading to dysregulation of central CRF-CRFR1-PKA signaling within the LHb. Although CRF significantly increased neuronal excitability in non-MD neurons in both the presence and absence of synaptic blockade (Figure 1), CRF's action on neuronal excitability was blunted in MD rats both in the presence and absence of fast synaptic transmission, given that there was no significant increase in LHb neuronal excitability in MD rats after adding CRF (Figure 8, A and B). Except for a significant increase in the amplitude of fAHP accompanied by a lower AP threshold after CRF, no significant changes were detected in mAHP and Rin by CRF in MD rats (table S1). Assuming sustained CRF signaling dysregulation by MD, we next examined the effects of MD on the mRNA abundance of *Crh* and *Crhr1* in LHb tissues. Relative to non-MD rats, no significant changes were detected in *Crh* or *Crhr1* mRNA expression in MD rats (Figure 8B).

Because we had established that CRF signaling affected the activity of postsynaptic SK channels and that the protein abundance of SK2 channels was diminished by MD, we sought to determine whether an impaired function in SK channels in MD rats underlies the blunted effect of CRF on LHb intrinsic excitability (in conditions in which synaptic transmission was blocked). The first piece of evidence to support this hypothesis was that bath application of apamin had a diminished effect on the intrinsic excitability of MD LHb neurons (Figure 8C). Except for a significant increase in fAHP by apamin, mAHP and Rin were unaltered after exposure in MD slices (table S2). We also tested the effects of another positive SK channel modulator, 1-EBIO, which is similar to CYPPA. Bath application of 1-EBIO significantly decreased intrinsic neuronal excitability of LHb neurons in slices from MD rats despite of the lower abundance of SK2 channel protein in MD rats (Figure 8D). This decrease in intrinsic excitability was associated with a significant increase in mAHP amplitude and a lower Rin whereas fAHP was unaffected (table S3). PKA-dependent regulation of SK channel activity and trafficking of SK2 plays an important role in neuronal excitability, and we had also observed a complete inhibition of CRF-induced increases in LHb intrinsic excitability by postsynaptic inhibition of PKA through modulation of AHPs (Figure 5D). Although mRNA expression of *Crhr1* was not significantly decreased by MD, we hypothesized that PKA abundance and/or function may have been persistently altered as a biochemical neuroadaptation in response to CRF-CRFR1 activation during MD contributing to increased LHb neuronal excitability. First we tested whether postsynaptic manipulation of PKA activity in LHb neurons could diminish LHb intrinsic excitability in MD rats through changes in mAHP. We detected a significant decrease in the number of APs triggered in response to depolarization within 15 min after initiation of the whole cell recording with PKI(6–22)-filled pipettes in LHb neurons recorded from MD rats which was

maximized and then stabilized at 30min from the start of our recordings. This significant decline in intrinsic excitability was not evident within the same time-frame of our recordings in LHb neurons with normal internal solution-filled pipettes from MD rats (Figure 8E) and in PKI(6–22) experiments performed in non-MD rats (figure S4). Although postsynaptic inhibition of PKA did not affect fAHPs, the decreased LHb intrinsic excitability of MD rats by PKA inhibition was associated with a significant increase in the amplitude of mAHP parallel with a lower Rin and AP threshold (table S4). Western blot analysis of LHb tissues from non-MD and MD rats revealed no significant difference in the abundance of PKA protein between non-MD and MD rats (Figure 8F).

We also tested the effects of acute CRF on mEPSCs and mIPSCs recorded from LHb neurons in slices from MD rats. Except for a significant leftward shift in the cumulative probability curve of mEPSC amplitude, CRF did not affect mEPSCs in MD rats (figure S8A), similar to our observations in non-MD rats. Similarly, acute bath application of CRF significantly reduced GABAergic synaptic transmission onto LHb neurons both pre- and post-synaptically in MD rats suggesting that the acute effects of CRF on synaptic transmission remained intact following MD (figure S8B). Consistently, E/I ratio measurements also did not change after CRF application in MD rats (figure S3B). Altogether, our data suggest that LHb intrinsic neuronal responsiveness to acute CRF was altered following MD due to changes in SK channel activity and abundance in addition to a persistent increase in PKA activity downstream to CRF-CRFR1 signaling.

Discussion

This study establishes a role for an extrahypothalamic CRF-CRFR1-PKA signaling in the LHb. We demonstrate that the stress neuropeptide CRF increased the excitability of LHb neurons and depressed GABAergic transmission onto LHb neurons through eCB signaling. We found that blockade of CRFR1 and postsynaptic Ca^{2+} or disruption of postsynaptic G-protein signaling prevented the excitatory effect of CRF. *Crhr1* and *Crhr2* gene expression often overlaps and *Crhr1* and *Crhr2* mRNAs have also been detected in the rodent habenula (21, 40). Our findings suggest that postsynaptic G-protein-mediated CRFR1 signaling requires intracellular Ca^{2+} signaling to increase LHb neuronal excitability. The increase in neuronal excitability induced by CRF was associated with a significant reduction in the amplitude of the mAHP with a parallel increase in input resistance (suggesting possible decreases in K^+ conductances underlying membrane resistance including leak K^+ currents in addition to SK channels by CRF). Consistently, we found that blocking SK channels mimicked and occluded CRF's effect while positive enhancement of SK channel function blocked CRF's effect. CRF increases neuronal excitability in several brain regions (41–43). Decreases in the amplitude of AHPs and K^+ channel function, including SK type, are associated with an increase in neuronal excitability (44, 45). On the other hand, CRF receptor signaling in VTA DA neurons potentiates the function of SK channels (31).

We also found that CRF increased the amplitude of fAHPs (mostly mediated by BK channels) and blocking BK channels prevented the effects of CRF on excitability, AHPs, and Rin. fAHPs may facilitate high-frequency firing of LHb neurons similar to that shown in CA1 hippocampal pyramidal neurons by limiting activation of other K^+ channels and Na^+

channel inactivation (35, 46–49). In fact, in addition to an increase in the amplitude of fAHP, CRF significantly lowered AP threshold suggesting that a hyperpolarized inter-spike membrane potential during high AP firing rates might be maintained by fAHPs to prevent voltage-gated Na⁺ channel inactivation and promote suppression of SK channel function by CRF. Activated CRFR1 primarily signals by Gs protein coupling, resulting in the induction of PKA activity and increase in hippocampal neuronal excitability (50) although CRFR1 can also affect protein kinase C (PKC) activity through Gq-coupling (51). We found that postsynaptic inhibition of PKA in LHb neurons significantly blocked CRF's action on LHb intrinsic excitability and AHPs suggesting that PKA-dependent modulation of AHPs through opposing actions on BK and SK channel activity underlie CRF's excitatory effects. It has been shown that PKA-mediated phosphorylation of the SK2 channel subunit significantly decreases membrane SK2 channels and SK channel function, whereas BK channel activity is augmented by PKA (36–39).

Other studies have shown that the action of CRF is dependent on an organism's previous exposure to stress (9–11). CRF modulates DA signaling and mesolimbic reward function (26, 27); thus, an alteration of CRF signaling within the LHb and subsequent changes in LHb function could significantly affect DA release from the VTA. Here we found that MD increased both basal and depolarization-induced AP firing of LHb neurons. In addition, the increase in LHb intrinsic excitability following MD was accompanied by a decrease in mAHP. Indeed, protein abundance of SK2 was lower in MD rats. Positive modulation of SK channels did not reduce firing rate of LHb neurons in non-MD rats suggesting that the basal activity of these channels may be relatively high. On the other hand, positive enhancement of SK activity normalized intrinsic excitability in LHb neurons of MD rats. A loss of apamin-sensitive SK channels (SK2 channels contribute to apamin sensitive portion of mAHP) could be related to increased PKA signaling in MD rats. MD was not associated with a significant increase in *Crh* mRNA expression or a decrease in *Crhr1* mRNA expression to suggest hyperactivation of CRF-CRFR1 signaling pathway in the LHb after MD. However, we found that the transduction signaling involving PKA is increased by MD, suggesting that this is an adaptation that persisted into adolescence. PKA inhibition was able to markedly attenuate the excitability of LHb neurons accompanied by a significant increase in mAHPs in MD rats with no significant alterations in BK channel abundance or activity. Since Western blot analysis of LHb tissues from non-MD and MD rats revealed no significant difference in the abundance of PKA protein (Figure 8F), this suggests a possible sustained rebound increase in PKA activity and signaling in LHb neurons. Therefore, the blunting of CRF's excitatory effects in the LHb of MD rats is partly due to MD-induced reduction in abundance of SK2 and also PKA-mediated changes in SK channel function and possibly trafficking.

CRF pre- or post-synaptically potentiates GABAergic transmission in some brain areas, such as dorsal raphe, central amygdala, and BNST (52–54), while we found that CRF presynaptically depressed GABAergic transmission onto LHb neurons in a CB1 receptor-dependent manner. Inhibition of CB1 receptors not only prevented the suppressing effect of CRF on presynaptic GABA release, but also led to an increase in the probability of presynaptic GABA release. These results point to the possible presence of a tonic eCB signaling and persistent CB1 receptor activation in the regulation of GABAergic

transmission in the LHb. Glutamatergic synaptic transmission, however, was unaffected by CRF. This indicates that CRF could increase LHb neuronal excitability not only through changes in membrane properties, but also by depressing GABAergic transmission. Similar to non-MD rats, CRF did not affect the synaptic excitatory to inhibitory balance of LHb neurons.

Both AMPA/NMDA ratios and the amplitude of mEPSCs were larger in LHb neurons from MD rats, suggesting an induction of a postsynaptic glutamatergic potentiation by MD. This glutamatergic potentiation was not associated with a change in AMPAR subunit composition. We assume that reduction in SK2 protein abundance by MD likely facilitated potentiation of glutamatergic synapses in LHb neurons. In fact, hippocampal CA1 long-term potentiation (LTP) was accompanied by a PKA-dependent endocytosis of SK2 channels (38). Similarly, potentiation of glutamatergic synapses and reduction of K⁺ conductance underlie LHb hyperexcitability after repeated injections of cocaine (55). MD also increased presynaptic glutamate release in LHb neurons similar to what was found in LHb neurons projecting to the VTA in learned helplessness animal models of depression (16). Although glutamatergic synaptic modifications induced by MD could support LHb hyperexcitability, MD was also associated with significant increases in GABAergic synaptic transmission both pre- and post-synaptically. Alteration in GABAergic synaptic strength could have been triggered in response to long-term changes in neuronal excitability induced by MD to stabilize neuronal activity (homeostatic synaptic scaling) (56). However, it is also possible that changes in LHb intrinsic excitability is a neuroadaptation triggered by MD-induced changes in GABAergic synaptic inputs to the LHb. Main GABAergic structures projecting to the LHb include the entopeduncular nucleus and the VTA, which are shown to co-release GABA and glutamate onto LHb neurons (15, 57). Therefore, an increase in glutamate and GABA release from such inputs onto LHb neurons following MD may elicit a homeostatic increase in LHb neuronal intrinsic excitability observed here. Yet, we found an imbalance of excitation to inhibition of synaptic inputs onto LHb neurons favoring excitation. The acute suppressing effects of CRF on GABAergic transmission remained intact in MD animals with no subsequent alteration in E/I balance in response to acute CRF. Although MD rats still showed higher E/I ratios compared to non-MD rats (Supplementary figure 3) promoting LHb excitability.

The role of CRF-CRFR1-PKA regulation of SK channels in MD rats is an important question that merits investigation given increasing interest in the role of the extrahypothalamic CRF system in health and disease. Recent advances in optogenetic interrogation of neuronal circuitries and the availability of promoter driven Cre mouse and rat lines will help us to elucidate the specific contributions of synaptic inputs to LHb dysfunction after MD and further delineate mechanisms underlying MD-induced dysregulation of CRF signaling in the LHb.

Materials and Methods

All experiments were carried out in accordance with the National Institutes of Health guidelines for the care and use of laboratory animals and were approved by the Uniformed

Services University Institutional Animal Care and Use Committee. All efforts were made to minimize animal suffering and reduce the number of animals used.

Maternal deprivation procedure

Half of the male pups of litter of Sprague–Dawley rats at P9 were isolated from the dam and their siblings at 10:00 a.m. for 24 hours (MD group). The isolated rats were placed in a separate quiet room and kept on a heating pad (34° C) and not disturbed until being returned to their home cage. The remaining non-separated male rat pups received the same amount of handling and served as the non-maternally deprived control group (non-MD group). Then starting at P21, each day through P28, two MD and non-MD (one from each condition) rats (age-matched) of the same litter were sacrificed to obtain LHb tissue and neurons for electrophysiology and Western Blotting analyses. With the exception of Figures 6 A–D and 7C, where two or three neurons were recorded per animal (n represents number of recorded cells per rat), one cell per animal was recorded to demonstrate that the MD effects observed were consistent across multiple animals and across the P21–28 time frame; therefore, all reported n values in those graphs represent the number of animals.

Slice preparation for electrophysiology

Rats were anesthetized with isoflurane and immediately decapitated. The brains were quickly dissected and placed into ice-cold artificial cerebrospinal fluid (ACSF) containing (in mM): 126 NaCl, 21.4 NaHCO₃, 2.5 KCl, 1.2 NaH₂PO₄, 2.4 CaCl₂, 1.00 MgSO₄, 11.1 glucose, 0.4 ascorbic acid, saturated with 95% O₂–5% CO₂. Sagittal slices containing the LHb were cut at 250 µm and incubated in ACSF at 34°C for at least 1 hour. Slices were then transferred to a recording chamber and perfused with ascorbic acid-free ACSF at 28°C.

Electrophysiology

Whole cell and cell-attached recordings were performed on LHb slices using a patch amplifier (Multiclamp 700B) under infrared-differential interference contrast microscopy. Data acquisition and analysis were carried out using DigiData 1440A, pClamp 10 (molecular devices, Union City, CA), Clampfit and Mini Analysis 6.0.3 (Synaptosoft Inc.). Signals were filtered at 3 kHz and digitized at 10 kHz. The recording ACSF was the same as the cutting solution except that it was ascorbic acid-free. Spontaneous activity was monitored using cell-attached voltage-clamp recordings of action potentials (APs) at $I = 0$ pA for 5 minutes. Number of APs was counted over 5 min and spike frequency was calculated. Neuronal excitability recordings in response to depolarization were performed in whole-cell current-clamp mode. The patch pipettes (3–6 MΩ) were filled with 130 mM K-gluconate, 15 mM KCl, 4 mM ATP-Na⁺, 0.3 mM GTP-Na⁺, 1 mM EGTA, and 5 mM HEPES (pH adjusted to 7.28 with KOH, osmolarity adjusted to 275–280 mOsm). LHb neurons were given increasingly depolarizing current steps at +10pA intervals ranging from +10pA to +100pA, allowing us to measure AP generation in response to membrane depolarization (5 sec duration). Current injections were separated by a 20s interstimulus interval and neurons were kept at –65 mV with manual direct current injection between pulses. Synaptic transmission blockade was achieved by adding 6,7-dinitroquinoxaline-2,3-dione (DNQX; 10 µM), picrotoxin (100µM) and APV (50 µM) to block AMPA, GABA_A and NMDA receptor-mediated synaptic transmission, respectively. The number of APs

induced by depolarization at each intensity was counted and averaged for each experimental group.

Because AHP amplitude increases with increasing number of APs, we measured AP threshold, mAHP and fAHP amplitudes at the current step that was sufficient to generate the first AP/s (58). AP threshold was measured at the beginning of the upward rise of the AP. mAHP was measured as the difference between AP threshold and the peak negative membrane potential at the end of the current step. fAHPs were calculated as the difference between AP threshold and the peak negative potential following the AP (Figure 1E). Resting membrane potential (RMP) was assessed at the beginning of the recording by quickly switching to $I=0$. Input resistance (R_{in}) was determined by injecting a small (50 pA) hyperpolarizing current pulse (5s) and calculated by dividing the steady-state voltage response by the current pulse amplitude. Reported AP half width, AP peak amplitude were obtained from measurements of AP characteristics in Clampfit.

Whole-cell recordings of GABA_AR-mediated mIPSCs were isolated in ACSF perfused with DNQX; 10 μ M, strychnine (1 μ M) and tetrodotoxin (TTX, 1 μ M). The patch pipettes (3–6 M Ω) were filled with 125 mM KCl, 2.8 mM NaCl, 2 mM MgCl₂, 2 mM ATP-Na⁺, 0.3 mM GTP-Na⁺, 0.6 mM EGTA, and 10 mM HEPES (pH adjusted to 7.28 with KOH, osmolarity adjusted to 275–280 mOsm). AMPAR-mediated mEPSCs were isolated in ACSF perfused with picrotoxin (100 μ M), D-APV (50 μ M) and TTX (1 μ M). Patch pipettes for mEPSC recordings were filled with 117 mM Cs-gluconate, 2.8 mM NaCl, 5 mM MgCl₂, 2 mM ATP-Na⁺, 0.3 mM GTP-Na⁺, 0.6 mM EGTA, and 20 mM HEPES (pH adjusted to 7.28 with CsOH, osmolarity adjusted to 275–280 mOsm). For both mIPSCs and mEPSCs, LHb neurons were voltage-clamped at -70 mV and recorded over 10 sweeps, each lasting 50 seconds.

EPSCs were evoked with a stimulating electrode placed in the stria medullaris. Evoked EPSCs were recorded in ACSF perfusion containing picrotoxin (100 μ M) while the cell was voltage-clamped at $+40$ mV. Internal solution for patch pipettes was similar to that used for mEPSC recordings (Cs-gluconate-based) but also included intracellular spermine (10 μ M). AMPAR-mediated currents were isolated with D-APV (50 μ M), a selective NMDA receptor antagonist. Isolated AMPAR-mediated currents were then subtracted from the combined EPSC to provide the NMDA receptor-mediated current and thus, the AMPA/NMDA ratio. AMPAR EPSCs were also recorded at holding potentials ranging from -65 to $+40$ mV in the presence of picrotoxin (100 μ M), APV (50 μ M) and intracellular spermine (10 μ M) included in Cs-gluconate-based internal. Normalized current-voltage ($I-V$) curves were then generated by dividing the AMPAR EPSC peak amplitudes by the mean of AMPAR EPSC peak amplitude recorded at -65 mV. AMPAR EPSC rectification was determined by dividing peak AMPAR EPSCs amplitudes recorded at -65 mV by those recorded at $+40$ mV.

The excitatory and inhibitory balance (EPSC/IPSC, E/I ratio,) was recorded with a Cs-gluconate-based internal solution similar to mEPSC recordings. Evoked EPSCs and IPSCs from LHb neurons were recorded in the same neuron in drug-free ACSF using a stimulating electrode placed in the stria medullaris. EPSCs were recorded at the reversal potential for GABA_A IPSCs (-55 mV), and IPSCs were recorded at the reversal potential for EPSCs ($+10$

mV) in the same LHB neuron. The E/I ratio was then calculated as EPSC/IPSC amplitude ratio by dividing the average peak amplitude of 10 consecutive sweeps of EPSCs or IPSCs from the same recording. The cell input resistance and series resistance were monitored through all the experiments and if these values changed by more than 10%, data were not included.

Antalarmin, CYPPA, 1-EBIO, WIN 55,212-2 and AM251 were prepared as stock solution in DMSO and were diluted (1:10000) to final concentration in ACSF of 1 μ M, 10 μ M, 300mM, 2 μ M, and 10 μ M, respectively. Stock solutions for apamin, rat CRF, iberiotoxin, and Antisauvagine-30 were prepared in distilled water and diluted (1:1000) to final concentration in ACSF of 100nM, 250nM, 100nM, and 25nM, respectively. The intra-pipette concentrations of BAPTA, GDP β s and PKI(6–22) were 30mM, 300 μ M, and 10 μ M, respectively. In some of our control interleaved experiments, AP recordings were continued for an hour without addition of drugs and no significant changes were observed over time. For all drug experiments, a baseline depolarization-induced AP recording/ mIPSC/mEPSC was recorded in each neuron, and then the appropriate drug was added to the slice by the perfusate and AP generation in response to depolarizing current steps was again tested 25–30 min after. For apamin and CYPPA experiments, a second baseline with apamin or CYPPA was obtained before the addition of CRF and AP generation was re-evaluated 25–30min after CRF application. CRF wash-out experiments in Supplementary figures were performed following 15min of bath application of CRF and the recordings continued at least for 90 min after the initiation of wash out of CRF.

Western Blotting

The LHB was dissected bilaterally from sagittal slices (300 μ M) of non-MD or MD rats in ACSF and then snap frozen in liquid nitrogen and stored at -80°C . Tissues were thawed, washed in ice-cold PBS and lysed in RIPA buffer containing protease inhibitors (Sigma). Samples were then sonicated, incubated on ice for 30 minutes and centrifuged at 10,000g for 20 min at 4°C . Protein concentration in the supernatant was determined by Pierce BCA Protein Assay Kit (Life Technologies). Equal amounts of protein (20 μ g) were combined with loading buffer, boiled for 5 min, and loaded onto 4–20% precast polyacrylamide gel (Bio-Rad Laboratories). Separated proteins were transferred onto nitrocellulose membranes, blocked with casein-based blocking reagent (I-Block, Life Technologies) for 60 minutes at room temperature and then incubated overnight at 4°C with rabbit antibodies recognizing SKs (Antibody against KCa2.1 for SK1, Antibody against KCa2.2 for SK2, and Antibody against KCa2.3 (C-term) for SK3, 1:300, Alomone Labs), antibody against PKA regulatory β 2 subunit (1:5000, Abcam ab75993), antibody against BK *slo-1* (α 1 subunit of BK) (1:1000, Millipore MADN70) and GAPDH (1:1000, Cell signaling 2118) or antibody against actin (1:10,000, Abcam, ab6276). Secondary antibodies used were HRP-linked specific for rabbit (1:2000, Cell signaling) and mouse (1:2000, Cell signaling) IgG. After incubation, the membranes were washed with PBS-T and exposed to the appropriate horseradish peroxidase-linked secondary antibody (Cell Signaling). Blots were developed with Clarity Western ECL Substrate (Bio-Rad Laboratories) and detected using a BioRad ChemiDoc Touch image acquisition system (BioRad Laboratories, Hercules, CA, USA). Data was analyzed using ImageJ software. Total abundance of target protein was normalized

to appropriate endogenous control. All data were normalized to non-MD group with summary data reported as fold change.

Quantitative RT-PCR

Brain tissue from the LHb was collected by dissection from both non-MD and MD rats. Total RNA isolation was achieved through Trizol (Life Technologies)/chloroform-based extraction. Samples were homogenized and then quantified using the Nanodrop spectrophotometer at 260 and 280 nm optical densities. Reverse Transcriptase amplification of complementary DNA (cDNA) from total RNA was conducted with a High-Capacity RNA-to-cDNA Kit (Applied Biosystems) and BioRad CFX96 Real-Time System (Hercules, CA). TaqMan gene expression assays (Thermo Scientific) with probes for corticotropin releasing hormone (*Crh*) (catalog no. Rn01462137_m1), corticotropin releasing hormone receptor 1 (*Crhr1*) (catalog no. Rn00578611_m1) and actin (*Actb*) (catalog no. Rn00667869_m1) as the endogenous control to quantify these target genes during amplification. Each sample was run in triplicate. The comparative Ct (C_T) was calculated by subtracting C_T averages of the target gene from the respective endogenous control. For each sample calculated, the changes in target gene expression are given as 2^{-C_T} . All data were normalized to the non-MD group and presented as fold change.

Data analysis

Values are presented as means $\pm$ SEM. Statistical significance was determined using unpaired or paired two tailed Student's t-test, Mann-Whitney test, two way ANOVA or repeated-measures ANOVA (RM-ANOVA) with Bonferroni post hoc analysis. The threshold for significance was set at $*P < 0.05$ for all analyses. The peak values of the evoked paired EPSCs were measured relative to the same baseline. A stable baseline value was considered in each sweep of paired pulses starting at 20–50ms right before the emergence of the EPSC current using p-Clamp 10 software. Mini Analysis software was used to detect and measure mIPSCs and mEPSCs using preset detection parameters of mIPSCs and mEPSCs with an amplitude cutoff of 5 pA. The Kolmogorov–Smirnov test (KS test) was performed for the statistical analyses of cumulative probability plots of mEPSCs and mIPSCs. All statistical analyses were performed using IBM SPSS Statistics 24, GraphPad Prism 7 or Origin 2016.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The opinions and assertions contained herein are the private opinions of the authors and are not to be construed as official or reflecting the views of the Uniformed Services University of the Health Sciences or the Department of Defense or the Government of the United States. We are grateful to Drs. David Lovinger, Veronica Alvarez, Brian Cox and Mr. Shawn Gouty for their critical discussions for the present work. **Funding:** This work was supported by the National Institutes of Health (NIH)–National Institute of Drugs of Abuse (NIDA) Grant#R01 DA039533, Brain & behavior research foundation (formerly known as NARSAD) grant and Department of Defense intramural grant from the Uniformed Services University to FSN. The funding agencies did not contribute to writing this article or deciding to submit it.

References

1. Valentino RJ, Lucki I, Van Bockstaele E. Corticotropin-releasing factor in the dorsal raphe nucleus: Linking stress coping and addiction. *Brain research*. 2010; 1314:29–37. [PubMed: 19800322]
2. Koob GF. The role of CRF and CRF-related peptides in the dark side of addiction. *Brain research*. 2010; 1314:3–14. [PubMed: 19912996]
3. Silberman Y, Winder DG. Emerging role for corticotropin releasing factor signaling in the bed nucleus of the stria terminalis at the intersection of stress and reward. *Front Psychiatry*. 2013; 4:42. [PubMed: 23755023]
4. Sanders J, Nemeroff C. The CRF System as a Therapeutic Target for Neuropsychiatric Disorders. *Trends Pharmacol Sci*. 2016; 37:1045–1054. [PubMed: 27717506]
5. Silberman Y, Matthews RT, Winder DG. A corticotropin releasing factor pathway for ethanol regulation of the ventral tegmental area in the bed nucleus of the stria terminalis. *J Neurosci*. 2013; 33:950–960. [PubMed: 23325234]
6. Grieder TE, et al. VTA CRF neurons mediate the aversive effects of nicotine withdrawal and promote intake escalation. *Nat Neurosci*. 2014; 12:1751–8.
7. Ungless MA, et al. Corticotropin-releasing factor requires CRF binding protein to potentiate NMDA receptors via CRF receptor 2 in dopamine neurons. *Neuron*. 2003; 39:401–407. [PubMed: 12895416]
8. Partridge JG, et al. Stress increases GABAergic neurotransmission in CRF neurons of the central amygdala and bed nucleus stria terminalis. *Neuropharmacology*. 2016; 107:239–250. [PubMed: 27016019]
9. Henry B, Vale W, Markou A. The effect of lateral septum corticotropin-releasing factor receptor 2 activation on anxiety is modulated by stress. *J Neurosci*. 2006; 26:9142–9152. [PubMed: 16957071]
10. Lemos JC, et al. Severe stress switches CRF action in the nucleus accumbens from appetitive to aversive. *Nature*. 2012; 490:402–406. [PubMed: 22992525]
11. Waselus M, Nazzaro C, Valentino RJ, Van Bockstaele EJ. Stress-induced redistribution of corticotropin-releasing factor receptor subtypes in the dorsal raphe nucleus. *Biol Psychiatry*. 2009; 66:76–83. [PubMed: 19362706]
12. Hikosaka O. The habenula: from stress evasion to value-based decision-making. *Nat Rev Neurosci*. 2010; 11:503–513. [PubMed: 20559337]
13. Maroteaux M, Mameli M. Cocaine evokes projection-specific synaptic plasticity of lateral habenula neurons. *J Neurosci*. 2012; 32:12641–12646. [PubMed: 22956853]
14. Neumann PA, et al. Increased excitability of lateral habenula neurons in adolescent rats following cocaine self-administration. *Int J Neuropsychopharmacol*. 2015; 18
15. Shabel SJ, Proulx CD, Piriz J, Malinow R. Mood regulation. GABA/glutamate co-release controls habenula output and is modified by antidepressant treatment. *Science (New York, N.Y.)*. 2014; 345:1494–1498.
16. Li B, et al. Synaptic potentiation onto habenula neurons in the learned helplessness model of depression. *Nature*. 2011; 470:535–539. [PubMed: 21350486]
17. Stamatakis AM, Stuber GD. Activation of lateral habenula inputs to the ventral midbrain promotes behavioral avoidance. *Nat Neurosci*. 2012; 15:1105–7. [PubMed: 22729176]
18. Root DH, et al. Norepinephrine activates dopamine D4 receptors in the rat lateral habenula. *J Neurosci*. 2015; 35:3460–3469. [PubMed: 25716845]
19. Chappell PB, et al. Alterations in corticotropin-releasing factor-like immunoreactivity in discrete rat brain regions after acute and chronic stress. *J Neurosci*. 1986; 6:2908–2914. [PubMed: 3020187]
20. Chen Y, Brunson KL, Muller MB, Cariaga W, Baram TZ. Immunocytochemical distribution of corticotropin-releasing hormone receptor type-1 (CRF(1))-like immunoreactivity in the mouse brain: light microscopy analysis using an antibody directed against the C-terminus. *J Comp Neurol*. 2000; 420:305–323. [PubMed: 10754504]
21. Justice NJ, Yuan ZF, Sawchenko PE, Vale W. Type 1 corticotropin-releasing factor receptor expression reported in BAC transgenic mice: implications for reconciling ligand-receptor

- mismatch in the central corticotropin-releasing factor system. *J Comp Neurol.* 2008; 511:479–496. [PubMed: 18853426]
22. Ellenbroek BA, Derks N, Park HJ. Early maternal deprivation retards neurodevelopment in Wistar rats. *Stress.* 2005; 8:247–257. [PubMed: 16423713]
 23. Pruessner JC, Champagne F, Meaney MJ, Dagher A. Dopamine release in response to a psychological stress in humans and its relationship to early life maternal care: a positron emission tomography study using [¹¹C]raclopride. *J Neurosci.* 2004; 24:2825–2831. [PubMed: 15028776]
 24. Nemeroff CB. *Paradise Lost: The Neurobiological and Clinical Consequences of Child Abuse and Neglect.* Neuron. 2016; 89:892–909. [PubMed: 26938439]
 25. Authement ME, et al. Histone Deacetylase Inhibition Rescues Maternal Deprivation-Induced GABAergic Metaplasticity through Restoration of AKAP Signaling. *Neuron.* 2015; 86:1240–1252. [PubMed: 26050042]
 26. Wanat MJ, Bonci A, Phillips PE. CRF acts in the midbrain to attenuate accumbens dopamine release to rewards but not their predictors. *Nat neurosci.* 2013; 16:383–385. [PubMed: 23416448]
 27. Macey DJ, Koob GF, Markou A. CRF and urocortin decreased brain stimulation reward in the rat: reversal by a CRF receptor antagonist. *Brain research.* 2000; 866:82–91. [PubMed: 10825483]
 28. Storm JF. Action potential repolarization and a fast after-hyperpolarization in rat hippocampal pyramidal cells. *J Physiol.* 1987; 385:733–759. [PubMed: 2443676]
 29. Lee SH, et al. Multiple Forms of Endocannabinoid and Endovanilloid Signaling Regulate the Tonic Control of GABA Release. *J Neurosci.* 2015; 35:10039–10057. [PubMed: 26157003]
 30. Behrends JC, ten Bruggencate G. Changes in quantal size distributions upon experimental variations in the probability of release at striatal inhibitory synapses. *J Neurophysiol.* 1998; 79:2999–3011. [PubMed: 9636103]
 31. Riegel AC, Williams JT. CRF facilitates calcium release from intracellular stores in midbrain dopamine neurons. *Neuron.* 2008; 57:559–570. [PubMed: 18304485]
 32. Dermizaki I, Tsatsanis C, Alexaki VI, Castanas E, Margioris AN. Roles of protein kinase A (PKA) and PKC on corticotropin-releasing hormone (CRH)-induced elevation of cytosolic calcium from extra- and intra-cellular sources. *Hormones (Athens).* 2004; 3:252–258. [PubMed: 16982600]
 33. Chang SY, Kim U. Ionic mechanism of long-lasting discharges of action potentials triggered by membrane hyperpolarization in the medial lateral habenula. *J Neurosci.* 2004; 24:2172–2181. [PubMed: 14999068]
 34. Stocker M. Ca(2+)-activated K⁺ channels: molecular determinants and function of the SK family. *Nat Rev Neurosci.* 2004; 5:758–770. [PubMed: 15378036]
 35. Gu N, Vervaeke K, Storm JF. BK potassium channels facilitate high-frequency firing and cause early spike frequency adaptation in rat CA1 hippocampal pyramidal cells. *J Physiol.* 2007; 580:859–882. [PubMed: 17303637]
 36. Ren Y, et al. Regulation of surface localization of the small conductance Ca²⁺-activated potassium channel, Sk2, through direct phosphorylation by cAMP-dependent protein kinase. *J Biol Chem.* 2006; 281:11769–11779. [PubMed: 16513649]
 37. Bond CT, et al. Small conductance Ca²⁺-activated K⁺ channel knock-out mice reveal the identity of calcium-dependent after hyperpolarization currents. *J Neurosci.* 2004; 24:5301–5306. [PubMed: 15190101]
 38. Lin MT, Lujan R, Watanabe M, Adelman JP, Maylie J. SK2 channel plasticity contributes to LTP at Schaffer collateral-CA1 synapses. *Nat Neurosci.* 2008; 11:170–177. [PubMed: 18204442]
 39. Kyle BD, Braun AP. The regulation of BK channel activity by pre- and post-translational modifications. *Front Physiol.* 2014; 5:316. [PubMed: 25202279]
 40. Eghbal-Ahmadi M, et al. The developmental profile of the corticotropin releasing factor receptor (CRF2) in rat brain predicts distinct age-specific functions. *Brain Res Dev Brain Res.* 1998; 107:81–90. [PubMed: 9602071]
 41. Giesbrecht CJ, Mackay JP, Silveira HB, Urban JH, Colmers WF. Countervailing modulation of Ih by neuropeptide Y and corticotrophin-releasing factor in basolateral amygdala as a possible mechanism for their effects on stress-related behaviors. *J Neurosci.* 2010; 30:16970–16982. [PubMed: 21159967]

42. Wanat MJ, Hopf FW, Stuber GD, Phillips PE, Bonci A. Corticotropin-releasing factor increases mouse ventral tegmental area dopamine neuron firing through a protein kinase C-dependent enhancement of Ih. *J Physiol.* 2008; 586:2157–2170. [PubMed: 18308824]
43. Qiu DL, et al. Corticotrophin-releasing factor augments the I(H) in rat hypothalamic paraventricular nucleus parvocellular neurons in vitro. *J Neurophysiol.* 2005; 94:226–234. [PubMed: 15800070]
44. Sanchez G, Rodriguez MJ, Pomata P, Rela L, Murer MG. Reduction of an after hyperpolarization current increases excitability in striatal cholinergic interneurons in rat parkinsonism. *J Neurosci.* 2011; 31:6553–6564. [PubMed: 21525296]
45. Matthews EA, Weible AP, Shah S, Disterhoft JF. The BK-mediated fAHP is modulated by learning a hippocampus-dependent task. *Proc Natl Acad Sci USA.* 2008; 105:15154–15159. [PubMed: 18799739]
46. Brenner R, et al. BK channel beta4 subunit reduces dentate gyrus excitability and protects against temporal lobe seizures. *Nat Neurosci.* 2005; 8:1752–1759. [PubMed: 16261134]
47. Montgomery JR, Meredith AL. Genetic activation of BK currents in vivo generates bidirectional effects on neuronal excitability. *Proc Natl Acad Sci USA.* 2012; 109:18997–19002. [PubMed: 23112153]
48. Shruti S, Clem RL, Barth AL. A seizure-induced gain-of-function in BK channels is associated with elevated firing activity in neocortical pyramidal neurons. *Neurobiol Dis.* 2008; 30:323–330. [PubMed: 18387812]
49. Wang B, et al. Knockout of the BK beta4-subunit promotes a functional coupling of BK channels and ryanodine receptors that mediate a fAHP-induced increase in excitability. *J Neurophysiol.* 2016; 116:456–465. [PubMed: 27146987]
50. Blank T, et al. Corticotropin-releasing factor receptors couple to multiple G-proteins to activate diverse intracellular signaling pathways in mouse hippocampus: role in neuronal excitability and associative learning. *J Neurosci.* 2003; 23:700–707. [PubMed: 12533630]
51. Nimitvilai S, et al. Dopamine D2 receptor desensitization by dopamine or corticotropin releasing factor in ventral tegmental area neurons is associated with increased glutamate release. *Neuropharmacology.* 2014; 82:28–40. [PubMed: 24657149]
52. Kirby LG, et al. Corticotropin-releasing factor increases GABA synaptic activity and induces inward current in 5-hydroxytryptamine dorsal raphe neurons. *J Neurosci.* 2008; 28:12927–12937. [PubMed: 19036986]
53. Lowery-Gionta EG, et al. Corticotropin releasing factor signaling in the central amygdala is recruited during binge-like ethanol consumption in C57BL/6J mice. *J Neurosci.* 2012; 32:3405–3413. [PubMed: 22399763]
54. Kash TL, Winder DG. Neuropeptide Y and corticotropin-releasing factor bi-directionally modulate inhibitory synaptic transmission in the bed nucleus of the stria terminalis. *Neuropharmacology.* 2006; 51:1013–1022. [PubMed: 16904135]
55. Meye FJ, et al. Cocaine-evoked negative symptoms require AMPA receptor trafficking in the lateral habenula. *Nat Neurosci.* 2015; 18:376–378. [PubMed: 25643299]
56. Nelson SB, Turrigiano GG. Strength through diversity. *Neuron.* 2008; 60:477–482. [PubMed: 18995822]
57. Root DH, et al. Single rodent mesohabenular axons release glutamate and GABA. *Nat Neurosci.* 2014; 17:1543–1551. [PubMed: 25242304]
58. Tricoire L, et al. A blueprint for the spatiotemporal origins of mouse hippocampal interneuron diversity. *J Neurosci.* 2011; 31:10948–10970. [PubMed: 21795545]

Figure 1.

CRF increased the excitability of Lhb neurons. **A:** Whole cell patch clamp recording of action potentials (APs) in Lhb neurons from non-MD rats in response to depolarizing current injections (I) and CRF (250nM) bath application in slices, with synaptic transmission intact ($n=7$ cells from 7 rats; $F_{(1,6)}=12.3$). **B:** Sample AP recordings of Lhb neurons in response to a 50pA depolarizing current step before (baseline, black) and after CRF (red) application (calibration bar: 20mV/1s) from data in (A). **C:** Whole cell patch clamp recordings of APs as described in (A), with fast synaptic transmission blocked. ($n=10$ cells from 10 rats; $F_{(1,9)}=13.901$). **D:** Sample AP recordings as described in (B) with synaptic

blockade. **E**: Schematic of the method to measure AP threshold, mAHP and fAHP in sample AP recordings in response to the lowest depolarizing current step that generated the first AP/s before (black) and after CRF (red) application (calibration bar: 20mV/1s). Input resistance (R_{in}) was calculated from the steady-state voltage deflections generated in response to a 50pA hyperpolarizing current (calibration bar: 10mV/1s). **F**: Average amplitude of fAHP, mAHP and R_{in} derived from recordings in (C). Data are means $\pm$ SEM from cells for each condition (one cell from each rat); * $P < 0.05$, *** $P < 0.001$, by two-way RM-ANOVA (A and C) or paired Student's t-test (F).

Figure 2.

CRF depressed GABAergic transmission without affecting glutamatergic synaptic transmission. **A:** Top: Representative AMPAR-mediated mEPSC traces recorded from LHb neurons from non-MD rats before (baseline) and after CRF (250 nM) application (calibration bars: 30 pA/5s). Graphs below: Average mEPSC amplitude and frequency, and the cumulative probability plots of amplitude and frequency (inter-event interval) for all mEPSCs before and after CRF application. (n=10 cells from 10 non-MD rats). **B:** As described in (A) for GABA_AR-mediated mIPSCs before and after CRF application (calibration bars: 50 pA/5s). (n=6 cells from 6 non-MD rats). Data are means ± SEM from

cells for each condition (one cell from each rat). * $P < 0.05$, **** $P < 0.0001$ by paired Student's t -tests, or KS tests for cumulative distribution curves.

Figure 3.

CRF suppressed GABA release from GABAergic terminals through retrograde eCB signaling. (A–C): Average mIPSC amplitude and frequency, and cumulative probability plots of amplitude and frequency (inter-event interval) for all mIPSCs before and after the application of (A) CRF (250 nM) with GDPβS (300 μM) present in the patch pipette (n=10 cells from 10 non-MD rats), (B) WIN55,212-2 (2 μM; n=8 cells from 8 non-MD rats), or (C) CRF with AM251 (10 μM) present in the perfusate (n=7 cells recorded from 7 non-MD rats). Data are means ± SEM from cells for each condition (one cell from each rat). *P<

0.05, *** $P < 0.001$, **** $P < 0.001$, by paired Student's t -tests, or KS tests for the cumulative distribution curves.

Figure 4.

CRF increased LHb intrinsic excitability through G-protein-dependent CRFR1 signaling and requires intracellular Ca^{2+} . (**A to D**) Whole cell patch clamp recordings of APs in LHb neurons from non-MD rats in response to depolarizing step currents (left), with representative AP traces (right) in response to a 100pA depolarizing current step before (baseline, black) and after application of CRF (250 nM, red), CRFR1 antagonist antalarmin (A; 1 μM ; $F_{(1,6)}=0.222$), CRFR2 antagonist antisauvagine (B; 25 nM; $n=6$ cells from 6 non-MD rats, $F_{(1,5)}=5.057$), G-protein inhibitor GDP β S (C; 300 μM ; $n=4$ cells from 4 non-MD rats, $F_{(1,3)}=0.86$), and calcium chelator BAPTA (D; 30 mM; $n=6$ cells from 6 non-MD rats,

$F_{(1,5)} = 0.327$) was included in the patch pipette or perfusate). All recordings performed with fast synaptic transmission blocked. Data are means $\pm$ SEM from cells for each condition (one cell from each rat); ** $P < 0.01$ by two-way RM-ANOVA.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 5.

Pharmacological modulation of SK and BK channels altered CRF-induced increases in LHb intrinsic excitability in a PKA-dependent manner. (A–D): AP recordings in response to depolarizing current steps with representative AP traces in LHb neurons from non-MD rats in response to a 100pA current step before (baseline, black) and after (A) SK channel blocker apamin (100nM, blue; $n=11$ cells from 11 non-MD rats, $F_{(2,20)}=9.127$), (B) SK channel positive modulator CYPPA (10μM, green; $n=9$ cells from 9 non-MD rats, $F_{(2,16)}=2.058$), (C) BK channel blocker iberiotoxin (100nM, teal blue; $n=7$ cells from 7 non-MD rats, $F_{(1,6)}=3.039$), or (D) PKI(6–22) (10 μM, sky blue; $n=6$ cells from 6 non-MD rats,

$F_{(1,5)}=2.773$) alone or with subsequent CRF (250 nM, red) bath application. Right: Amplitudes of fAHP, mAHP and Rin derived from the AP recordings. All recordings were performed with fast synaptic transmission blocked. Data are means $\pm$ SEM from cells for each condition (one cell from each rat). * $P<0.05$, *** $P<0.001$, by two-way RM-ANOVA (left panels) or paired Student's t-tests (right panels).

Figure 6.

MD increased the excitability of LHb neurons. **A**. Top: Representative spontaneous AP recordings of LHb neurons from non-MD and MD rats using cell-attached voltage clamp recordings. Bottom: Percent of spontaneously active LHb neurons (left) and mean AP frequency (right) from each group (n= 33 cells from 18 rats or 29 cells from 19 rats, respectively). **B**. Representative traces (top) and number of APs (bottom) recorded from LHb neurons using whole cell patch clamp recording, with synaptic transmission intact, in slices from non-MD and MD rats in response to depolarizing current steps (calibration bar: 20mV/1s) [n= 24 cells from 11 rats (non-MD) or 24 cells from 9 rats (MD), $F_{(1,460)}=1.69$]. **C**.

As described in (B), with fast synaptic transmission blocked [n= 25 cells from 17 rats (non-MD) or 31 cells from 18 rats (MD), $F_{(1,600)}=35.84$]. **D:** Average amplitude of fAHP, mAHP and Rin derived from AP recordings in (C). **E:** Representative Western blots and quantitation of SK2 abundance (β -actin: loading control) in LHb tissue homogenates from non-MD and MD rats (n= 6 biological replicates). Data are means $\pm$ SEM. * $P<0.05$, ** $P<0.01$ by unpaired Student's t-test; **** $P<0.0001$ by two-way ANOVA.

Figure 7.

MD potentiated both glutamatergic and GABAergic synaptic transmission onto LHb neurons and increased E/I ratio. **A:** Representative AMPAR-mediated mEPSC traces from non-MD and MD rats (calibration bars: 30 pA/5s). Average mEPSC amplitude and frequency (left) and cumulative probability plots of amplitude and frequency (inter-event interval) (right) in non-MD and MD rats ($n=20$ cells from 20 rats or 17 cells from 17 rats, respectively). **B:** As described in (A) for GABA_AR-mediated mIPSC traces from non-MD and MD rats (calibration bars: 50 pA/5s; $n=13$ cells from 13 rats or 9 cells from 9 rats, respectively). **C:** Summary of E/I ratios obtained from non-MD and MD rats ($n=16$ cells

from 8 rats or 12 cells from 9 rats, respectively), with representative traces of evoked EPSCs (black, recorded at -55 mV holding potential) and IPSCs (red, recorded at $+10$ mV holding potential) in response to the stria medullaris stimulation for LHb neurons. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.0001$ by unpaired Student's t -tests, or KS tests for the cumulative distribution curves.

Figure 8.

The effects of CRF and apamin on LHB intrinsic excitability were blunted following MD in a PKA-dependent manner. **A:** Whole cell patch clamp AP recordings in LHB neurons in slices from MD rats in response to depolarizing current steps with representative AP traces (in response to a 100pA current step) before (baseline, black) and after application of CRF (250 nM, red), with fast synaptic transmission intact (top) or blocked (bottom). $n = 6$ cells from 6 MD rats; $F_{(1,5)} = 0.38$ and $F_{(1,5)} = 2.335$, respectively. **B:** Relative expression of *Crh* and *Crhr1* mRNA in the LHB from non-MD and MD rats (*Crh*: $n = 8$ and 10 biological replicates, respectively; *Crhr1*: $n = 9$ and 12 biological replicates, respectively). Data are

normalized to expression in the non-MD group and are means $\pm$ SEM. **C:** AP recordings in LHb neurons from MD rats in response to depolarizing current steps with representative AP traces (in response to a 100pA current step) before (baseline, black) and after application of (C) apamin (100nM, blue) or (D) 1-EBIO (300 mM, green), with fast synaptic transmission blocked. n=12 cells from 12 rats (C; $F_{(1,11)}=1.816$) or 7 cells from 7 rats (D; $F_{(1,6)}=5.677$). **E:** AP recordings in response to depolarizing current steps with representative AP traces (in response to a 100pA current step) in LHb neurons from MD rats with patch pipettes filled with normal internal solution (black) or along with PKI(6–22) (10 μ M; sky blue). Each condition n=8 cells from 8 rats; $F_{(1,140)}=45.02$. Data are means $\pm$ SEM. * $P<0.05$ by two-way RM-ANOVA; **** $P<0.0001$ by two-way ANOVA. **F:** Representative Western blots and quantitative data of PKA (GAPDH: loading control) in LHb tissue homogenates from non-MD and MD rats (n= 11 biological replicates).