Hall and his successors for recovery from apparent death—the establishment of physiological and pathological laboratories at home and abroad, and the issue of their reports—and the whole modern body of military medicine, surgery, and hygiene gained from the experience of the long series of wars which have followed in quick succession on the campaign in the Crimea.”

We cannot conclude this short notice without congratulating the Society and Mr. Wheatley, the librarian, on the successful issue of their arduous undertaking.

On Deafness, Giddiness, and Noises in the Head. By
EDWARD WOAKES, M.D. London: H. K. Lewis. 1879.

THIS treatise, a considerable portion of which has appeared at different times in the Medical Journals, is a valuable addition to aural literature. The first part of the work treats of ear complications during dentition, and in the exanthemata; also of ear-cough, and the laryngeal complications dependent on diseases of the ear. The greater part, however, is occupied with the consideration of labyrinthine giddiness and noises in the head, phenomena which receive very inadequate notice in the usual text books; certainly they are not generally treated of in a way commensurate with their frequency and their distressing character. Dr. Woakes has, in the volume before us, examined these symptoms, especially with regard to their causation, in a scientific spirit, and with a thoroughness which is truly commendable; and he has undoubtedly helped to elucidate a confessedly difficult subject. He finds a clue to the causation of these phenomena in the sources of the vessels which supply the affected region, and in the nerves which control these vessels. The vaso-motor nervous connections of the ear are traced most carefully, in order to discover alliances between parts histologically distinct, near or remote, which might explain vascular disturbance in the organ of hearing. His explanations are based on the fact that “in numerous instances in the economy the local connections of vaso-motor innervation are such that given tracts of tissue, not otherwise connected, are thereby kept in nutritive correlation one with another.” The references to the anatomical relations of the nerves of the ear are very interesting and instructive; and, while we may not always agree with his hypotheses, or admit that his deductions are always warranted by the facts, we must

concede that the views promulgated are both original and suggestive.

In the first part of his book, Dr. Woakes very properly insists on the importance of looking to the ear when brain symptoms develop themselves in an infant, and points out how readily, by the open petro-squamosal fissure in the child, inflammation in the tympanic cavity may extend to the membranes of the brain. On this subject he says, "I am more and more convinced that many obscure and often fatal illnesses in children which, from their affinities, are regarded as acute cases of hydrocephalus or brain fever, would, if examined from an aural point of view, be found either to originate in this organ, or be due to its implication in a reflex way; and I cannot too strongly insist on the importance of such an examination being made whenever brain symptoms arise, and the cause of them is doubtful, or in any respects obscure."

That common affection of infantile years, purulent otitis, has unquestionably often its source in a carious tooth, or in the process of dentition. Dr. Woakes finds an explanation in the circumstance that the inferior dental nerve and the nervi vasorum, round the temporal branch of the carotid, are brought into communication at the otic ganglion. The extraction of the carious tooth, or incision of the inflamed gums would, he says, probably cure the inflamed tympanum by "diminishing the waves of irritation."

We are familiar with the fact that a fit of coughing may be provoked by a mechanical irritation of the meatus of the ear. May not a blast of cold air playing on the external auditory canal—Dr. Woakes asks—produce other laryngeal disturbances, if long enough continued? The chain of events is as follows: "The irritation of the sensitive fibres of the auriculo-pneumo-gastricus distributed in the meatus, is reflected along the motor fibres of the superior laryngeal nerve, exciting in the larynx the act of coughing, by causing contraction of the crico-thyroid muscle, and in some very susceptible subjects, of vomiting also. If the stimulus be sufficiently prolonged, other conditions ensue—viz., tissue changes which make the evidence of participation in the process of the vaso-motor elements quite unimpeachable. We then see functional derangement becoming structural lesion, not by varying the experiment, but simply by prolonging it." In suggesting that a draught of cold air falling upon the ear may occasionally, in this way, excite spasmodic croup it is proposed "to provide against exposing the ears to the access of cold winds in

children predisposed to this affection, a provision which can be readily arranged by allowing the hair to fall over the ears, which it would do naturally if not prevented, as well as protecting the chest and throat, to which regions attention is at present entirely confined."

A most important section of the book is devoted to tinnitus aurium, the *bête noire* of aural surgery as it may be called, which forms a prominent feature in many diseases of the auditory apparatus, and is often more insufferable to the patient than the deafness with which it is generally associated. The continuity and monotony of the sound sometimes become intolerable, distracting the mind in the duties of the day, and making sleep almost impossible by night, bringing about a state of nervousness and irritability bordering upon mental disease, and, in cases where a predisposition exists, leading up to insanity itself. Deaf to the sounds of the outer world, he is tormented with the most dreadful noises within, and the unfortunate sufferer has been known to seek relief from his affliction in self-destruction. Any additional light which might aid in dispelling the obscurity still enveloping this subject is welcome, and it can be safely said that Dr. Woakes, in dealing with the subject of noises in the head, has done good service in that direction. While the causes on which the different forms of tinnitus aurium depend are not yet clear, we think that the following scheme, by Dr. Woakes, is a useful classification, and likely to help the student of this subject:—

LOCAL MORBID CONDITION.			CHARACTER OF SOUNDS.		
1. Labyrinthine congestion—arterial,	-	-	-	Pulsating noises.	
Do. do. venous,	-	-	-	Rushing noises.	
2. Anæmia, aneurism—extra aural,	-	-	-	Pulsating noises.	
3. Fluid in tympanum or Eustachian tubes,	} Catarrh of middle ear,			{ Bubbling and gurgling noises.	
4. Congestion of membrana tympani, Eustachian cavity of tympanum, or Eustachian tube, or all of these together,	{ Chronic non-suppurative inflammation of ear,			{ Tidal noises (the sound heard when a shell is held to the ear.)	
	inducing				
5. Undue contraction of intrinsic muscles,	{ Reflex irritation of tensor tympani nerve from otic ganglion,			{ Singing and tidal.	

In discussing the sounds due to labyrinthine congestion, which are pulsating or rushing, according as the congestion is arterial or venous, Dr. Woakes places emphasis on the fact that the internal ear, or perceptive part of the organ, gets its supply of blood from the vertebral artery which, rising from the sub-

clavian in the neck, is quite distinct from the blood supply of the more external parts. The calibre of this vessel is regulated by the *nervi vasorum* from the inferior cervical ganglion of the sympathetic, and is in this way brought into important relationships. When the inhibitory power of the nerve is removed, the ramifications of the terminal branches of the vessel in the labyrinth are not only surcharged with blood, but also press upon the delicate nervous arrangements, and so it is not surprising that the sounds of the ear are synchronous with the pulse. This is the class of cases for which the author would use hydrobromic acid, a drug the use of which was suggested by Dr. Fothergill, from the consideration of its effects in counteracting the aural symptoms produced by large doses of quinine. Dr. Woakes considers that cinchona and the hydrobromic salts have a special influence on the area supplied by the vertebral artery, that while the former paralyses the vessels, thus causing dilatation, the latter increases the inhibitory control of these vessels, and so brings about contraction. This is one of the instances in this work where an attractive theory, requiring more evidence before we can give it complete acceptance, is advanced as an explanation of the causation of a symptom. In like manner, iodine is held to have a specific influence on the circulation of the middle ear, and "the salts of ammonia owe their adaptability to the object in view to the physiological property which they possess of causing the peripheral vessels to contract their calibre." The truth is, it is impossible to define the class of cases improved by hydrobromic acid; while some cases seem benefited by it, unfortunately, in many cases of pulsating tinnitus, it is quite useless. Neither its *modus operandi*, nor the precise indications for its employment are at all clearly settled. With respect to the use of electricity in the treatment of noises in the head, Dr. Woakes rightly points out the importance of selecting the proper kind and strength of current, the mode of applying it, and the proper intervals between each application. He would employ the Faradic form when the tinnitus is presumed to be due to pressure of the ossicular chain upon the membrane of the foramen ovale, from defective action of the tensor tympani or stapedius muscle, or abnormal fixing of the ossicles from adhesions or other causes. In such cases, gentle Faradism applied at weekly intervals, by stimulating the stapedius and thus pulling out the stapes from the fenestra ovalis, or by increasing the mobility of the ossicles, might tend to remedy some of the causes on which the noises in the head may depend.

Our experience has not at all led us to share in Dr. Woakes' opinion of the usefulness of forcing vapour into the tympanum by the Valsalva method for the relief of disease of the middle ear. Many people, with Eustachian tubes normally permeable, are quite unable to inflate their tympana in this way; while, should there be any obstruction, such as a thickened mucous membrane, the patient's efforts are futile, and they are, at the same time, very likely to aggravate the hyperæmic condition which may be present. We are astonished to find the author recommending the bivalve form of speculum, an instrument which we thought was now discarded as cumbrous and unsafe, besides being constructed to effect an impossibility—namely, the dilatation of a passage having unyielding osseous walls.

We have done with fault finding, and would only add that to all who are interested in aural diseases, Dr. Woakes' work will, from its really scientific efforts to explain little understood phenomena, amply repay perusal; while, at the same time, it is far from being devoid of instruction to the general practitioner of medicine, because it has an interest beyond the particular organ of which it treats.

REPORTS OF HOSPITAL AND PRIVATE PRACTICE.

GLASGOW ROYAL INFIRMARY.

REPORTS UNDER THE SUPERVISION OF DR. J. WALLACE ANDERSON.

FROM DR. MACEWEN'S WARDS.

RADICAL CURE OF HERNIA: LIGATURE OF INTERNAL INGUINAL RING BY CHROMIC ACID SUTURES.—A. S., æt. 29, was admitted 29th September, suffering from inguinal hernia on the left side, of two years' duration. This had caused her very considerable annoyance, and she had frequent attacks of vomiting with great pain. These attacks occurred at intervals when the hernia was down, and lasted for some time after it was reduced. She was therefore determined to have it operated upon. On examination, Dr. Macewen found an inguinal hernia on the left side, the ring of which was large, and the hernia protruded whenever the patient rose out of bed. She wore a truss, and stated that she had done so for a couple of years.