Published in final edited form as:

React Oxyg Species (Apex). 2018 May ; 5(15): 145–158.

Oxidant Stress and Lipid Peroxidation in Acetaminophen Hepatotoxicity

Hartmut Jaeschke and Anup Ramachandran

Department of Pharmacology, Toxicology & Therapeutics, University of Kansas Medical Center, Kansas City, Kansas, USA

Abstract

Acetaminophen (APAP) overdose is the most frequent cause of liver injury and acute liver failure in many western countries. The mechanism of APAP-induced hepatocyte necrosis has been investigated extensively. The formation of a reactive metabolite and its binding to cellular proteins was initially thought to be responsible for cell death. A competing hypothesis was introduced that questioned the relevance of protein binding and instead suggested that P450-derived oxidant stress and lipid peroxidation causes APAP-induced liver injury. However, work over the last 15 years has reconciled some of these apparent contradictory hypotheses. This review summarizes the present state of knowledge on the role of reactive oxygen species (ROS) in APAP hepatotoxicity. Detailed investigations into the sources and relevance of the oxidant stress have clearly shown the critical role of the electron transport chain of mitochondria as main source of the oxidant stress. Other potential sources of ROS such as cytochrome P450 enzymes or NADPH oxidase on phagocytes are of limited relevance. The mitochondria-derived superoxide and peroxynitrite formation is initiated by the binding of the reactive metabolite to mitochondrial proteins and the amplification by mitogen activated protein kinases. The consequences of this oxidant stress are the opening of the mitochondrial membrane permeability transition pore with cessation of ATP synthesis, nuclear DNA fragmentation and ultimately cell necrosis. Lipid peroxidation is not a relevant mechanism of cell death but can be a marker of ROS formation. These mechanistic insights suggest that targeting mitochondrial oxidant stress is a promising therapeutic option for APAP hepatotoxicity.

Keywords

Acetaminophen hepatotoxicity; reactive oxygen species; mitochondria; cytochrome P450; neutrophils; lipid peroxidation

1. INTRODUCTION

Acetaminophen (APAP) is a widely used analgesic and anti-pyretic drug. It is considered safe at therapeutic doses but can cause liver injury and even acute liver failure and death when an overdose is consumed [1,2]. Some people overdose by taking a single large

For Correspondence: Hartmut Jaeschke, PhD., University of Kansas Medical Center, Department of Pharmacology, Toxicology & Therapeutics, 3901 Rainbow Blvd, MS 1018, Kansas City, KS, 66160, USA., Tel. +1 913 588 7969, hjaeschke@kumc.edu.

CONFLICT OF INTEREST DISCLOSURE

The authors declare that there is no conflict of interest.

overdose in an attempt to commit suicide others overdose unintentionally by taking multiple overdoses because APAP is present in numerous over-the-counter medications. As such, APAP is one of the most consumed drugs but is also one of the most important causes of drug-induced liver injury in the clinic. For this reason, the mechanisms of APAP hepatotoxicity have been investigated in animal models and in humans for more than 40 years [3,4]. The first hypothesis to explain the toxicity was that a small part of APAP is being metabolized by cytochrome P450 enzymes to form a reactive metabolite, which can be detoxified by glutathione (GSH), but when the capacity of hepatic GSH is exhausted, it also covalently binds to cellular proteins and causes cell death [5–7]. This early insight into the mechanism of cell death led to the development of N-acetylcysteine (NAC) as clinical antidote [8]. However, the covalent binding hypothesis was also challenged by an oxidant stress and lipid peroxidation (LPO) hypothesis [9], which created a significant controversy at the time [10,11]. This review discusses how this oxidant stress hypothesis has evolved and the current view of the co-existence of covalent binding and oxidant stress in the pathophysiology of APAP-induced cell death.

2. SOURCES OF OXIDANT STRESS: CYTOCHROME P450 ENZYMES

The first evidence of an oxidant stress after APAP overdose was reported in the late 1970s when it was shown that mice exhaled LPO products such as ethane and pentane [9]. Because it was newly recognized at the time that in a regenerating system *in vitro*, microsomal cytochrome P450 enzymes can produce superoxide and hydrogen peroxide [12], it was hypothesized that the metabolism of APAP triggered the oxidant stress [9]. This idea was supported by subsequent experiments showing reduced LPO and protection against APAP hepatotoxicity with P450 inhibitors and the opposite effects with inducers [13]. Later studies indicated that the main P450 enzyme involved in oxidative APAP metabolism is Cyp2E1 [14], which also participates in ethanol metabolism where it has been postulated to be a major source of ROS in ethanol-induced liver injury [15]. Both leakage of ROS during ethanol metabolism and induction of Cyp2E1 expression by ethanol with subsequent ROS formation through uncoupling have been shown [15]. Based on this insight from ethanol toxicity, it was assumed that the Cyp2E1-dependent metabolism of APAP is responsible for the oxidant stress [16]. However, very early after a role of ROS was first shown, the relevance of a metabolism-dependent oxidant stress was questioned because no direct evidence for ROS formation was found. If superoxide and hydrogen peroxide is formed during the metabolism of APAP, some of it would be detoxified by copper-zinc superoxide dismutase (SOD1) and glutathione peroxidase, which would lead to glutathione disulfide (GSSG) formation in the cytosol. Such an oxidant stress would cause an increase in hepatic GSSG levels and release of GSSG into bile and plasma as was shown for exposure to known oxidants such as t-butyl hydroperoxide [17,18] or the redox-cycling agent diquat [19]. However, APAP metabolism did neither cause elevated GSSG levels in the liver nor enhanced GSSG release into bile or plasma indicating that there is no relevant oxidant stress during the metabolism phase both in rats [17] and in mice [20]. In addition, when the formation of ROS was measured by 2',7'-dichlorofluorescein (DCF) fluorescence in isolated mouse hepatocytes, an oxidant stress was observed only after GSH depletion at 3–4 h after APAP exposure [21]. This again suggests that there is no oxidant stress during the

main metabolism phase of APAP in cultured hepatocytes [21]. Furthermore, when rats were exposed to a high overdose of APAP there was extensive depletion of hepatic GSH levels and formation of protein adducts indicating oxidative metabolism of APAP through the P450 system [22]. However, despite this extensive metabolism, no evidence of an oxidant stress or liver injury was detected in these rats *in vivo* [22]. Taken together, attempts to detect direct evidence for ROS formation during the metabolism phase of APAP toxicity consistently failed to find any support in rats or mice and in cultured hepatocytes. The early observations that LPO and toxicity correlated with P450 activities [13] does not contradict these findings because the formation of the reactive metabolite NAPQI and covalent binding to proteins is the initiating event in the toxicity. Thus, cytochrome P450 enzymes including Cyp2E1 are unlikely a relevant source of the oxidant stress observed during APAP toxicity.

3. SOURCES OF OXIDANT STRESS: MITOCHONDRIA

During the controversial discussion whether an oxidant stress actually occurs during APAP hepatotoxicity, it was discovered that hepatic GSSG levels increased substantially after the metabolism of APAP is over and GSH levels start to recover [23]. Interestingly, there was no GSSG release into bile suggesting that GSSG must be formed in a compartment where it cannot be released into the cytosol [23]. Mitochondria are cell organelles that can only exchange GSH but not GSSG between the matrix and the cytosol [24]. Therefore, when mitochondria were isolated and GSSG was measured in these organelles at various times after APAP, it was estimated that almost all of the GSSG detected in the whole liver is actually located within mitochondria [23,25]. This mitochondrial oxidant stress was confirmed in cultured hepatocytes using MitoSox Red [26]. In addition, after it was reported that the potent oxidant peroxynitrite was formed from superoxide and nitric oxide during APAP hepatotoxicity [27], it was shown that this peroxynitrite was also generated predominantly inside mitochondria [28]. Consistent with this observation, a loss of mitochondrial DNA and of SOD2, both located in the mitochondrial matrix, was found during APAP toxicity [28,29]. Thus, there is extensive, direct evidence that the formation of superoxide, hydrogen peroxide and peroxynitrite during APAP-induced liver injury occurs mainly within mitochondria.

The critical question whether this mitochondrial oxidative and nitrosative stress is actually relevant for the pathophysiology was subsequently addressed in a number of experiments. Support for the role of peroxynitrite was provided by the delayed treatment with sulfhydryl reagents GSH or NAC, which promoted the recovery of hepatic and mitochondrial GSH levels [30–32]. This mitochondrial GSH directly scavenged peroxynitrite [30] and detoxified ROS [30,32] and protected against APAP-induced liver injury [30–32]. In addition, the SOD mimetic Mito-TEMPO effectively protected against APAP-induced liver injury by dismutating superoxide and preventing peroxynitrite formation [33]. The fact that the mitochondria-targeted Mito-TEMPO was an order of magnitude more potent than TEMPO, which does not accumulate selectively in mitochondria, further supports the critical role of mitochondrial oxidant stress [33]. The phenolic compound resveratrol, which is a known antioxidant [34], effectively protected against APAP hepatotoxicity by scavenging peroxynitrite and preventing mitochondrial dysfunction [35]. In addition to the pharmacological intervention to scavenge ROS, mice with a partial deficiency of

mitochondrial SOD2 proved to be much more susceptible to APAP toxicity [36]. The enhanced susceptibility to APAP-induced liver injury in these mice was correlated with enhanced levels of GSSG, protein carbonyls, and nitrotyrosine adducts in the mitochondria [36]. More recently, several studies provided evidence that inhibiting ROS formation in mitochondria may be equally effective as scavenging them. Complex I of the mitochondrial electron transport chain is known to be an important source of ROS formation [37] and biguanides such as metformin are known to arrest complex I in the deactivated state and thereby preventing ROS leakage [38]. Consistent with these observations, metformin profoundly protected against APAP hepatotoxicity by inhibiting complex I thereby substantially attenuating the mitochondrial oxidant stress [39]. MCJ (methylation-controlled J protein), an endogenous regulator of complex I, reduces its activity [40]. Deficiency of MCJ in CD8+ T cells increased complex I activity and mitochondrial respiration but not mitochondrial ROS formation by promoting the formation of electron transport chain supercomplexes [41]. Interestingly, mice with MCJ-deficiency were highly protected against APAP hepatotoxicity, which correlated with improved mitochondrial function and reduced mitochondrial oxidant stress [42]. In summary, the direct evidence for a mitochondrial oxidant stress and peroxynitrite formation and the consistent beneficial effects when ROS are either scavenged or their formation is prevented, provides convincing evidence for the critical role of mitochondrial ROS derived from the electron transport chain in the pathophysiology of APAP-induced liver injury.

One aspect that has not been investigated in great detail is the question why the electron transport chain is leaking more electrons after APAP exposure. Early studies have shown that APAP can lead to inhibition of mitochondrial respiration [43]. Importantly, the mitochondrial dysfunction correlates with protein adduct formation in mitochondria [44]. A hepatotoxic dose of APAP caused extensive protein adducts formation in the whole liver and in mitochondria [44]. However, *N*-acetyl-*m*-aminophenol (AMAP), a non-toxic analog of APAP in mice, triggered extensive protein adducts formation only in the whole liver but not in mitochondria [44]. Interestingly, AMAP caused cytotoxicity in human hepatocytes, which correlated with mitochondrial protein adducts formation [45]. Although the specific adducted proteins in mitochondria that induce the ROS leakage from the electron transport chain have yet to be identified, the current data support the concept that the critical initiating event is the formation of the reactive metabolite *N*-acetyl-*p*-benzoquinone imine (NAPQI) and its reaction with mitochondrial proteins.

Although the initial protein adduct formation peaks by about 1–2 h after APAP treatment in the mouse [46,47], the full blown mitochondrial oxidant stress and parenchymal cell necrosis occurs several hours later [48]. This suggests the initial oxidant stress is insufficient to trigger the mitochondrial permeability transition pore (MPTP) opening and cell death. More recently it was recognized that the mitogen activated protein kinase (MAPK) c-Jun *N*-terminal kinase (JNK) is activated (phosphorylated) and P-JNK translocates to the mitochondria [49]. P-JNK binds to Sab (SH3 domain-binding protein that preferentially associates with Btk) in the outer mitochondrial membrane and triggers through several additional proteins a further impairment of the electron transport chain and enhanced ROS formation [50]. Thus, the JNK activation pathway *in vivo* clearly represents an amplification cycle for oxidant stress and peroxynitrite formation in the mitochondria [51]. Although it

was hypothesized that JNK is activated by oxidant stress, JNK is not a redox-sensitive kinase. Apoptosis signal-regulating kinase 1 (ASK1), which is sequestered by thioredoxin and can be activated by thioredoxin oxidation [52,53] and mixed-lineage protein kinase 3 (MLK3) [54] are redox-sensitive MAP kinases, which activate JNK through the MAP2 kinase MKK4 [55]. Inhibition or deletion of each of these kinases attenuates APAP-induced liver injury suggesting that this amplification loop for the oxidant stress is critical for cell death *in vivo* [52–55].

4. SOURCES OF OXIDANT STRESS: NADPH OXIDASE

The extensive necrosis during APAP-induced liver injury leads to the release of damage-associated molecular patterns (DAMPs), which trigger a sterile inflammatory response with formation of cytokines and chemokines and the activation and recruitment of phagocytes in the liver including Kupffer cells, neutrophils and blood monocytes [56–58]. All these inflammatory cells contain NADPH oxidase (NOX2), which can generate superoxide and hydrogen peroxide into the phagosome or to the extracellular space [59]. In addition, the neutrophil contains myeloperoxidase, which can form the potent oxidant and chlorinating agent hypochlorous acid from hydrogen peroxide and chloride [59]. In previous studies it was clearly demonstrated that a Kupffer cell-induced oxidant stress (mainly hydrogen peroxide) contributes to the early phase of hepatic ischemia-reperfusion injury [60]. Neutrophils are responsible for a second injury phase mainly by generating hypochlorite [61]. Although the ROS are generated outside of hepatocytes, due to the close proximity of the inflammatory cells to the target, these oxidants enter the cell and generate an intracellular oxidant stress in hepatocytes [62]. The presence of neutrophil-derived oxidants was demonstrated by increased intracellular GSSG levels and chlorotyrosine protein adducts in hepatocytes during the neutrophil-induced injury phase in different models [61,63–65]; this intracellular oxidant stress could be eliminated by inhibiting the neutrophil cytotoxicity [65,66]. A moderate extracellular-derived oxidant stress results in mitochondrial dysfunction, mitochondrial oxidant stress, MPTP opening and necrotic cell death [67,68]. Thus, in order to cause tissue injury, the phagocytes need to be recruited into the tissue and generate these oxidants in sufficient quantities and long enough close to the target cell in order to be able to disturb the intracellular homeostasis of the cells and generate mitochondrial dysfunction leading to the MPTP opening and cell death [62].

The first report of Kupffer cell activation and their potential impact on the injury after APAP overdose was reported in rats [69]. However, these findings were never confirmed and are unlikely to be relevant as rats show a high resistance to APAP and are a poor model for the human pathophysiology [22]. Nevertheless, a second report using a mouse model suggested that inactivation of Kupffer cells with gadolinium chloride (GdCl₃) treatment eliminated nitrotyrosine staining and liver injury [70]. These findings, however, were questioned for a number of reasons. First, the most active Kupffer cells are located in the periportal region [71], which makes it unlikely that ROS and peroxynitrite generated by these cells can selectively cause cell death in the centrilobular area. Second, deficiency of NADPH oxidase activity (in gp91phox knockout mice), the principle enzyme responsible for ROS formation in Kupffer cells, did not prevent the oxidant stress or injury [31]. Third, elimination of Kupffer cells with clodronate liposomes did not protect but actually even increased the

injury [72]. Together these data do not support the hypothesis that a Kupffer cell-derived oxidant stress can be responsible for APAP-induced liver injury.

Neutrophils are recruited into the liver after the early injury phase [73]. Necrotic cells release DAMPs including high mobility group box 1 (HMGB1) protein, mitochondrial DNA and nuclear DNA fragments [74,75], which activate toll like receptors, e.g. TLR9 [57,58] leading to activation of neutrophils by formation of cytokines and chemokine [57,58]. The controversial question is whether these neutrophils actually cause additional injury. In support of a neutrophil-mediated injury it was shown that neutropenia attenuates APAP hepatotoxicity [76,77]. However, it was argued that early pretreatment with a neutropenia-inducing antibody actually causes a preconditioning effect that is responsible for the protection rather than the absence of neutrophils [78]. In addition, there is no neutrophil activation and priming for ROS formation during the early injury phase in mice or humans [79,80], there is no direct evidence for a neutrophil-induced oxidant stress (chlorotyrosine staining) [81] and interventions that reduce neutrophil transmigration (anti-CD18 antibodies, CD18- or ICAM-1-deficiency) [73,79,81] have no effect on the oxidant stress and injury during APAP hepatotoxicity. Although not every result published in favor of a neutrophil-mediated injury phase can be readily explained as being caused by an off-target effect, it is difficult to overlook the fact gp91phox KO mice, which have neutrophils that are incapable of generating an oxidant stress and cause cytotoxicity, show the same oxidant stress and peroxynitrite formation and liver injury after APAP overdose as wild type animals [31,80]. Thus, the preponderance of experimental evidence does not support a neutrophil-induced injury during APAP hepatotoxicity but indicates that the main purpose of neutrophil recruitment into the injured liver is to assist in clean-up of necrotic tissue in preparation of regeneration [56,57].

Blood-derived monocytes are also recruited into the liver of animals with APAP-induced liver injury. They accumulate mainly at later time points, i.e. after the peak of injury, and preferentially localize in the areas of necrosis [82,83]. Although these cells contain NADPH oxidase and can generate ROS under certain conditions [84], in the case of APAP they develop an anti-inflammatory macrophage phenotype (M2) [82,85,86]. The recruitment occurs mainly through monocyte chemoattractant protein-1 (MCP-1, CCL2), which can be generated by macrophages but also by injured hepatocytes [82]. The receptor for MCP-1 on monocytes/macrophages is CCR2. Mice with CCR2 deficiency are not protected against APAP toxicity but show reduced regeneration and repair of the necrotic lesions [82,83,87,88]. Similarly, animals deficient of MCP-1 experience the same injury as wild type animals but again have a delayed repair [82]. These results indicate that blood-derived monocytes/macrophages are important for the recovery from the injury and do not likely participate in the injury process. Furthermore, the fact that gp91phox knock out mice, which have a non-functional NADPH oxidase, show a similar recovery as wild type animals suggest that the removal of necrotic cell debris by these phagocytes is not dependent of ROS formation [80].

In summary, all phagocytes have the capacity to generate ROS and all have been shown to be involved in various inflammatory tissue injury models including hepatic ischemia-reperfusion injury, obstructive cholestasis and endotoxin shock [89]. However, extensive investigations into the role of these inflammatory in APAP hepatotoxicity did not provide

any convincing evidence that inflammatory cell-derived oxidant stress is a relevant factor in the pathophysiology.

5. CONSEQUENCES OF OXIDANT STRESS: LIPID PEROXIDATION

Lipid peroxidation (LPO) is a frequently, even today, invoked as a mechanism of cell death during APAP hepatotoxicity [e.g. 90]. LPO is a free radical reaction process that is initiated by hydroxyl radical formation from hydrogen peroxide (Fe^{2+} -dependent Fenton reaction) and the generation of lipid radicals leading to the destruction of polyunsaturated fatty acids in lipid membranes [91]. LPO can cause rapid catastrophic breakdown of the membrane potential and ion gradients leading to cell necrosis. The lipid soluble vitamin α -tocopherol (vitamin E) is a highly effective radical scavenger that can interrupt the radical chain reaction and block LPO [91]. Early studies on LPO in the APAP mouse model used ethane and pentane exhalation and hepatic malondialdehyde as indicators of LPO [9,13]. Massive LPO led to severe injury in these animals within 4 h [9,13] and vitamin E pretreatment effectively inhibited LPO and injury induced by APAP [92]. Although these studies clearly support the hypothesis that LPO as the cause and not a consequence of the injury, one aspect that is frequently overlooked when discussing these experiments is that Wendel and coworkers used animals fed a vitamin E-deficient diet with a high content of polyunsaturated fatty acids [9,13,92]. However, a more recent follow-up study using animals on a regular diet found only minimal evidence for LPO after APAP and treatment with d-alpha-tocopheryl acetate, which increased hepatic tocopherol levels by almost 7-fold, had no effect on liver injury [93]. These data suggest that animals on a regular diet have enough lipid-soluble antioxidants to prevent extensive LPO and thus LPO is quantitatively not a relevant injury mechanism in APAP-induced liver injury [94]. However, the much higher susceptibility of animals on a diet low in vitamin E but high in polyunsaturated fatty acids indicates a potential danger under special circumstances of the oxidant stress generated after APAP overdose. On the other hand, these data also suggest that the mechanism of cell death even *in vivo* can be changed based on the experimental conditions. This means that special attention needs to be paid to the translational relevance of any experimental design.

6. OXIDANT STRESS AND MITOCHONDRIAL PERMEABILITY TRANSITION PORE OPENING

The mitochondrial permeability transition (MPT) is a phenomenon whereby the mitochondrial inner membrane becomes permeable to molecules of less than 1500 da [95], resulting in release of mitochondrial proteins into the cytosol and subsequent mitochondrial dysfunction. It has been well established that APAP overdose causes activation of the MPT, which results in dissipation of the proton gradient across the inner mitochondrial membrane and resulting loss of mitochondrial membrane potential, a collapse of ATP generation, leading to hepatocyte necrosis [95,96]. The regulation of the MTP has been shown to be influenced by matrix calcium and free radicals, and is also dependent on electron flux through the electron transport chain, especially from complex I [97]. This is relevant in light of the earlier mentioned, beneficial effect of metformin against APAP hepatotoxicity [39], which inhibited complex I activity. The molecular composition of the mitochondrial

permeability transition pore (MPTP) has been under intense investigation for decades and the current consensus implicates the proteins Bax and Bak as its components on the outer mitochondrial membrane [98,99], which function along with subunits of the ATP synthase on the inner mitochondrial membrane [94]. The concept that ATP synthase may form a component of the MPTP originated from the identification of the interaction of its lateral stalk (the Oligomycin Sensitivity Conferring Protein, OSCP) with cyclophilin D, the best characterized component of the MPTP [100]. Inhibition of cyclophilin D was shown to provide partial protection against APAP induced hepatocyte necrosis *in vitro* [95]. However, this effect depends on concentration of APAP *in vivo*, with protection against liver injury at moderate 200mg/kg concentrations [101], but no effect at a higher concentration of 600mg/kg [102]. This dose dependence of APAP concentrations on the MPT was also seen using intravital microscopy, where a dose of 150mg/kg APAP resulted in transient loss of membrane potential without cell necrosis [103]. In addition to calcium and reactive oxygen species, cellular iron has also been shown to induce the MPT [104]. The ferric chelator deferoxamine (Desferal) was shown to delay APAP hepatotoxicity in mice [105] and an increase in cytosolic chelatable ferrous iron, presumably from disrupted lysosomes [106], which is taken up into polarized mitochondria to trigger the MPT and hepatocyte necrosis was observed after APAP treatment of hepatocytes *in vitro* [107]. Further characterization of the lysosome to mitochondrial iron transfer revealed that lysosomally targeted iron chelation or inhibition of the mitochondrial calcium uniporter provided significant protection against mitochondrial depolarization and APAP hepatotoxicity [108] suggesting that cellular iron acts in concert with calcium and reactive oxygen species to influence the MPT.

7. SUMMARY AND CONCLUSIONS

ROS and peroxynitrite are critical mediators of APAP-induced cell death (Figure 1). The electron transport chain in mitochondria is the principal source of the oxidant stress. Binding of NAPQI to mitochondrial proteins initiates the oxidant stress, which is further amplified through JNK activation. The dependence of the oxidant stress on reactive metabolite formation and protein binding reconciles previously competing hypotheses. Other sources of ROS such as cytochrome P450 and NADPH oxidase appear to be of limited relevance in the pathophysiology (Figure 1). The major consequence of the mitochondrial oxidant stress is the opening of the MPTP, which leads to cessation of ATP synthesis, nuclear DNA fragmentation and ultimately necrotic cell death. LPO is quantitatively insufficient to cause cell death but can be considered as evidence for oxidant stress. Based on this mechanistic insight, potential therapeutic targets are interventions to prevent the mitochondria oxidant stress, to scavenge superoxide or peroxynitrite inside mitochondria, to prevent the MPTP formation and to remove damaged mitochondria by autophagy and replace them by biogenesis.

Acknowledgments

This work was supported in part by a National Institutes of Health grant R01 DK102142, and by grants from the National Institute of General Medical Sciences (P20 GM103549 and P30 GM118247) of the National Institutes of Health.

ABBREVIATIONS

AMAP	N-acetyl-m-aminophenol
APAP	acetaminophen
ASK1	apoptosis signal-regulating kinase 1
DAMPs	damage-associated molecular patterns
GSH	glutathione
GSSG	glutathione disulfide
HMGB1	high mobility group box 1
JNK	c-jun N-terminal kinase
LPO	lipid peroxidation
MAPK	mitogen activated protein kinase
MCP-1	monocyte chemoattractant protein-1
MLP3	mixed-lineage protein kinase 3
MPTP	mitochondrial permeability transition pore
N-acetylcysteine	
NAPQI	N-acetyl-p-benzoquinone imine
ROS	reactive oxygen species
SOD	superoxide dismutase

References

1. Larson AM. Acetaminophen hepatotoxicity. Clin Liver Dis. 2007 Aug; 11(3):525–48. DOI: 10.1016/j.cld.2007.06.006 [PubMed: 17723918]
2. Jaeschke H. Acetaminophen: Dose-Dependent Drug Hepatotoxicity and Acute Liver Failure in Patients. Dig Dis. 2015; 33(4):464–71. DOI: 10.1159/000374090 [PubMed: 26159260]
3. Jaeschke H, Xie Y, McGill MR. Acetaminophen-induced Liver Injury: from Animal Models to Humans. J Clin Transl Hepatol. 2014 Sep; 2(3):153–61. DOI: 10.14218/JCTH.2014.00014 [PubMed: 26355817]
4. Ramachandran A, Jaeschke H. Mechanisms of acetaminophen hepatotoxicity and their translation to the human pathophysiology. J Clin Transl Res. 2017 Feb; 3(Suppl 1):157–169. DOI: 10.18053/jctres.03.2017S1.002 [PubMed: 28670625]
5. Mitchell JR, Jollow DJ, Potter WZ, Davis DC, Gillette JR, Brodie BB. Acetaminophen-induced hepatic necrosis. I. Role of drug metabolism. J Pharmacol Exp Ther. 1973 Oct; 187(1):185–94. [PubMed: 4746326]
6. Jollow DJ, Mitchell JR, Potter WZ, Davis DC, Gillette JR, Brodie BB. Acetaminophen-induced hepatic necrosis. II. Role of covalent binding in vivo. J Pharmacol Exp Ther. 1973 Oct; 187(1):195–202. [PubMed: 4746327]

7. Mitchell JR, Jollow DJ, Potter WZ, Gillette JR, Brodie BB. Acetaminophen-induced hepatic necrosis. IV. Protective role of glutathione. *J Pharmacol Exp Ther*. 1973 Oct; 187(1):211–7. [PubMed: 4746329]
8. Rumack BH, Bateman DN. Acetaminophen and acetylcysteine dose and duration: past, present and future. *Clin Toxicol (Phila)*. 2012 Feb; 50(2):91–8. DOI: 10.3109/15563650.2012.659252 [PubMed: 22320209]
9. Wendel A, Feuerstein S, Konz KH. Acute paracetamol intoxication of starved mice leads to lipid peroxidation in vivo. *Biochem Pharmacol*. 1979 Jul 1; 28(13):2051–5. [PubMed: 475847]
10. Mitchell JR, Smith CV, Hughes H, Lauterburg BH, Horning MG. Overview of alkylation and peroxidation mechanisms in acute lethal hepatocellular injury by chemically reactive metabolites. *Semin Liver Dis*. 1981 May; 1(2):143–50. [PubMed: 7347018]
11. Smith CV, Mitchell JR. Acetaminophen hepatotoxicity in vivo is not accompanied by oxidant stress. *Biochem Biophys Res Commun*. 1985 Nov 27; 133(1):329–36. [PubMed: 4074372]
12. Kuthan H, Tsuji H, Graf H, Ullrich V. Generation of superoxide anion as a source of hydrogen peroxide in a reconstituted monooxygenase system. *FEBS Lett*. 1978 Jul 15; 91(2):343–5. [PubMed: 210047]
13. Wendel A, Feuerstein S. Drug-induced lipid peroxidation in mice--I. Modulation by monooxygenase activity, glutathione and selenium status. *Biochem Pharmacol*. 1981 Sep 15; 30(18):2513–20. [PubMed: 7306203]
14. Lee SS, Buters JT, Pineau T, Fernandez-Salguero P, Gonzalez FJ. Role of CYP2E1 in the hepatotoxicity of acetaminophen. *J Biol Chem*. 1996 May 17; 271(20):12063–7. [PubMed: 8662637]
15. Cederbaum AI. Role of Cytochrome P450 and Oxidative Stress in Alcohol-Induced Liver Injury. *Reactive Oxygen Species*. 2017; 4(11):303–319. DOI: 10.20455/ros.2017.851
16. Kass GE. Mitochondrial involvement in drug-induced hepatic injury. *Chem Biol Interact*. 2006 Oct 27; 163(1–2):145–59. DOI: 10.1016/j.cbi.2006.06.007 [PubMed: 16889760]
17. Lauterburg BH, Smith CV, Hughes H, Mitchell JR. Biliary excretion of glutathione and glutathione disulfide in the rat. Regulation and response to oxidative stress. *J Clin Invest*. 1984 Jan; 73(1):124–33. DOI: 10.1172/JCI111182 [PubMed: 6690473]
18. Adams JD Jr, Lauterburg BH, Mitchell JR. Plasma glutathione and glutathione disulfide in the rat: regulation and response to oxidative stress. *J Pharmacol Exp Ther*. 1983 Dec; 227(3):749–54. [PubMed: 6655568]
19. Smith CV, Hughes H, Lauterburg BH, Mitchell JR. Oxidant stress and hepatic necrosis in rats treated with diquat. *J Pharmacol Exp Ther*. 1985 Oct; 235(1):172–7. [PubMed: 3930698]
20. Smith CV, Jaeschke H. Effect of acetaminophen on hepatic content and biliary efflux of glutathione disulfide in mice. *Chem Biol Interact*. 1989; 70(3–4):241–8. [PubMed: 2743472]
21. Bajt ML, Knight TR, Lemasters JJ, Jaeschke H. Acetaminophen-induced oxidant stress and cell injury in cultured mouse hepatocytes: protection by N-acetyl cysteine. *Toxicol Sci*. 2004 Aug; 80(2):343–9. DOI: 10.1093/toxsci/kfh151 [PubMed: 15115886]
22. McGill MR, Williams CD, Xie Y, Ramachandran A, Jaeschke H. Acetaminophen-induced liver injury in rats and mice: comparison of protein adducts, mitochondrial dysfunction, and oxidative stress in the mechanism of toxicity. *Toxicol Appl Pharmacol*. 2012 Nov 1; 264(3):387–94. DOI: 10.1016/j.taap.2012.08.015 [PubMed: 22980195]
23. Jaeschke H. Glutathione disulfide formation and oxidant stress during acetaminophen-induced hepatotoxicity in mice in vivo: the protective effect of allopurinol. *J Pharmacol Exp Ther*. 1990 Dec; 255(3):935–41. [PubMed: 2262912]
24. Olafsdottir K, Reed DJ. Retention of oxidized glutathione by isolated rat liver mitochondria during hydroperoxide treatment. *Biochim Biophys Acta*. 1988 Mar 17; 964(3):377–82. [PubMed: 3349102]
25. Tirmenstein MA, Nelson SD. Acetaminophen-induced oxidation of protein thiols. Contribution of impaired thiol-metabolizing enzymes and the breakdown of adenine nucleotides. *J Biol Chem*. 1990 Feb 25; 265(6):3059–65. [PubMed: 2303440]

26. Yan HM, Ramachandran A, Bajt ML, Lemasters JJ, Jaeschke H. The oxygen tension modulates acetaminophen-induced mitochondrial oxidant stress and cell injury in cultured hepatocytes. *Toxicol Sci*. 2010 Oct; 117(2):515–23. DOI: 10.1093/toxsci/kfq208 [PubMed: 20616211]
27. Hinson JA, Pike SL, Pumford NR, Mayeux PR. Nitrotyrosine-protein adducts in hepatic centrilobular areas following toxic doses of acetaminophen in mice. *Chem Res Toxicol*. 1998 Jun; 11(6):604–7. DOI: 10.1021/tx9800349 [PubMed: 9625727]
28. Cover C, Mansouri A, Knight TR, Bajt ML, Lemasters JJ, Pessayre D, Jaeschke H. Peroxynitrite-induced mitochondrial and endonuclease-mediated nuclear DNA damage in acetaminophen hepatotoxicity. *J Pharmacol Exp Ther*. 2005 Nov; 315(2):879–87. DOI: 10.1124/jpet.105.088898 [PubMed: 16081675]
29. Agarwal R, MacMillan-Crow LA, Rafferty TM, Saba H, Roberts DW, Fifer EK, James LP, Hinson JA. Acetaminophen-induced hepatotoxicity in mice occurs with inhibition of activity and nitration of mitochondrial manganese superoxide dismutase. *J Pharmacol Exp Ther*. 2011 Apr; 337(1):110–6. DOI: 10.1124/jpet.110.176321 [PubMed: 21205919]
30. Knight TR, Ho YS, Farhood A, Jaeschke H. Peroxynitrite is a critical mediator of acetaminophen hepatotoxicity in murine livers: protection by glutathione. *J Pharmacol Exp Ther*. 2002 Nov; 303(2):468–75. DOI: 10.1124/jpet.102.038968 [PubMed: 12388625]
31. James LP, McCullough SS, Knight TR, Jaeschke H, Hinson JA. Acetaminophen toxicity in mice lacking NADPH oxidase activity: role of peroxynitrite formation and mitochondrial oxidant stress. *Free Radic Res*. 2003 Dec; 37(12):1289–97. [PubMed: 14753753]
32. Saito C, Zwingmann C, Jaeschke H. Novel mechanisms of protection against acetaminophen hepatotoxicity in mice by glutathione and N-acetylcysteine. *Hepatology*. 2010 Jan; 51(1):246–54. DOI: 10.1002/hep.23267 [PubMed: 19821517]
33. Du K, Farhood A, Jaeschke H. Mitochondria-targeted antioxidant Mito-Tempo protects against acetaminophen hepatotoxicity. *Arch Toxicol*. 2017 Feb; 91(2):761–773. DOI: 10.1007/s00204-016-1692-0 [PubMed: 27002509]
34. Holthoff JH, Woodling KA, Doerge DR, Burns ST, Hinson JA, Mayeux PR. Resveratrol, a dietary polyphenolic phytoalexin, is a functional scavenger of peroxynitrite. *Biochem Pharmacol*. 2010 Oct 15; 80(8):1260–5. DOI: 10.1016/j.bcp.2010.06.027 [PubMed: 20599800]
35. Du K, McGill MR, Xie Y, Bajt ML, Jaeschke H. Resveratrol prevents protein nitration and release of endonucleases from mitochondria during acetaminophen hepatotoxicity. *Food Chem Toxicol*. 2015 Jul; 81:62–70. DOI: 10.1016/j.fct.2015.04.014 [PubMed: 25865938]
36. Ramachandran A, Lebofsky M, Weinman SA, Jaeschke H. The impact of partial manganese superoxide dismutase (SOD2)-deficiency on mitochondrial oxidant stress, DNA fragmentation and liver injury during acetaminophen hepatotoxicity. *Toxicol Appl Pharmacol*. 2011 Mar 15; 251(3):226–33. DOI: 10.1016/j.taap.2011.01.004 [PubMed: 21241727]
37. Grivennikova VG, Kapustin AN, Vinogradov AD. Catalytic activity of NADH-ubiquinone oxidoreductase (complex I) in intact mitochondria. evidence for the slow active/inactive transition. *J Biol Chem*. 2001 Mar 23; 276(12):9038–44. DOI: 10.1074/jbc.M009661200 [PubMed: 11124957]
38. Matsuzaki S, Humphries KM. Selective inhibition of deactivated mitochondrial complex I by biguanides. *Biochemistry*. 2015 Mar 24; 54(11):2011–21. DOI: 10.1021/bi501473h [PubMed: 25719498]
39. Du K, Ramachandran A, Weemhoff JL, Chavan H, Xie Y, Krishnamurthy P, Jaeschke H. Editor's Highlight: Metformin Protects Against Acetaminophen Hepatotoxicity by Attenuation of Mitochondrial Oxidant Stress and Dysfunction. *Toxicol Sci*. 2016 Dec; 154(2):214–226. DOI: 10.1093/toxsci/kfw158 [PubMed: 27562556]
40. Hatle KM, Gummadidala P, Navasa N, Bernardo E, Dodge J, Silverstrim B, Fortner K, Burg E, Suratt BT, Hammer J, Radermacher M, Taatjes DJ, Thornton T, Anguita J, Rincon M. MCJ/DnaJC15, an endogenous mitochondrial repressor of the respiratory chain that controls metabolic alterations. *Mol Cell Biol*. 2013 Jun; 33(11):2302–14. DOI: 10.1128/MCB.00189-13 [PubMed: 23530063]
41. Champagne DP, Hatle KM, Fortner KA, D'Alessandro A, Thornton TM, Yang R, Torralba D, Tomás-Cortázar J, Jun YW, Ahn KH, Hansen KC, Haynes L, Anguita J, Rincon M. Fine-Tuning of CD8(+) T Cell Mitochondrial Metabolism by the Respiratory Chain Repressor MCJ Dictates

- Protection to Influenza Virus. Immunity. 2016 Jun 21; 44(6):1299–311. DOI: 10.1016/j.immuni.2016.02.018 [PubMed: 27234056]
42. Barbier-Torres L, Iruzubieta P, Fernández-Ramos D, Delgado TC, Taibo D, Guitiérrez-de-Juan V, Varela-Rey M, Azkargorta M, Navasa N, Fernández-Tussy P, Zubiete-Franco I, Simon J, Lopitz-Otsoa F, Lachiondo-Ortega S, Crespo J, Masson S, McCain MV, Villa E, Reeves H, Elortza F, Lucena MI, Hernández-Alvarez MI, Zorzano A, Andrade RJ, Lu SC, Mato JM, Anguita J, Rincón M, Martínez-Chantar ML. The mitochondrial negative regulator MCJ is a therapeutic target for acetaminophen-induced liver injury. Nat Commun. 2017 Dec 12.8(1):2068.doi: 10.1038/s41467-017-01970-x [PubMed: 29233977]
43. Meyers LL, Beierschmitt WP, Khairallah EA, Cohen SD. Acetaminophen-induced inhibition of hepatic mitochondrial respiration in mice. Toxicol Appl Pharmacol. 1988 May; 93(3):378–87. [PubMed: 3368917]
44. Tirmenstein MA, Nelson SD. Subcellular binding and effects on calcium homeostasis produced by acetaminophen and a nonhepatotoxic regioisomer, 3'-hydroxyacetanilide, in mouse liver. J Biol Chem. 1989 Jun 15; 264(17):9814–9. [PubMed: 2524496]
45. Xie Y, McGill MR, Du K, Dorko K, Kumer SC, Schmitt TM, Ding WX, Jaeschke H. Mitochondrial protein adducts formation and mitochondrial dysfunction during N-acetyl-m-aminophenol (AMAP)-induced hepatotoxicity in primary human hepatocytes. Toxicol Appl Pharmacol. 2015 Dec 1; 289(2):213–22. DOI: 10.1016/j.taap.2015.09.022 [PubMed: 26431796]
46. Roberts DW, Bucci TJ, Benson RW, Warbritton AR, McRae TA, Pumford NR, Hinson JA. Immunohistochemical localization and quantification of the 3-(cystein-S-yl)-acetaminophen protein adduct in acetaminophen hepatotoxicity. Am J Pathol. 1991 Feb; 138(2):359–71. [PubMed: 1992763]
47. McGill MR, Lebofsky M, Norris HR, Slawson MH, Bajt ML, Xie Y, Williams CD, Wilkins DG, Rollins DE, Jaeschke H. Plasma and liver acetaminophen-protein adduct levels in mice after acetaminophen treatment: dose-response, mechanisms, and clinical implications. Toxicol Appl Pharmacol. 2013 Jun 15; 269(3):240–9. DOI: 10.1016/j.taap.2013.03.026 [PubMed: 23571099]
48. Knight TR, Kurtz A, Bajt ML, Hinson JA, Jaeschke H. Vascular and hepatocellular peroxynitrite formation during acetaminophen toxicity: role of mitochondrial oxidant stress. Toxicol Sci. 2001 Aug; 62(2):212–20. [PubMed: 11452133]
49. Hanawa N, Shinohara M, Saberi B, Gaarde WA, Han D, Kaplowitz N. Role of JNK translocation to mitochondria leading to inhibition of mitochondria bioenergetics in acetaminophen-induced liver injury. J Biol Chem. 2008 May 16; 283(20):13565–77. DOI: 10.1074/jbc.M708916200 [PubMed: 18337250]
50. Win S, Than TA, Min RW, Aghajan M, Kaplowitz N. c-Jun N-terminal kinase mediates mouse liver injury through a novel Sab (SH3BP5)-dependent pathway leading to inactivation of intramitochondrial Src. Hepatology. 2016 Jun; 63(6):1987–2003. DOI: 10.1002/hep.28486 [PubMed: 26845758]
51. Saito C, Lemasters JJ, Jaeschke H. c-Jun N-terminal kinase modulates oxidant stress and peroxynitrite formation independent of inducible nitric oxide synthase in acetaminophen hepatotoxicity. Toxicol Appl Pharmacol. 2010 Jul; 246(1–2):8–17. DOI: 10.1016/j.taap.2010.04.015 [PubMed: 20423716]
52. Nakagawa H, Maeda S, Hikiba Y, Ohmae T, Shibata W, Yanai A, Sakamoto K, Ogura K, Noguchi T, Karin M, Ichijo H, Omata M. Deletion of apoptosis signal-regulating kinase 1 attenuates acetaminophen-induced liver injury by inhibiting c-Jun N-terminal kinase activation. Gastroenterology. 2008 Oct; 135(4):1311–21. DOI: 10.1053/j.gastro.2008.07.006 [PubMed: 18700144]
53. Xie Y, Ramachandran A, Breckenridge DG, Liles JT, Lebofsky M, Farhood A, Jaeschke H. Inhibitor of apoptosis signal-regulating kinase 1 protects against acetaminophen-induced liver injury. Toxicol Appl Pharmacol. 2015 Jul 1; 286(1):1–9. DOI: 10.1016/j.taap.2015.03.019 [PubMed: 25818599]
54. Sharma M, Gadang V, Jaeschke A. Critical role for mixed-lineage kinase 3 in acetaminophen-induced hepatotoxicity. Mol Pharmacol. 2012 Nov; 82(5):1001–7. DOI: 10.1124/mol.112.079863 [PubMed: 22918968]

55. Zhang J, Min RWM, Le K, Zhou S, Aghajani M, Than TA, Win S, Kaplowitz N. The role of MAP2 kinases and p38 kinase in acute murine liver injury models. *Cell Death Dis.* 2017 Jun 29;8(6):e2903.doi: 10.1038/cddis.2017.295 [PubMed: 28661486]
56. Jaeschke H, Williams CD, Ramachandran A, Bajt ML. Acetaminophen hepatotoxicity and repair: the role of sterile inflammation and innate immunity. *Liver Int.* 2012 Jan; 32(1):8–20. DOI: 10.1111/j.1478-3231.2011.02501.x [PubMed: 21745276]
57. Woolbright BL, Jaeschke H. Role of the inflammasome in acetaminophen-induced liver injury and acute liver failure. *J Hepatol.* 2017 Apr; 66(4):836–848. DOI: 10.1016/j.jhep.2016.11.017 [PubMed: 27913221]
58. Kubes P, Mehal WZ. Sterile inflammation in the liver. *Gastroenterology.* 2012 Nov; 143(5):1158–72. DOI: 10.1053/j.gastro.2012.09.008 [PubMed: 22982943]
59. El-Benna J, Hurtado-Nedelec M, Marzaioli V, Marie JC, Gougerot-Pocidalo MA, Dang PM. Priming of the neutrophil respiratory burst: role in host defense and inflammation. *Immunol Rev.* 2016 Sep; 273(1):180–93. DOI: 10.1111/imr.12447 [PubMed: 27558335]
60. Jaeschke H, Farhood A. Neutrophil and Kupffer cell-induced oxidant stress and ischemia-reperfusion injury in rat liver. *Am J Physiol.* 1991 Mar; 260(3 Pt 1):G355–62. [PubMed: 2003603]
61. Hasegawa T, Malle E, Farhood A, Jaeschke H. Generation of hypochlorite-modified proteins by neutrophils during ischemia-reperfusion injury in rat liver: attenuation by ischemic preconditioning. *Am J Physiol Gastrointest Liver Physiol.* 2005 Oct; 289(4):G760–7. DOI: 10.1152/ajpgi.00141.2005 [PubMed: 15994427]
62. Jaeschke H. Mechanisms of Liver Injury II. Mechanisms of neutrophil-induced liver cell injury during hepatic ischemia-reperfusion and other acute inflammatory conditions. *Am J Physiol Gastrointest Liver Physiol.* 2006 Jun; 290(6):G1083–8. DOI: 10.1152/ajpgi.00568.2005 [PubMed: 16687579]
63. Jaeschke H, Ho YS, Fisher MA, Lawson JA, Farhood A. Glutathione peroxidase-deficient mice are more susceptible to neutrophil-mediated hepatic parenchymal cell injury during endotoxemia: importance of an intracellular oxidant stress. *Hepatology.* 1999 Feb; 29(2):443–50. DOI: 10.1002/hep.510290222 [PubMed: 9918921]
64. Hasegawa T, Ito Y, Wijeweera J, Liu J, Malle E, Farhood A, McCuskey RS, Jaeschke H. Reduced inflammatory response and increased microcirculatory disturbances during hepatic ischemia-reperfusion injury in steatotic livers of ob/ob mice. *Am J Physiol Gastrointest Liver Physiol.* 2007 May; 292(5):G1385–95. DOI: 10.1152/ajpgi.00246.2006 [PubMed: 17307725]
65. Gujral JS, Farhood A, Bajt ML, Jaeschke H. Neutrophils aggravate acute liver injury during obstructive cholestasis in bile duct-ligated mice. *Hepatology.* 2003 Aug; 38(2):355–63. DOI: 10.1053/jhep.2003.50341 [PubMed: 12883479]
66. Gujral JS, Hinson JA, Farhood A, Jaeschke H. NADPH oxidase-derived oxidant stress is critical for neutrophil cytotoxicity during endotoxemia. *Am J Physiol Gastrointest Liver Physiol.* 2004 Jul; 287(1):G243–52. DOI: 10.1152/ajpgi.00287.2003 [PubMed: 15044177]
67. Nieminen AL, Saylor AK, Tesfai SA, Herman B, Lemasters JJ. Contribution of the mitochondrial permeability transition to lethal injury after exposure of hepatocytes to t-butylhydroperoxide. *Biochem J.* 1995 Apr 1; 307(Pt 1):99–106. [PubMed: 7718000]
68. Nieminen AL, Byrne AM, Herman B, Lemasters JJ. Mitochondrial permeability transition in hepatocytes induced by t-BuOOH: NAD(P)H and reactive oxygen species. *Am J Physiol.* 1997 Apr; 272(4 Pt 1):C1286–94. DOI: 10.1152/ajpcell.1997.272.4.C1286 [PubMed: 9142854]
69. Laskin DL, Gardner CR, Price VF, Jollow DJ. Modulation of macrophage functioning abrogates the acute hepatotoxicity of acetaminophen. *Hepatology.* 1995 Apr; 21(4):1045–50. [PubMed: 7705777]
70. Michael SL, Pumford NR, Mayeux PR, Niesman MR, Hinson JA. Pretreatment of mice with macrophage inactivators decreases acetaminophen hepatotoxicity and the formation of reactive oxygen and nitrogen species. *Hepatology.* 1999 Jul; 30(1):186–95. DOI: 10.1002/hep.510300104 [PubMed: 10385655]
71. Bautista AP, Mészáros K, Bojta J, Spitzer JJ. Superoxide anion generation in the liver during the early stage of endotoxemia in rats. *J Leukoc Biol.* 1990 Aug; 48(2):123–8. [PubMed: 2164555]

72. Ju C, Reilly TP, Bourdi M, Radonovich MF, Brady JN, George JW, Pohl LR. Protective role of Kupffer cells in acetaminophen-induced hepatic injury in mice. *Chem Res Toxicol*. 2002 Dec; 15(12):1504–13. [PubMed: 12482232]
73. Lawson JA, Farhood A, Hopper RD, Bajt ML, Jaeschke H. The hepatic inflammatory response after acetaminophen overdose: role of neutrophils. *Toxicol Sci*. 2000 Apr; 54(2):509–16. [PubMed: 10774834]
74. Antoine DJ, Williams DP, Kipar A, Jenkins RE, Regan SL, Sathish JG, Kitteringham NR, Park BK. High-mobility group box-1 protein and keratin-18, circulating serum proteins informative of acetaminophen-induced necrosis and apoptosis in vivo. *Toxicol Sci*. 2009 Dec; 112(2):521–31. DOI: 10.1093/toxsci/kfp235 [PubMed: 19783637]
75. McGill MR, Sharpe MR, Williams CD, Taha M, Curry SC, Jaeschke H. The mechanism underlying acetaminophen-induced hepatotoxicity in humans and mice involves mitochondrial damage and nuclear DNA fragmentation. *J Clin Invest*. 2012 Apr; 122(4):1574–83. DOI: 10.1172/JCI59755 [PubMed: 22378043]
76. Liu ZX, Han D, Gunawan B, Kaplowitz N. Neutrophil depletion protects against murine acetaminophen hepatotoxicity. *Hepatology*. 2006 Jun; 43(6):1220–30. DOI: 10.1002/hep.21175 [PubMed: 16729305]
77. Marques PE, Amaral SS, Pires DA, Nogueira LL, Soriani FM, Lima BH, Lopes GA, Russo RC, Avila TV, Melgaço JG, Oliveira AG, Pinto MA, Lima CX, De Paula AM, Cara DC, Leite MF, Teixeira MM, Menezes GB. Chemokines and mitochondrial products activate neutrophils to amplify organ injury during mouse acute liver failure. *Hepatology*. 2012 Nov; 56(5):1971–82. DOI: 10.1002/hep.25801 [PubMed: 22532075]
78. Jaeschke H, Liu J. Neutrophil depletion protects against murine acetaminophen hepatotoxicity: another perspective. *Hepatology*. 2007 Jun; 45(6):1588–9. DOI: 10.1002/hep.21549 [PubMed: 17539019]
79. Williams CD, Bajt ML, Farhood A, Jaeschke H. Acetaminophen-induced hepatic neutrophil accumulation and inflammatory liver injury in CD18-deficient mice. *Liver Int*. 2010 Oct; 30(9):1280–92. DOI: 10.1111/j.1478-3231.2010.02284.x [PubMed: 20500806]
80. Williams CD, Bajt ML, Sharpe MR, McGill MR, Farhood A, Jaeschke H. Neutrophil activation during acetaminophen hepatotoxicity and repair in mice and humans. *Toxicol Appl Pharmacol*. 2014 Mar 1; 275(2):122–33. DOI: 10.1016/j.taap.2014.01.004 [PubMed: 24440789]
81. Cover C, Liu J, Farhood A, Malle E, Waalkes MP, Bajt ML, Jaeschke H. Pathophysiological role of the acute inflammatory response during acetaminophen hepatotoxicity. *Toxicol Appl Pharmacol*. 2006 Oct 1; 216(1):98–107. DOI: 10.1016/j.taap.2006.04.010 [PubMed: 16781746]
82. Dambach DM, Watson LM, Gray KR, Durham SK, Laskin DL. Role of CCR2 in macrophage migration into the liver during acetaminophen-induced hepatotoxicity in the mouse. *Hepatology*. 2002 May; 35(5):1093–103. DOI: 10.1053/jhep.2002.33162 [PubMed: 11981759]
83. Holt MP, Cheng L, Ju C. Identification and characterization of infiltrating macrophages in acetaminophen-induced liver injury. *J Leukoc Biol*. 2008 Dec; 84(6):1410–21. DOI: 10.1189/jlb.0308173 [PubMed: 18713872]
84. Arthur MJ, Kowalski-Saunders P, Wright R. *Corynebacterium parvum*-elicited hepatic macrophages demonstrate enhanced respiratory burst activity compared with resident Kupffer cells in the rat. *Gastroenterology*. 1986 Jul; 91(1):174–81. [PubMed: 3011580]
85. Antoniadis CG, Quaglia A, Taams LS, Mitry RR, Hussain M, Abeles R, Possamai LA, Bruce M, McPhail M, Starling C, Wagner B, Barnardo A, Pomplun S, Auzinger G, Bernal W, Heaton N, Vergani D, Thursz MR, Wendon J. Source and characterization of hepatic macrophages in acetaminophen-induced acute liver failure in humans. *Hepatology*. 2012 Aug; 56(2):735–46. DOI: 10.1002/hep.25657 [PubMed: 22334567]
86. Laskin DL, Sunil VR, Gardner CR, Laskin JD. Macrophages and tissue injury: agents of defense or destruction? *Annu Rev Pharmacol Toxicol*. 2011; 51:267–88. DOI: 10.1146/annurev.pharmtox.010909.105812 [PubMed: 20887196]
87. Hogaboam CM, Bone-Larson CL, Steinhäuser ML, Matsukawa A, Gosling J, Boring L, Charo IF, Simpson KJ, Lukacs NW, Kunkel SL. Exaggerated hepatic injury due to acetaminophen challenge in mice lacking C-C chemokine receptor 2. *Am J Pathol*. 2000 Apr; 156(4):1245–52. DOI: 10.1016/S0002-9440(10)64995-4 [PubMed: 10751350]

88. You Q, Holt M, Yin H, Li G, Hu CJ, Ju C. Role of hepatic resident and infiltrating macrophages in liver repair after acute injury. *Biochem Pharmacol.* 2013 Sep 15; 86(6):836–43. DOI: 10.1016/j.bcp.2013.07.006 [PubMed: 23876342]
89. Jaeschke H. Reactive oxygen and mechanisms of inflammatory liver injury: Present concepts. *J Gastroenterol Hepatol.* 2011 Jan; 26(Suppl 1):173–9. DOI: 10.1111/j.1440-1746.2010.06592.x [PubMed: 21199529]
90. Yoshioka H, Aoyagi Y, Fukuishi N, Gui MY, Jin YR, Li XW, Adachi Y, Ohno N, Takeya K, Hitotsuyanagi Y, Miura N, Nonogaki T. Suppressive effect of kamebakaurin on acetaminophen-induced hepatotoxicity by inhibiting lipid peroxidation and inflammatory response in mice. *Pharmacol Rep.* 2017 Oct; 69(5):903–907. DOI: 10.1016/j.pharep.2017.04.004 [PubMed: 28624597]
91. Jaeschke, H., Ramachandran, A. Antioxidant Defense Mechanisms. In: McQueen, CA., editor. *Comprehensive Toxicology.* 3. Vol. 2. Oxford: Elsevier Ltd; 2018. p. 277-295.
92. Werner C, Wendel A. Hepatic uptake and antihepatotoxic properties of vitamin E and liposomes in the mouse. *Chem Biol Interact.* 1990; 75(1):83–92. [PubMed: 2364459]
93. Knight TR, Fariss MW, Farhood A, Jaeschke H. Role of lipid peroxidation as a mechanism of liver injury after acetaminophen overdose in mice. *Toxicol Sci.* 2003 Nov; 76(1):229–36. DOI: 10.1093/toxsci/kfg220 [PubMed: 12944590]
94. Amodeo GF, Torregrosa MES, Pavlov EV. From ATP synthase dimers to C-ring conformational changes: unified model of the mitochondrial permeability transition pore. *Cell Death Dis.* 2017 Dec 4.8(12):1.doi: 10.1038/s41419-017-0042-3 [PubMed: 29233966]
95. Kon K, Kim JS, Jaeschke H, Lemasters JJ. Mitochondrial permeability transition in acetaminophen-induced necrosis and apoptosis of cultured mouse hepatocytes. *Hepatology.* 2004 Nov; 40(5):1170–9. DOI: 10.1002/hep.20437 [PubMed: 15486922]
96. Masubuchi Y, Suda C, Horie T. Involvement of mitochondrial permeability transition in acetaminophen-induced liver injury in mice. *J Hepatol.* 2005 Jan; 42(1):110–6. DOI: 10.1016/j.jhep.2004.09.015 [PubMed: 15629515]
97. Bernardi P, Krauskopf A, Basso E, Petronilli V, Blachly-Dyson E, Di Lisa F, Forte MA. The mitochondrial permeability transition from in vitro artifact to disease target. *FEBS J.* 2006 May; 273(10):2077–99. DOI: 10.1111/j.1742-4658.2006.05213.x [PubMed: 16649987]
98. Karch J, Kwong JQ, Burr AR, Sargent MA, Elrod JW, Peixoto PM, Martinez-Caballero S, Osinska H, Cheng EH, Robbins J, Kinnally KW, Molkentin JD. Bax and Bak function as the outer membrane component of the mitochondrial permeability pore in regulating necrotic cell death in mice. *Elife.* 2013 Aug 27.2:e00772.doi: 10.7554/eLife.00772 [PubMed: 23991283]
99. Karch J, Molkentin JD. Identifying the components of the elusive mitochondrial permeability transition pore. *Proc Natl Acad Sci U S A.* 2014 Jul 22; 111(29):10396–7. DOI: 10.1073/pnas.1410104111 [PubMed: 25002521]
100. Biasutto L, Azzolini M, Szabò I, Zoratti M. The mitochondrial permeability transition pore in AD 2016: An update. *Biochim Biophys Acta.* 2016 Oct; 1863(10):2515–30. DOI: 10.1016/j.bbamcr.2016.02.012 [PubMed: 26902508]
101. Ramachandran A, Lebofsky M, Baines CP, Lemasters JJ, Jaeschke H. Cyclophilin D deficiency protects against acetaminophen-induced oxidant stress and liver injury. *Free Radic Res.* 2011 Feb; 45(2):156–64. DOI: 10.3109/10715762.2010.520319 [PubMed: 20942566]
102. LoGuidice A, Boelsterli UA. Acetaminophen overdose-induced liver injury in mice is mediated by peroxynitrite independently of the cyclophilin D-regulated permeability transition. *Hepatology.* 2011 Sep 2; 54(3):969–78. DOI: 10.1002/hep.24464 [PubMed: 21626531]
103. Hu J, Ramshesh VK, McGill MR, Jaeschke H, Lemasters JJ. Low Dose Acetaminophen Induces Reversible Mitochondrial Dysfunction Associated with Transient c-Jun N-Terminal Kinase Activation in Mouse Liver. *Toxicol Sci.* 2016 Mar; 150(1):204–15. DOI: 10.1093/toxsci/kfv319 [PubMed: 26721299]
104. Rauen U, Petrat F, Sustmann R, de Groot H. Iron-induced mitochondrial permeability transition in cultured hepatocytes. *J Hepatol.* 2004 Apr; 40(4):607–15. DOI: 10.1016/j.jhep.2003.12.021 [PubMed: 15030976]

105. Schnellmann JG, Pumford NR, Kusewitt DF, Bucci TJ, Hinson JA. Deferoxamine delays the development of the hepatotoxicity of acetaminophen in mice. *Toxicol Lett.* 1999 May 20; 106(1): 79–88. DOI: 10.1016/S0378-4274(99)00021-1 [PubMed: 10378453]
106. Woolbright BL, Ramachandran A, McGill MR, Yan HM, Bajt ML, Sharpe MR, Lemasters JJ, Jaeschke H. Lysosomal instability and cathepsin B release during acetaminophen hepatotoxicity. *Basic Clin Pharmacol Toxicol.* 2012 Dec; 111(6):417–25. DOI: 10.1111/j.1742-7843.2012.00931.x [PubMed: 22900545]
107. Kon K, Kim JS, Uchiyama A, Jaeschke H, Lemasters JJ. Lysosomal iron mobilization and induction of the mitochondrial permeability transition in acetaminophen-induced toxicity to mouse hepatocytes. *Toxicol Sci.* 2010; 117(1):101–8. DOI: 10.1093/toxsci/kfq175 [PubMed: 20584761]
108. Hu J, Kholmukhamedov A, Lindsey CC, Beeson CC, Jaeschke H, Lemasters JJ. Translocation of iron from lysosomes to mitochondria during acetaminophen-induced hepatocellular injury: Protection by starch-desferal and minocycline. *Free Radic Biol Med.* 2016 Aug; 97:418–426. DOI: 10.1016/j.freeradbiomed.2016.06.024 [PubMed: 27345134]

Figure 1. Reactive oxygen species and APAP hepatotoxicity

APAP is metabolized through the cytochrome P450 system to generate the reactive metabolite NAPQI, in a step that is unlikely to produce significant oxidant stress. NAPQI forms mitochondrial protein adducts, which induces mitochondrial oxidant stress and peroxynitrite formation and activates the MAPK JNK, which translocates to mitochondria and amplifies reactive oxygen species production within mitochondria. This also causes lysosomal instability and release of free iron, which is taken up by mitochondria. Along with translocation of Bax to the mitochondrial outer membrane (OMM) this initiates the mitochondrial permeability transition (MPT) mediated by Bax on the OMM and components of the ATP synthase on the inner mitochondrial membrane (IMM). MPT activation subsequently causes cellular necrosis and release of DAMPs, which then result in neutrophil

infiltration. Neutrophils contain NADPH oxidase and myeloperoxidase, which could then generate reactive oxygen species outside the hepatocyte.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript