Published in final edited form as:

Free Radic Biol Med. 2018 May 01; 119: 45–55. doi:10.1016/j.freeradbiomed.2018.01.025.

Circadian redox rhythms in the regulation of neuronal excitability

Mia Y. Bothwell¹ and Martha U. Gillette^{1,2,3,4,5,†}

¹Department of Molecular and Integrative Physiology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

²Department of Cell and Developmental Biology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

³Neuroscience Program, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

⁴Department of Bioengineering, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

⁵Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

Abstract

Oxidation-reduction reactions are essential to life as the core mechanisms of energy transfer. A large body of evidence in recent years presents an extensive and complex network of interactions between the circadian and cellular redox systems. Recent advances show that cellular redox state undergoes a ~24-h (circadian) oscillation in most tissues and is conserved across the domains of life. In nucleated cells, the metabolic oscillation is dependent upon the circadian transcription-translation machinery and, vice versa, redox-active proteins and cofactors feed back into the molecular oscillator. In the suprachiasmatic nucleus (SCN), a hypothalamic region of the brain specialized for circadian timekeeping, redox oscillation was found to modulate neuronal membrane excitability. The SCN redox environment is relatively reduced in daytime when neuronal activity is highest and relatively oxidized in nighttime when activity is at its lowest. There is evidence that the redox environment directly modulates SCN K⁺ channels, tightly coupling metabolic rhythms to neuronal activity. Application of reducing or oxidizing agents produces rapid changes in membrane excitability in a time-of-day-dependent manner. We propose that this reciprocal interaction may not be unique to the SCN. In this review, we consider the evidence for circadian redox oscillation and its interdependencies with established circadian timekeeping mechanisms. Furthermore, we will investigate the effects of redox on ion-channel gating dynamics and membrane excitability. The susceptibility of myriad different ion channels to

[†]To whom correspondence should be addressed: Telephone: 217-244-1355; Fax: 217-244-1648; mgillett@illinois.edu.

Author Disclosure Statement

No competing financial interests exist.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

modulation by changes in the redox environment suggests that circadian redox rhythms may play a role in the regulation of all excitable cells.

Graphical Abstract

Keywords

brain; cellular redox; circadian rhythms; hypothalamus; ion channels; K^+ channels; neuronal excitability; neurophysiology; redox; suprachiasmatic nucleus

1. Introduction

The suprachiasmatic nucleus (SCN) of the hypothalamus is the master regulator of the circadian rhythms in mammals. It generates the daily rhythms of behavior, metabolism, and other important physiological processes. The main driver of the circadian clock is a transcription-translation feedback loop of core circadian genes. However, emerging evidence suggest that metabolic oscillators also play a crucial role in the generation of circadian rhythms. Circadian rhythms in cycles of oxidation and reduction have been reported in a broad array of mammalian tissues and cell types and are conserved across the domains of life [1]. The discovery of a near-24-h oscillation of redox state in the SCN also revealed that cellular metabolic state could modulate neuronal excitability, an integral component of SCN timekeeping, via modification of redox-sensitive K^+ channels [2]. These studies demonstrate that redox homeostasis is dynamic, displays circadian characteristics, and may play a role in the regulation of daily rhythms of electrically excitable cells.

Since the pioneering work of Hodgkin and Huxley [3], the scientific community has developed deep insights into neuronal membrane dynamics. Neuronal excitability is linked directly to ion channel activity. A change in permeability of ions across the plasma membrane can lead to significant changes in resting membrane potential. The electrical properties of neurons and other excitable cells rely on many different types of voltage-gated, ligand-gated, and leak channels that are permeable to ions such as Na^+ , K^+ , Cl^- , and Ca^{2+} .

Membrane potential (V_m) is determined by the differential distribution of these ions. Ion channels are regulated by a vast number of ligands, post-translational modifications, and other mechanisms. They are susceptible to modulation by phosphorylation, second messengers, gaseous signaling molecules such as carbon monoxide (CO) and nitric oxide (NO), and by the redox environment [2, 4–7].

Cellular redox encompasses the dynamic regulation of reactive oxygen species (ROS), antioxidants, and redox-sensitive metabolic cofactors. ROS have been viewed historically as toxic. There are many studies regarding their detrimental effects to the body and contributions to disease and aging. However, there is also increasing evidence that ROS signaling is integral to a multitude of normal cellular processes and signaling pathways [8–10]. ROS generation and redox homeostasis are no longer only relevant as sources of oxidative stress. Mounting evidence in the past few years shows circadian rhythms in redox state are an intrinsic and dynamic feature of all cell types and may contribute to daily regulatory processes.

Intrinsic circadian oscillations of signaling molecules exist in the SCN and hippocampus, opening windows of excitability and susceptibility. Cyclic changes that gate activity have been termed “iterative metaplasticity” to describe states permissive for plasticity mechanisms that are expressed as daily cycles [11]. In nucleated cells, the molecular circadian clock is reciprocally connected to the redox system. These interactions produce daily rhythms in redox state which can then modulate neuronal activity via regulation of ion channels [2]. Thus, day/night differences in redox state may play a role in the generation of daily changes in brain states that underlie the potential to establish long-lasting changes in brain function that we know as memory. Indeed, changes in the redox state have been found to modulate cognitive decline [12,13].

In this review, we examine fundamental features of circadian rhythms, the role of the SCN as master circadian clock, and explore the reciprocal connections between circadian timekeeping and the cellular redox oscillation, including the redox modulation of neuronal excitability. Then, we consider the many reports of redox modification of ion channel activity. We propose that circadian regulation of neuronal excitability via redox-sensitive ion channels may not be unique to the SCN.

2. Circadian rhythms

The 24-h cycle of day and night generated by the Earth’s rotation has accompanied and driven the evolution of most organisms. As a result, myriad life forms, from some prokaryotes to all eukaryotes, have developed intrinsic daily rhythms in cellular processes, behavior, and metabolism. The self-sustained circadian oscillation provides an evolutionary advantage as it allows organisms to coordinate their internal states and anticipate changes in the timing and duration between night and day so that cellular, physiological, and behavioral events occur at appropriate times. Misalignment of the internal clock with the external environment can disrupt these functions and lead to disease [14–17].

3. The mammalian suprachiasmatic nucleus as master clock

The mammalian circadian system is organized hierarchically into a master oscillator and secondary oscillators in the brain and body. The suprachiasmatic nucleus (SCN) of the hypothalamus is the master circadian clock in mammals that synchronizes peripheral clocks in other brain regions and organ systems [18–20]. The SCN is a pair of small nuclei on either side of the third ventricle, directly above the optic chiasm in the anterior hypothalamus (Figure 1) [21]. Each nucleus is composed of approximately 10,000 tightly compacted cells whose collective activity is considered to be the central “pacemaker” of circadian rhythm [22]. Destruction of the SCN results in the loss of daily rhythms in sleep-wake cycle, body temperature, locomotor activity, drinking, and endocrine release [23,24].

3.1. Oscillation of clock genes and proteins in the SCN

Circadian rhythms are generated by a transcription-translation feedback loop of clock genes and proteins that form an oscillatory molecular clock [25]. This core molecular clock consists of a heterodimeric complex of protein products of the genes *CLOCK* and *BMAL1*, which positively regulate the expression of *Period* (*Per1*, *2*, and *3*) and *Cryptochrome* (*Cry1* and *2*) genes. The accumulated protein products form their own heterodimeric transcriptional repressor complex of PER and CRY that represses the activity of *CLOCK* and *BMAL1*. The transcriptional-translational loop of these core clock genes repeats with a period of approximately 24 h and is the basis for the mammalian molecular clock.

An additional interlocking feedback loop involves the *BMAL1/CLOCK*-mediated transcription of nuclear receptor genes, *Rev-Erba/β* and *RAR-related orphan receptor alpha* (*Rora*). To complete the circadian loop, REV-ERB and ROR proteins then compete for binding sites within the promoters of *Bmal1* and *CLOCK*, where REV-ERB inhibits transcription and ROR initiates transcription. Additionally, casein kinase 1 (CK1)-mediated phosphorylation contributes to timekeeping through the destabilization of PER proteins [26]. Post-translational modifiers like protein kinases and small molecule messengers such as cAMP and Ca^{2+} play important roles to determine and modify the intrinsic circadian rhythm [27–32]. Together, these modifiers work to synchronize firing rate, gene expression, and secretion across the SCN.

3.2. Oscillation of neuronal activity in the SCN

Circadian rhythms in SCN electrical activity were first revealed via *in vivo* recordings by Inouye and Kawamura in 1979 [33]. The activity of the SCN neuronal population oscillates: it is nearly electrically silent at night but exhibits significantly higher frequency during the day, peaking at midday [2,34,35]. This rhythm exists endogenously and will persist for days in an SCN brain slice without input from other brain regions or the external environment [34,36,37]. Interestingly, this high level of spontaneous action potentials occurs during the day regardless of whether the species is nocturnal or diurnal [38]. The SCN is able to generate these neuronal activity rhythms *in vivo*, in slice, and in dispersed cell culture [36,39,40].

Circadian oscillation in neuronal electrical activity is generated at the level of individual neurons. The spontaneous firing of SCN neurons emerges from modulation of a combination of intrinsic currents. These currents can be divided into those that provide excitatory drive and depolarization during the day, those that maintain a regular pattern of action potentials, and those that produce the hyperpolarization that renders SCN neurons silent at night [38]. Underlying activity rhythms are day/night changes in resting membrane potential. Studies have shown the resting membrane potential of neurons in the SCN is most depolarized during midday and most hyperpolarized in the early night [2]. The daytime depolarization of SCN neurons is accompanied by a decrease in overall K^+ conductance while a larger K^+ conductance at night results in hyperpolarization [41,42]. The differences in membrane potential and ionic conductances during the day and night persist even in the absence of synaptic activity and are likely clock-driven [43].

Whereas neuronal activity in the SCN could be interpreted as an output of the molecular clock that mediates communication with other brain regions, research from the past 20 years has established that neuronal activity is an important component of rhythmic gene expression. A decade ago, a landmark study showed that electrically silencing *Drosophila melanogaster* pacemaker neurons stops the free-running rhythms of core molecular clock components PERIOD and TIMELESS, as well as behavioral activity rhythms [44]. Furthermore, a recent study showed that optogenetic activation and suppression of SCN neuronal firing in mice can reset the phase and alter the period of the molecular clock, respectively [45]. These conceptual advances establish that the molecular rhythm and neuronal activity rhythm are, in fact, linked as core clock mechanisms.

4. Extra-SCN clocks

Rhythms of core clock genes are found in cells and tissues throughout the body and in regions of the brain outside of the SCN. These molecular rhythms persist when cells or tissues are maintained in culture in the absence of the SCN. Because non-SCN cells also contain these endogenous molecular oscillators, the molecular circadian clock is now recognized as a fundamental component of all cells [46,47]. However, the SCN alone behaves as the master clock. A key difference between the SCN and peripheral oscillators with respect to emergent behavior is in their network properties. SCN explants are capable of sustaining synchrony via synaptic and neuropeptidergic coupling mechanisms. While isolated cells and cells within cultured peripheral tissue can oscillate independently of one another [48], they are coupled within tissues in the body [47].

In complex organisms, the circadian system is comprised of multiple oscillators. Peripheral oscillators are not directly entrained by light, so they rely on the SCN for synchronization within tissues and to the external environment. Circadian oscillations have been identified within various structures of the central nervous system. Robust rhythms in core clock gene expression and electrical activity have been observed in extra-SCN hypothalamic nuclei, the olfactory bulb, amygdala, cerebellum, cerebral cortex, and hippocampus [49–51]. These areas differ in phase, ability to sustain rhythmicity, and their relationship to the SCN. The SCN entrains other oscillators partly through efferent projections, but primarily through release of diffusible factors such as peptides and hormones [52,53]. Circadian oscillations in

the liver and hippocampus can be entrained by external cues other than light. Restricted feeding is a salient signal for entraining the circadian clocks of these peripheral oscillators, whereas the SCN is relatively unaffected [54].

5. Circadian rhythms and metabolism

A large body of evidence has demonstrated circadian rhythmicity in metabolism. Plasma concentrations of glucose and insulin show a day/night pattern in mammals and the SCN is required for this rhythmicity [55]. In addition, glucose utilization in the SCN is rhythmic, with the highest level of uptake in the subjective day when SCN neuronal activity peaks [56]. Furthermore, circadian rhythms and metabolism are inextricably linked. Food acts as a powerful *zeitgeber* (time-giver, German) and the circadian clock contributes to metabolic homeostasis by shifting physiological processes to support changes in activity [57]. Animal models that lack a functional circadian clock show not only a lack of rhythmicity, but also abnormal metabolic phenotypes [58].

Transcriptomic studies reveal that nearly half of the mouse genome oscillates with a near-24-h rhythm in an organ-specific manner [59]. Strikingly, many of the genes that show circadian oscillation are involved in metabolic functions [60,61]. Metabolomic screens of mice and humans have found that large amounts of metabolites are rhythmic even when conducted in constant environmental conditions. Furthermore, nuclear hormone receptors that participate in fatty acid and glucose metabolism as well as metabolic hormones, like glucagon and insulin, display circadian oscillation [62,63]. Recently, work from Feeney *et al.* demonstrated that in eukaryotes, the level of intracellular Mg^{2+} displays circadian rhythmicity and can feed back to regulate clock periodicity [64]. What makes this finding exciting is that Mg^{2+} is a necessary cofactor for ATP to be biologically active, thus giving circadian significance to global cellular energetics and vice versa. These combined data strongly suggest that metabolism is influenced by circadian rhythm and there is increasing evidence that these metabolic fluctuations can feed back to regulate elements of the clock.

6. Cellular redox signaling

Energy metabolism is linked to redox homeostasis. On a cellular level, metabolic state is represented by the redox state. Redox is the potential to donate or receive electrons for biochemical processes and redox state can be defined by the balance of oxidizing and reducing potential. This can be evaluated biochemically by the ratio of small molecule cellular redox pairs, such as glutathione disulfide (GSSG)/glutathione (GSH), nicotinamide adenine dinucleotide (NAD^+ /NADH), nicotinamide adenine dinucleotide phosphate ($NADP^+$ /NADPH) or dehydroascorbic acid (DHA)/ascorbic acid (AA) (Figure 2). These redox molecule pairs are sensitive to electron flow and to fluctuations in the reducing/oxidizing potential of the cellular environment.

Cells possess multiple regulators of redox state. GSH is the most abundant free thiol in cells and is primarily responsible for maintaining a physiological intracellular redox environment. Reduced GSH is oxidized to GSSG, and the GSH/GSSG ratio is frequently used to assess overall redox state [65]. The thioredoxin (Trx) system is another ubiquitous thiol-reducing

complex. Like GSH, thioredoxins serve as electron donors to reversibly reduce intracellular protein disulfides and maintain redox homeostasis [66]. Peroxiredoxins (Prxs) are a family of peroxidases that scavenge and reduce hydrogen peroxide (H_2O_2) as regulators of the cellular redox environment. Mammals express six different Prx isoforms (Prx1–Prx6) with distinct subcellular distributions in different brain regions and cell types. Upon encountering H_2O_2 , Prxs go through several levels of oxidation.

The reduced thiol ($-\text{SH}$) reacts to form a sulphenic acid ($-\text{SOH}$), then sulphinic acid ($-\text{SO}_2\text{H}$), and a fully oxidized sulphonic acid ($-\text{SO}_3\text{H}$) [67].

Relative to its size, the brain consumes a disproportionately large amount of O_2 and is highly sensitive to oxidative stress and damage. Reactive oxygen species (ROS), such as H_2O_2 , superoxide anion (O_2^-), and hydroxyl radical ($\text{HO}^\bullet$), are frequently implicated in oxidative damage and pathogenesis of disease. O_2^- is generated primarily as a byproduct of mitochondrial oxidative phosphorylation and also by NADPH oxidase (NOX) enzyme complexes [68]. Only moderately reactive, O_2^- is rapidly converted to H_2O_2 by superoxide dismutases. H_2O_2 is more inert and capable of diffusing through cellular membranes [69]. In low levels, H_2O_2 can act as a local signaling molecule and, in high levels, can have cytotoxic effects [70]. H_2O_2 is catalytically broken down to H_2O by scavenging enzymes such as catalase (Prxs), or glutathione peroxidases (GPx), which keep ROS levels contained. More reactive than either O_2^- or H_2O_2 is $\text{HO}^\bullet$, which can be generated from H_2O_2 in the presence of metal ions via the Fenton reaction. A careful overall balance in production and neutralization of oxidants is maintained using a variety of mechanisms. When ROS levels rise significantly and overwhelm the antioxidant defense system, oxidative stress can occur and cause damage to cellular structures [71]. However, small changes in ROS are integral to many biochemical processes. Accumulating evidence suggests that ROS also play important roles in intracellular signal transduction and many physiological processes [72–74].

Protein amino acid residues are major targets of cellular oxidants. Many of the effects of signaling via ROS are through the reversible modification of exposed cysteine and methionine residues of redox-sensitive proteins. Cysteines and methionines have reactive sulfur-containing side chains that make them more susceptible to redox modifications than other amino acids. The thiol ($-\text{SH}$) functional groups of cysteine residues make them especially susceptible to oxidation. Upon exposure to H_2O_2 , the redox-sensitive thiol can be reversibly oxidized to reactive sulphenic acid ($-\text{SOH}$) or further oxidized to sulfinic acid ($-\text{SO}_2\text{H}$) and irreversibly to sulfonic acid ($-\text{SO}_3\text{H}$) [75]. Non-enzymatic and reversible disulfide bond formation with a nearby cysteine is the most common product of $-\text{SOH}$ cysteine oxidation [76].

The redox state of sulfur-containing amino acids is tightly regulated by GSH. GSH is a major source of cellular cysteine. Its linkage to redox-sensitive cysteine residues within proteins is termed S-glutathionylation. This transient incorporation of GSH into cellular protein via reactive thiols is a metabolically driven form of post-translational modification [77]. Redox-sensitive methionine residues do not form disulfide bonds and, instead, are oxidized to methionine sulfoxide (MetO) by a variety of different ROS [78]. The reduction of oxidized methionine is catalyzed by the enzyme methionine sulfoxide reductase (MsrA)

[79]. The redox modification of these sulfur-containing amino acids can result in functional changes to protein structure and/or activity.

7. Redox-molecular clockwork interactions

There is compelling evidence that redox state may play a regulatory role in the circadian clockwork. Reduced forms of the redox cofactors NAD(H) and NADP(H) have been shown to enhance DNA-binding activity of the core clock proteins BMAL1 and CLOCK, while their oxidized forms inhibit it. Even minute changes in cellular redox state can affect binding activity of these circadian transcriptional activators [80]. The BMAL1/CLOCK heterodimer regulates the expression of nicotinamide phosphoribosyltransferase (NAMPT), a rate-limiting enzyme in the NAD⁺ salvage pathway [81,82]. This relationship is the driving force for rhythmic levels of NAD⁺, which in turn activate NAD⁺-dependent histone deacetylases, sirtuin 1 (SIRT1) and sirtuin 3 (SIRT3) [82,83]. SIRT1, an important element of metabolic control, displays circadian oscillatory activity, and alters PER2 stabilization and CLOCK function [84–86]. SIRT1 is localized in the mitochondrial matrix where it mediates the deacetylation of metabolic enzymes (Figure 3).

Heme, a protein cofactor that acts as a sensor of cellular redox state, is another component in the metabolism-circadian relationship. Heme reversibly binds to the circadian nuclear receptors REV-ERB α and REV-ERB β in a redox state-dependent manner and modulates their activity, providing a link between energy sensing and an arm of the molecular clock [87–89]. Heme also binds to neuronal PAS domain protein 2 (NPAS2), a CLOCK paralog that also binds to BMAL, and modulates its DNA-binding activity [90]. Another link between metabolism and circadian systems comes from the clock's sensitivity to the AMP/ATP ratio through AMP-activated protein kinase (AMPK)-mediated phosphorylation of CRY and casein kinase I (CK1) [91].

Phosphorylation of CRY results in its degradation, and phosphorylation of CK1 ϵ increases its activity and accelerates the degradation of PER2. Most recently, Hirano et al. showed that in mammalian cells, CRY proteins are stabilized by the redox cofactor flavin adenine dinucleotide (FAD) [92].

The first evidence of a molecular clock-independent circadian oscillation came in the form of redox rhythms [93]. Both sulphinic and sulphonic forms of peroxiredoxin are considered “hyper-oxidized.” A circadian oscillation of hyper-oxidized Prx proteins has been observed in red blood cells. The presence of this rhythm in cells that lack a nucleus established that circadian metabolic cycles can exist independently of the core molecular clock based on transcription-translation. However, in embryonic fibroblastic cells from *mCry1/2* $-/-$ mice, hyper-oxidized Prx rhythms are altered [93]. Therefore, in nucleated cells, there is interplay between a non-translational redox-based oscillator and the clock gene-dependent oscillator. This redox cycle of Prxs is conserved broadly across domains of life, from prokaryotes to the rodent SCN [1]. These findings raise the possibility that rhythms in metabolism may be an ancient timekeeping oscillator [94].

Recent studies have identified the pentose phosphate pathway (PPP) as a regulator of circadian redox rhythms. Glucose utilization through the PPP is essential for replenishing the cytoplasmic NADPH content as well as for the generation of nucleotide and amino acid precursors. NADPH availability is required for the regeneration of cellular antioxidants, such as GSH, and prevention of oxidative stress. By pharmacological and genetic inhibition of the PPP, Rey and colleagues demonstrated that NADPH availability is able to alter both the phase and period of the molecular clock in human osteosarcoma cells [95]. A similar study found that PPP inhibition affects only clock gene expression amplitude and phase with no changes in periodicity [96]. These inconsistencies may be explained by the differences in inhibitors and cell types employed. Taken together, these data suggest that metabolic and redox mechanisms can modulate the core circadian clock.

This is by no means an exhaustive list of known circadian-metabolism interactions. However, these findings provide strong support for metabolic regulation of the core molecular clock and vice versa. In mammalian systems, redox rhythms work in concert with molecular circadian timekeeping and have both input and output roles.

8. Circadian redox rhythms and SCN neuronal excitability

Corresponding circadian oscillations of redox state and neuronal excitability were observed in the SCN of rats and mice [2]. Through measurement of FAD/NAD(P)H, glutathiolation, and DHA/AA ratios, Wang et al. reported that the SCN redox environment is at its most reduced in the mid-subjective day and most oxidized in the early subjective night. Strikingly, midday corresponds to the peak in spontaneous action potentials and most depolarized state of membrane potential (V_m), assessed by patch clamp recording of SCN neurons. Early night corresponds to the lowest levels of spontaneous action potentials and V_m of SCN neurons is most hyperpolarized. The circadian redox rhythm is not present in arrhythmic *Bmal1*^{-/-} mice, suggesting that it is dependent upon a functional circadian molecular clock [2].

Imposed changes in redox state cause immediate changes in excitability [32]. Thus, unlike the transcription-translational relationship of changes in the molecular clock to SCN physiology, redox modulates neuronal excitability by tight coupling. A parallel relationship between redox state and membrane potential of SCN neurons suggests metabolic state has a modulatory role in membrane excitability. Patch-clamp electrophysiology showed that redox state regulates SCN neuronal excitability via post-translational modulation of K^+ channels. The relatively reduced daytime environment decreases hyperpolarizing K^+ currents, which results in membrane depolarization and increased neuronal activity. Conversely, the oxidized nighttime environment potentiates K^+ currents, resulting in membrane hyperpolarization and reduced neuronal activity. Application of pharmacological blockers suggests that both leak and A-type K^+ channels in SCN neurons are sensitive to redox environment and mediate changes in membrane excitability (Figure 4) [2]. These findings established a new connection between redox state and neuronal excitability in the SCN: metabolic state can be a modulator, rather than only the result, of neuronal activity.

Extending beyond the central nervous system (CNS), endogenous redox oscillations can have modulatory effects on other excitable cells in the body. For instance, there is considerable research on how the heart responds to ROS. The redox environment can affect cardiac excitability via modulation of Ca^{2+} , Na^+ , and K^+ ion channels [97]. Furthermore, a growing number of studies are focused on the relationship between redox and pain pathways, specifically, the effects of redox modulation of ion channels and excitability in peripheral sensory neurons [98–100].

9. Redox modulation of K^+ channel activity

Ion channels permeable to K^+ are components of the plasma membrane of most mammalian cells. K^+ channels play central roles in the regulation of neuronal excitability via the resting membrane potential, the repolarization phase of action potentials, and firing frequency [7]. Alterations in their activity can lead to changes in many aspects of neuronal activity. Four major classes of K^+ channels can be identified when grouped by common characteristics and function: voltage-gated, Ca^{2+} -activated, inward-rectifying, and leak K^+ channels. In this section, we will discuss the redox sensitivity of voltage-gated, Ca^{2+} -activated, and inward-rectifying K^+ channels, which have been more thoroughly studied.

9.1. Voltage-gated K^+ channels

Voltage-gated potassium (Kv) channels form a large and evolutionarily conserved family of ion channels that play essential roles in the generation and propagation of electrical impulses in excitable cells. Each Kv channel contains four α -subunits that form the pore structure and may also contain auxiliary β -subunits that modify channel activity [102]. Kv channels can be physiologically divided based on inactivation properties into fast-inactivation A-type K^+ channels and delayed-rectifier K^+ channels. Rapid inactivation occurs via the “ball and chain” mechanism where the N-terminus of an α - or β -subunit blocks the pore and inhibits ion flow [103]. This is also sometimes referred to as “N-type inactivation.”

Several rapidly inactivating (A-type) Kv channels have been found to be sensitive to the redox environment. Ruppertsberg et al. showed that A-type K^+ channels cloned from mammalian brain can be modulated by intracellular GSH. In an oxidized state, reversible formation of a disulfide bond forms in the inactivating gate, preventing it from closing/inactivation. Application of reduced GSH rapidly restored fast-inactivation of the channel [104]. This modulation of inactivation properties may lead to changes in hyperpolarization or depolarization of the membrane. Similar redox modulation of β -subunit cysteine residues can determine inactivation states [103,105]. Additionally, subsequent studies have shown that the oxidation of methionine in Kv channels rapidly reverses the inactivated state of the channel [106].

Several types of delayed-rectifier K^+ channels are expressed in the CNS with distinct channels localized to specific areas [107]. In freshly dissociated CA1 neurons from the rodent hippocampus, application of H_2O_2 inhibits delayed rectifier K^+ currents. The effect is completely abolished by the addition of membrane-permeable thiol-reducing agents, indicating the involvement of sulfhydryl groups [108]. Furthermore, oxidation by H_2O_2 increases M-type K^+ current in superior cervical ganglion (SCG) neurons and, consequently,

decreases neuronal resting membrane potential [109]. These effects result in reduced neuronal excitability and are reversed by the reducing action of dithiothreitol (DTT). The mammalian Shaker family channels (Kv1) can undergo inactivation from their associated β subunits (Kv β). Studies have shown that the Kv β subunits are oxidoreductase enzymes that bind and utilize NADPH, coupling channel activity with cellular redox [105]. Oxidation of Kv β -bound NADPH limits N-type inactivation by obstructing the N-terminus from blocking the channel, leading to a potentiated K⁺ current (Figure 4) [101].

9.2. Ca²⁺-activated K⁺ channels

The large conductance Ca²⁺- and voltage-activated K⁺ (BK) channels contribute to a variety of functions, including the regulation of membrane excitability and intracellular Ca²⁺ concentration. BK channels are activated during action potentials by membrane depolarization and Ca²⁺ entry through voltage-sensitive Ca²⁺ channels. Fast BK activation leads to hyperpolarization and closure of Ca²⁺ channels [110]. Under physiological conditions, intracellular Mg²⁺ also is capable of enhancing BK channel activity with low affinity independent of the high affinity Ca²⁺ binding site [111–113]. BK channels are essential for circadian pacemaker neuronal firing in the SCN with peak activity that suppresses spontaneous firing at night [114]. Daily expression of SCN BK channels shows circadian rhythmicity in the absence of light and is dependent on the intrinsic molecular clock [115–117]. Although BK channel expression is decreased during the day and BK activity is sensitive to the intracellular Ca²⁺ concentration, N-type inactivation of BK channels also contributes to the decrease in daytime K⁺ current that modulates SCN firing rhythms [114].

Similar to voltage-gated K⁺ channels, BK channels contain a tetrameric pore formed of α subunits that may be associated with auxiliary β subunits that confer various functional properties to the channel [118]. BK channels have been shown to be directly modulated by redox state with somewhat inconsistent results. Direct application of reducing and oxidizing agents modulates activity of human Ca²⁺-activated K⁺ (BK) channels expressed in *Xenopus* oocytes and human embryonic kidney cells. The thiol-reducing agent DTT enhances and stabilizes channel activity while the oxidizing agent H₂O₂ decreases the channel open probability [119].

In dissociated adult rat hippocampal pyramidal neurons, application of GSSG, oxidized glutathione, increases channel open probability and thus activity, while application of GSH, reduced glutathione, reverses the effects [120]. Another study using cultured neonatal rat hippocampal neurons reported the opposite results: GSSG decreases channel activity and GSH activates the channels [121]. These conflicting results may be explained by the heterogeneity of ion channels due to the variety of accessory subunits. BK channels have four known β subunits and each modulates channel function in different ways [122]. Additionally, BK channels have both cysteine and methionine residues that are redox sensitive; oxidation of the different amino acids appears to have opposite effects on channel activity [123–125].

9.3 Inward rectifying K⁺ channels

Unlike most K⁺ channels, inwardly rectifying K⁺ (K_{ir}) channels are characterized by greater inward flow rather than outward flow. They are an important class of K⁺ channels that contribute to regulation of membrane potential and activity in excitable cells. Structurally, K_{ir} channels are tetramers with subunits of two transmembrane domains and cytoplasmic N- and C-terminal domains linked by a conserved pore region [126–128]. Under physiological conditions, these channels display greater conductance when the membrane potential more negative than the reversal potential of K⁺ and permit less flow when membrane potential is more positive [129–131]. Inward flow of ions through K_{ir} channels is due to interactions with intracellular Mg²⁺ and polyamines, which physically block the cytoplasmic region preventing ion efflux [132–134].

High-resolution kinetic analyses demonstrate that K_{ir} channels are susceptible to redox regulation. Ruppersberg and Fakler first observed that presence of the reducing agent DTT in the buffer solution altered the kinetics of K_{ir}2.1 channels expressed in *Xenopus* oocytes [135]. Furthermore, application of DTT reversibly increased channel activity of G protein-coupled inwardly rectifying K⁺ (GIRK) channels K_{ir}3.1/3.4 [136]. This modulatory activity depends upon a specific cysteine residue in the N-terminus because mutation of this cytoplasmic residue abolishes redox-mediated effects. Beyond redox modifications, the dependence of K_{ir} and BK channel activity on cytoplasmic Mg²⁺ availability raises the interesting question of whether circadian oscillation in intracellular Mg²⁺ concentration can also exert rhythmic modulatory effects on ion-channel kinetics. Thus, there are multiple potential avenues for circadian metabolic regulation of neuronal excitability.

10. Redox modulation of Ca²⁺ channel activity

Calcium has the unique role of contributing to the membrane electrical potential as well as acting as a second messenger in the cytoplasm. Intracellular Ca²⁺ is an important regulator of neurotransmitter release, ion channel activation, gene regulation, and action potential generation [137]. Elevated cytosolic Ca²⁺ in neurons comes from the endoplasmic reticulum (ER) or the extracellular space.

10.1. Voltage-gated Ca²⁺ channels

Ca²⁺ influx from the extracellular space is achieved primarily through several voltage-gated Ca²⁺ channels and Ca²⁺-permeable cation channels. Voltage-dependent Ca²⁺ channels can be classified based on their sensitivity to activation into high-voltage-activated (HVA) and low-voltage-activated (LVA) channels. LVA channels are also known as “transient” T-type channels. HVA channels are further subdivided into L-, N-, P, Q-, and R-type channels. Both L-type and T-type Ca²⁺ currents contribute to the excitatory drive of SCN neurons in the morning, and the L-type current amplitude shows a diurnal rhythm [38].

T-type Ca²⁺ channels have been widely implicated in redox sensitivity of nociceptive neurons [138]. A study using dissociated primary sensory dorsal root ganglion (DRG) neurons demonstrated that T-type currents are significantly enhanced in the presence of a sulfhydryl-reducing agent and inhibited in the presence of a sulfhydryl-oxidizing agent,

suggesting that T-type Ca^{2+} channels are redox sensitive [139]. Interestingly, the same study by Todorovic et al. showed that while redox reagents affected the T current in DRG cells, none had an effect on the HVA current, even at high concentrations. Furthermore, the presence of reducing agents enhances neuronal excitability of peripheral nociceptors through T-type channel modulation [98,99]. Studies show that the $\text{Ca}_v3.2$ isoform of T-type Ca^{2+} channels is the target of redox modulation [139–141]. Consistent with these results, $\text{Ca}_v3.2$ T-type currents in the thalamus are enhanced by thiol-reducing agents L-cysteine and DTT, while the oxidizing agent 5,5-dithio-bis-(2-nitrobenzoic acid) (DTNB) inhibits the currents [142].

Many studies have demonstrated that L-type Ca^{2+} currents are affected by reducing and oxidizing agents. However, most of these were conducted in non-CNS cells. Thiol-specific oxidizing and reducing agents have been shown to modulate L-type Ca^{2+} channels in cultured cardiomyocytes and smooth muscle cells. Oxidizing agents increased the basal L-type current and reducing agents decreased the current in some cells whereas oxidizing agents inhibited the current in other cells [131,143,144]. There is evidence that changes to current activity in the presence of H_2O_2 and thiol-oxidizing agents is due to S-glutathionylation of the L-type Ca^{2+} channel [145,146]. These conflicting effects of oxidation may reflect differences in subunit composition, cell type, and membrane voltage.

10.2. Intracellular Ca^{2+} channels

Intracellular Ca^{2+} stores are released from the endoplasmic reticulum (ER) by the inositol triphosphate (IP_3) receptor in IP_3 -mediated release and Ca^{2+} -induced Ca^{2+} release via the ryanodine receptor (RyR) on the ER. Ample evidence demonstrates that the activity of RyR is sensitive to the cellular redox environment. RyRs are Ca^{2+} channels embedded in the membrane of the endoplasmic reticulum and are responsible for release of Ca^{2+} from intracellular stores. RyR are susceptible to many posttranslational modifications, with redox modification being one of them. The sulfhydryl groups of RyR cysteine residues are subject to reversible S-nitrosylation and S-glutathionylation, which can affect channel structure and activity. Additionally, oxidizing agents increase RyR opening probability while reducing agents have the opposite effect [147–149]. The RyR is most active under oxidizing conditions, making redox state a potential regulator of intracellular Ca^{2+} signaling. It is noteworthy that the activity of the RyR in the SCN undergoes tight circadian regulation. RyR are sensitive to activation in early night, when the oxidation state of the SCN is maximal [2,150].

11. Redox modulation of Na^+ channel activity

Voltage-dependent sodium channels underlie the generation of action potentials in many excitable cell types. The fast-inactivating Na^+ current initiates action potentials, however, the Na^+ current also has a voltage-dependent non-inactivating component in some cells [151]. This persistent Na^+ current is a key component of the excitatory drive for SCN neurons to be spontaneously active [38]. Like K^+ and Ca^{2+} channels, Na^+ channels are susceptible to redox modulation. Wang and colleagues observed that application of Chloramine-T (ChT), an oxidant that targets both cysteine and methionine residues, inhibits channel inactivation in

toad myelinated nerve fibers and squid giant axons [152,153]. Further investigation revealed that it is the oxidation of methionine residues by ChT that attenuates fast inactivation of some Na_v isoforms [154]. These studies show oxidation of several methionine residues suppresses inactivation; however, the cysteine residues of Na⁺ channels are also targets of redox modulation. In cultured neuroblastoma cells, thiol oxidizers significantly inhibited Na⁺ currents, which could be fully reversed by thiol-reducing agents DTT and intracellular GSH [155].

12. Circadian redox modification of membrane excitability in *Drosophila melanogaster*

The *Drosophila melanogaster* circadian system provides perhaps the most direct example for the interconnections between redox status, neuronal excitability, and the core components of the circadian clock. Approximately 150 neurons have been identified as the circadian clock circuit in the *Drosophila* CNS. These neurons are organized into clusters that include the large and small lateral ventral neurons (ILNvs and sLNvs), which play important roles in timekeeping [156]. Electrical properties of these pacemaker neurons show diurnal variation with higher resting membrane potential and greater frequency of spontaneous firing during the subjective day than the subjective night [157]. Similar to the SCN, this oscillation in excitability helps maintain circadian rhythmicity and also is subject to circadian regulation [44,158,159].

In this system, Cryptochrome (CRY) is photosensitive and functions as a blue light photoreceptor that regulates circadian entrainment [160,161]. CRY is expressed in clock neurons and interacts with the core circadian machinery by binding to Timeless (TIM) upon light activation to initiate its degradation. Light-activated CRY is also known to regulate the firing rate of ILNv arousal neurons through modification of K⁺ channel conductance [162]. In fact, CRY is coupled to neuronal electrical activity by redox modulation of ion channels. Upon light activation, the CRY chromophore flavin adenine dinucleotide (FAD) is reduced. This change in redox state is coupled to redox-sensitive β subunits of the voltage gated K⁺ channels, leading to rapid membrane depolarization [163].

13. Conclusion

The balance between generation of ROS and availability of antioxidant defense systems in the cell is reflected in the redox state. There is increasing evidence that small changes in ROS can regulate diverse cellular processes and pathways. Many proteins, including ion channels, can be modified by oxidizing or reducing reactions. The discovery of endogenous and conserved circadian redox oscillations brings time-of-day significance to all cellular processes that are susceptible to redox regulation. Wang and colleagues demonstrated that circadian redox environment of the SCN can modulate K⁺ ion channel activity and thus neuronal excitability [2]. This is the first and, to our knowledge, only example of the modulation of neuronal excitability by redox state in a circadian-dependent manner in mammals.

The SCN is unique among brain regions in that synchronized neuronal firing is independent of synaptic input, depending instead upon modulation of multiple intrinsic currents for daytime excitatory drive and nightly silencing. It is not difficult to appreciate how circadian redox regulation of key ion channels can rhythmically modulate SCN neuronal activity. However, it is challenging to infer the relationship between redox state and neuronal activity in other brain regions where studies have focused primarily on high-resolution relationships between redox and ion channel kinetics; insight on cellular changes in membrane potential and excitability due to redox modulation of specific ion channels in these regions is lacking. We predict that this relationship may vary across brain regions on a case-by-case basis depending on neuronal and circuit properties. This important area remains to be explored.

Numerous reports have demonstrated redox regulation of ion channels that contribute to cellular excitability. Thus, it is likely that circadian rhythms in redox state extend beyond the SCN to diverse brain regions where they may contribute to regulation of neuronal excitability. Furthermore, the circadian timing system is present in all tissues and cellular redox state has emerged as a contributor to tissue-specific rhythmicity. The redox oscillation has the potential to orchestrate the activation state of ion channels that underlie daily oscillations of electrical activity in all excitable cells throughout the body, enabling a nuanced fluctuation of excitable states. Redox-driven modulation of ion channels may be a fundamental form of regulation of physiological processes. The likelihood of this scenario raises the urgency of understanding the cellular and molecular groundwork and timing mechanisms that generate the metabolic redox oscillator and their relationship to the transcription-translation oscillator. Establishing these interconnections will shed light on the general role of cellular redox state in circadian timekeeping and in physiological regulation broadly.

Acknowledgments

The authors thank Jennifer Mitchell for provocative discussions and contributions to the manuscript, Ian Bothwell and Raj Iyer for contributions to some figures, and Ann Benefiel for preparing the manuscript. We are grateful for support from the National Institutes of Health (RO1HL086870, R21MH101655, and U01 MH109062) and the National Science Foundation (DBI 1450962 EAGER and IOS 1354913). Content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH or the NSF.

Abbreviations

AA	ascorbic acid
BK	large conductance Ca^{2+} - and voltage-activated K^{+} channel
ChT	chloramine-T
CK1	casein kinase 1
CNS	central nervous system
CO	carbon monoxide
CRY	cryptochrome
DHA	dehydroascorbic acid

DRG	dorsal root ganglion
DTNB	5,5-dithio-bis-(2-nitrobenzoic acid)
DTT	dithiothreitol
ER	endoplasmic reticulum
FAD	flavin adenine dinucleotide
GPx	glutathione peroxidase
GSH	glutathione
GSSG	glutathione disulfide
Kv	voltage-gated potassium channel
HVA	high-voltage-activated
IP₃	inositol triphosphate
LVA	low-voltage-activated
MetO	methionine sulfoxide
MsrA	methionine sulfoxide reductase
NADH	nicotinamide adenine dinucleotide
NADPH	nicotinamide adenine dinucleotide phosphate
NAMPT	nicotinamide phosphoribosyltransferase
NO	nitric oxide
NOX	NADPH oxidase
NPAS2	neuronal PAS domain protein 2
PER	period
PPP	pentose phosphate pathway
Prx	peroxiredoxin
ROR	RAR-related orphan receptor
ROS	reactive oxygen species
RyR	ryanodine receptor
SCN	suprachiasmatic nucleus
SIRT1	sirtuin 1
SIRT3	sirtuin 3

Trx	thioredoxin
VIP	vasoactive intestinal peptide
V_m	membrane potential

References

1. Edgar RS, Green EW, Zhao Y, van Ooijen G, Olmedo M, Qin X, Xu Y, Pan M, Valekunja UK, Feeney KA, Maywood ES, Hastings MH, Baliga NS, Merrow M, Millar AJ, Johnson CH, Kyriacou CP, O'Neill JS, Reddy AB. Peroxiredoxins are conserved markers of circadian rhythms. *Nature*. 2012; 485doi: 10.1038/nature11088
2. Wang TA, Yu YV, Govindaiah G, Ye X, Artinian L, Coleman TP, Sweedler JV, Cox CL, Gillette MU. Circadian rhythm of redox state regulates excitability in suprachiasmatic nucleus neurons. *Science*. 2012; 337:839–842. DOI: 10.1126/science.1222826 [PubMed: 22859819]
3. Hodgkin AL, Huxley AF, Katz B. Measurement of current-voltage relations in the membrane of the giant axon of *Loligo*. *J Physiol*. 1952; 116:424–448. DOI: 10.1113/jphysiol.1952.sp004716 [PubMed: 14946712]
4. Hille B. Modulation of ion-channel function by G-protein-coupled receptors. *Trends Neurosci*. 1994; 17:531–6. [PubMed: 7532338]
5. Misonou H, Mohapatra DP, Park EW, Leung V, Zhen D, Misonou K, Anderson AE, Trimmer JS. Regulation of ion channel localization and phosphorylation by neuronal activity. *Nat Neurosci*. 2004; 7:711–718. DOI: 10.1038/nn1260 [PubMed: 15195093]
6. Peers C, Boyle JP, Scragg JL, Dallas ML, Al-Owais MM, Hettiarachichi NT, Elies J, Johnson E, Gamper N, Steele DS. Diverse mechanisms underlying the regulation of ion channels by carbon monoxide. *Br J Pharmacol*. 2015; 172:1546–1556. DOI: 10.1111/bph.12760 [PubMed: 24818840]
7. Patel R, Sesti F. Oxidation of ion channels in the aging nervous system. *Brain Res*. 2016; 1639:174–185. DOI: 10.1016/j.brainres.2016.02.046 [PubMed: 26947620]
8. Knapp LT, Klann E. Role of reactive oxygen species in hippocampal long-term potentiation: contributory or inhibitory? *J Neurosci Res*. 2002; 70:1–7. DOI: 10.1002/jnr.10371 [PubMed: 12237859]
9. Borquez DA, Urrutia PJ, Wilson C, van Zundert B, Nunez MT, Gonzalez-Billault C. Dissecting the role of redox signaling in neuronal development. *J Neurochem*. 2016; 137:506–517. DOI: 10.1111/jnc.13581 [PubMed: 26875993]
10. Prieto-Bermejo R, Hernandez-Hernandez A. The importance of NADPH oxidases and redox signaling in angiogenesis. *Antioxidants (Basel, Switzerland)*. 2017; 6doi: 10.3390/antiox6020032
11. Iyer R, Wang TA, Gillette MU. Circadian gating of neuronal functionality: a basis for iterative metaplasticity. *Front Syst Neurosci*. 2014; 8doi: 10.3389/fnsys.2014.00164
12. Massaad C, Klann E. Reactive oxygen species in the regulation of synaptic plasticity and memory. *Antioxid Redox Signal*. 2011; 14:2013–2054. DOI: 10.1089/ars.2010.3208 [PubMed: 20649473]
13. Paula-Lima AC, Adasme T, Hidalgo C. Contribution of Ca²⁺ release channels to hippocampal synaptic plasticity and spatial memory: potential redox modulation. *Antioxid Redox Signal*. 2014; 21:892–914. DOI: 10.1089/ars.2013.5796 [PubMed: 24410659]
14. Davidson AJ, Sellix MT, Daniel J, Yamazaki S, Menaker M, Block GD. Chronic jet-lag increases mortality in aged mice. *Curr Biol*. 2006; 16doi: 10.1016/j.cub.2006.09.058
15. Barnard AR, Nolan PM. When clocks go bad: Neurobehavioural consequences of disrupted circadian timing. *PLoS Genet*. 2008; 4doi: 10.1371/journal.pgen.1000040
16. Menet JS, Rosbash M. When brain clocks lose track of time: Cause or consequence of neuropsychiatric disorders. *Curr Opin Neurobiol*. 2011; 21:849–857. DOI: 10.1016/j.conb.2011.06.008 [PubMed: 21737252]
17. Huang W, Ramsey KM, Marcheva B, Bass J. Circadian rhythms, sleep, and metabolism. *J Clin Invest*. 2011; 121:2133–2141. DOI: 10.1172/JCI46043 [PubMed: 21633182]

18. Moore RY, Eichler VB. Loss of a circadian adrenal corticosterone rhythm following suprachiasmatic lesions in the rat. *Brain Res.* 1972; 42:201–206. DOI: 10.1016/0006-8993(72)90054-6 [PubMed: 5047187]
19. Stephan FK, Zucker I. Circadian rhythms in drinking behavior and locomotor activity of rats are eliminated by hypothalamic lesions. *Proc Natl Acad Sci U S A.* 1972; 69:1583–6. DOI: 10.1073/pnas.69.6.1583 [PubMed: 4556464]
20. Lehman MN, Silver R, Gladstone WR, Kahn RM, Gibson M, Bittman EL. Circadian rhythmicity restored by neural transplant. Immunocytochemical characterization of the graft and its integration with the host brain. *J Neurosci.* 1987; 7:1626–1638. [PubMed: 3598638]
21. Schwartz WJ, Reppert SM, Eagan SM, Moore-Ede MC. In vivo metabolic activity of the suprachiasmatic nuclei: a comparative study. *Brain Res.* 1983; 274:184–187. DOI: 10.1016/0006-8993(83)90538-3 [PubMed: 6684493]
22. Ralph M, Foster R, Davis F, Menaker M. Transplanted suprachiasmatic nucleus determines circadian period. *Science.* 1990; 247:975–978. DOI: 10.1126/science.2305266 [PubMed: 2305266]
23. Eastman C, Rechtschaffen A. Circadian temperature and wake rhythms of rats exposed to prolonged continuous illumination. *Physiol Behav.* 1983; 31:417–427. DOI: 10.1016/0031-9384(83)90061-6 [PubMed: 6657763]
24. Meyer-Bernstein EL, Jetton AE, Matsumoto SI, Markuns JF, Lehman MN, Bittman EL. Effects of suprachiasmatic transplants on circadian rhythms of neuroendocrine function in golden hamsters. *Endocrinology.* 1999; 140:207–218. DOI: 10.1210/en.140.1.207 [PubMed: 9886827]
25. Partch CL, Green CB, Takahashi JS. Molecular architecture of the mammalian circadian clock. *Trends Cell Biol.* 2014; 24:90–99. DOI: 10.1016/j.tcb.2013.07.002 [PubMed: 23916625]
26. Caroline H, Ko JST. Molecular components of the mammalian circadian clock. *Hum Mol Genet.* 2006; 15(Spec No):R271–7. DOI: 10.1093/hmg/ddl207 [PubMed: 16987893]
27. Gillette MU, Prosser RA. Circadian rhythm of the rat suprachiasmatic brain slice is rapidly reset by daytime application of cAMP analogs. *Brain Res.* 1988; 474:348–52. [PubMed: 2850092]
28. Tamaru T, Okada M, Nagai K, Nakagawa H, Takamatsu K. Periodically fluctuating protein kinases phosphorylate CLOCK, the putative target in the suprachiasmatic nucleus. *J Neurochem.* 1999; 72:2191–2197. DOI: 10.1046/j.1471-4159.1999.0722191.x [PubMed: 10217302]
29. Lundkvist GB, Kwak Y, Davis EK, Tei H, Block GD. A calcium flux is required for circadian rhythm generation in mammalian pacemaker neurons. *J Neurosci.* 2005; 25:7682–6. DOI: 10.1523/JNEUROSCI.2211-05.2005 [PubMed: 16107654]
30. O'Neill JS, Maywood ES, Chesham JE, Takahashi JS, Hastings MH. cAMP-dependent signaling as a core component of the mammalian circadian pacemaker. *Science.* 2008; 320:949–953. DOI: 10.1126/science.1152506 [PubMed: 18487196]
31. Atkinson SE, Maywood ES, Chesham JE, Wozny C, Colwell CS, Hastings MH, Williams SR. Cyclic AMP signaling control of action potential firing rate and molecular circadian pacemaking in the suprachiasmatic nucleus. *J Biol Rhythms.* 2011; 26:210–220. DOI: 10.1177/0748730411402810 [PubMed: 21628548]
32. Gillette MU, Wang TA. Brain circadian oscillators and redox regulation in mammals. *Antioxid Redox Signal.* 2014; 20:1–47. DOI: 10.1089/ars.2013.5598 [PubMed: 23725018]
33. Inouye ST, Kawamura H. Persistence of circadian rhythmicity in a mammalian hypothalamic “island” containing the suprachiasmatic nucleus. *Proc Natl Acad Sci.* 1979; 76:5962–5966. DOI: 10.1073/pnas.76.11.5962 [PubMed: 293695]
34. Prosser RA, Gillette MU. The mammalian circadian clock in the suprachiasmatic nuclei is reset in vitro by cAMP. *J Neurosci.* 1989; 9:1073–81. [PubMed: 2538580]
35. Belle MDC, Diekmann CO, Forger DB, Piggins HD. Daily electrical silencing in the mammalian circadian clock. *Science.* 2009; 326:281–284. DOI: 10.1126/science.1169657 [PubMed: 19815775]
36. Green DJ, Gillette R. Circadian rhythm of firing rate recorded from single cells in the rat suprachiasmatic brain slice. *Brain Res.* 1982; 245:198–200. [PubMed: 6889453]

37. Groos G, Hendriks J. Circadian rhythms in electrical discharge of rat suprachiasmatic neurones recorded in vitro. *Neurosci Lett*. 1982; 34:283–288. DOI: 10.1016/0304-3940(82)90189-6 [PubMed: 6298675]
38. Colwell CS. Linking neural activity and molecular oscillations in the SCN. *Nat Rev Neurosci*. 2011; 12:553–69. DOI: 10.1038/nrn3086 [PubMed: 21886186]
39. Meijer JH, Watanabe K, Schaap J, Albus H, Détári L. Light responsiveness of the suprachiasmatic nucleus: long-term multiunit and single-unit recordings in freely moving rats. *J Neurosci*. 1998; 18:9078–87. [PubMed: 9787011]
40. Welsh DK, Logothetis DE, Meister M, Reppert SM. Individual neurons dissociated from rat suprachiasmatic nucleus express independently phased circadian firing rhythms. *Neuron*. 1995; 14:697–706. [PubMed: 7718233]
41. Jiang ZG, Yang Y, Liu ZP, Allen CN. Membrane properties and synaptic inputs of suprachiasmatic nucleus neurons in rat brain slices. *J Physiol*. 1997; 141–59.
42. de Jeu M, Hermes M, Pennartz C. Circadian modulation of membrane properties in slices of rat suprachiasmatic nucleus. *Neuroreport*. 1998; 9:3725–9. [PubMed: 9858386]
43. Kuhlman SJ, McMahon DG. Rhythmic regulation of membrane potential and potassium current persists in SCN neurons in the absence of environmental input. *Eur J Neurosci*. 2004; 20:1113–1117. DOI: 10.1111/j.1460-9568.2004.03555.x [PubMed: 15305881]
44. Nitabach MN, Blau J, Holmes TC. Electrical silencing of *Drosophila* pacemaker neurons stops the free-running circadian clock. *Cell*. 2002; 109:485–495. DOI: 10.1016/S0092-8674(02)00737-7 [PubMed: 12086605]
45. Jones JR, Tackenberg MC, McMahon DG. Manipulating circadian clock neuron firing rate resets molecular circadian rhythms and behavior. *Nat Neurosci*. 2015; 18:1–5. DOI: 10.1038/nn.3937 [PubMed: 25547471]
46. Balsalobre A, Damiola F, Schibler U. A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell*. 1998; 93:929–937. DOI: 10.1016/S0092-8674(00)81199-X [PubMed: 9635423]
47. Yoo S-H, Yamazaki S, Lowrey PL, Shimomura K, Ko CH, Buhr ED, Slepka SM, Hong H-K, Oh WJ, Yoo OJ, Menaker M, Takahashi JS. PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci*. 2004; 101:5339–5346. DOI: 10.1073/pnas.0308709101 [PubMed: 14963227]
48. Welsh DK, Yoo S-H, Liu AC, Takahashi JS, Kay SA. Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression. *Curr Biol*. 2004; 14:2289–2295. DOI: 10.1016/j.cub.2004.11.057 [PubMed: 15620658]
49. Abe M, Herzog ED, Yamazaki S, Straume M, Tei H, Sakaki Y, Menaker M, Block GD. Circadian rhythms in isolated brain regions. *J Neurosci*. 2002; 22:350–6. [PubMed: 11756518]
50. Granados-Fuentes D. The suprachiasmatic nucleus entrains, but does not sustain, circadian rhythmicity in the olfactory bulb. *J Neurosci*. 2004; 24:615–619. DOI: 10.1523/JNEUROSCI.4002-03.2004 [PubMed: 14736846]
51. Guilding C, Piggins HD. Challenging the omnipotence of the suprachiasmatic timekeeper: Are circadian oscillators present throughout the mammalian brain? *Eur J Neurosci*. 2007; 25:3195–3216. DOI: 10.1111/j.1460-9568.2007.05581.x [PubMed: 17552989]
52. Silver R, LeSauter J, Tresco PA, Lehman MN. A diffusible coupling signal from the transplanted suprachiasmatic nucleus controlling circadian locomotor rhythms. *Nature*. 1996; 382:810–813. DOI: 10.1038/382810a0 [PubMed: 8752274]
53. Mohawk JA, Green CB, Takahashi JS. Central and peripheral circadian clocks in mammals. *Ann Rev Neurosci*. 2013; 35:445–462. DOI: 10.1146/annurev-neuro-060909-153128.CENTRAL
54. Loh DH, Jami SA, Flores RE, Truong D, Ghiani CA, O'Dell TJ, Colwell CS. Misaligned feeding impairs memories. *Elife*. 2015; 4:1–16. DOI: 10.7554/eLife.09460
55. la Fleur SE, Kalsbeek A, Wortel J, Fekkes ML, Buijs RM. A daily rhythm in glucose tolerance: a role for the suprachiasmatic nucleus. *Diabetes*. 2001; 50:1237–43. [PubMed: 11375322]
56. Schwartz WJ, Gainer H. Suprachiasmatic nucleus: use of ¹⁴C-labeled deoxyglucose uptake as a functional marker. *Science*. 1977; 197:1089–91. [PubMed: 887940]

57. Brown SA, Azzi A. Peripheral circadian oscillators in mammals. *Handb Exp Pharmacol*. 2013; 217:45–66. DOI: 10.1007/978-3-642-25950-0-3
58. Turek FW, Joshu C, Kohsaka A, Lin E, Ivanova G, McDearmon E, Laposky A, Losee-Olson S, Easton A, Jensen DR, Eckel RH, Takahashi JS, Bass J. Obesity and metabolic syndrome in circadian Clock mutant mice. *Science*. 2005; 308:1043–5. DOI: 10.1126/science.1108750 [PubMed: 15845877]
59. Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB. A circadian gene expression atlas in mammals: Implications for biology and medicine. *Proc Natl Acad Sci*. 2014; 111:16219–16224. DOI: 10.1073/pnas.1408886111 [PubMed: 25349387]
60. Akhtar RA, Reddy AB, Maywood ES, Clayton JD, King VM, Smith AG, Gant TW, Hastings MH, Kyriacou CP. Circadian cycling of the mouse liver transcriptome, as revealed by cDNA microarray, is driven by the suprachiasmatic nucleus. *Curr Biol*. 2002; 12:540–550. DOI: 10.1016/S0960-9822(02)00759-5 [PubMed: 11937022]
61. Reddy AB, Karp NA, Maywood ES, Sage EA, Deery M, O'Neill JS, Wong GKY, Chesham J, Odell M, Lilley KS, Kyriacou CP, Hastings MH. Circadian orchestration of the hepatic proteome. *Curr Biol*. 2006; 16:1107–1115. DOI: 10.1016/j.cub.2006.04.026 [PubMed: 16753565]
62. Hansen AP, Johansen K. Diurnal patterns of blood glucose, serum free fatty acids, insulin, glucagon and growth hormone in normals and juvenile diabetics. *Diabetologia*. 1970; 6:27–33. DOI: 10.1007/BF00425888 [PubMed: 5434734]
63. Yang X, Downes M, Yu RT, Bookout AL, He W, Straume M, Mangelsdorf DJ, Evans RM. Nuclear receptor expression links the circadian clock to metabolism. *Cell*. 2006; 126:801–810. DOI: 10.1016/j.cell.2006.06.050 [PubMed: 16923398]
64. Feeney KA, Hansen LL, Putker M, Olivares-Yañez C, Day J, Eades LJ, Larrondo LF, Hoyle NP, O'Neill JS, van Ooijen G. Daily magnesium fluxes regulate cellular timekeeping and energy balance. *Nature*. 2016; 532:375–379. DOI: 10.1038/nature17407 [PubMed: 27074515]
65. Schafer FQ, Buettner GR. Redox environment of the cell as viewed through the redox state of the glutathione disulfide/glutathione couple. *Free Radic Biol Med*. 2001; 30:1191–1212. DOI: 10.1016/S0891-5849(01)00480-4 [PubMed: 11368918]
66. Arnér ES, Holmgren A. Physiological functions of thioredoxin and thioredoxin reductase. *Eur J Biochem*. 2000; 267:6102–6109. DOI: 10.1046/j.1432-1327.2000.01701.x [PubMed: 11012661]
67. Rhee SG. Overview on peroxiredoxin. *Mol Cells*. 2016; 39:1–5. DOI: 10.14348/molcells.2016.2368 [PubMed: 26831451]
68. Schulz E, Wenzel P, Munzel T, Daiber A. Mitochondrial redox signaling: Interaction of mitochondrial reactive oxygen species with other sources of oxidative stress. *Antioxid Redox Signal*. 2014; 20:308–324. DOI: 10.1089/ars.2012.4609 [PubMed: 22657349]
69. Bienert GP, Moller ALB, Kristiansen KA, Schulz A, Moller IM, Schjoerring JK, Jahn TP. Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J Biol Chem*. 2007; 282:1183–1192. DOI: 10.1074/jbc.M603761200 [PubMed: 17105724]
70. Veal E, Day A. Hydrogen peroxide as a signaling molecule. *Antioxid Redox Signal*. 2011; 15:147–151. DOI: 10.1089/ars.2011.3968 [PubMed: 21375475]
71. Cross CE, Halliwell B, Borish ET, Pryor WA, Ames BN, Saul RL, McCord JM, Harman D. Oxygen radicals and human disease. *Ann Intern Med*. 1987; 107:526–545. DOI: 10.7326/0003-4819-107-4-526 [PubMed: 3307585]
72. Torres M, Forman HJ. Redox signaling and the MAP kinase pathways. *Biofactors*. 2003; 17:287–296. DOI: 10.1002/biof.5520170128 [PubMed: 12897450]
73. Forman HJ, Fukuto JM, Torres M. Redox signaling: thiol chemistry defines which reactive oxygen and nitrogen species can act as second messengers. *Am J Physiol Cell Physiol*. 2004; 287:C246–56. DOI: 10.1152/ajpcell.00516.2003 [PubMed: 15238356]
74. Finkel T. Signal transduction by reactive oxygen species. *J Cell Biol*. 2011; 194:7–15. DOI: 10.1083/jcb.201102095 [PubMed: 21746850]
75. Comini MA. Measurement and meaning of cellular thiol:disulfide redox status. *Free Radic Res*. 2016; 50:246–271. DOI: 10.3109/10715762.2015.1110241 [PubMed: 26695718]

76. Rehder DS, Borges CR. Cysteine sulfenic acid as an intermediate in disulfide bond formation and nonenzymatic protein folding. *Biochemistry*. 2010; 49:7748–7755. DOI: 10.1021/bi1008694 [PubMed: 20712299]
77. Grek CL, Zhang J, Manevich Y, Townsend DM, Tew KD. Causes and consequences of cysteine s-glutathionylation. *J Biol Chem*. 2013; 288:26497–26504. DOI: 10.1074/jbc.R113.461368 [PubMed: 23861399]
78. Levine RL, Mosoni L, Berlett BS, Stadtman ER. Methionine residues as endogenous antioxidants in proteins. *Proc Natl Acad Sci U S A*. 1996; 93:15036–40. DOI: 10.1073/pnas.93.26.15036 [PubMed: 8986759]
79. Zhang X-H, Weissbach H. Origin and evolution of the protein-repairing enzymes methionine sulfoxide reductases. *Biol Rev Camb Philos Soc*. 2008; 83:249–257. DOI: 10.1111/j.1469-185X.2008.00042.x [PubMed: 18557976]
80. Rutter J. Regulation of Clock and NPAS2 DNA binding by the redox state of NAD cofactors. *Science*. 2001; 293:510–514. DOI: 10.1126/science.1060698 [PubMed: 11441146]
81. Ramsey KM, Yoshino J, Brace CS, Abrassart D, Kobayashi Y, Marcheva B, Hong H-K, Chong JL, Buhr ED, Lee C, Takahashi JS, Imai S-i, Bass J. Circadian clock feedback cycle through NAMPT-mediated NAD⁺ biosynthesis. *Science*. 2009; 324:651–654. DOI: 10.1126/science.1171641 [PubMed: 19299583]
82. Nakahata Y, Sahar S, Astarita G, Kaluzova M, Sassone-Corsi P. Circadian control of the NAD⁺ salvage pathway by CLOCK-SIRT1. *Science*. 2009; 324:654–657. DOI: 10.1126/science.1170803 [PubMed: 19286518]
83. Peek CB, Affinati AH, Ramsey KM, Kuo H-Y, Yu W, Sena LA, Ilkayeva O, Marcheva B, Kobayashi Y, Omura C, Levine DC, Bacsik DJ, Gius D, Newgard CB, Goetzman E, Chandel NS, Denu JM, Mrksich M, Bass J. Circadian clock NAD⁺ cycle drives mitochondrial oxidative metabolism in mice. *Science*. 2013; 342:1243417–1243417. DOI: 10.1126/science.1243417 [PubMed: 24051248]
84. Asher G, Gatfield D, Stratmann M, Reinke H, Dibner C, Kreppel F, Mostoslavsky R, Alt FW, Schibler U. SIRT1 regulates circadian clock gene expression through PER2 deacetylation. *Cell*. 2008; 134:317–328. DOI: 10.1016/j.cell.2008.06.050 [PubMed: 18662546]
85. Belden WJ, Dunlap JC. SIRT1 is a circadian deacetylase for core clock components. *Cell*. 2008; 134:212–214. DOI: 10.1016/j.cell.2008.07.010 [PubMed: 18662537]
86. Bellet MM, Orozco-Solis R, Sahar S, Eckel-Mahan K, Sassone-Corsi P. The time of metabolism: NAD⁺, SIRT1, and the circadian clock. *Cold Spring Harb Symp Quant Biol*. 2011; 76:31–38. DOI: 10.1101/sqb.2011.76.010520 [PubMed: 22179986]
87. Carter EL, Gupta N, Ragsdale SW. High affinity heme binding to a heme regulatory motif on the nuclear receptor Rev-erb β leads to its degradation and indirectly regulates its interaction with nuclear receptor corepressor. *J Biol Chem*. 2016; 291:2196–2222. DOI: 10.1074/jbc.M115.670281 [PubMed: 26670607]
88. Carter EL, Ramirez Y, Ragsdale SW. The heme-regulatory motif of nuclear receptor Rev-erb β is a key mediator of heme and redox signaling in circadian rhythm maintenance and metabolism. *J Biol Chem*. 2017; 292:11280–11299. DOI: 10.1074/jbc.M117.783118 [PubMed: 28500133]
89. Gerhart-Hines Z, Lazar MA. Rev-erba and the circadian transcriptional regulation of metabolism. *Diabetes Obes Metab*. 2015; 17:12–16. DOI: 10.1111/dom.12510 [PubMed: 26332963]
90. Dioum EM, Rutter J, Tuckerman JR, Gonzalez G, Marie-Alda G-G, L MS. {NPAS2:} a gas-responsive transcription factor. *Science*. 2002; 298:2385–2387. DOI: 10.1126/science.1078456 [PubMed: 12446832]
91. Lee Y, Kim E-K. AMP-activated protein kinase as a key molecular link between metabolism and clockwork. *Exp Mol Med*. 2013; 45:e33.doi: 10.1038/emm.2013.65 [PubMed: 23887727]
92. Hirano A, Braas D, Fu Y-H, Ptacek LJ. FAD Regulates CRYPTOCHROME protein stability and circadian clock in mice. *Cell Rep*. 2017; 19:255–266. DOI: 10.1016/j.celrep.2017.03.041 [PubMed: 28402850]
93. O'Neill JS, Reddy AB. Circadian clocks in human red blood cells. *Nature*. 2011; 469:498–503. DOI: 10.1038/nature09702 [PubMed: 21270888]

94. Causton HC, Feeney KA, Ziegler CA, O'Neill JS. Metabolic cycles in yeast share features conserved among circadian rhythms. *Curr Biol.* 2015; 25:1056–1062. DOI: 10.1016/j.cub.2015.02.035 [PubMed: 25866393]
95. Rey G, Valekunja UK, Feeney KA, Wulund L, Milev NB, Stangherlin A, Ansel-Bollepalli L, Velagapudi V, O'Neill JS, Reddy AB. The pentose phosphate pathway regulates the circadian clock. *Cell Metab.* 2016; 24:462–473. DOI: 10.1016/j.cmet.2016.07.024 [PubMed: 27546460]
96. Putker M, Crosby P, Feeney KA, Hoyle NP, Costa ASH, Gaude E, Frezza C, O'Neill JS. Mammalian circadian period, but not phase and amplitude, is robust against redox and metabolic perturbations. *Antioxid Redox Signal.* 2017; doi: 10.1089/ars.2016.6911
97. Aggarwal NT, Makielski JC. Redox control of cardiac excitability. *Antioxid Redox Signal.* 2013; 18:432–68. DOI: 10.1089/ars.2011.4234 [PubMed: 22897788]
98. Nelson MT, Joksovic PM, Perez-Reyes E, Todorovic SM. The endogenous redox agent L-cysteine induces T-type Ca^{2+} channel-dependent sensitization of a novel subpopulation of rat peripheral nociceptors. *J Neurosci.* 2005; 25:8766–8775. DOI: 10.1523/JNEUROSCI.2527-05.2005 [PubMed: 16177046]
99. Jagodic MM, Pathirathna S, Nelson MT, Mancuso S, Joksovic PM, Rosenberg ER, Bayliss DA, Jevtovic-Todorovic V, Todorovic SM. Cell-specific alterations of T-type calcium current in painful diabetic neuropathy enhance excitability of sensory neurons. *J Neurosci.* 2007; 27:3305–3316. DOI: 10.1523/JNEUROSCI.4866-06.2007 [PubMed: 17376991]
100. Evans JG, Todorovic SM. Redox and trace metal regulation of ion channels in the pain pathway. *Biochem J.* 2015; 470:275–280. DOI: 10.1042/BJ20150522 [PubMed: 26341484]
101. Pan Y, Weng J, Levin EJ, Zhou M. Oxidation of NADPH on Kvbeta1 inhibits ball-and-chain type inactivation by restraining the chain. *Proc Natl Acad Sci U S A.* 2011; 108:5885–5890. DOI: 10.1073/pnas.1100316108 [PubMed: 21436029]
102. Gonzalez C, Baez-Nieto D, Valencia I, Oyarzun I, Rojas P, Naranjo D, Latorre R. $\text{K}(+)$ channels: function-structural overview. *Compr Physiol.* 2012; 2:2087–2149. DOI: 10.1002/cphy.c110047 [PubMed: 23723034]
103. Rettig J, Heinemann SH, Wunder F, Lorra C, Parcej DN, Dolly JO, Pongs O. Inactivation properties of voltage-gated K^{+} channels altered by presence of beta-subunit. *Nature.* 1994; 369:289–294. DOI: 10.1038/369289a0 [PubMed: 8183366]
104. Ruppersberg JP, Stocker M, Pongs O, Heinemann SH, Frank R, Koenen M. Regulation of fast inactivation of cloned mammalian $\text{IK}(\text{A})$ channels by cysteine oxidation. *Nature.* 1991; 352:711–714. DOI: 10.1038/352711a0 [PubMed: 1908562]
105. Bähring R, Milligan CJ, Vardanyan V, Engeland B, Young BA, Dannenberg J, Waldschutz R, Edwards JP, Wray D, Pongs O. Coupling of voltage-dependent potassium channel inactivation and oxidoreductase active site of $\text{Kv}\beta$ subunits. *J Biol Chem.* 2001; 276:22923–22929. DOI: 10.1074/jbc.M100483200 [PubMed: 11294861]
106. Ciorba MA, Heinemann SH, Weissbach H, Brot N, Hoshi T. Modulation of potassium channel function by methionine oxidation and reduction. *Proc Natl Acad Sci U S A.* 1997; 94:9932–9937. [PubMed: 9275229]
107. Vacher H, Mohapatra DP, Trimmer JS. Localization and targeting of voltage-dependent ion channels in mammalian central neurons. *Physiol Rev.* 2008; 88:1407–1447. DOI: 10.1152/physrev.00002.2008 [PubMed: 18923186]
108. Hasan SMK, Redzic ZB, Alshuaib WB. Hydrogen peroxide-induced reduction of delayed rectifier potassium current in hippocampal neurons involves oxidation of sulfhydryl groups. *Brain Res.* 2013; 1520:61–69. DOI: 10.1016/j.brainres.2013.05.019 [PubMed: 23688542]
109. Gamper N, Zaika O, Li Y, Martin P, Hernandez CC, Perez MR, Wang AYC, Jaffe DB, Shapiro MS. Oxidative modification of M-type K^{+} channels as a mechanism of cytoprotective neuronal silencing. *EMBO J.* 2006; 25:4996–5004. DOI: 10.1038/sj.emboj.7601374 [PubMed: 17024175]
110. Lee US, Cui J. BK channel activation: Structural and functional insights. *Trends Neurosci.* 2010; 33:415–423. DOI: 10.1016/j.tins.2010.06.004 [PubMed: 20663573]
111. Golowasch J, Kirkwood A, Miller C. Allosteric effects of Mg^{2+} on the gating of Ca^{2+} -activated K^{+} channels from mammalian skeletal muscle. *J Exp Biol.* 1986; 124:5–13. [PubMed: 2428908]

112. Zhang X, Solaro CR, Lingle CJ. Allosteric regulation of BK channel gating by Ca(2+) and Mg(2+) through a nonselective, low affinity divalent cation site. *J Gen Physiol.* 2001; 118:607–36. DOI: 10.1085/jgp.118.5.607 [PubMed: 11696615]
113. Shi J, Cui J. Intracellular Mg(2+) Enhances the function of BK-type Ca(2+)-activated K(+) channels. *J Gen Physiol.* 2001; 118:589–606. [PubMed: 11696614]
114. Whitt JP, Montgomery JR, Meredith AL. BK channel inactivation gates daytime excitability in the circadian clock. *Nat Commun.* 2016; 7:10837.doi: 10.1038/ncomms10837 [PubMed: 26940770]
115. Panda S, Antoch MP, Miller BH, Su AI, Schook AB, Straume M, Schultz PG, Kay SA, Takahashi JS, Hogenesch JB. Coordinated transcription of key pathways in the mouse by the circadian clock. *Cell.* 2002; 109:307–320. [PubMed: 12015981]
116. Meredith AL, Wiler SW, Miller BH, Takahashi JS, Fodor AA, Ruby NF, Aldrich RW. BK calcium-activated potassium channels regulate circadian behavioral rhythms and pacemaker output. *Nat Neurosci.* 2006; 9:1041–1049. DOI: 10.1038/nn1740 [PubMed: 16845385]
117. Pitts GR, Ohta H, McMahon DG. Daily rhythmicity of large-conductance Ca2+-activated K+ currents in suprachiasmatic nucleus neurons. *Brain Res.* 2006; 1071:54–62. DOI: 10.1016/j.brainres.2005.11.078 [PubMed: 16412396]
118. Torres YP, Morera FJ, Carvacho I, Latorre R. A marriage of convenience: beta-subunits and voltage-dependent K+ channels. *J Biol Chem.* 2007; 282:24485–24489. DOI: 10.1074/jbc.R700022200 [PubMed: 17606609]
119. DiChiara TJ, Reinhart PH. Redox modulation of hsls Ca2+-activated K+ channels. *J Neurosci.* 1997; 17:4942–4955. [PubMed: 9185532]
120. Gong L, Gao TM, Huang H, Tong Z. Redox modulation of large conductance calcium-activated potassium channels in CA1 pyramidal neurons from adult rat hippocampus. *Neurosci Lett.* 2000; 286:191–194. [PubMed: 10832017]
121. Soh H, Jung W, Uhm DY, Chung S. Modulation of large conductance calcium-activated potassium channels from rat hippocampal neurons by glutathione. *Neurosci Lett.* 2001; 298:115–118. [PubMed: 11163291]
122. Brenner R, Perez GJ, Bonev AD, Eckman DM, Kosek JC, Wiler SW, Patterson AJ, Nelson MT, Aldrich RW. Vasoregulation by the beta1 subunit of the calcium-activated potassium channel. *Nature.* 2000; 407:870–876. DOI: 10.1038/35038011 [PubMed: 11057658]
123. Zeng X-H, Xia X-M, Lingle CJ. Redox-sensitive extracellular gates formed by auxiliary beta subunits of calcium-activated potassium channels. *Nat Struct Biol.* 2003; 10:448–454. DOI: 10.1038/nsb932 [PubMed: 12740608]
124. Santarelli LC, Wassef R, Heinemann SH, Hoshi T. Three methionine residues located within the regulator of conductance for K+ (RCK) domains confer oxidative sensitivity to large-conductance Ca2+-activated K+ channels. *J Physiol.* 2006; 571:329–348. DOI: 10.1113/jphysiol.2005.101089 [PubMed: 16396928]
125. Zhang G, Xu R, Heinemann SH, Hoshi T. Cysteine oxidation and rundown of large-conductance Ca2+-dependent K+ channels. *Biochem Biophys Res Commun.* 2006; 342:1389–1395. DOI: 10.1016/j.bbrc.2006.02.079 [PubMed: 16516848]
126. Ho K, Nichols CG, Lederer WJ, Lytton J, Vassilev PM, Kanazirska MV, Hebert SC. Cloning and expression of an inwardly rectifying ATP-regulated potassium channel. *Nature.* 1993; 362:31–38. DOI: 10.1038/362031a0 [PubMed: 7680431]
127. Kubo Y, Baldwin TJ, Nung Jan Y, Jan LY. Primary structure and functional expression of a mouse inward rectifier potassium channel. *Nature.* 1993; 362:127–133. DOI: 10.1038/362127a0 [PubMed: 7680768]
128. Kuo A, Gulbis JM, Antcliff JF, Rahman T, Lowe ED, Zimmer J, Cuthbertson J, Ashcroft FM, Ezaki T, Doyle DA. Crystal structure of the potassium channel KirBac1.1 in the closed state. *Science.* 2003; 300:1922–1926. DOI: 10.1126/science.1085028 [PubMed: 12738871]
129. Hagiwara S, Takahashi K. The anomalous rectification and cation selectivity of the membrane of a starfish egg cell. *J Membr Biol.* 1974; 18:61–80. DOI: 10.1007/BF01870103 [PubMed: 4854650]

130. Miyazaki S-I, Takahashi K, Tsuda K, Yoshii M. Analysis of non-linearity observed in the current—voltage relation of the tunicate embryo. *J Physiol.* 1974; 238:55–77. DOI: 10.1113/jphysiol.1974.sp010510 [PubMed: 4838798]
131. Lacampagne A, Duittoz A, Bolanos P, Peineau N, Argibay JA. Effect of sulfhydryl oxidation on ionic and gating currents associated with L-type calcium channels in isolated guinea-pig ventricular myocytes. *Cardiovasc Res.* 1995; 30:799–806. [PubMed: 8595629]
132. Vandenberg CA. Inward rectification of a potassium channel in cardiac ventricular cells depends on internal magnesium ions. *Physiol Sci.* 1987; 84:2560–2564. DOI: 10.1073/pnas.84.8.2560
133. Matsuda H, Saigusa A, Irisawa H. Ohmic conductance through the inwardly rectifying K channel and blocking by internal Mg²⁺ Nature. 1987; 325:156–159. DOI: 10.1038/325156a0 [PubMed: 2433601]
134. Lopatin AN, Makhina EN, Nichols CG. Potassium channel block by cytoplasmic polyamines as the mechanism of intrinsic rectification. *Nature.* 1994; 372:366–369. DOI: 10.1038/372366a0 [PubMed: 7969496]
135. Ruppersberg JP, Fakler B. Complexity of the regulation of Kir2.1 K⁺ channels. *Neuropharmacology.* 1996; 35:887–893. [PubMed: 8938719]
136. Zeidner G, Sadjia R, Reuveny E. Redox-dependent gating of G protein-coupled inwardly rectifying K⁺ channels. *J Biol Chem.* 2001; 276:35564–70. DOI: 10.1074/jbc.M105189200 [PubMed: 11466316]
137. Frazier HN, Maimaiti S, Anderson KL, Brewer LD, Gant JC, Porter NM, Thibault O. Calcium's role as nuanced modulator of cellular physiology in the brain. *Biochem Biophys Res Commun.* 2017; 483:981–987. DOI: 10.1016/j.bbrc.2016.08.105 [PubMed: 27553276]
138. Todorovic SM, Jevtovic-Todorovic V. Redox regulation of neuronal voltage-gated calcium channels. *Antioxid Redox Signal.* 2014; 21:880–891. DOI: 10.1089/ars.2013.5610 [PubMed: 24161125]
139. Todorovic SM, Jevtovic-Todorovic V, Meyenburg A, Mennerick S, Perez-Reyes E, Romano C, Olney JW, Zorumski CF. Redox modulation of T-type calcium channels in rat peripheral nociceptors. *Neuron.* 2001; 31:75–85. [PubMed: 11498052]
140. Nelson MT, Joksovic PM, Su P, Kang H-W, Van Deusen A, Baumgart JP, David LS, Snutch TP, Barrett PQ, Lee J-H, Zorumski CF, Perez-Reyes E, Todorovic SM. Molecular mechanisms of subtype-specific inhibition of neuronal T-type calcium channels by ascorbate. *J Neurosci.* 2007; 27:12577–12583. DOI: 10.1523/JNEUROSCI.2206-07.2007 [PubMed: 18003836]
141. Voisin T, Bourinet E, Lory P. Genetic alteration of the metal/redox modulation of Cav3.2 T-type calcium channel reveals its role in neuronal excitability. *J Physiol.* 2016; 594:3561–3574. DOI: 10.1113/JP271925 [PubMed: 26931411]
142. Joksovic PM, Nelson MT, Jevtovic-Todorovic V, Patel MK, Perez-Reyes E, Campbell KP, Chen C-C, Todorovic SM. CaV3.2 is the major molecular substrate for redox regulation of T-type Ca²⁺ channels in the rat and mouse thalamus. *J Physiol.* 2006; 574:415–430. DOI: 10.1113/jphysiol.2006.110395 [PubMed: 16644797]
143. Hool LC. Hypoxia increases the sensitivity of the L-type Ca(2+) current to beta-adrenergic receptor stimulation via a C2 region-containing protein kinase C isoform. *Circ Res.* 2000; 87:1164–1171. DOI: 10.1161/01.RES.87.12.1164 [PubMed: 11110774]
144. Yamaoka K, Yakehiro M, Yuki T, Fujii H, Seyama I. Effect of sulfhydryl reagents on the regulatory system of the L-type Ca channel in frog ventricular myocytes. *Pflugers Arch.* 2000; 440:207–215. DOI: 10.1007/s004249900242 [PubMed: 10898520]
145. Tang H, Viola HM, Filipovska A, Hool LC. Ca(v)1.2 calcium channel is glutathionylated during oxidative stress in guinea pig and ischemic human heart. *Free Radic Biol Med.* 2011; 51:1501–1511. DOI: 10.1016/j.freeradbiomed.2011.07.005 [PubMed: 21810465]
146. Johnstone VPA, Hool LC. Glutathionylation of the L-type Ca²⁺ channel in oxidative stress-induced pathology of the heart. *Int J Mol Sci.* 2014; 15:19203–19225. DOI: 10.3390/ijms151019203 [PubMed: 25340983]
147. Marengo JJ, Hidalgo C, Bull R. Sulfhydryl oxidation modifies the calcium dependence of ryanodine-sensitive calcium channels of excitable cells. *Biophys J.* 1998; 74:1263–1277. DOI: 10.1016/S0006-3495(98)77840-3 [PubMed: 9512024]

148. Xu L, Eu JP, Meissner G, Stamler JS. Activation of the cardiac calcium release channel (ryanodine receptor) by poly-S-nitrosylation. *Science*. 1998; 279:234–7. [PubMed: 9422697]
149. Sun J, Yamaguchi N, Xu L, Eu JP, Stamler JS, Meissner G. Regulation of the cardiac muscle ryanodine receptor by O(2) tension and S-nitrosoglutathione. *Biochemistry*. 2008; 47:13985–90. DOI: 10.1021/bi8012627 [PubMed: 19053230]
150. Ding JM, Buchanan GF, Tischkau Sa, Chen D, Kuriashkina L, Faiman LE, Alster JM, McPherson PS, Campbell KP, Gillette MU. A neuronal ryanodine receptor mediates light-induced phase delays of the circadian clock. *Nature*. 1998; 394:381–384. DOI: 10.1038/28639 [PubMed: 9690474]
151. Crill WE. Persistent sodium current in mammalian central neurons. *Annu Rev Physiol*. 1996; 58:349–362. DOI: 10.1146/annurev.ph.58.030196.002025 [PubMed: 8815799]
152. Wang GK. Irreversible modification of sodium channel inactivation in toad myelinated nerve fibres by the oxidant chloramine-T. *J Physiol*. 1984; 346:127–141. [PubMed: 6321714]
153. Wang GK, Brodwick MS, Eaton DC. Removal of sodium channel inactivation in squid axon by the oxidant chloramine-T. *J Gen Physiol*. 1985; 86:289–302. DOI: 10.1085/jgp.86.2.289 [PubMed: 2413161]
154. Kassmann M, Hansel A, Leipold E, Birkenbeil J, Lu S-Q, Hoshi T, Heinemann SH. Oxidation of multiple methionine residues impairs rapid sodium channel inactivation. *Pflügers Arch*. 2008; 456:1085–1095. DOI: 10.1007/s00424-008-0477-6 [PubMed: 18369661]
155. Evans JR, Bielefeldt K. Regulation of sodium currents through oxidation and reduction of thiol residues. *Neuroscience*. 2000; 101:229–236. [PubMed: 11068151]
156. Yoshii T, Hermann-Luibl C, Helfrich-Förster C. Circadian light-input pathways in *Drosophila*. *Commun Integr Biol*. 2016; 9doi: 10.1080/19420889.2015.1102805
157. Sheeba V, Gu H, Sharma VK, O'Dowd DK, Holmes TC. Circadian- and light-dependent regulation of resting membrane potential and spontaneous action potential firing of *Drosophila* circadian pacemaker neurons. *J Neurophysiol*. 2008; 99:976–88. DOI: 10.1152/jn.00930.2007 [PubMed: 18077664]
158. Nitabach MN. Electrical hyperexcitation of lateral ventral pacemaker neurons desynchronizes downstream circadian oscillators in the fly circadian circuit and induces multiple behavioral periods. *J Neurosci*. 2006; 26:479–489. DOI: 10.1523/JNEUROSCI.3915-05.2006 [PubMed: 16407545]
159. Sheeba V, Fogle KJ, Kaneko M, Rashid S, Chou YT, Sharma VK, Holmes TC. Large ventral lateral neurons modulate arousal and sleep in *Drosophila*. *Curr Biol*. 2008; 18:1537–1545. DOI: 10.1016/j.cub.2008.08.033 [PubMed: 18771923]
160. Emery P, So WV, Kaneko M, Hall JC, Rosbash M. CRY, a *Drosophila* clock and light-regulated cryptochrome, is a major contributor to circadian rhythm resetting and photosensitivity. *Cell*. 1998; 95:669–679. [PubMed: 9845369]
161. Emery P, Stanewsky R, Helfrich-Forster C, Emery-Le M, Hall JC, Rosbash M. *Drosophila* CRY is a deep brain circadian photoreceptor. *Neuron*. 2000; 26:493–504. [PubMed: 10839367]
162. Fogle KJ, Parson KG, Dahm NA, Holmes TC. CRYPTOCHROME is a blue-light sensor that regulates neuronal firing rate. *Science*. 2011; 331:1409–1413. DOI: 10.1126/science.1199702 [PubMed: 21385718]
163. Fogle KJ, Baik LS, Houl JH, Tran TT, Roberts L, Dahm NA, Cao Y, Zhou M, Holmes TC. CRYPTOCHROME-mediated phototransduction by modulation of the potassium ion channel beta-subunit redox sensor. *Proc Natl Acad Sci U S A*. 2015; 112:2245–2250. DOI: 10.1073/pnas.1416586112 [PubMed: 25646452]

Highlights

- Cellular redox state undergoes a ~24-h (circadian) oscillation
- Circadian redox oscillation in the SCN regulates neuronal excitability
- Redox mechanisms are involved in the regulation of ion channels
- Redox oscillation may play a role in global regulation of cellular excitability

Figure 1.

The mammalian circadian clock is located in the suprachiasmatic nucleus (SCN) of the anterior hypothalamus. *Upper:* In this fresh coronal section of the brain of a rat, the SCN lies near the base (within the dashed box). *Lower:* Vasoactive intestinal peptide (VIP) immunoreactive staining shows the pair of small hypothalamic nuclei is positioned medially, on either side of the third ventricle (3V), and directly superior to the heavily myelinated optic chiasm (OC).

Figure 2.

Cellular redox state is determined by several redox cofactor pairs. Redox state can be evaluated by the balance of small molecule redox cofactor pairs that are sensitive to the oxidizing/reducing potentials of the cell. These reversible reactions are regenerated enzymatically. Depicted are glutathione disulfide (GSSG)/glutathione (GSH), nicotinamide adenine dinucleotide (NAD⁺/NADH), nicotinamide adenine dinucleotide phosphate (NADP⁺/NADPH), and dehydroascorbic acid (DHA)/ascorbic acid (AA).

Figure 3.

Interactions between cellular metabolism, the circadian molecular clock, and ion channels. The core molecular clock consists of positive elements, such as circadian locomotor output kaput (CLOCK) and brain and muscle ARNT-like 1 (BMAL1), that positively regulate the expression of PERs and CRYs, which comprise negative elements, repressing the activity of CLOCK and BMAL1 [26]. An additional feedback loop involves the BMAL1/CLOCK-mediated transcription of REV-ERB and RAR-related orphan receptor (ROR) proteins that inhibit and initiate *Bmal1* transcription, respectively. Casein kinase 1 (CK1) enzymes contribute to timekeeping through phosphorylation and destabilization of PER. Reduced forms of NAD(H) and NADP(H) enhance DNA-binding activity of BMAL1 and CLOCK, while oxidized NAD⁺ and NADP⁺ inhibit their DNA binding [80]. Nicotinamide phosphoribosyltransferase (NAMPT) is a rate-limiting enzyme in the NAD⁺ biosynthetic pathway and is the driving force for rhythmic levels of cellular NAD⁺ which, in turn, activate NAD⁺-dependent histone deacetylases, sirtuin 1 (SIRT1) and sirtuin 3 (SIRT3) [82,83]. SIRT1 has been found to exhibit circadian oscillatory activity and alter clock gene expression. SIRT3 is located in the mitochondria where it mediates metabolic processes. Heme is a known sensor of cellular redox state that binds to nuclear receptors REV-ERB α and REV-ERB β and modulates their activity [87,88]. AMP-activated protein kinase (AMPK) is an important sensor of cellular energy state activated by a high AMP/ATP ratio. AMPK mediates phosphorylation of CRY and CK1 [91]. Phosphorylation of CRY marks it for degradation and phosphorylation of CK1 increases its activity and accelerates degradation of PER2. The balance between ROS levels and antioxidant defense systems, such as peroxiredoxins (Prx) and glutathione (GSH), contribute to the overall redox state. Ca²⁺, K⁺, and Na⁺ ion channels have been shown to be sensitive to redox modifications.

Figure 4.

Day-night oscillation of the redox environment modulates SCN neuronal excitability. In the relatively reduced daytime environment, more K⁺ channels close and the neuronal membrane becomes more depolarized, bringing the membrane potential closer to threshold for action potentials. In the relatively oxidized nighttime environment, more K⁺ channels open and the neuronal membrane becomes hyperpolarized [2]. Some voltage-gated K⁺ channels have been shown to have oxidoreductase Kvβ subunits that bind the redox cofactor NADPH, which determines gating dynamics [101]. This confers sensitivity to the local redox environment.