Published in final edited form as:

Cardiol Young. 2018 May ; 28(5): 697–701. doi:10.1017/S1047951118000045.

Early Post-Operative Remodeling Following Repair of Tetralogy of Fallot Utilizing Unsedated Cardiac Magnetic Resonance: a Pilot Study

Michael P DiLorenzo^a, Elizabeth Goldmuntz^a, Susan C Nicolson^b, Mark A Fogel^a, and Laura Mercer-Rosa^a

^aDivision of Cardiology, Department of Pediatrics, The Children's Hospital of Philadelphia, 3401 Civic Center Blvd, Philadelphia, PA 19104, USA

^bDepartment of Anesthesia, The Children's Hospital of Philadelphia, 3401 Civic Center Blvd, Philadelphia, PA 19104, USA

Abstract

Introduction—The right ventricular adaptations early after surgery in infants with tetralogy of Fallot are important to understand the changes that occur later on in life; this physiology has not been fully delineated. We sought to assess early post-operative right ventricular remodeling in patients with tetralogy of Fallot by cardiac magnetic resonance imaging.

Materials and Method—Subjects with tetralogy of Fallot under 1 year of age were recruited following complete surgical repair for tetralogy of Fallot. Protocol-based cardiac magnetic resonance imaging to assess anatomy, function, and flows was performed before hospital discharge using the Feed and Sleep technique, an unsedated imaging technique.

Results—Magnetic resonance imaging was completed in 16 subjects at a median age of 77 days (Interquartile range 114). There was normal ventricular ejection fraction and indexed right ventricular end-diastolic volume (48 ± 13 cc/m²), but elevated right ventricular mass (Z-score 6.2 ± 2.4). Subjects requiring a transannular patch or right ventricle to pulmonary artery conduit had moderate pulmonary insufficiency (Regurgitant fraction 27 ± 16).

Discussion—Early right ventricular remodeling after surgical repair for tetralogy of Fallot is characterized by significant pulmonary regurgitation, right ventricular hypertrophy, and lack of dilation. Performing cardiac magnetic resonance imaging using the Feed and Sleep technique is

Corresponding Author: Michael DiLorenzo, MD MSCE, 3959 Broadway, 2 North, New York, NY 10032, USA. Tel: (917) 370-5255, mpd2001@columbia.edu.

CONFLICTS OF INTEREST

None.

ETHICAL STANDARDS

The authors assert that all procedures contributing to this work comply with the ethical standards of Ethical Advisory Board and the National Institute of Health, and with the Helsinki Declaration of 1975, as revised in 2008, and has been approved by the institutional committee (Institutional Review Board, Children's Hospital of Philadelphia). There was no animal experimentation in this study.

Statement on the welfare of animals

This article does not contain any studies with animals performed by any of the authors.

Informed consent

Informed consent was obtained from all individual participants included in the study.

feasible in infants younger than five months. These results might open new avenues to study longitudinal right ventricular changes in tetralogy of Fallot and to further explore the utility of unsedated magnetic resonance imaging in patients with other types of congenial heart defects.

Keywords

Tetralogy of Fallot; Congenital heart disease; Cardiac Magnetic Resonance Imaging; Right ventricle

INTRODUCTION

Despite excellent long-term survival rates,^{1,2} most patients with tetralogy of Fallot are left with substantial pulmonary regurgitation following surgical repair with its consequent long-term deleterious effects.^{3,4} Cardiac MRI is the gold standard for evaluating anatomy, ventricular function, right ventricular mass, and valvar insufficiency following surgical repair for tetralogy of Fallot and has been utilized mostly during long-term follow up to guide the decision to intervene on the pulmonary valve.⁵⁻¹⁰ Although perioperative outcomes have been described in tetralogy of Fallot, there is little information about the early ventricular adaptations, including pulmonary regurgitation, which occur immediately following surgical repair and may be important to understanding the changes which occur later in life and put the magnitude of these alterations into perspective.¹¹⁻¹³ The Feed and Sleep technique is an anesthesia-free method for performing cardiac MRI, and has been used in select patients with proven utility in the assessment of intra- and extra-cardiac anatomy in certain congenital heart defects.¹⁴⁻¹⁶

The objectives of our study were to investigate right ventricle remodeling in terms of right ventricular function, pulmonary regurgitation, pulmonary artery anatomy, and residual lesions in the early post-operative period in patients operated for tetralogy of Fallot and to investigate the feasibility of the Feed and Sleep technique to obtain this information in post-operative neonates and infants.

MATERIALS AND METHOD

Subjects were recruited sequentially following complete surgical repair for tetralogy of Fallot between March 2013 and January 2016. Institutional Review Board approval and parental informed consent were obtained prior to cardiac MRI exams. The medical history was reviewed to assess patient safety prior to undergoing unsedated cardiac MRI. Inclusion criteria included the diagnosis of tetralogy of Fallot and under one year of age at time of enrollment. Patients requiring mechanical ventilation, medically unstable at scheduled time of cardiac MRI, or those with contraindications to cardiac MRI, such as pacemakers, were excluded. Complete surgical repair was defined as ventricular septal defect closure with or without right ventricular outflow tract intervention (transannular patch or right ventricle to pulmonary artery conduit). Protocol-based cardiac MRIs were performed with the Feed and Sleep technique just prior to hospital discharge, at a time when the patient was deemed stable to undergo the study without sedation. If the patient was discharged on a weekend prior to scheduling the cardiac MRI, the test was performed as soon as possible after hospital discharge.

The cardiac MRI protocol used has been previously described.¹³ Briefly, subjects were fed 3 to 4 hours prior to the planned initiation of the procedure and fasted up until their cardiac MRI. Immediately prior to the procedure, electrocardiographic leads were placed, the patient was fed, swaddled in an infant blanket, and placed in a vacuum immobilizer (MedVac bag, CFI Medical Solutions/Contour Fabricators, Fenton, MI, USA). After a pulse oximeter probe and ear muffs were applied, the lights were dimmed in the scanner to minimize stimuli. The study proceeded once the patient fell asleep or remained motionless. Cardiac MRI sequences were prioritized to assess anatomy (static steady state free precession axial imaging), ventricular function (cine steady state free precession 4-chamber and short axis), and flow assessments using phase contrast cardiac MRI across the aortic root, main and branch pulmonary arteries. An exam was considered complete if the intracardiac anatomy, right and left ventricular ejection fraction, and pulmonary regurgitant fraction could be assessed. Contrast was not utilized as part of the study protocol since subjects were preparing for hospital discharge and therefore did not have intravenous lines. Pulmonary regurgitation was considered moderate if the regurgitant fraction was > 20% and severe if the regurgitant fraction was >40%. Right ventricular mass Z-score was calculated based on normative data in children and main, right, and left pulmonary artery dimensions were indexed to (Body Surface Area)^{0.5} based on the linear relationship between vascular size and this indexing method.^{17,18}

Statistical analysis

Continuous variables were described using mean with standard deviation or median with interquartile range, if not normally distributed. Categorical variables were described with counts and percentages. For all analyses, statistical significance was indicated by a two-sided p value <0.05. Pearson or Spearman correlation coefficients were determined to assess the association between cardiac MRI variables.

RESULTS

Baseline data

Cardiac MRI was attempted in 21 subjects. Baseline demographics for these patients are listed in Table 1. Five patients did not complete the study due to inability to remain motionless for the entire exam. Sixteen patients (76%) completed the cardiac MRI protocol using the Feed and Sleep technique. Of those, 9 (60%) were asleep for the entire exam, 2 (13%) were asleep for part of the exam, and 4 (27%) were awake throughout the exam. One patient received sedation for another procedure prior to cardiac MRI, which was then performed during recovery. The median age at cardiac MRI for the entire group (n=21) was 123 days (Interquartile range 121) with a mean weight of 5.2 ± 1.7 kg. Those that could not complete the study were older at the time of attempted cardiac MRI as compared to those who could complete the study [174 days (Interquartile range 23) vs. 77 (Interquartile range 114), respectively, p=0.03].

Right ventricular remodeling

Among the 16 patients who completed the study, there was overall normal ventricular function, indexed right ventricular volumes, and right ventricular volume Z-scores. There

was on average moderate pulmonary regurgitation among the 11 subjects requiring right ventricular outflow tract intervention (transannular patch or right ventricle to pulmonary artery conduit) (Regurgitant fraction $27 \pm 16\%$, Table 2). Of this group, four (36%) had moderate regurgitation and two (11%) had severe regurgitation. Subjects requiring only ventricular septal defect closure had mild regurgitation. There was overall significant right ventricular hypertrophy (Right ventricular mass Z-score 6.2 ± 2.4 , Table 2). Subjects had a well-balanced circulation with a pulmonary to systemic blood flow ratio of 0.9 ± 0.2 . A small residual ventricular septal defect was present in 3 of 10 subjects. There was no association between pulmonary regurgitant fraction and indexed right ventricular end diastolic volume ($r=0.19$, $p=0.5$) or right ventricular mass Z-score ($r=-0.19$, $p=0.5$).

DISCUSSION

Pulmonary regurgitation resulting from surgical repair is a main determinant of outcome in tetralogy of Fallot, in particular with regard to its deleterious effect on the right ventricle. Characterization of right ventricular remodeling by cardiac MRI early after surgical repair for tetralogy of Fallot has not been described. In this study we sought to investigate the early right ventricular remodeling in patients operated for tetralogy of Fallot and to test the feasibility of unsedated cardiac MRI to obtain this information. The study found that neonates and infants have moderate pulmonary regurgitation and increased right ventricular mass with normal volumes and function. The Feed and Sleep technique allows for complete cardiac assessment in tetralogy of Fallot patients less than five months of age in the early post-operative period.

Right ventricular remodeling

There was overall moderate pulmonary regurgitation among those requiring a transannular patch or right ventricle to pulmonary artery conduit. Other reports have also demonstrated moderate pulmonary regurgitation in children and adolescents with tetralogy of Fallot.^{5,8,19,20} The determinants of pulmonary regurgitation are multifactorial,^{21,22} and include right ventricular compliance, pulmonary vascular resistance, heart rate and orifice size. While we cannot comment on the longitudinal changes in pulmonary regurgitation based on these comparisons, our findings would suggest that the degree of pulmonary regurgitation might be established early post-operatively. Longitudinal assessment of pulmonary regurgitation is thus necessary to further elaborate changes long-term.

Right ventricular mass was significantly increased in the early post-operative period, which is a reflection of insufficient time for right ventricular remodeling in response to relief of the outflow tract obstruction; Right ventricular hypertrophy is thus most likely a reflection of the pre-operative physiology. Right ventricular mass could be an important parameter to monitor in this patient population, and follow-up studies to determine the longitudinal changes in right ventricular mass, and physiologic drivers of these changes, are necessary.

Right ventricular volumes were within normal limits in our cohort, most likely due to the early interval between surgery and cardiac MRI. As such, there was no association between right ventricular end-diastolic volume and pulmonary regurgitant fraction. This may suggest

that the right ventricle does not dilate immediately, but rather remodels over the course of a number of years.

Even though on average mild residual disease was present early postoperatively, some patients demonstrated significant pulmonary regurgitation, and they may be at risk for accelerated right ventricular dilation and early dysfunction and should therefore be carefully monitored.

This study determined a number of parameters that can be used as a reference. Pulmonary artery size and architecture was obtained, which can be utilized as a baseline for future assessments; branch pulmonary artery stenosis is a potentially manageable residual lesion that has deleterious effects if not addressed.^{22,23} Right ventricular function was also obtained early after surgery. Better understanding of the early right ventricular remodeling after tetralogy of Fallot repair may provide insight into post-operative management strategies and guide follow-up. Early post-operative cardiac MRI could be of potential clinical use, as it can establish the burden of residual lesions, define the branch pulmonary artery anatomy, and help identify patients that could benefit from early follow up and interventional procedures.

Feed and Sleep technique

The Feed and Sleep technique provides a complete cardiac assessment in post-operative neonates and infants with tetralogy of Fallot. This technique was previously shown to be feasible in brain and extra-cardiac MRIs, as well as in cardiac MRIs that had specific anatomic questions.^{14–16,24} To our knowledge, our study is the first attempt to extend the use of the Feed and Sleep technique to perform a comprehensive cardiac MRI evaluation in tetralogy of Fallot. A complete dataset was obtained in approximately 75% of cardiac MRIs in this study. The failure rate is multifactorial, and includes greater time to obtain a full protocol, attempting Feed and Sleep in an older age group, and having all of our patients scanned prior to discharge. Indeed, three of the infants who failed Feed and Sleep had an attempted cardiac MRI within 5 days of surgery, so it is possible that post-operative pain may have contributed to an inability to fall asleep or remain motionless during the scan. Because our study protocol sought to perform cardiac MRI prior to discharge, the window for performing a study for research purposes was quite narrow and dependent on both the patient's stability and schedule availability.

This study demonstrates that the Feed and Sleep technique is feasible in post-operative patients that are younger than 5 months of age. One advantage of this technique is that sedation is not required, and although sedation is generally safe, it does carry some risks. In addition, the Feed and Sleep allows for normal physiology to be present (which could be distorted by sedation with or without intubation and mechanical ventilation), is relatively fast, and complete datasets can be obtained using the duration of the child's nap.^{24,25} Although we did not investigate the cost-effectiveness of this technique, a non-sedated protocol is probably cost-effective, given that the required support staff is limited to the bedside nurse and cardiac MRI personnel, and post-procedure recovery is unnecessary.¹⁶

Limitations

This was a prospective pilot study; therefore, we are limited by a small sample size. Thus, we cannot comment on the statistical significance of differences in pulmonary regurgitation or right ventricular volume based on anatomy or type of surgical repair due to a lack of power. However, the goal of the study was to demonstrate the right ventricular changes that occur early after tetralogy of Fallot repair and to test the feasibility of the Feed and Sleep technique. Further studies with larger sample sizes are necessary to address these questions. Because this analysis was restricted to patients who were clinically stable and ready for discharge, our sample may not have included those with more significant disease who may not have been stable for examination, or those who had less significant disease and early hospital discharge.

Conclusions

Early right ventricular remodeling after surgical repair for tetralogy of Fallot is characterized by significant pulmonary regurgitation, right ventricular hypertrophy, and lack of dilation. Longitudinal examination in this cohort could provide further insight into right ventricular remodeling and help identify a subset of children at particular risk for deleterious consequences. Unsedated cardiac MRI is a feasible approach for infants less than 5 months of age and could be considered in other forms of congenital heart defects.

Acknowledgments

None

FINANCIAL SUPPORT

This work was supported by the National Institutes of Health [M.P.D, NIH grants T32 HL007915; L.M.R, K01HL125521], and Pulmonary Hypertension Association supplement to K01HL125521 (L.M.R).

References

1. Murphy JG, Gersh BJ, Mair DD, et al. Long-term outcome in patients undergoing surgical repair of tetralogy of Fallot. *N Engl J Med*. 1993; 329:593–599. [PubMed: 7688102]
2. Nollert G, Fischlein T, Bouterwek S, Böhmer C, Klinner W, Reichart B. Long-term survival in patients with repair of tetralogy of Fallot: 36-year follow-up of 490 survivors of the first year after surgical repair. *J Am Coll Cardiol*. 1997; 30:1374–1383. [PubMed: 9350942]
3. Bouzas B, Kilner PJ, Gatzoulis MA. Pulmonary regurgitation: not a benign lesion. *Eur Heart J*. 2005; 26:433–439. [PubMed: 15640261]
4. Gatzoulis MA, Balaji S, Webber SA, et al. Risk factors for arrhythmia and sudden cardiac death late after repair of tetralogy of Fallot: a multicentre study. *Lancet*. 2000; 356:975–981. [PubMed: 11041398]
5. Geva T, Sandweiss BM, Gauvreau K, Lock JE, Powell AJ. Factors associated with impaired clinical status in long-term survivors of tetralogy of Fallot repair evaluated by magnetic resonance imaging. *J Am Coll Cardiol*. 2004; 43:1068–1074. [PubMed: 15028368]
6. Kilner PJ, Geva T, Kaemmerer H, Trindade PT, Schwitter J, Webb GD. Recommendations for cardiovascular magnetic resonance in adults with congenital heart disease from the respective working groups of the European Society of Cardiology. *Eur Heart J*. 2010; 31:794–805. [PubMed: 20067914]

7. Knauth AL, Gauvreau K, Powell AJ, et al. Ventricular size and function assessed by cardiac MRI predict major adverse clinical outcomes late after tetralogy of Fallot repair. *Heart*. 2008; 94:211–216. [PubMed: 17135219]
8. Niezen RA, Helbing WA, Van der Wall EE, Van der Geest RJ, Rebergen SA, De Roos A. Biventricular systolic function and mass studied with MR imaging in children with pulmonary regurgitation after repair for tetralogy of Fallot. *Radiology*. 1996; 201:135–140. [PubMed: 8816534]
9. Villafañe J, Feinstein JA, Jenkins KJ, et al. Hot Topics in Tetralogy of Fallot. *J Am Coll Cardiol*. 2013; 62:2155–2166. [PubMed: 24076489]
10. Rebergen SA, Chin JG, Ottenkamp J, Van der Wall EE, De Roos A. Pulmonary regurgitation in the late postoperative follow-up of tetralogy of Fallot. Volumetric quantitation by nuclear magnetic resonance velocity mapping. *Circulation*. 1993; 88:2257–2266. [PubMed: 8222120]
11. Pigula FA, Khalil PN, Mayer JE, Pedro J, Jonas RA. Repair of tetralogy of Fallot in neonates and young infants. *Circulation*. 1999; 100:II–157. [PubMed: 10567300]
12. Sachdev MS, Bhagyavathy A, Varghese R, Coelho R, Kumar RS. Right Ventricular Diastolic Function After Repair of Tetralogy of Fallot. *Pediatr Cardiol*. 2006; 27:250–255. [PubMed: 16391981]
13. Mercer-Rosa L, Pinto N, Yang W, Tanel R, Goldmuntz E. 22q11.2 Deletion syndrome is associated with perioperative outcome in tetralogy of Fallot. *J Thorac Cardiovasc Surg*. 2013; 146:868–873. [PubMed: 23312975]
14. Fogel MA, Pawlowski TW, Harris MA, et al. Comparison and Usefulness of Cardiac Magnetic Resonance Versus Computed Tomography in Infants Six Months of Age or Younger With Aortic Arch Anomalies Without Deep Sedation or Anesthesia. *Am J Cardiol*. 2011; 108:120–125. [PubMed: 21679782]
15. Shariat M, Mertens L, Seed M, et al. Utility of Feed-and-Sleep Cardiovascular Magnetic Resonance in Young Infants with Complex Cardiovascular Disease. *Pediatr Cardiol*. 2015; 36:809–812. [PubMed: 25527227]
16. Windram J, Grosse-Wortmann L, Shariat M, Greer M-L, Crawford MW, Yoo S-J. Cardiovascular MRI without sedation or general anesthesia using a feed-and-sleep technique in neonates and infants. *Pediatr Radiol*. 2012; 42:183–187. [PubMed: 21861089]
17. Buechel E, Kaiser T, Jackson C, Schmitz A, Kellenberger CJ. Normal right- and left ventricular volumes and myocardial mass in children measured by steady state free precession cardiovascular magnetic resonance. *J Cardiovasc Magn Reson*. 2009; 11:19. [PubMed: 19545393]
18. Sluysmans T. Theoretical and empirical derivation of cardiovascular allometric relationships in children. *J Appl Physiol*. 2005; 99:445–457. [PubMed: 15557009]
19. Lee C, Lee C-H, Kwak JG, et al. Factors associated with right ventricular dilatation and dysfunction in patients with chronic pulmonary regurgitation after repair of tetralogy of Fallot: analysis of magnetic resonance imaging data from 218 patients. *J Thorac Cardiovasc Surg*. 2014; 148:2589–2595. [PubMed: 25173118]
20. Mercer-Rosa L, Paridon SM, Fogel MA, et al. 22q11.2 Deletion Status and Disease Burden in Children and Adolescents With Tetralogy of Fallot. *Circ Cardiovasc Genet*. 2015; 8:74–81. [PubMed: 25561045]
21. Chaturvedi RR, Redington AN. Pulmonary regurgitation in congenital heart disease. *Heart*. 2007; 93:880–889. [PubMed: 17569817]
22. Redington AN. Determinants and Assessment of Pulmonary Regurgitation in Tetralogy of Fallot: Practice and Pitfalls. *Cardiol Clin*. 2006; 24:631–639. [PubMed: 17098516]
23. Harris MA, Whitehead KK, Gillespie MJ, et al. Differential Branch Pulmonary Artery Regurgitant Fraction Is a Function of Differential Pulmonary Arterial Anatomy and Pulmonary Vascular Resistance. *JACC Cardiovasc Imaging*. 2011; 4:506–513. [PubMed: 21565739]
24. Rangamani S, Varghese J, Li L, et al. Safety of cardiac magnetic resonance and contrast angiography for neonates and small infants: a 10-year single-institution experience. *Pediatr Radiol*. 2012; 42:1339–1346. [PubMed: 22885603]
25. Dorfman A, Odegard K, Powell A, Laussen P, Geva T. Risk Factors for Adverse Events During Cardiovascular Magnetic Resonance in Congenital Heart Disease. *J Cardiovasc Magn Reson*. 2007; 9:793–798. [PubMed: 17891617]

Table 1

Baseline demographic data

n = 21*	n (%)	Mean ± SD/Median (IQR)
Male	15 (71)	
Anatomy		
Pulmonary Stenosis	17 (81)	
Pulmonary Atresia	4 (19)	
Genetic Syndrome	9 (43)	
Age at surgery (days)		114 (128)
Age at cardiac MRI (days)		123 (121)
Time from surgery to cardiac MRI (days)		8 (15)
Weight at cardiac MRI (kg)		5.2 ± 1.7
Surgical repair- RV outflow tract		
Transannular Patch	10 (48)	
RV to pulmonary artery conduit	3 (14)	
None	8 (38)	

MRI, Magnetic Resonance Imaging; RV, Right ventricle; SD, Standard Deviation; IQR, Interquartile Range

* Includes one patient recovering from sedation during the start of the cardiac MRI

Table 2

Cardiac MRI data.

n = 16	n (%)	Mean $\pm$ SD/Median (IQR)
Length of cardiac MRI scan (minutes)	16	38 (8)
RV ejection fraction (%)	16	62 $\pm$ 10
Transannular patch	8 (50)	64 $\pm$ 10
RV to pulmonary artery conduit	3 (19)	54 $\pm$ 3
Ventricular septal defect only	5 (31)	64 $\pm$ 10
LV ejection fraction (%)	16	70 $\pm$ 8
Transannular patch	8 (50)	71 $\pm$ 11
RV to pulmonary artery conduit	3 (19)	69 $\pm$ 4
Ventricular septal defect only	5 (31)	69 $\pm$ 5
Indexed RV end-diastolic volume (cc/m²)	16	48 $\pm$ 13
Transannular patch	8 (50)	49 $\pm$ 15
RV to pulmonary artery conduit	3 (19)	48 $\pm$ 3
Ventricular septal defect only	5 (31)	45 $\pm$ 14
RV end-diastolic volume Z-score	16	0.8 $\pm$ 2.4
Transannular patch	8 (50)	1.4 $\pm$ 3
RV to pulmonary artery conduit	3 (19)	1.1 $\pm$ 0.7
Ventricular septal defect only	5 (31)	-0.4 $\pm$ 2.8
Indexed RV end-systolic volume (cc/m²)	16	18 $\pm$ 8
Indexed LV end-diastolic volume (cc/m²)	16	39 $\pm$ 12
Transannular patch	8 (50)	36 $\pm$ 6
RV to pulmonary artery conduit	3 (19)	35 $\pm$ 5
Ventricular septal defect only	5 (31)	47 $\pm$ 19
Indexed RV mass (grams/m²)	16	25.6 $\pm$ 9.6
RV mass Z-score	16	6.2 $\pm$ 2.4
Transannular patch	8 (50)	6.3 $\pm$ 2.4
RV to pulmonary artery conduit	3 (19)	7.3 $\pm$ 3.9
Ventricular septal defect only	5 (31)	5.3 $\pm$ 1.0
Pulmonary Regurgitant fraction	16	
Transannular patch	8 (50)	29 $\pm$ 18
RV to pulmonary artery conduit	3 (19)	21 $\pm$ 9.6
Ventricular septal defect only	5 (31)	1 $\pm$ 1.6
Right pulmonary artery flow (% of total)	16	58 $\pm$ 14
Pulmonary to systemic blood flow ratio	16	0.9 $\pm$ 0.2

n = 16	n (%)	Mean ± SD/Median (IQR)
Indexed pulmonary artery size (mm/BSA^{0.5})	16	
Right pulmonary artery		9.3 ± 4.7
Left pulmonary artery		7 ± 4.4
Main pulmonary artery		20 ± 6.1

MRI, Magnetic Resonance Imaging; RV, Right ventricle; LV, Left Ventricle; BSA, Body Surface Area; SD, Standard Deviation; IQR, Interquartile range