CORRESPONDENCE

Fitness to Drive in Cardiovascular Disease

by Prof. Dr. med. Hermann H. Klein, Prof. Dr. med. Udo Sechtem, and Prof. Dr. med. Hans-Joachim Trappe in issue 41/2017

The Expert's Duty of Confidentiality

The well structured and informative article on fitness to drive by Klein et al. (1) unfortunately contains an incorrect and at best misleading comment on medical confidentiality where an expert report is required.

At the end of the first paragraph on page 696 of the article (1), the authors say: "If the patient asks the physician to produce a report on his or her fitness to drive, the obligation to maintain confidentiality is of course removed." This is not the case. The patient as the person who commissions the report remains in charge. If, for example, the expert report contains information that can have a negative effect for the patient, and he or she decides not to present this report to third parties, the doctor's duty of confidentiality remains unchanged. In such a scenario, the doctor writing the report is not authorized to inform third parties—for example, the responsible authorities—of the negative conclusion concerning for instance the patient's driving ability in his/her report. The patient remains at leisure to find a new expert, who—in borderline cases—might come to another conclusion that is more favorable for the patient.

DOI: 10.3238/arztebl.2018.0188a

References

1. Klein HH, Sechtem U, Trappe HJ: Fitness to drive in cardiovascular disease. *Dtsch Arztebl Int* 2017; 114: 692–702.

Dr. med. Thomas Schneider
Seehausen

Conflict of interest statement

The author declares that no conflict of interest exists.

Fitness to Drive After Sternotomy

The CME article (1) provides a risky recommendation for patients after cardiac surgery involving sternotomy in Table 4. The table says that after cardiac surgery, patients are fit to drive 2–4 weeks after the intervention. This may be sufficient for a cardiovascular system that is healing without complications, but it is not a sufficiently long period for the sternum to heal. The time period in the table is not long enough to guarantee a sufficiently high degree of stability of the bone for the driver to look over their shoulder.

I know several patients who drove their car to a hospital for follow-up treatment/rehabilitation between 14 and 21 days after cardiac surgery. Later they returned with similar narratives: "Driving was not a problem, but parking while looking over my shoulder in the rehab venue's car park caused a cracking noise in my

thorax, and afterwards the two halves of my breastbone were loose and dislodged against each other."

Some such patients will have to undergo repeat surgery, whereas in others, conservative attempts can be made by using thoracic bandages to treat the problem. Yet others put up with a permanently loose and mobile sternum.

Three months after cardiac surgery, almost no breastbone will have completely healed, according to CT criteria (CT score for sternal healing in [2]; CT, computed tomography). After 6 months, a maximum of 7 in every 10 breast bones shows complete healing on CT examination (2).

From a clinical perspective, many such breastbones that have not completely healed according to the CT criteria are none the less strong and stable, and the patients have no symptoms. Still, the recommendation should be not to look over one's shoulder—which involves twisting the upper body—for a minimum of 6 weeks after cardiac surgery involving median sternotomy. This is a physical restriction that—according to what I know about driving lessons—does not allow for active driving.

In my long years' experience, 6 weeks are the absolute lower limit for a possible look over the shoulder—and even this only where the bone has been observed to be safely clinically stable during the healing period.

DOI: 10.3238/arztebl.2018.0118b

References

1. Klein HH, Sechtem U, Trappe HJ: Fitness to drive in cardiovascular disease. *Dtsch Arztebl Int* 2017; 114: 692–702.
2. Raman J, Lehmann S, Zehr K, et al.: Sternal closure with rigid plate fixation versus wire closure: a randomized controlled multicenter trial. *Ann Thorac Surg* 2012; 94: 1854–61.

Dr. med. Arnd Afflerbach

Klinik für Herz-, Thorax- und Gefäßchirurgie, St.-Johannes-Hospital
amd.afflerbach@joho-dortmund.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank both our correspondents for their constructive contributions.

Dr. Schneider points out that the sentence "If the patient asks the physician to produce a report on his or her fitness to drive, the obligation to maintain confidentiality is of course removed" is misleading. In all likelihood, Dr Schneider refers to the guidelines for expert reports issued by the Federal Highway Research Institute (*Bundesanstalt für Straßenwesen*, BASt), which

says under section 2.3, “Legal position of the expert” that the person to be assessed/reported on is the expert’s client, contractual partner, and party liable for costs. The expert report may be passed to the authorities or third parties only with the explicit agreement of that person—otherwise medical confidentiality will prevent this.

In our article on this subject, we complied with the rules and regulations for the licensing of drivers in Germany (*Fahrerlaubnisverordnung*, FeV). These regulate the approach of the licensing authorities for assessing the affected person’s fitness to drive in § 11, “Fitness.” If the licensing authorities are in doubt about a person’s fitness to drive, they can commission a report to help them come to a decision. The affected person is obliged to commission at their own expense and within a defined time frame an expert report on their own fitness to drive. They have to inform the licensing authorities about which institution has been commissioned to provide the report. If the requested report is not presented by the deadline, the authorities may conclude that the affected person is unfit to drive. The person is entitled to view the transmitted documents. Since the affected person is the one who commissioned the expert report for the licensing authorities, s/he therefore agrees with this approach. It is not difficult to see that the requirements of the BASt and the FeV regarding expert reports for the licensing authorities diverge substantially. As far as we can find out, this discrepancy has existed for years and can be resolved only by the Ministry of Transport, which is responsible for the guidelines on expert reports of both the FeV and the BASt. From a legal perspective, the FeV as a statutory instrument is superior to the guidelines of the BASt.

Matters are different if a client (a private company or authority) requests an expert report on an employee’s fitness to drive. In this scenario, the expert should be released from their duty of confidentiality by the affected person if the expert report is sent to the client.

In his letter, Dr. Afflerbach points out that in Table 4 in the article, drivers in group 1 are described as fit to drive, 2–4 weeks after sternotomy. He claims that they will be fit to drive after 6 weeks at the earliest.

Unpublished guidelines for expert reports on fitness to drive from the European Union (New Standards for Driving and Cardiovascular Diseases) of 2013 state that no specific European guidelines exist that stipulate when a patient is fit to drive after cardiac surgery. Dr. Afflerbach’s correspondence was extremely compelling, and it therefore seems to make sense to assume fitness to drive only 6–8 weeks after successful cardiac surgery. The likelihood with which other participants in road traffic will be harmed if someone returns to driving—appropriate to their condition—as early as 2–4 weeks after sternotomy is impossible to answer. Furthermore, such patients will mostly be in rehabilitation clinics.

DOI: 10.3238/arztebl.2018.0188c

References

1. Klein HH, Sechtem U, Trappe HJ: Fitness to drive in cardiovascular disease. *Dtsch Arztebl Int* 2017; 114: 692–702.

On behalf of the authors

Prof. Dr. med. Hermann H. Klein

Idar-Oberstein
hermann@klein-devries.de

Conflict of interest statement

Prof Klein has received travel expenses and speaker honoraria from Kel-Con.

CLINICAL SNAPSHOT

Leukemia Cutis

A 66-year-old man complained of skin changes that had not responded to treatment. He had first noticed them three years before; they had progressed in recent months, and his general condition had declined. He was under treatment with systemic glucocorticoids and other drugs for suspected sarcoidosis. On physical examination, there were multiple indolent erythematous nodules at many sites on his skin. Histopathological examination of a biopsy specimen revealed an interstitial infiltrate of atypical lymphocytes; laboratory testing revealed a massive leukocytosis (96 000/ μ L). We were able to cytogenetically demonstrate the Philadelphia chromosome and to establish the initial diagnosis of chronic myeloid leukemia (CML). Cutaneous findings in patients with CML may include nodules, plaques, vesicles, or ulcers. The simultaneous cutaneous and systemic manifestation of the disease has been described in as many as 23% of patients; approximately 7% of patients with cutaneous infiltration do not have leukemia in the blood. In such cases, skin biopsy is essential for the establishment of the diagnosis.

Prof. Dr. med. Joachim Dissemond, Klinik und Poliklinik für Dermatologie, Venerologie und Allergologie, Universitätsklinikum Essen, joachim.dissemond@uk-essen.de

Dr. med. Elvir Cesko, Praxis, Essen

Conflict of interest statement: The authors state that they have no conflict of interest.

Cite this as: Dissemond J, Cesko E: Leukemia cutis. *Dtsch Arztebl Int* 2018; 115: 189. DOI: 10.3238/arztebl.2018.0189

Translated from the original German by Ethan Taub, M.D.