Published in final edited form as:

J Neurogenet. 2016 ; 30(3-4): 259–275. doi:10.1080/01677063.2016.1255212.

K⁺ Channel Reorganization and Homeostatic Plasticity during Postembryonic Development: Biophysical and Genetic Analyses in Acutely Dissociated *Drosophila* Central Neurons

Taixiang Saur^{1,2,4,a}, I-Feng Peng^{3,b}, Peng Jiang^{1,2,c}, Neng Gong^{1,2}, Wei-Dong Yao^{4,d}, Tian-Le Xu^{1,2,e}, and Chun-Fang Wu^{1,2,3}

¹Department of Neurobiology and Biophysics, School of Life Sciences, University of Science and Technology of China, Hefei 230027, China

²Institute of Neuroscience and Key Laboratory of Neurobiology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China

³Department of Biology, University of Iowa, Iowa City, IA 52242 USA

⁴New England Primate Research Center, Harvard Medical School, Southborough, MA 01772 USA

Abstract

Intrinsic electric activities of neurons play important roles in establishing and refining neural circuits during development. However, how the underlying ionic currents undergo postembryonic reorganizations remains largely unknown. Using acutely dissociated neurons from larval, pupal, and adult *Drosophila* brains, we show drastic re-assemblies and compensatory regulations of voltage-gated (I_{Kv}) and Ca^{2+} -activated ($I_{K(Ca)}$) K^+ currents during postembryonic development. Larval and adult neurons displayed prominent fast inactivating I_{Kv} , mediated by the *Shaker* (*Sh*) channel to a large extent, while in the same neurons $I_{K(Ca)}$ was far smaller in amplitude. In contrast, pupal neurons were characterized by large sustained I_{Kv} and prominent $I_{K(Ca)}$, encoded predominantly by the *slowpoke* (*slo*) gene. Surprisingly, deletion of *Sh* in the *Sh^M* null mutant removed inactivating, transient I_{Kv} from large portions of neurons at all stages. Interestingly, elimination of *Sh* currents was accompanied by upregulation of non-*Sh* transient I_{Kv} . In comparison, the *slo^I* mutation abolished the vast majority of $I_{K(Ca)}$, particularly at the pupal stage. Strikingly, the deficiency of $I_{K(Ca)}$ in *slo* pupae was compensated by the transient component of I_{Kv} mediated by *Sh* channels. Thus, $I_{K(Ca)}$ appears to play critical roles in pupal development and its absence induces functional compensations from a specific transient I_{Kv} current. While mutants lacking either *Sh* or *slo* currents survived normally, *Sh*;*slo* double mutants deficient in both failed to survive through pupal metamorphosis. Together, our data highlight significant reorganizations

Correspondence to: Wei-Dong Yao; Tian-Le Xu; Chun-Fang Wu.

^aMcLean Hospital, Harvard Medical School, Belmont, MA 02478, USA

^bJoekai Biotechnology LLC, Kunshan City, Jiangsu Province 215300, China

^cDepartment of Developmental Neuroscience, Munroe-Meyer Institute, Holland Regenerative Medicine Program, University of Nebraska Medical Center, Omaha, NE 68198, USA

^dDepartments of Psychiatry and Neuroscience, State University of New York, Health Science Center at Syracuse, Upstate Medical University, Syracuse, NY 13210, USA

^eDiscipline of Neuroscience and Department of Anatomy, Histology, and Embryology, Collaborative Innovation Center for Brain Science, Shanghai Jiao Tong University School of Medicine, Shanghai 200025, China

and homeostatic compensations of K^+ currents during postembryonic development and uncover previously unrecognized roles for *Sh* and *slo* in this plastic process.

Introduction

The intrinsic membrane properties of a neuron, mediated by various voltage- and ligand-activated ion currents that determine its overall excitability state and firing patterns, give the neuron its functional identity. At the circuit level, intrinsic properties dictate the translation of dendritic input to axonal output, which can guide the establishment and remodeling of connections in neural circuits (Goodman and Shatz, 1993) while mediating directly the functioning of the circuits appropriate for specific developmental stages (Getting, 1989; Harris-Warrick and Marder, 1991). *Drosophila* metamorphosis offers a particularly extreme case of postembryonic development since during pupation, the central nervous system (CNS) undergoes a drastic reconstruction via processes that include elimination of certain larval neurons, modifications of the persisting ones, and proliferation and differentiation of adult-specific neurons to accommodate profound changes in body functions and behaviors (Truman, 1990; Kimura and Truma, 1990; Consoulas et al., 2002; Hebbar et al., 2006). Thus, *Drosophila* metamorphosis provides a unique system to investigate the genetic control of the transforming physiological properties of CNS neurons during this morphologically defined postembryonic developmental process.

K^+ currents are a key element of intrinsic properties regulating excitability, frequency coding and firing patterns of a neuron (Hille, 2001). K^+ currents in *Drosophila* neurons and muscle cells are characterized by two voltage-gated (I_{Kv}) currents, I_A (inactivating transient A-type current) and I_K (persistent delayed rectifier) (Salkoff & Wyman, 1983; Solc and Aldrich, 1988; Baker and Salkoff, 1990; Saito and Wu, 1991), and two Ca^{2+} -activated ($I_{K(Ca)}$) currents, fast I_{CF} and slow I_{CS} (Singh and Wu, 1989; Saito and Wu, 1991). Five genes are known to encode these currents in *Drosophila*: *Shaker* (*Sh*, I_A), *Shal* (I_A), *Shab* (slowly inactivating I_K), *Shaw* (non-inactivating I_K), and *slowpoke* (*slo*, $I_{K(Ca)}$) (Iverson et al., 1988; Timpe et al., 1988; Butler et al., 1989; Wei et al., 1990; Atkinson et al., 1991). Previous studies of *Sh* and *Shal* mutants suggest that *Sh* channels mediate I_A in embryonic, larval, and adult muscle (Salkoff and Wyman, 1981; Haugland and Wu, 1990; Brodie and Bate, 1993), whereas *Shal* encodes most I_{Kv} in embryonic, and possibly pupal, neurons, with little contribution from *Sh* (Baker and Salkoff, 1990; Tsunoda and Salkoff, 1995).

Electrophysiological analyses of cultured neurons from *Drosophila* embryos (Byerly and Leung, 1988; Saito and Wu, 1991; Tsunoda and Salkoff, 1995; Zhao et al., 1995; Yao and Wu, 1999; Peng and Wu, 2007a), larvae (Solc and Aldrich, 1988; Wright and Zhong, 1995), pupae (Baker and Salkoff, 1990), and adult brains (Wu et al., 2001) have provided in-depth understandings of neuronal K^+ current properties at the specific stages. However, this approach does not ensure a direct “profiling” of K^+ currents across distinct metamorphic stages due to the “extra” in vitro culturing periods and potential complications derived from different culture and growth conditions. Only limited studies have investigated the contribution of *Sh* to K^+ currents in larval and adult neurons in vivo (Choi et al., 2004;

Gasque et al., 2005). Additionally, the molecular identities that mediate neuronal $I_{K(Ca)}$ at the various stages still remain to be determined.

We addressed these questions using acutely dissociated neurons from *Drosophila* larval, pupal, and adult brains. This approach allowed direct comparisons of K^+ currents expressed in neurons from these distinct developmental stages. We show drastic reorganizations of both I_{K_V} and $I_{K(Ca)}$ during metamorphosis. Our data indicate that the Sh^M mutation affects a transient I_{K_V} in most (>60%) of larval and adult neurons, which was accompanied by an upregulation of remaining I_{K_V} , suggesting an unexpectedly large role for *Sh* in larval and adult neurons. In contrast, the slo^I mutation eliminated the vast majority of neuronal $I_{K(Ca)}$ from all stages, and the near complete loss of $I_{K(Ca)}$ was nevertheless compensated by increased I_{K_V} in pupal neurons. Our data provide the first “profile” of neuronal K^+ currents during *Drosophila* postembryonic development, assign contributions for *Sh* and *slo* to larval, pupal and adult K^+ currents, and highlight a form of robust neuronal plasticity that governs K^+ current re-assemblies and compensations during postembryonic development.

Materials and methods

Fly stocks

Drosophila stocks were reared on cornmeal-yeast-dextrose media at 21–23°C and ~ 60% relative humidity with 12 hr dark/light cycles. Canton S (CS) and Oregon R (OR) strains were used as wild type (WT) control. The *slowpoke* (slo^I) fly, which is an ethyl methanesulfonate (EMS)-induced temperature-sensitive mutant, becomes paralyzed at 38°C, as described previously (Elkins et al., 1986). The *Shaker* mutant allele, Sh^M , isolated in an X-ray-irradiation screen, is a null mutation due to an insertion of a 2.2 kb DNA fragment which eliminates all *Shaker* gene products (Kamb et al., 1987; Zhao et al., 1995). The Sh^M flies exhibit vigorous leg-shaking under ether anesthesia (Kaplan and Trout, 1969). The UAS-GFP; OK107-GAL4 flies, in which green fluorescent protein (GFP) expression is driven by a mushroom body (MB)-specific driver OK107-GAL4, were utilized to label MB neurons in order to examine the cell type-specific development of ionic currents in MB neurons. All strains were obtained from the Department of Biological Sciences, University of Iowa, Iowa, U.S.A. except that the UAS-GFP; OK107-GAL4 line (Li et al., 2013) was obtained from Dr. A.-K. Guo's laboratory at Institute of Neuroscience, Chinese Academy of Sciences, Shanghai.

Solution

The Ca^{2+} -, Mg^{2+} -free solution contained (mg/100ml): 800 NaCl, 20 KCl, 5 NaH_2PO_4 , 100 $NaHCO_3$ and 100 glucose; was pHed to 7.2–7.3 with 1 M HCl. Recording solution contained (mM): 130 NaCl, 5 KCl, 2.5 $CaCl_2$, 6 $MgCl_2$, 10 HEPES and 10 glucose. The composition of the patch pipette solution was (mM): 130 KCl, 4 $MgCl_2$, 10 EGTA, 10 HEPES, 1 $CaCl_2$ and 2 ATP. To isolate voltage-gated K^+ currents, 0.1 μM tetrodotoxin (TTX) was added to the Ca^{2+} -free solution (mM) (2.5 $CaCl_2$ was substituted by 2.5 $MgCl_2$, plus 0.5 EGTA) to eliminate voltage-gated Na^+ current (I_{Na}), voltage-gated Ca^{2+} current (I_{Ca}) and Ca^{2+} -activated K^+ current, respectively. Voltage-gated Ba^{2+} current (I_{Ba})-recording solution consisted of (mM): 40 $BaCl_2$, 50 NMDG, 40 TEA-Cl, 10 4-AP, 6 $MgCl_2$, 10

HEPES and 10 glucose. I_{Na}-recording solution contained (mM): 90 NaCl, 40 TEA-Cl, 10 4-AP, 6 MgCl₂, 10 HEPES and 10 glucose. All the recording solutions were adjusted to pH of 7.2–7.3 with Tris. The CsCl patch pipette solution contained (mM): 130 CsCl, 4 MgCl₂, 10 HEPES, 10 EGTA, 1 CaCl₂, and 2 ATP. All chemicals were obtained from Sigma.

Acute dissociation of *Drosophila* brain neurons

The developmental stages of *Drosophila* were defined by the criteria of Bainbridge and Bownes (Bainbridge and Bownes, 1981). L-III, EP, MP, LP and Ad represent third-instar larva, early-pupa (P5–8), middle-pupa (P8–11), late-pupa (P11–15) and adult (1–4d) stages, respectively. For the preparation of all stages, only the fly brains were dissected to generate the acutely dissociated neurons as reported previously (Wu et al., 2001) with some minor modifications. Briefly, the fly brains were dissected carefully with a pair of sharp twisters and were then incubated in Ca²⁺/Mg²⁺-free solution with 0.05 mg/ml collagenase IA for 15 min at 25°C. The treated brains were rinsed twice with recording solution and transferred into a culture dish, where the brains were mounted with a U-shaped grid made of metal filament and nylon fibers. A vibrator with a glass-dissecting micropipette was attached onto a manipulator, and the dissecting pipette was lightly placed on the surface of the mounted fly brains under a dissecting microscope (XTL-II, Tech Instrument Co. Ltd, Beijing, China). The neurons were dissociated by a series of horizontal movements (5 Hz, 100 µm) of the pipette tip, which were generated from a pulse generator (EE 1641B, Nanjing, China). GFP-labeled neurons derived from MBs were identified under the fluorescence microscope. Drugs were applied via a ‘Y-tube’ method in all experiments. The tip of the ‘Y-tube’ was positioned approximately 100 µm away from the neurons studied, allowing a complete exchange of external solution surrounding the neuron within 20 ms (Wu et al., 2001).

Whole-cell recordings

Whole-cell recordings were carried out under voltage-clamp conditions at 22–25°C. The small size of the neurons enabled us to attain giga-ohm seals (>1 GΩ) with or without a gentle suction. The resistance between the recording electrode filled with pipette solution and the reference electrode was 7–9 MΩ. Serial resistance compensation was not essential in these experiments, as reported in earlier studies on cultured larval neurons (Solc and Aldrich, 1988; Wright and Zhong, 1995; Yu et al., 1999). Data were collected with an Axopatch 200B patch-clamp amplifier (Axon Instruments, Foster City, CA) equipped with Digidata 1320A and Clampex (version 8.0.1, Axon Instruments). Clampfit 8.0.1 and Origin 7.0 (OriginLab) were used to analyze and plot the data. The currents were sampled at 5 kHz and filtered at 2 kHz. Unless otherwise mentioned, the membrane potential was held at –70 mV. Statistical analysis was performed using Student’s *t*-test or one-way ANOVA, as appropriate.

Survival rates

Flies of *Sh^M/XX/Y;;slo⁹⁸/TM6* genotype were allowed to lay eggs for one day. The genotypes of all F1 3rd instar larvae were determined. For each genotype, 25 male, 3rd instar larvae were collected (four repeats) for the examination of survivability through the pupal stage. In *Sh;;slo* double mutants, only about 10% of the larvae survived through the pupal stage to adulthood.

Real time PCR

An on-column DNase digestion was performed during RNA preparation to remove potential genomic DNA contamination (RNeasy Mini Kit Qiagen, Valencia, CA). Reverse transcription was carried out using the SuperScript III Kit (Invitrogen, Carlsbad, CA) with random primers. The copies of dsDNAs containing the conserved domain corresponding to the individual K⁺ channel probes were amplified in PCR in proportion to the initial abundance. Using Power SYBR Green PCR Master Mix (Qiagen), the fluorescent reaction products were quantified optically with an ABI Prism 7000 machine (ABI, Foster City, CA). For each reaction, a single PCR product was indicated by a characteristic dsDNA dissociation temperature. Triplicates of PCR runs were performed for each gene. The abundance of transcripts for individual K⁺ channel genes was determined by normalizing their optical measurements to that of GAPDH from the same genotypes. The sequences for the primers used in this study and additional experimental details have been described previously (Peng and Wu, 2007b).

Results

Whole-cell currents in dissociated *Drosophila* central neurons at various developmental stages

We dissociated neurons from the third-instar larval (L-III), early (EP), mid (MP) and late pupal (LP), and adult (Ad) *Drosophila* brains. The acutely dissociated neurons were classified into three types according to their shape and size (Wu et al., 1983; Wu et al., 2001): large (> 8 μm), round neuroblast-like type I neurons, small (2–5 μm) type II neurons, and intermediate (5–8 μm) type III neurons. Type II and III neurons accounted for the majority of neurons and thus were used in the present study (Fig. 1A). Despite the diverse morphologies of the neurons recorded, there was no correlation between cell morphologies and ionic current phenotypes. Furthermore, compared to the previously described mechanical dissociation method (Wu et al., 2001), the addition of enzymatic treatment in the present protocol produced a larger number of neurons with longer processes. The majority of the recorded neurons (~91% in a total of 677 neurons) had processes one to three times the diameter of their soma (Fig. 1A).

We performed whole-cell voltage-clamp recordings on the soma of type II/III neurons. Consistent with previous work (Solc and Aldrich, 1988; Saito and Wu, 1991; Wu et al., 2001), the outward K⁺ currents that are composed of both voltage-dependent and Ca²⁺-activated components are the major electrophysiological signature of *Drosophila* CNS neurons. The outward currents often masked a small inward voltage-gated Ca²⁺ current (I_{Ca}) in the total whole-cell K⁺ currents (WCCs). The small I_{Ca} could be observed after the blockade of K⁺ channels (data not shown, cf. Saito and Wu, 1997; Peng and Wu, 2007b). No obvious macroscopic voltage-gated Na⁺ current (I_{Na}) was recorded in most of the cell body among neurons examined.

To better dissect the distinct components that make up the WCCs, we categorized WCCs into four distinct types based on their inactivation kinetics and the relative amplitudes of the sustained components (Fig. 1B). Specifically, types A and D currents inactivated slowly and

were well-fitted with a single exponential function. The sustained component of type D reached more than half of its peak current amplitude, while that of type A did not, accounting for the major difference between the two types. On the other hand, types B and C inactivated in a bi-exponential manner. Type C differs from type B in that the sustained component of type C was more than 50% of its peak current amplitude. The four WCC types presumably reflect differential compositions of distinct K^+ current components, which in *Drosophila* include two voltage-gated K^+ currents (VGCs), the inactivating transient I_A and the delayed rectifier component (I_K) (Solc and Aldrich, 1988; Baker and Salkoff, 1990; Saito and Wu, 1991; Tsunoda and Salkoff, 1995a; 1995b; Yao and Wu, 1999; Peng and Wu, 2007a), and two types of Ca^{2+} -activated K^+ currents (CACs), fast I_{CF} and slow I_{CS} (Singh and Wu, 1989; Peng and Wu, 2007b).

We first examined how the four biophysically different WCC types varied during postembryonic development (Fig. 1C). The percentage of type A WCC remained largely unaltered throughout development; in contrast, those of types B and D displayed most dynamic regulations. Specifically, type B was observed in the majority of larval and adult neurons, while type D dominated pupal stages. In addition, types B and D displayed progressive changes during pupal development, with the percentage of type B increasing and that of type D decreasing, respectively, as pupation progressed. Finally, type C WCC was observed in minimal number of neurons at all stages. In summary, types A and B WCCs largely accounted for the transient and sustained whole-cell K^+ currents in larval and adult neurons, whereas the current types that constitute the WCCs in pupal neurons were more heterogeneous and developmentally regulated. Our data reveal previously unappreciated dynamic features that characterize the K^+ current profiles during pupal development.

Voltage-gated K^+ currents (VGCs), I_A and I_K

WCCs are composed of transient and sustained components of voltage-gated K^+ currents (VGCs) and Ca^{2+} -activated K^+ currents (CACs). To investigate how VGCs and CACs undergo differential developmental regulations, we separated the two currents by electrophysiological and pharmacological means. VGCs were isolated from WCCs in a Ca^{2+} -free external solution to remove both I_{Ca} and CACs, and were classified into four subtypes (Fig. 2A) by the same criteria used for WCC categorizations. The percentages of VGC types observed at each stage differed somewhat from those of corresponding WCC types, especially during pupal development, which may reflect increased contributions of CACs to the total WCCs in pupae (see below). However, the four VGC types showed similar progression of increase or decline compared to the four WCC types across these stages (Fig. 2B). These data are consistent with the idea of more predominant contribution of VGCs than CACs to WCCs in *Drosophila* larval and adult neurons.

To further quantify the transient and sustained components of VGCs, the inactivating I_A and sustained I_K components of VGCs were segregated with a prepulse paradigm (Fig. 2C). It is well established that I_A is inactivated by a conditioning prepulse of -40 mV, whereas I_K is relatively unaffected (Solc and Aldrich, 1988). Subtraction of I_K (Fig. 2C2) from the VGCs evoked from a prepulse of -90 mV (Fig. 2C1) generated I_A (Fig. 2C1-2). I-V curve analyses revealed substantial variations of I_A and I_K current densities (pooled from all types) at

different stages (Fig. 2D, E). To better illustrate the developmental progression of these voltage-dependent transient and sustained components, we examined the relative current densities of I_A and I_K (normalized to the L-III stage) at 10 mV. As shown in Fig. 2F, I_A and I_K displayed distinct developmental profiles: I_A declined from L-III to reach a minimum at EP, gradually recovered throughout pupation, and reached the maximum at Ad; in contrast, I_K reached a plateau at MP and stayed relatively unaltered afterwards. I_A and I_K isolated from different VGC types also grossly followed these developmental profiles (Fig. S1). These data suggest that I_A and I_K are differentially regulated during development.

Ca²⁺-activated K⁺ currents (CACs): I_{CF} , I_{CS}

We next investigated how CACs change developmentally. CACs could be removed by abolishing I_{Ca} with Ca²⁺-free external solution or by applying 0.2 mM CdCl₂ to isolate VGCs (Saito and Wu, 1997; Peng and Wu, 2007b). Therefore, after obtaining WCCs (Fig. 3A1), subtraction of VGCs (Fig. 3A2) resulted in CACs (Fig. 3A1-2). The CACs isolated by this method were mediated by both outward K⁺ currents and a smaller inward I_{Ca} , which was masked by K⁺ currents of greater amplitudes (Fig. 3A and see Ba²⁺ current measurements below). Two subtypes of CACs, types A and D (Fig. 3A), were identified using the same criteria employed for WCC and VGC classifications. The inactivation of both CAC subtypes was well-fitted by one-exponential function. In particular, the property of type A is similar to that of I_{CF} , with the sustained component constituting less than 50% of the peak CACs, whereas type D resembles I_{CS} , whose sustained components account for more than 50% of the peak CACs. The kinetics and pharmacological properties of the type A and D CACs in *Drosophila* brain neurons were consistent with the notion that they consisted of largely I_{CF} and I_{CS} , respectively, with characteristic features of these $I_{K(Ca)}$ reported for *Drosophila* embryonic neurons (Saito and Wu, 1991), larval muscles (Singh and Wu, 1989; 1990; Komatsu et al. 1990) and adult flight muscles (Elkins et al., 1986). Therefore, type A and D currents can serve as useful indicators for the major underlying component of I_{CF} and I_{CS} , respectively. A third CAC subtype, type E, was characterized by its minor current density, with the peak amplitude less than 10 pA/pF (not shown).

The distributions of the three CAC types are shown in Fig. 3B. At the Ad and L-III stages, most neurons did not show appreciable amount of CACs, as suggested by the high percentage of type E CAC. In contrast, types A and D CACs dominated in pupal neurons. The I-V curves of I_{CF} and I_{CS} derived from the inactivating and sustained components of both types of CACs (Fig. 3C, D) are in line with the distribution of CAC subtypes during development. Specifically, I_{CF} and I_{CS} , which displayed similar profiles across developmental stages, are low at the larval stage, increased to the peak at pupal stages and subsequently declined at the adult stage (Fig. 3E). I_{CF} and I_{CS} isolated from individual CAC subtypes (A, D, and E) also grossly followed these developmental profiles (Fig. S1). Together, these data suggest that Ca²⁺-activated K⁺ currents make a relatively small contribution to the total K⁺ currents in the larval and adult neurons, but contribute substantially at the pupal stage.

Ba²⁺ current (I_{Ba}) through voltage-gated Ca²⁺ channels

The increased CACs at pupal stages suggest that Ca²⁺-dependent processes may play a particularly important role in pupal development. Since the major source of Ca²⁺ for triggering CACs is most likely the influx through Ca²⁺ channels, we reasoned that these CAC and Ca²⁺ channels may be developmentally coupled. To test this idea, we measured ionic currents mediated by voltage-gated Ca²⁺ channels by replacing external Ca²⁺ with Ba²⁺ (40 mM) to avoid triggering the large outward CACs activated by Ca²⁺ (Saito and Wu, 1997; Peng and Wu, 2007b). I_{Ba} measurements also provided a more robust determination of Ca²⁺ channel development because Ba²⁺ is a more effective charge carrier for Ca²⁺ channels (Hille, 1992) and circumvented the rapid Ca²⁺-dependent inactivation of Ca²⁺ channels limiting I_{Ca} amplitudes (Kramer and Zucker, 1985).

With a 50-ms depolarization step, I_{Ba} inactivated slowly (Fig. 4A). This sustained, inward I_{Ba} was blocked by 0.2 mM CdCl₂ and was undetectable in the absence of extracellular Ca²⁺ and Ba²⁺, indicating that the current was carried through voltage-activated Ca²⁺ channels. The I-V curves showed that I_{Ba} activated at approximately -30 mV and reached a peak near 10 mV (Fig. 4B). I_{Ba} was first detected at L-III, peaked at EP, and subsequently declined slightly and plateaued throughout MP, LP and AD stages (Fig. 4C). This developmental profile resembled that displayed by I_{CF} and I_{CS} during larval and pupal stages (Fig. 3E), and to a lesser degree, I_K (Fig. 2F). This profile contrasts with the U-shaped developmental time course of I_A. Our data suggest that the developmental onset of Ca²⁺-dependent K⁺ and Ca²⁺ currents may be coupled, and that Ca²⁺-dependent processes may play an unusually important role during pupal development.

Ion current development in GFP-labeled mushroom body neurons

Dissociated neurons from the whole brain as employed in this study contained heterogeneous neuronal populations. The various types of WCCs, VGCs, and CACs observed in these mixed populations and the differential distributions of these current types at different developmental stages could therefore reflect differences in population cell types, rather than overall regulation of current types at these stages. This possibility, while unlikely, was tested in an identified population of central neurons in the mushroom body (MB), the locus of olfactory learning in *Drosophila* (Heisenberg, 2003). MB neurons were prepared from the UAS-GFP; OK107 flies and were identified using GFP fluorescence microscopy (Fig. 5A). Although the current densities in MB neurons were generally smaller than those at other stages (Fig. 5C, Table 1), similar types of WCCs, VGCs, CACs, and I_{Ba} were recorded from this subpopulation of neurons (data not shown). In addition, the distributions of different WCC and VGC types at Ad, MP, and L-III stages (Fig. 5B) paralleled those of whole brain neurons (Fig. 2B), most evidenced by the abundance of the type B WCC and VGC in Ad and L-III neurons and the enrichment of type A and type D WCC and VGC in MP neurons. Finally, the bell-shaped I_{CF}, I_{CS}, and I_{Ba} profiles as well as the V-shaped I_A profile observed in the MB neurons (Fig. 5D) also resembled those of mixed populations (Fig. 2F, 3E, 4C). Although different MB neurons cannot be considered molecularly and functionally identical, our results suggest that it is unlikely that the different distributions of WCCs, VGCs, and CACs types at different developmental stages were due primarily to sampling biases in the neuronal populations examined.

Correlations of WCCs, VGCs and CACs from the same neurons

The distributions of WCC, VGC and CAC subtypes across development suggest that biophysically distinct K^+ current types make varied contributions to total K^+ currents in neurons at different postembryonic stages (Fig. 1C; 2B). In addition, our data indicate that VGCs make up a major portion of total K^+ currents in adult and larval neurons, while CACs contribute more significantly to the total currents in pupal neurons (Fig. 3B). To examine how the various subtypes of VGCs and CACs combined to make up the specific WCC types in individual neurons sampled at different stages, we correlated WCC, VGC, and CAC subtypes from the same neurons. At both L-III and Ad stages, the vast majority of WCCs are type B (Fig. 6A L-III, Ad), which, following pharmacological separations of CACs from VGCs, were correlated with type B VGC and type E CAC. These data indicate that whole-cell K^+ currents from most larval and adult neurons consist of voltage-dependent type B current with minimal contributions from Ca^{2+} -activated K^+ current. Most of the remaining neurons expressed type A WCC that was supported by either type A or type B VGC which were linked to type A, D or E CAC. At pupal stages, whole-cell K^+ currents are less dominated by any specific type (Fig. 6A MP), although type B became the minority and were outnumbered by type A and type D WCCs. Notably, Ca^{2+} -activated K^+ currents, both type A and D, were prominent in pupal neurons and associated with varied WCC/VGC subtypes, presenting a sharp transformation from the predominant populations in Ad/L-III neurons that displayed only small-amplitude type E CACs. Although at the MP stage neurons with type B or type D WCCs still correlated well with type B and type D VGCs, respectively, the degree of correlation was reduced when compared to the adult and larval stages. The diminished correlation is likely due to increased abundance of type A and D CACs that display temporal characteristics comparable to type A and D VGCs and WCCs, which may somewhat distort the correlation. Our data reveal an unexpected re-organization of functional compositions of whole-cell K^+ currents at the successive developmental stages and again highlight an important role for Ca^{2+} -dependent currents in pupal neurons.

Functional role of *Sh* I_A and compensatory effects of *Sh^M* on K^+ currents during development

To investigate the identity of channels that underlies the differential contributions of VGCs to the total WCCs at different developmental stages, we examined the mutational effects of *Sh* and *slo*, two well-characterized genes that encode a voltage-gated I_A and a Ca^{2+} -activated I_{CF} K^+ channel, respectively. Figures 6–9 show results based on *Sh^M*, a null mutation that removes the A-type K^+ current mediated by the *Sh* gene product (Kamb et al., 1987; Zhao et al., 1995). Our analyses indicate that the *Sh* I_A channel mediates a substantial portion of VGCs in both larval and adult neurons, but to a lesser degree in pupal neurons, and that removal of *Sh* I_A may induce a compensatory up-regulation of a different transient K^+ current in larval and adult neurons.

Most notably, *Sh^M* markedly reduced the incidences of type B (and to a lesser degree type C) VGC, with concomitant increases in type A and D VGCs in neurons from all stages (Fig. 7A; Table 1). In addition, direct correlations of WCCs, VGCs, and CACs from the same *Sh^M* neurons also revealed substantial loss of type B VGC but increased abundance of types A and D VGCs (Fig. 6B). In contrast, the occurrence and distributions of CAC subtypes

(Fig. 6B), the current densities of I_{CF} and I_{CS} (Table 2) and I_{Ba} (data not shown) were not affected in Sh^M at all stages. These data suggest that the *Sh* channel mediates a significant portion of type B and C VGCs at all three postembryonic stages, and that the absence of the *Sh* channel apparently led to re-distributions of the other VGC subtypes.

To gain insight into the apparent reorganization of VGC subtypes in Sh^M neurons, we investigated the properties of the remaining transient I_A in Sh^M , using the prepulse inactivation protocol to extract I_A (Fig. 2C). We compared the current densities of the inactivating transient I_A and non-inactivating sustained I_k between wild-type and Sh^M neurons at different stages. As expected, at both L-III and Ad stages, the type B I_A current densities were significantly reduced in Sh^M (Fig. 7B, D). Unexpectedly, coupled with an increased frequency of occurrence of type A VGC, I_A current densities for type A VGC were significantly increased in adult and larval Sh^M neurons (Table 1; Fig. 7B, C). However, when current densities were pooled from all VGC subtypes, the inactivating I_A in Sh^M was still significantly decreased compared to WT, whereas I_k was not affected (Fig. S2). Our results suggest that voltage-gated K^+ currents may undergo functional compensations at larval and adult stages when *Sh*-mediated currents are absent, although these compensations appeared suboptimal, as indicated by a less than WT total I_A density (Fig. S2).

At the mid-pupal stage, we found that the I_A densities for both the type A and type B VGCs were reduced in Sh^M neurons (Table 1), but the pooled I_A and I_k densities were unaltered in Sh^M MP neurons (Fig. S2), suggesting a different pattern of VGC redistribution within a more complex WCC phenotypes during pupation (Fig. 6).

The notion of redistribution of K^+ channel subtypes is further indicated by altered kinetics and voltage dependence of the remaining I_A in Sh^M . The time-dependent activation (assessed by time to peak, T_p) of I_A differed substantially for type A and B VGC, with I_A in neurons displaying type A VGC activating more slowly than that showing type B (Fig. 8, Table 1). Sh^M significantly prolonged the T_p of I_A in type A current at all voltages (Fig. 8B), but did not affect that in type B (Fig. 8C, Table 1) for both L-III and Ad neurons. Sh^M had little effect on T_p of either type A or type B currents in MP neurons, consistent with the idea that different patterns of VGC redistribution occur during metamorphosis (Table 1, Figs. 6–7). As shown in Table 1, the kinetics of I_A recovery from inactivation were not affected when the remaining I_A in both VGCs of type A and B in Sh^M neurons was examined with a twin-pulse protocol (10 ms depolarization to 30 mV from -90 mV, with varying inter-pulse intervals).

However, the voltage dependence of I_A inactivation was altered for the remaining I_A in VGCs of type B, but not type A, in Sh^M neurons (Fig. 9 and Table 1), as examined by fitting the steady-state inactivation of I_A with a Boltzmann equation (Fig. 9, cf. Peng and Wu, 2007a). The half-inactivation voltage ($V_{1/2}$) was more hyperpolarized for I_A in type A compared with that of type B VGC in WT neurons. Sh^M negatively shifted the inactivation curve of I_A of type B, i.e. the remaining non-*Sh* (Fig. 9B3), without affecting that of type A current (Fig. 9B2, Table 1) at both L-III and Ad stages (Table 1), although the slope of the steady-state inactivation curve was not altered by Sh^M at any stages (data not shown). This is in contrast with the more depolarized $V_{1/2}$ for I_A in *Sh* mutants reported previously based on

ensemble results from dissociated neuronal cultures (Baker and Salkoff 1990; Gasque et al., 2005; but see Peng and Wu, 2007a). Again, *Sh^M* did not affect the $V_{1/2}$ for either type A or type B currents in MP neurons.

Taken together, the simplest interpretation of our results is that *Sh*-dependent currents represent a key component of voltage-gated K^+ currents in larval and adult stages but contribute less to pupal I_A currents. *Sh* I_A represents a major component of the type B VGC. Further, the loss of *Sh* results in significant functional compensations of VGCs. The compensations are more prominent in adult and larval than in pupal neurons, consistent with the larger contribution of (*Sh*-containing) type B VGC in adult and larval neurons. Such compensation could be achieved via upregulation of other voltage-gated K^+ channels with properties consistent with those characterized for the remaining I_A in *Sh^M* neurons (Figs. 8 and 9, Table 1).

Effects of *slo¹* mutation on CACs and VGCs

Previous studies on *Drosophila* muscle have shown that the *slo¹* mutation impairs BK-type Ca^{2+} -activated K^+ currents I_{CF} (Elkins et al., 1986; Singh and Wu, 1989; Komatsu et al., 1990). To determine the varying role of CACs at different developmental stages, we examined CACs and the underlying I_{CF} and I_{CS} in the *slo¹* mutant. Our results (Figs. 6C and 10) reveal that *slo* mediates the vast majority of all the Ca^{2+} -activated K^+ currents in postembryonic neurons, and that absence of *slo* induced profound functional compensations from transient voltage-gated K^+ currents, particularly in pupal neurons.

As shown in Fig. 10A, *slo¹* displayed a Ca^{2+} -activated K^+ current profile dominated by the small type E CAC in the majority of neurons at all stages (Fig. 10A), which contrasts to the more prominent type A and D CACs found in subsets of adult and larval, and most of pupal, neurons from the wild-type CNS (Fig. 3A). To reveal how *slo¹* affects different CAC types, we isolated I_{CF} and I_{CS} as described in Figure. 3. In contrast to previous studies that show a preferential elimination of I_{CF} in *Drosophila* embryonic, larval and adult muscles (Elkins et al., 1986; Gho and Mallart, 1986; Singh and Wu, 1989; Komatsu et al., 1990; Broadie and Bate, 1993), we found that current densities of both I_{CF} and I_{CS} were drastically diminished in *slo¹* neurons throughout postembryonic development (Fig. 10B,C,D; Fig. S3; Table 2). Consistently, correlation analyses of WCCs, VGCs, and CACs from the same *slo¹* neurons showed near complete loss of neurons expressing type A and type D CACs (Fig. 6C). Finally, there were near perfect correlations between WCC and VGC subtypes, exhibiting that WCCs in the mutant neurons are predominantly VGCs (Fig. 6C).

To investigate how the loss of *slo* might alter the expression of voltage-gated K^+ currents, we measured the I_A and I_K of VGCs in *slo¹* at different stages. While *slo¹* had no effect on either I_A or I_K of type A and type B VGCs at adult and larval stages, the mutation significantly increased the I_A (Fig. 11A, but not I_K , not shown) in pupal neurons, leading to increased current densities (in pA/pF) of both type A and type B VGCs at the pupal stage (Table 1). In addition, type B VGCs were found in a higher percentage of *slo¹* pupal neurons (Fig. 11B) and in the majority of L-III and Ad neurons (Fig. 6C). These data suggest that voltage-gated I_A components associated with type B VGCs were globally upregulated to compensate for the loss of Ca^{2+} -activated I_{CF} and I_{CS} at the pupal stage. Consistently, in *slo¹*

pupal brains we found a marked increase in the mRNAs for *Sh*, and a less prominent increase for *Shal* (Fig. 11C). This result provides a mechanism for the previous report (Lee et al., 2008) that *Sh* K⁺ channels are required for the synaptic strength homeostasis to compensate for the loss of Ca²⁺-activated K⁺ channels in *slo* neuromuscular junctions. These data demonstrated that functional deficiency of a Ca²⁺-activated channel type could recruit compensatory adjustments from completely different channel families. The pupa-specific VGC reorganizations associated with *slo^l* mirrors the important roles of CACs in developing pupal neurons.

***Sh*;*slo* double mutants failed to survive through adulthood**

The significant contributions of *Sh* and *slo* to K⁺ currents, as well as the compensatory regulation at their absence, suggest important roles of these channels in *Drosophila* development. We thus constructed *Sh^{l33};;slo^l* and *Sh^M;;slo⁹⁸* (kept over X or 3rd chromosome balancers) to compare with *Sh* or *slo* single mutants for effects on post-embryonic developments (Fig. 11D). The mutant *slo⁹⁸* is an independent allele of *slo* that is similar to *slo^l* in many respects of neuromuscular junction and behavioral phenotypes (Lee and Wu, 2010; Atkinson et al., 2000). We determined the survival rates of the single and double mutants through the larval and the pupal stages. Our data indicated that while single *Sh* (*Sh^M* and *Sh^{l33}*) or *slo* (*slo^l* and *slo⁹⁸*) mutants survived nearly normally through early adulthood, *Sh^M;;slo⁹⁸* double mutants displayed substantially reduced survival through the larval stages, and virtually all failed to eclose to reach adulthood (Fig. 11D). The decreased survival rates in double mutants compared to single mutants were consistent with the homeostatic compensations across channel families between *Sh* and *slo* shown above. The data suggest that *Sh* can serve as a major backup for the missing *slo*-dependent Ca²⁺-activated K⁺ currents, and that absence of both in double mutants is catastrophic. Our findings thus identify a central, but previously unrecognized roles for *Sh* and *slo* channels in *Drosophila* development.

Discussion

How individual neurons participate in circuitry functions that mediate behaviors depends, to a great degree, on their intrinsic properties in situ. In this study, we adapted a neuronal preparation that allowed comprehensive and quantitative biophysical investigations of ion currents from soma of freshly isolated neurons from the CNS of *Drosophila* (Wu et al., 2001). These neurons were devoid of long neurites and complex arbors, avoiding interferences from uncontrolled remote processes during voltage-clamp recording. In addition, this acute dissociation approach permitted immediate assessments of neuronal ion channels in their natural settings without further culturing, which may alter channel expression and assembly during prolonged in vitro growth or re generation. Here we provide a complete profile of the various K⁺ and Ca²⁺ currents during *Drosophila* postembryonic development. Analyses of mutants that are deficient in *Sh*- and/or *slo*-encoded channels uncovered their major contributions to the composition, reorganization, and homeostatic plasticity of neuronal K⁺ currents at different postembryonic stages.

Postembryonic development of *Drosophila* K⁺ currents

K⁺ currents in *Drosophila* neurons have been the subject of intense study (Byerly and Leung, 1988; Solc and Aldrich, 1988; Baker and Salkoff, 1990; Saito and Wu, 1991; Tsunoda and Salkoff, 1995; Wright and Zhong, 1995; Zhao et al., 1995; Baines and Bate, 1998; Yao and Wu, 1999; Peng and Wu, 2007a; 2007b; Srinivasan et al., 2012). However, a detailed understanding of how K⁺ currents develop, especially at postembryonic phases, has been hampered due to the electrophysiological inaccessibility to most of postembryonic neurons. Using acutely dissociated neurons from different postembryonic stages, we provide the first comprehensive, quantitative biophysical assessments of K⁺ current phenotypes in *Drosophila* central neurons at larval, pupal, and adult stages. We show that I_A displays a U-shaped developmental profile, minimizing at an early pupal stage, whereas both I_{CF} and I_{CS} exhibit inverted U profiles, peaking at the same early/mid pupal stages. In contrast, I_K shows a steady increase as metamorphosis progresses, reaching a plateau beyond the mid-pupal stage. Similar profiles were also observed in morphologically and functionally more homogeneous neurons dissociated from the mushroom bodies. Thus, these profiles likely reflect a global program of regulated expression and/or reassembly of different channel subunits in CNS neurons during postembryonic development.

This postembryonic I_A profile extends the sequential and progressive development reported for I_A in embryonic neurons (Baines and Bate, 1998) and muscles (Broadie and Bate, 1993) in *Drosophila*. Such monotonic increase of I_A is found in embryonic neurons in vertebrates as well (O'Dowd et al., 1988; Rivera and Spitzer, 1990; Spitzer 1994). However, the onset of metamorphosis, i.e. early pupa, is characterized by a vigorous remodeling of CNS at both the individual cell and the circuit levels, during which larval dendritic and axonal branches prune, followed by the growth of adult-specific processes (Williams and Truman, 2005). Notably, transient I_A has been negatively correlated to axonal arborization and growth. For example, *Drosophila* hyperexcitability mutants that lack both *Sh* and *ether à go-go* (*eag*) display increased larval motoneuron terminal branching (Budnik et al., 1990; Zhong et al., 1992), whereas overexpression of *Sh*-like channels leads to a reduction in morphologically differentiated neurons in vertebrate cell cultures (Jones and Ribera, 1994). In this context, suppression of I_A at early pupal stages (Fig. 2) may increase nerve excitability and promote axonal arborization, activity-dependent synaptic maturation, and neural circuit reorganization during early metamorphosis.

The delayed rectifier I_K contributes to membrane repolarization and the maintenance of repetitive nature of action potential firing (Hille, 2001). Our study demonstrates a progressive increase of I_K during postembryonic development (Fig. 2), which further extends the gradual increase of I_K in the development of both *Drosophila* embryos (Baines and Bate, 1998) and vertebrate embryonic cultures (O'Dowd et al., 1988; Ribera and Spitzer, 1990). A larger I_K is expected to limit the number of spikes. Consistent with this theme, mutation of the *Shab* channel in *Drosophila* converts characteristic repetitive firing patterns into run-away spikes and high-frequency oscillations in mutant “giant” neuron cultures (Peng and Wu, 2007a). In an identified motoneuron in *Manduca*, I_K gradually increases during postembryonic metamorphosis, in parallel to the continuing reduction of membrane excitability during its transformation to adult form (Dutch and Levine, 2000).

Ca^{2+} -dependent $\text{I}_{\text{K}(\text{Ca})}$ contributes to action potential repolarization and hyperpolarizing after potentials, and hence neuronal membrane excitability and firing patterns (for review, see e.g. Vergara et al., 1998). Therefore, Ca^{2+} influx through voltage-gated Ca^{2+} channels not only orchestrates complex intracellular Ca^{2+} signaling processes, but also triggers $\text{I}_{\text{K}(\text{Ca})}$ that directly regulates membrane repolarization and in turn limits further Ca^{2+} influx. We found that these two functionally coupled channel types, voltage-gated Ca^{2+} channels and Ca^{2+} -activated K^{+} channels, also displayed similar developmental profiles, suggesting that their expression is also developmentally coupled during metamorphosis. During insect CNS metamorphosis, extensive axonal and dendritic pruning, synaptogenesis, and synaptic modification occur, all of which may depend on spatially and temporally regulated Ca^{2+} signaling. The surges of both I_{Ca} and $\text{I}_{\text{K}(\text{Ca})}$ at the onset of metamorphosis may reflect a drastic change in neuronal Ca^{2+} signaling and excitability requirements for the CNS remodeling.

Composition, reorganization, and homeostatic plasticity of postembryonic K^{+} currents: role of *Sh* and *slo*

Our demonstration of a significant contribution of *Sh* channels to I_{A} in *Drosophila* CNS neurons contrasts to the previous genetic and electrophysiological studies of I_{K} in cultured embryonic and larval central neurons that have led to the idea that *Shal* and *Shab* mediate nearly all of I_{A} and I_{K} , respectively, and that *Sh* does not make significant contributions to I_{A} in these neurons (Baker and Salkoff, 1990; Tsunoda and Salkoff, 1995). However, prolonged culturing of dissociated neurons in vitro could alter intrinsic gene expression programs in studies employing cultured neurons. It is interesting to speculate that processes of neuronal re-growth in culture might recapitulate the condition of CNS remodeling initiation in early pupal development where I_{A} expression is minimal among the postembryonic stages (Figs. 2F and S1).

Our analyses of *Sh^M* neurons demonstrated that the null mutation drastically reduced incidences of the fast inactivating type B VGCs in all three developmental stages, including larval (70.2 to 20%), pupal (31.8 to 21.4%), and adult (70.8 to 20.8%), and curtailed the current densities of this VGC type by 49%, and 41%, and 52% in mutant larval, pupal, and adult neurons, respectively (Figs. 6B, 7 and Table 1). These results demonstrate that *Sh* makes significantly larger than expected contributions to I_{A} in a majority of CNS neurons. *Sh* contributions in different developmental stages are not equal, being more prominent in larval and adult neurons and less in pupal neurons, consistent with the reported two expression peaks of *Sh* transcripts, at the larval and adult stages, respectively (Rogerio and Tejedor, 1995). Together, our study on populations of wild-type and *Sh^M* neurons from various postembryonic stages uncover surprisingly greater roles of *Sh* in larval and adult neurons.

In *Drosophila* embryonic, larval, and adult muscle cells, *slo* eliminates the transient, but not the sustained components, of $\text{I}_{\text{K}(\text{Ca})}$ (Elkins et al., 1986; Singh and Wu, 1989, 1990; Broadie and Bate, 1993), indicating that *slo* mediates I_{CF} in muscles. The molecular identities of $\text{I}_{\text{K}(\text{Ca})}$ in neurons are less clear, although a previous study using cultured embryonic *Drosophila* “giant” neurons suggests that *slo* appears to affect the late as well as the transient components of $\text{I}_{\text{K}(\text{Ca})}$ (Saito and Wu, 1991). Our study on neurons freshly dissociated from

various postembryonic brains demonstrated that both I_{CF} and I_{CS} were strikingly reduced by *slo*¹ (Fig. 10), suggesting that *slo* mediates the vast majority of somatic I_{CF} and I_{CS} in central neurons during all postembryonic stages. However, *slo*¹ did not completely eliminate I_{CF} and I_{CS} in at least a subset of neurons and non-*slo* $I_{K(Ca)}$ channels were present. This is indicated by small populations of remaining type A and D CAC neurons, and most strikingly during metamorphosis, when a great majority of type A and D CACs was converted into type E CAC of small amplitudes in *slo* pupal neurons (Figs. 6C and 10A). The possibility that these remaining CACs may be mediated by channels encoded by other genes awaits further investigations. In summary, in the *Drosophila* nervous system, a single gene, *slo*, appears to encode the majority of the Ca^{2+} -activated K^{+} currents across the entire embryonic and postembryonic phases.

Homeostatic maintenance of neuronal activity represents a form of plasticity by which neurons re-adjust to maintain their membrane excitability in response to chronic internal (e.g. genetic) or external perturbations (Spitzer, 1999; Turrigiano, 1999, 2008). This can be achieved through reorganization by re-adjusting the proportions of different ion channel subtypes. For instance, overexpression of *Shal*-mediated I_A in lobster neurons induces a compensatory increase of hyperpolarization-activated inward I_h (MacLean et al., 2003).

Our study presents two cases of channel homeostatic plasticity in response to genetic perturbations in *Drosophila*. In *Sh*^M, there was a specific increase of *Sh*-independent I_A , but not I_K , that was particularly evident in larval and adult neurons, a compensation that could be due to upregulation of other homologous genes in the *Sh* family. The properties of the remaining voltage-gated K^{+} currents after *Sh* channel removal in *Sh*^M neurons reveal some clues to the candidate voltage-gated K^{+} channels for the homeostatic compensation. The dominant type A VGC in *Sh*^M, presumably converted from the type B majority in WT (Figs. 6B and 7A), exhibited a much longer time to peak, indicating slower activation kinetics (Fig. 8), and the remaining non-*Sh* I_A in Type B VGC showed a hyperpolarizing shift in steady-state inactivation (a more negative $V_{1/2}$, Fig. 9A,C and Table 1). These findings indicate that the *Shal* channel could be a good candidate based on the kinetic and voltage-dependent properties (Baker and Salkoff, 1990; Tsunoda and Salkoff, 1995; Ryglewski and Duch, 2009). Further studies will be required to determine this conjecture.

In contrast, *slo*¹ displayed a specific upregulation of I_A , but not I_K , mediated largely by an increased *Sh* expression and most evident in mutant pupal neurons. The contrasting stage-dependent compensations of K^{+} currents in *Sh*^M and *slo*¹ mutants mirrors their more prominent contributions in the corresponding postembryonic stages: *Sh* for larval and adult and *slo* for pupal stages.

The upregulation of *Sh* currents in *slo*¹ pupal neurons provide a unique example of cross-family homeostatic regulation of K^{+} currents. This is presumably because I_A and I_{CF} have similar kinetic and other biophysical properties (Elkins and Ganetzky, 1988; Singh and Wu, 1990) and can provide some functional redundancies. This notion is consistent with the previous study in larval neuro-muscular junctions (Lee et al., 2008) indicating that *Sh* K^{+} channels are up-regulated to maintain synaptic strength homeostasis in *slo* mutants.

Similar cross-family compensation is observed in the *Drosophila* Ca^{2+} channel mutant, *cacophony* (*cac*). In *cac*, the majority of Ca^{2+} -activated K^+ currents are lost (resulting from the near complete elimination of Ca^{2+} currents in mutant neurons), but an upregulation of the *Sh* I_A is induced to partially compensate for the reduction of $\text{I}_{\text{K}(\text{Ca})}$ (Peng and Wu, 2007). In both cases, interestingly, the homeostatic regulation appeared to be unidirectional since I_A was upregulated in *slo* and *cac* but neither Ca^{2+} currents nor $\text{I}_{\text{K}(\text{Ca})}$ were altered in K^+ channel mutants. Our data indicate that *Sh* can serve as a functional backup for other channels with similar physiological properties, e.g. *slo* channels. This novel compensation mechanism is indispensable during pupal development, as *Sh;;slo* double mutants failed to enclose to become adults. Thus, although *Sh* mediates only a minor portion of I_A in pupal neurons during metamorphosis, its role as a “plasticity” gene represents an essential feature in *Drosophila* postembryonic development.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We dedicate this manuscript to the special issue in honor of the lifetime achievements of Dr. Barry Ganetzky. His early pioneering work on *Sh* and *slo* has engendered several active research areas that continues to produce exciting new findings. This study based on hundreds of *Sh* and *slo* neurons is a testimony of how he profoundly influenced the development of this important research field in the past decades.

This study was supported by the Knowledge Innovation Project from the Chinese Academy of Sciences (KSCX2-YW-R-35) and the National Basic Research Program of China (No. 2006CB500803) to T.-L. X., US NIH Grants GM088804 and AG051513 to C.-F. W., DA032283 to W.-D.Y., and OD011103 to New England Primate Research Center, Harvard University. C.-F. W. was also supported by the Wang Kuancheng Scholarship of the Chinese Academy of Sciences while conducting research at the Shanghai Institutes for Biological Sciences. We thank Dr. Z.-R. Wang for comments on the manuscript, Q. Wang and H. Lu for technical assistance, and T. Patience for assistance in manuscript preparation.

Frequently used abbreviations

WCC	Whole-Cell Current
VGC	Voltage-Gated K^+ Current
CAC	Ca^{2+} -Activated K^+ Current
L-III	third-instar larva
EP,MP,LP	Early, Middle, and Late Pupa
Ad	Adult
I_A, I_K	transient and sustained (inactivating and non-inactivating) components of VGCs
ICF, ICS	fast, inactivating and slow, sustained components of CACs

References

- Atkinson NS, Robertson GA, Ganetzky B. A component of calcium-activated potassium channels encoded by the *Drosophila* slo locus. *Science*. 1991; 253:551–555. [PubMed: 1857984]
- Atkinson NS, Brenner R, Chang W, Larimer JL, Yu J. Molecular separation of two behavioral phenotypes by a mutation affecting the promoters of a Ca-activated K channel. *J Neurosci*. 2000; 20:2988–2993. [PubMed: 10751451]
- Bainbridge SP, Bownes M. Staging the metamorphosis of *Drosophila melanogaster*. *J Embryol Exp Morphol*. 1981; 66:57–80. [PubMed: 6802923]
- Baines RA, Bate M. Electrophysiological development of central neurons in the *Drosophila* embryo. *J Neurosci*. 1998; 18:4673–4683. [PubMed: 9614242]
- Baker K, Salkoff L. The *Drosophila* Shaker gene codes for a distinctive K^+ current in a subset of neurons. *Neuron*. 1990; 4:129–40. [PubMed: 2310571]
- Broadie KS, Bate M. Development of larval muscle properties in the embryonic myotubes of *Drosophila melanogaster*. *J Neurosci*. 1993a; 13:167–180. [PubMed: 8093714]
- Budnik V, Zhong Y, Wu CF. Morphological plasticity of motor axons in *Drosophila* mutants with altered excitability. *J Neurosci*. 1990; 10:3754–3768. [PubMed: 1700086]
- Butler A, Wei AG, Baker K, Salkoff L. A family of putative potassium channel genes in *Drosophila*. *Science*. 1989; 243:943–947. [PubMed: 2493160]
- Byerly L, Leung HT. Ionic currents of *Drosophila* neurons in embryonic cultures. *J Neurosci*. 1988; 8:4379–4393. [PubMed: 2460598]
- Choi JC, Park D, Griffith LC. Electrophysiological and morphological characterization of identified motor neurons in the *Drosophila* third instar larva central nervous system. *J Neurophysiol*. 2004; 91:2353–2365. [PubMed: 14695352]
- Consoulas C, Restifo LL, Levine RB. Dendritic remodeling and growth of motoneurons during metamorphosis of *Drosophila melanogaster*. *J Neurosci*. 2002; 22:4906–4917. [PubMed: 12077188]
- Duch C, Levine RB. Remodeling of membrane properties and dendritic architecture accompanies the postembryonic conversion of a slow into a fast motoneuron. *J Neurosci*. 2000; 20:6950–6961. [PubMed: 10995839]
- Elkins T, Ganetzky B. The role of potassium currents in *Drosophila* flight muscles. *J Neurosci*. 1988; 8:428–434. [PubMed: 3339425]
- Elkins T, Ganetzky B, Wu CF. A *Drosophila* mutation that eliminates a calcium-dependent potassium current. *Proc Natl Acad Sci U S A*. 1986; 83:8415–8419. [PubMed: 2430288]
- Gasque G, Labarca P, Reynaud E, Darszon A. Shal and shaker differential contribution to the K^+ currents in the *Drosophila* mushroom body neurons. *J Neurosci*. 2005; 25:2348–2358. [PubMed: 15745961]
- Getting PA. Emerging principles governing the operation of neural networks. *Annu Rev Neurosci*. 1989; 12:185–204. [PubMed: 2648949]
- Gho M, Mallart A. Two distinct calcium-activated potassium currents in larval muscle fibres of *Drosophila melanogaster*. *Pflugers Arch*. 1986; 407:526–533. [PubMed: 2431384]
- Goodman CS, Shatz CJ. Developmental mechanisms that generate precise patterns of neuronal connectivity. *Cell*. 1993; 72(Suppl):77–98. [PubMed: 8428376]
- Harris-Warrick RM, Marder E. Modulation of neural networks for behavior. *Annu Rev Neurosci*. 1991; 14:39–57. [PubMed: 2031576]
- Haugland NH, Wu C-F. A voltage-clamp analysis of gene-dosage effects of the Shaker locus on larval muscle potassium currents in *Drosophila*. *Journal of Neuroscience*. 1990; 10:1357–1371. [PubMed: 2109786]
- Hebbard S, Hall RE, Demski SA, Subramanian A, Fernandes JJ. The adult abdominal neuromuscular junction of *Drosophila*: A model for synaptic plasticity. *Dev Neurobiol*. 2006; 66:1140–1155.
- Heisenberg M. Mushroom body memoir: from maps to models. *Nat Rev Neurosci*. 2003; 4:266–275. [PubMed: 12671643]
- Hille, B. *Ionic Channels of Excitable Membranes*. Sunderland, MA: Sinauer; 1992.

- Hille, B. Ion channels of excitable membranes. Sunderland, MA: Sinauer Associates, Inc; 2001.
- Iverson LE, Tanouye MA, Lester HA, Davidson N, Rudy B. A-type potassium channels expressed from Shaker locus cDNA. *Proc Natl Acad Sci U S A*. 1988; 85:5723–7. [PubMed: 2456579]
- Jones SM, Ribera AB. Overexpression of a potassium channel gene perturbs neural differentiation. *J Neurosci*. 1994; 14:2789–2799. [PubMed: 8182440]
- Kamb A, Iverson LE, Tanouye MA. Molecular characterization of *Shaker*, a *Drosophila* gene that encodes a potassium channel. *Cell*. 1987; 50:405–413. [PubMed: 2440582]
- Kaplan WD, Trout WE 3rd. The behavior of four neurological mutants of *Drosophila*. *Genetics*. 1969; 61:399–409. [PubMed: 5807804]
- Kimura K, Truman J. Postmetamorphic cell death in the nervous and muscular systems of *Drosophila melanogaster*. *J Neurosci*. 1990; 10:403–411. [PubMed: 2106014]
- Komatsu A, Singh S, Rathe P, Wu CF. Mutational and gene dosage analysis of calcium-activated potassium channels in *Drosophila*: correlation of micro- and macroscopic currents. *Neuron*. 1990; 4:313–321. [PubMed: 2106331]
- Kramer RH, Zucker RS. Calcium-induced inactivation of calcium current causes the inter-burst hyperpolarization of Aplysia bursting neurones. *J Physiol*. 1985; 362:131–160. [PubMed: 2410598]
- Lee J, Ueda A, Wu CF. Pre- and post- synaptic mechanisms of synaptic strength homeostasis revealed by *Slowpoke* and *Shaker* K⁺ channel mutations in *Drosophila*. *Neuroscience*. 2008; 154:1283–1296. [PubMed: 18539401]
- Lee J, Wu CF. Orchestration of stepwise synaptic growth by K⁺ and Ca²⁺ channels in *Drosophila*. *J Neurosci*. 2010; 30:15821–15833. [PubMed: 21106821]
- Li H, Li Y, Lei Z, Wang K, Guo A. Transformation of odor selectivity from projection neurons to single mushroom body neurons mapped with dual-color calcium imaging. *PNAS*. 2013; 110:12084–12089. [PubMed: 23818618]
- MacLean JN, Zhang Y, Johnson BR, Harris-Warrick RM. Activity-independent homeostasis in rhythmically active neurons. *Neuron*. 2003; 37:109–120. [PubMed: 12526777]
- O'Dowd DK, Ribera AB, Spitzer NC. Development of voltage-dependent calcium, sodium, and potassium currents in *Xenopus* spinal neurons. *J Neurosci*. 1988; 8:792–805. [PubMed: 3346722]
- Peng IF, Wu CF. Differential contributions of Shaker and Shab K⁺ currents to neuronal firing patterns in *Drosophila*. *J Neurophysiol*. 2007a; 97:780–794. [PubMed: 17079336]
- Peng IF, Wu CF. *Drosophila* cacophony channels: a major mediator of neuronal Ca²⁺ currents and a trigger for K⁺ channel homeostatic regulation. *J Neurosci*. 2007b; 27:1072–1081. [PubMed: 17267561]
- Ribera AB, Spitzer NC. Differentiation of IKA in amphibian spinal neurons. *J Neurosci*. 1990; 10:1886–1891. [PubMed: 2355255]
- Rogero O, Tejedor FJ. Immunohistochemical characterization and developmental expression of *Shaker* potassium channels from the nervous system of *Drosophila*. *J Biol Chem*. 1995; 270:25746–25751. [PubMed: 7592756]
- Ryglewski S, Duch C. Shaker and Shal mediate transient calcium-independent potassium current in a *Drosophila* flight motoneuron. *J Neurophysiol*. 2009; 102:3673–3688. [PubMed: 19828724]
- Saito M, Wu CF. Expression of ion channels and mutational effects in giant *Drosophila* neurons differentiated from cell division-arrested embryonic neuroblasts. *J Neurosci*. 1991; 11:2135–2150. [PubMed: 1712379]
- Salkoff L, Wyman R. Genetic modification of potassium channels in *Drosophila*. *Shaker* Mutants, *Nature*. 1981; 293:228–230.
- Salkoff L, Wyman R. Ion currents in *Drosophila* flight muscles. *J. Physiol. Paris*. 1983; 80:275–282.
- Singh S, Wu CF. Complete separation of four potassium currents in *Drosophila*. *Neuron*. 1989; 2:1325–1329. [PubMed: 2516727]
- Singh S, Wu CF. Properties of potassium currents and their role in membrane excitability in *Drosophila* larval muscle fibers. *J Exp Biol*. 1990; 152:59–76. [PubMed: 2121887]
- Solc CK, Aldrich RW. Voltage-gated potassium channels in larval CNS neurons of *Drosophila*. *J Neurosci*. 1988; 8:2556–2570. [PubMed: 3249242]

- Spitzer, NC. Development of voltage-dependent and ligand-gated channels in excitable membranes. In: van Pelt, J. Corner, MA. Uylings, HBM., Lopes da Silva, FH., editors. Progress in brain research. Amsterdam: Elsevier; 1994. p. 169-179.
- Spitzer NC. New dimensions of neuronal plasticity. *Nat Neurosci.* 1999; 2:489–491. [PubMed: 10448206]
- Srinivasan S, Lance K, Levine RB. Contribution of EAG to excitability and potassium currents in *Drosophila* larval motoneurons. *J Neurophysiol.* 2012; 107:2660–2671. [PubMed: 22323637]
- Timpe LC, Jan YN, Jan LY. Four cDNA clones from the Shaker locus of *Drosophila* induce kinetically distinct A-type potassium currents in *Xenopus* oocytes. *Neuron.* 1988; 1:659–67. [PubMed: 3272184]
- Truman JW. Metamorphosis of the central nervous system of *Drosophila*. *J Neurobiol.* 1990; 21:1072–1084. [PubMed: 1979610]
- Tsunoda S, Salkoff L. Genetic analysis of *Drosophila* neurons: *Shal*, *Shaw*, and *Shab* encode most embryonic potassium currents. *J Neurosci.* 1995; 15:1741–1754. [PubMed: 7891132]
- Turrigiano GG. Homeostatic plasticity in neuronal networks: the more things change, the more they stay the same. *Trends Neurosci.* 1999; 22:221–227. [PubMed: 10322495]
- Turrigiano GG. The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell.* 2008; 135:422–435. [PubMed: 18984155]
- Vergara C, Latorre R, Marrion NV, Adelman JP. Calcium-activated potassium channels. *Curr Opin Neurobiol.* 1998; 8:321–329. [PubMed: 9687354]
- Wei A, Covarrubias M, Butler A, Baker K, Pak M, Salkoff L. K^+ current diversity is produced by an extended gene family conserved in *Drosophila* and mouse. *Science.* 1990; 248:599–603. [PubMed: 2333511]
- Williams DW, Truman JW. Remodeling dendrites during insect metamorphosis. *J Neurobiol.* 2005; 64:24–33. [PubMed: 15884009]
- Wright NJ, Zhong Y. Characterization of K^+ currents and the cAMP-dependent modulation in cultured *Drosophila* mushroom body neurons identified by lacZ expression. *J Neurosci.* 1995; 15:1025–1034. [PubMed: 7869080]
- Wu CF, Suzuki N, Poo MM. Dissociated neurons from normal and mutant *Drosophila* larval central nervous system in cell culture. *J Neurosci.* 1983; 3:1888–1899. [PubMed: 6310066]
- Wu CF, Ganetzky B, Jan LY, Jan YN. A *Drosophila* mutant with a temperature-sensitive block in nerve conduction. *Proc Natl Acad Sci U S A.* 1978; 75:4047–4051. [PubMed: 211514]
- Wu LJ, Lu Y, Xu TL. A novel mechanical dissociation technique for studying acutely isolated maturing *Drosophila* central neurons. *J Neurosci Methods.* 2001; 108:199–206. [PubMed: 11478979]
- Yao WD, Wu CF. Auxiliary Hyperkinetic beta subunit of K^+ channels: regulation of firing properties and K^+ currents in *Drosophila* neurons. *J Neurophysiol.* 1999; 81:2472–2484. [PubMed: 10322082]
- Yu D, Feng C, Guo A. Altered outward $K(+)$ currents in *Drosophila* larval neurons of memory mutants *rutabaga* and *amnesiac*. *J Neurobiol.* 1999; 40:158–170. [PubMed: 10413447]
- Zhao ML, Sable EO, Iverson LE, Wu CF. Functional expression of *Shaker* K^+ channels in cultured *Drosophila* “giant” neurons derived from *Sh* cDNA transformants: distinct properties, distribution, and turnover. *J Neurosci.* 1995; 15:1406–1418. [PubMed: 7869107]
- Zhong Y, Budnik V, Wu CF. Synaptic plasticity in *Drosophila* memory and hyperexcitable mutants: role of cAMP cascade. *J Neurosci.* 1992; 12:644–651. [PubMed: 1371316]

Fig. 1. Morphology and whole-cell K^+ current phenotypes of acutely dissociated *Drosophila* brain neurons at various postembryonic stages

A, Morphology of acutely dissociated neurons from larval, pupal and adult brains of *Drosophila*. L-III, EP, MP, LP and Ad represent third-instar larva, early (P5–8), middle (P8–11) and late (P11–15) pupa, and adult, respectively. Arrows indicate neuronal processes. **B**, Four subtypes of whole-cell K^+ currents (WCCs), types A–D, were identified based on the inactivation kinetics and the relative amplitude of sustained currents at 10 mV. After hyperpolarizing to -90 mV for 200 ms, WCCs were elicited by 20 mV depolarizing steps for 500 ms from -70 to 70 mV. Holding potential was -70 mV. **C**, Distribution histogram of WCC types at different developmental stages. Cell numbers are indicated in the parentheses for corresponding developmental stages.

Fig. 2. Voltage-gated K^+ currents (VGCs) at different developmental stages

A, Four subtypes of VGCs were identified as in Fig. 1B. **B**, Comparisons of percentage of WCCs and VGCs at different developmental stages. **C**, Isolation of I_A and I_K . VGCs were recorded in the external solution in which Ca^{2+} was replaced with Mg^{2+} to eliminate both the inward Ca^{2+} currents and the Ca^{2+} -activated K^+ currents. Two components of the VGCs, I_A and I_K were separated with a prepulse of -40 mV that inactivate the transient components (I_A) but left the sustained component I_K intact (**C2**). I_A was isolated by subtracting I_K from total VGCs (**C1–C2**). **D**, **E**, I-V curves of I_A and I_K currents (pA/pF, normalized to membrane capacitance in pA/pF) at different developmental stages. I_A was measured at the peak of derived transients, whereas I_K was measured at the end of the 500-ms depolarization pulses. **F**, Developmental progression of I_A and I_K . Current densities of different

components at 10 mV were normalized to those of L-III neurons. In this and subsequent figures, data points and error bars represent means and SEM.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Fig. 3. Development of Ca^{2+} -activated K^+ currents (CACs), I_{CF} and I_{CS}

A, Two subtypes of CACs, type A and D, were classified according to the criteria for WCCs and VGCs. CACs (traces 1–2) were obtained by subtracting the currents recorded in the absence of Ca^{2+} (traces 2, VGCs) from those recorded in the presence of 2.5 mM Ca^{2+} (traces 1, WCCs). I_{CS} , the sustained component, was measured at the end of CACs (traces 1–2), whereas I_{CF} was obtained by subtracting I_{CS} from the peak. **B**, Bar graph showing the percentage of three CACs at different postembryonic stages. An additional type, type E was characterized by its low current density (less than 10 pA/pF). **C**, **D**, I-V curves of I_{CF} and I_{CS} currents at different stages as indicated. **E**, Developmental progression of I_{CF} and I_{CS} . Current densities of different components at 10 mV were normalized to those of L-III neurons.

Fig. 4. Developmental profiles of Ca^{2+} channel-mediated Ba^{2+} current

A, An example of voltage-gated Ba^{2+} current (I_{Ba}) generated in response to 50 ms voltage steps in 20 mV increments, ranging from -70 to 50 mV, after a 200 ms prepulse of -90 mV. I_{Ba} was isolated using I_{Ba} -solution, in which I_{Na} was blocked by 0.1 μM TTX, and K^{+} currents were blocked by 20 mM TEA and 10 mM 4-AP. KCl in the pipette solution was substituted by same millimolar CsCl. *B*, I-V relationships of I_{Ba} at different developmental stages. *C*, Developmental progression of I_{Ba} . Current densities at 10 mV were normalized to those of L-III neurons.

Fig. 5. Development of various current components in mushroom body (MB) neurons
A, Morphology of acutely dissociated MB neurons. GFP-labeled MB neurons were identified in the UAS-GFP; OK107 line under a fluorescence microscope. Arrows indicate neuronal processes. **B**, Percentages of different WCC and VGC types at LIII, MP, and Ad stages. **C**, Mean densities of various current components at 10 mV during development. **D**, Developmental progression of different current components. I_A , I_K , I_{CF} and I_{CS} were recorded from 15 L-III, 18 MP and 18 Ad neurons, respectively. For I_{Ba} , 8 neurons were analyzed for each developmental stage.

Fig. 6. Correlating between WCCs, VGCs and CACs in the same neurons from WT, Sh^M , and slo^1 mutants

WCCs, VGCs, and CACs were recorded from individual neurons, classified into different types using criteria described in Text, and correlated. Three stages, L-III, MP, and Ad are shown for WT (A), Sh^M (B), and slo^1 (C) neurons. Lines connecting different specific subtypes of WCC, VGC, and CAC represent individual neurons.

Fig. 7. Effects of *Sh^M* on I_A and I_K

A, Percentage of neurons expressing distinctive VGCs in *Sh^M*, compared to WT, flies at L-III, MP and Ad stages. *B*, I_A densities (at 10 mV) from type A and type B VGCs from LIII and Ad WT and *Sh^M* neurons. Histogram bars show the averaged values. *C*, *D*, Mean I_A and I_K densities pooled from all VGC types in WT and *Sh^M* neurons. *E*, *F*, I-V curves I_A for type A and B I_A in WT and *Sh^M* neurons at L-III and adult stages. * $P < 0.05$, ** $P < 0.01$ (student's t-test).

Fig. 8. Effects of Sh^M on kinetics and voltage-dependence of I_A activation

A, Effects of Sh^M on time to peak (Tp) of I_A (VGCs). Bar graphs showing average Tp for type A and type B I_A at 10 mV in LIII and Ad neurons from WT and Sh^M . The inset shows a typical recording from type A subpopulation illustrating different activation kinetics of I_A in WT and Sh^M neurons.

B–C, Voltage dependence of Tp for type A (**B**) and type B (**C**) I_A in WT and Sh^M . * $P < 0.05$, ** $P < 0.01$ (student's t-test.).

Fig. 9. Effects of Sh^M on steady-state inactivation

A, Effects of Sh^M on the steady-state inactivation of I_A . Bar graphs summarizing average half-inactivation voltage ($V_{1/2}$) of I_A for type A and type B currents in LIII and Ad neurons from WT and Sh^M . The inserted current traces were obtained in response to a test potential of 40 mV after a prepulse to a range of membrane potentials from -120 to -30 mV in 10 mV increments.

B–C, I-V curves of steady-state inactivation of I_A for type A (**B**) and type B (**C**) subpopulations in WT (solid) and Sh^M (dashed) neurons. Peak I_A amplitudes were normalized and plotted as a function of the prepulse potentials. Curves were fitted with a Boltzmann function and the dotted lines show the half-inactivation points.

Fig. 10. Effects of *slo*¹ on I_{CF} and I_{CS}

A, Percentage of distinctive CACs at different stages in WT and *slo*¹ neurons.

B, I_{CF} and I_{CS} densities (V_h at 10 mV) in WT and *slo*¹ at different developmental stages. * $P < 0.05$, ** $P < 0.01$ (student's t-test.). C, D, I-V curves of I_{CF} (C) and I_{CS} (D) for CACs in WT and *slo*¹ MP neurons. Type A and D subpopulations of CACs showed voltage-dependent activation with type A peaking at 10 mV, while type D exhibiting delayed activation and peaking at voltage more positive than 30 mV.

Fig. 11. Effects of *slo* on *I_A* and *Sh* expression

A, Increased *I_A* current density in *slo*¹ pupal neurons. The increase was more pronounced in neurons expressing type B than type A VGCs (cf. Table 1). *B*, Increased percentage of neurons expressing type B VGCs in *slo*¹ pupal neurons (cf. Table 1). *C*, Increased expression of *Sh* mRNA determined by qPCR using adult WT and *slo*¹ heads. *D*, Survival to adulthood in *slo*⁹⁸, *Sh*^M and double mutants. Flies of *Sh*^M/XX/Y;;*slo*⁹⁸/TM6 genotype were used to determine the eclosion rate of the single and double mutants. Four repeats of 25 male, 3rd instar larvae of each genotype were collected for the examination.

Table 1

Summary of I_A properties of type A and type B VGCs in WT and Sh^M neurons at L-III, MP, and Ad stages.

Stage	VGCs	Stock	n	%	pA/pF	T_p (ms)	$V_{1/2}$ (mV)	τ_{rec} (ms)
L-III	A	WT	10	17.5	68.3 (24.4)	4.2 (0.4)	-82.9 (1.0)	23.7 (1.2)
		Sh^M	14	56.0	95.1 (21.8)**	6.2 (0.3)*	-84.4 (1.2)	21.5 (1.3)
		slo^I	5	15.4	65.7 (21.2)	4.1 (0.4)	-83.7 (1.3)	24.5 (1.2)
	B	WT	40	70.2	111.6 (12.9)	3.1 (0.2)	-77.1 (0.6)	28.8 (1.2)
		Sh^M	5	20.0	57.4 (18.9)**	3.0 (0.8)	-84.6 (1.3)*	28.2 (1.6)
		slo^I	19	73.1	107.1 (14.8)	2.9 (0.2)	-79.3 (0.8)	29.5 (1.5)
MP	A	WT	3	6.9	102.4 (11.8)	4.0 (0.7)	-77.5 (1.7)	23.8 (1.1)
		Sh^M	6	17.9	81.8 (26.4)*	3.7 (0.5)	-80.4 (1.7)	21.2 (1.4)
		slo^I	3	4.8	121.7 (26.5)#	4.2 (0.4)	-80.8 (1.2)	24.3 (1.1)
	B	WT	11	31.8	58.5 (5.3)	3.0 (0.1)	-78.8 (0.9)	28.8 (1.0)
		Sh^M	6	21.4	34.3 (11.3)*	3.0 (0.1)	-78.0 (1.1)	26.5 (0.8)
		slo^I	13	61.9	89.6 (12.3)##	2.9 (0.4)	-76.3 (0.9)	27.1 (1.2)
Ad	A	WT	7	12.5	97.7 (26.0)	4.0 (0.3)	-82.1 (1.2)	21.7 (0.6)
		Sh^M	9	37.5	115.9 (21.9)*	7.5 (1.0)**	-83.1 (1.1)	22.2 (0.4)
		slo^I	3	5.3	110.9 (13.8)	4.0 (0.3)	-83.4 (1.4)	23.7 (0.7)
	B	WT	35	70.8	139.0 (16.1)	2.6 (0.2)	-75.1 (0.8)	27.9 (1.6)
		Sh^M	5	20.8	66.8 (27.5)**	3.2 (0.4)	-82.6 (1.5)*	27.1 (1.2)
		slo^I	10	52.6	125.2 (16.6)	2.8 (0.3)	-78.8 (1.0)	26.4 (1.0)

Values are means $\pm$ S.E.M. n and %, sample size and % cells. T_p , time to peak. $V_{1/2}$, half-inactivation voltage of steady-state inactivation fitted by Boltzmann function. τ_{rec} , recovery time constant obtained by one exponential fits.

* $P < 0.05$,

** $P < 0.01$, Sh^M VS WT;

$P < 0.05$,

$P < 0.01$, slo^I VS WT.

Table 2

Effects of *slo^f* and *Sh^M* on I_{CF} and I_{CS} of different CAC subtypes at L-III, MP, and Ad stages.

	Type A CACs			Type D CACs			Type E CACs			
	n	I _{CF}	I _{CS}	n	I _{CF}	I _{CS}	n	I _{CF}	I _{CS}	
L-III	WT	11	33.2 (5.6)	19.8 (5.3)	6	11.6 (4.4)	23.3 (7.1)	40	3.6 (0.6)	2.1 (0.3)
	<i>Sh^M</i>	5	39.7 (9.1)	16.6 (3.8)	4	14.2 (4.4)	27.7 (20.)	16	3.2 (0.4)	1.7 (0.4)
	<i>slo^f</i>	3	22.8 (0.3)	1.1 (0.3)	0			25	5.3 (0.6)	1.1 (0.4)
MP	WT	24	56.6 (6.3)	20.7 (2.1)	18	52.3 (13.8)	76.3 (17.1)	2	0.2 (1.6)	-0.12 (1.6)
	<i>Sh^M</i>	16	65.2 (9.3)	30.3 (6.5)	11	54.4 (9.7)	73.0 (18.3)	1	1.2	0.3
	<i>slo^f</i>	1	20.0	8.6	0			21	2.5 (0.7)	1.5 (0.5)
Ad	WT	12	41.5 (11.0)	15.8 (4.0)	7	38.4 (12.0)	46.9 (19.2)	28	2.7 (0.5)	2.0 (0.4)
	<i>Sh^M</i>	4	40.3 (9.0)	13.6 (6.4)	4	40.7 (8.6)	44.4 (16.9)	16	2.6 (0.6)	3.7 (0.9)
	<i>slo^f</i>	1	32.0	18.1	0			18	3.8 (0.6)	1.1 (0.5)

Values are means ± S.E.M.