Published in final edited form as:

J Trauma Nurs. 2017 ; 24(6): 365–370. doi:10.1097/JTN.0000000000000327.

An Outcome Analysis of Nurse Practitioners in Acute Care Trauma Services

Anna Holliday, MSN, FNP-BC,

Charleston Area Medical Center, Charleston, WV

Damayanti Samanta, MS,

Center for Health Services and Outcomes Research, Charleston Area Medical Center,
Charleston, WV

Julie Budinger, MSN, ACNP-BC, GNP-BC,

Charleston Area Medical Center, Charleston, WV

Jessica Hardway, DNP, NP-C, and

Charleston Area Medical Center, Charleston, WV

Audis Bethea, PharmD, BCPS

Center for Health Services and Outcomes Research, Charleston Area Medical Center,
Charleston, WV

Abstract

The Department of Trauma at a Level 1 trauma center sought to improve outcomes by enhancing the continuity of care for patients admitted to trauma services. Departmental leadership explored opportunities to improve this aspect of patient care through expansion of existing trauma Nurse Practitioner (NP) services. The restructured trauma NP service model was implemented in September 2013. A retrospective study was conducted with patients who presented at the trauma center between September 2012 and August 2015. Patients with at least a 24 hour hospital length

Corresponding Author: Audis Bethea, PharmD, BCPS, Clinical Research Scientist/Pharmacy Specialist, Trauma/Surgery, Research Scientist, West Virginia Clinical and Translational Science Institute, 3211 MacCorkle Ave. SE, Charleston, WV 25304, Phone 304-388-3653, Fax 304-388-7790, audis.bethea@camc.org.

Conflict of Interest: Anna Holliday has no conflict of interest to report.

Damayanti Samanta has no conflict of interest to report.

Julie Budinger has no conflict of interest to report.

Jessica Hardway has no conflict of interest to report.

Audis Bethea has no conflict of interest to report.

Financial Disclosure: Anna Holliday has no financial disclosures.

Julie Budinger has no financial disclosures.

Jessica Hardway has no financial disclosures.

Author Contribution: Anna Holliday conceived the study, provided critical revision of the manuscript and approved the final version of the manuscript.

Damayanti Samanta contributed to data review and analysis, writing and revision of this manuscript, and approved the final version of the manuscript.

Julie Budinger was involved in writing of the manuscript, provided critical revision of the manuscript, and approved the final version of the manuscript.

Jessica Hardway was involved in writing of the manuscript, provided critical revision of the manuscript, and approved the final version of the manuscript.

Audis Bethea, PharmD, BCPS provided oversight for all processes pertaining to the current study including design, data analysis, writing, critical revision of the manuscript, and approved the final version of the manuscript

of stay were separated into three comparator groups by 12-month increments. Twelve months pre-, 12 months during and 12 months post implementation. Data revealed improvement in hospital length of stay, Intensive Care Unit (ICU) length of stay, time to place rehabilitation consultation, 30-day readmission, missed injury(s), and placement of discharge orders before noon. A significant decline in the rate of complications including pneumonia and deep vein thrombosis were also noted. Accordingly, expansion of the trauma NP model resulted in significant improvements in patient, and process of care outcomes. This model for NP services may prove to be beneficial for acute care settings at other hospitals with high volume trauma services.

Keywords

Nurse practitioner; trauma; outcomes

Background/Significance

Within the last 20 years, hospitals are increasingly relying on advanced practice nurses to help meet the demand of patient care (Morris et al., 2012). According to the American Association of Nurse Practitioners ("NP Fact Sheet," 2017), there are more than 222,000 licensed nurse practitioners (NP) in the U.S. The number of NPs have escalated due to legislative amendments that sanctioned an expanded scope of practice for NPs, increased number of NP trainees per year, limitations on resident work hours, shortage of physicians, optimization of human and capital resources, and the growing and changing patient population (Morris et al., 2012; Rejtar, Rangstrom, & Allcox, 2016; Sise et al., 2011). The role of NPs has evolved over time and they have transitioned from the realm of primary care on to becoming an integral part of more challenging acute care services (Resler, Hackworth, Mayo, & Rouse, 2014).

Nurse practitioners are considered to be an economical and practical approach in coordinating patient care in many trauma services (Collins et al., 2014). Trauma care is affected by high costs associated with severe injuries and increasing number of patients without adequate insurance coverage. Rapid flow of patients through a trauma system is critical to operational efficiency (Dutton et al., 2003). Charleston Area Medical Center, a 821 bed tertiary care center in West Virginia provides Level 1 trauma services to southwestern West Virginia and areas of bordering states. In an attempt to improve outcomes through enhancing the continuity of care for patients admitted to trauma services, the Department of Trauma explored opportunities to expand existing trauma NP services. Efforts focused on processes that would augment continuity for patients as they transitioned through various levels of care during their respective hospitalization. The original NP model had 2 full-time NPs and focused primarily on providing care through the expedition of discharges. The expansion of services required strategically positioning NPs in multiple patient care settings, facilitating their involvement globally throughout all components of patients' admission; including ICU, floor, and post-acute clinic visits.

The revised trauma NP service model was implemented in September 2013. Services were expanded to meet identified needs and address perceived areas of weakness within the

current structure of the institution's trauma services. Accomplishing this task required NP services to establish a presence and practice in several new venues. The NPs became a daily participant in multidisciplinary rounds in the ICU, which served to provide a consistent team member presence as patients transitioned from the ICU to the floor. Under the supervision of an attending surgeon, the NP team also began managing stable, non-critical traumas in order to improve continuity and progression of care in this sub-population of trauma patients. A NP clinic was also created in order to continue continuity and follow up for these patients following their acute phase of care. To aid in maintaining the integrity of patients' care, the resident service team provided support to the NP efforts on an as needed basis while the trauma attending performed bedside rounds with the NPs on a daily basis. The evolution of the NP service required expansion of the team from 2 to 5 full-time NPs. The 3 additional NPs were hired at different time points within a period of one year from September 2013. The original service model provided in hospital coverage Monday–Friday; however, under the new model NPs provide in hospital service coverage 7 days a week. Moving forward, the aim is to sustain the current service and expand provider numbers into the future.

Purpose

With the institution's commitment to significant expansion of the NP service it was crucial to quantify its impact on patient care. The current study endeavored to comprehensively assess trends in outcomes of patients admitted to trauma services over a period of three years that included a year prior to, a year during, and a year post implementation of the NP service model (Morris et al., 2012; Sise et al., 2011; Resler et al., 2014; Collins et al., 2014; Dutton et al., 2003; Haan et al., 2007; Spisso, O'Callaghan, McKennan, & Holcroft, 1990; Christmas et al., 2005).

Research Questions/hypothesis

Expansion of nurse practitioner services facilitated an improvement in outcomes associated with trauma admissions.

Methods

After approval by the Institutional Review Board, a retrospective review of patients admitted to an 821 bed tertiary care, level 1 trauma center was conducted between September 2012 and August 2015. Study patients were identified by the institution's trauma registry. Identified patients were included if they were admitted to trauma services with a hospital length of stay of at least 24 hours. All patients meeting the study criteria were separated into three cohorts: 12 months prior to implementation of the NP service (September 2012 to August 2013), during implementation of the NP service (September 2013 to August 2014), and 12 months following the implementation of the NP service (September 2014 to August 2015). These three years of data were collected to evaluate changes in patient outcomes.

Following approval from the Institution's Review Board, patients were identified and data were obtained through the institution's trauma registry. Data included patient characteristics and outcomes: age, Injury Severity Score (ISS), Glasgow Comma Scale (GCS) score at the

scene of injury, Abbreviated Injury Severity (AIS) scores, pre-existing conditions, hospital length of stay (LOS), Intensive Care Unit (ICU) LOS, time to place rehabilitation consultation, 30-day readmission, missed injury rate, unplanned ICU admissions, discharge order placed before noon, and complications during hospital stay. Evaluated complications included pneumonia, urinary tract infection (UTI), surgical site infection/deep surgical infection (SSI/DSI), acute renal failure, deep vein thrombosis (DVT), pulmonary embolism (PE), and sepsis.

Descriptive statistics were computed for each variable to describe the patient population. Continuous variables examining patient characteristics were reported as means and standard deviations and compared using one-way ANOVA. Categorical variables were presented as proportions and compared using chi-square or Fisher's Exact test. Outcome variables were analyzed with ANCOVA or binary logistic regression adjusting for differences in patient characteristics. An exception was made for ISS as differences in average ISS over the study period were of minimal clinical relevance. All comparisons were performed at a level of significance of $p = 0.05$. Data analysis was performed using SPSS Version 22.0 (IBM Corp, Armonk, NY).

Results

A total of 3,284 trauma patients were included in the analysis; 1,088 prior to implementation, 1,009 during the 12 months of implementation, and 1,187 in the first 12 months following implementation of the NP service. Average age and GCS of patients remained comparable, however, ISS significantly declined over the study period (13.09 vs. 12.08 vs. 11.77, $p=0.000$). In addition to ISS as an injury severity marker, AIS for the six body regions were examined and they remained equivalent over the study period. With regard to comorbidities, the proportion of patients with pre-existing conditions of neurologic disease (6.3% vs. 9.9% vs. 9.0%, $p=0.006$) and diabetes (12.3% vs. 12.3% vs. 15.3%, $p=0.050$) increased significantly over time. The percentage of patients with pre-existing pulmonary conditions, on the other hand, significantly decreased (5.5% vs. 7.8% vs. 2.4%, $p=0.000$), (Table 1).

Hospital LOS significantly declined over time. During the year of NP service implementation, hospital LOS decreased by 0.04 days and it further declined by 0.98 days in the first year following implementation (6.92 vs. 6.88 vs. 5.94, $p=0.007$). Similarly, ICU LOS significantly decreased (4.64 vs. 4.27 vs. 3.56, $p=0.001$) during the study periods. The time from admission to placement of rehabilitation consultations also significantly improved following NP service implementation (4.88 vs. 2.66 vs. 3.63, $p=0.000$). Although the time to rehabilitation consult increased during the first year post-implementation, it remained significantly lower than the pre-implementation time (4.88 vs. 3.63, $p=0.001$). Discharge orders placed before noon also significantly improved over the course of the study (47.0% vs. 50.3% vs. 64.3%, $p=0.000$). Changes in thirty-day readmission (3.8% vs. 3.1% vs. 2.0%, $p=0.078$) and missed injury rate (1.0% vs. 0.4% vs. 0.3%, $p=0.061$) improved, but did not reach statistical significance. Rates of unplanned ICU admissions did not significantly change (Table 2). Inpatient complication rates for pneumonia (8.6% vs. 8.3% vs. 5.1%,

p=0.002) and DVT (3.4% vs. 3.5% vs. 1.9%, p=0.028) also significantly decreased post-implementation of the NP service (Table 3).

Discussion

Trauma patients often present with complex injuries requiring coordination of care among various physician as well as non-physician services. Many trauma services have included NPs to improve continuity of patient care and overcome potential hindrances created by the reduction in resident hours (Haan et al., 2007). The Level 1 trauma center at the current study's institution expanded their existing NP model in September 2013 to promote continuity of patient care and improve outcomes.

Following expansion of the NP service outcomes of hospital and ICU LOS significantly decreased over time. The decrease in average hospital LOS by 0.98 days from 12 months prior, to 24 months post-implementation of the service equated to a decrease of approximately \$1.1 million in hospital charges per year. These outcomes are consistent with the findings of previous studies that have examined the impact of the addition of NPs to patient care models (Sise et al., 2011; Collins et al., 2014; Spisso et al., 1990; Christmas et al., 2005). Collins et al evaluated the impact of a pilot program in which NPs were assigned to provide care 5 days a week in the stepdown area of their Level 1 trauma center. The implementation of this NP practice model resulted in a decrease in overall trauma service hospital LOS by 0.8 days (Collins et al., 2014).⁶ Similar findings were also reported by Spisso et al. (1990) in a study comparing outcomes pre- and post-implementation of NPs in their trauma service. In this study an average decrease in hospital LOS of 1.05 days was seen in patients admitted to their service one year prior to, and post-implementation of the NP service. Two additional studies conducted by Christmas et al. (2005) and Sise et al. (2011) also demonstrated significant decline in ICU LOS following integration of NPs in their trauma care unit.

The current study also indicated that the expansion of NP services had a positive impact on the rate of missed injuries. Although missed injuries are not usually life threatening, they can result in long term disability for a patient (Kalemoglu et al., 2006). The rate of missed injuries decreased over the study period, but fell just short of achieving statistical significance. Similar findings were reported in a study conducted by Resler et al. (2014) in a pediatric trauma center. They reported that a significant percentage of missed injuries were identified with the addition of acute care trained pediatric nurse practitioners. The decline in the incidence of missed injuries in both studies was likely a result of improved in charting, consistency in the practitioners performing examinations, and improved tracking of injury lists.

Time to placement of rehabilitation consultation from admission and placement of discharge orders before noon were two additional outcomes that improved following implementation of the NP service at the study institution. These outcomes have not been previously examined in current literature assessing the impact of acute care NPs on inpatient outcomes. Identifying trauma patients with rehabilitation needs and early intervention likely serves to further expedite discharge from the acute care setting. In addition, previous literature

suggests that early involvement of rehabilitation services contributes to improved functional outcomes following traumatic injury (Lawrence, 2006). Similarly, placement of a discharge order(s) before noon allows for improvements in the flow of patients through the trauma system.

Post-admission complications of pneumonia and DVT were found to be significantly lower following the expansion of the NP model. Improvements in pneumonia rates were likely influenced by the significant decrease in the number of admitted patients with baseline pulmonary disease during the study period. Pneumonia rates may have also been impacted by the improvements in patients' LOS seen with the implementation of the expanded NP services model, thus decreasing the overall risk of nosocomial pneumonia. Significant decreases in DVT rates were likely influenced by routine NP evaluation of all trauma service patients for the presence or absence of pharmacologic DVT prophylaxis within the first 48 hours of admission. These select patients were followed for appropriate pharmacologic prophylaxis. This process continued until pharmacologic prophylaxis was initiated or a formal plan for the initiation of pharmacologic prophylaxis was developed and documented. Intuitively this process likely influenced DVT rates for the service by decreasing unnecessary delays in the initiation of pharmacologic DVT prophylaxis.

Readmission rates also declined over the course of the study. The overall readmission rate decreased by 1.8% resulting in 21 fewer patient readmissions during the final 12 month study period. While this improvement fell just short of statistical significance the decrease in hospital readmissions would have further contributed to the overall cost savings associated with the expansion of the NP model at the study institution. The impact on readmissions with NP involvement in the acute management of trauma patients has been evaluated in a previous study conducted by Morris et al. This study also conducted in a Level 1 trauma center, compared readmission rates between NP and resident coordinated management. This study, however, reported similar readmission rates between both patient care models (Morris et al., 2012).

Conclusion

This study provides a comprehensive review of outcomes related to the involvement of NPs in the continuum of care for trauma services patients. While there are several studies evaluating the impact of NPs in the provision of care in the trauma patient population, these studies have frequently been more limited in scope.^{1,5-6,9,10} Data from the current study, however, serves to comprehensively describe the impact of NPs involvement in provision of care to trauma patients. These findings support data from previous studies' by broadly evaluating morbidity and process of care variables that serve to directly impact LOS and ultimately an institutions' ability to provide quality care to a high volume of patients. Following expansion of the trauma NP model, improvement in patient outcomes including hospital LOS, ICU LOS, 30-day readmission, time to place rehabilitation consultation, missed injury, discharge order placement before noon, and complications of pneumonia and DVT were noted. The trauma NP service model developed at this study institution may prove beneficial in acute care settings at other institutions with high volume trauma services. The future goal for the NP service at the study institution is to continue to expand the model

to provide 24*7 days coverage. Success of the evolving model will rely on significant support from the medical, nursing, and administration teams.

Limitations

Although this study reports improved outcomes following expansion of the NP model, the study is not without limitations. The retrospective design of the study makes it reliant upon the identification, availability and completeness of patients' medical records. Accordingly, all patients admitted to the trauma center during the study period may not have been identified and included resulting in selection bias. In addition, the validity of the evaluated data depends on the accuracy with which it has been documented and subsequently collected. This serves to limit researchers' ability to practice quality control with the data. Also, this study does not seek to establish a causal relationship between the implementation of the NP service model and positive outcomes. It only describes the changes in outcomes reported prior to and post implementation of the model.

Acknowledgments

The authors would like to thank Teresa White MSN, FNP-BC, Tara Miller, MSN, FNP-BC, Kelly Jo Evans, BSN, RN, and Julton Tomanguillo MD for providing critical revision of the manuscript.

Audis Bethea is a research scientist for the West Virginia Clinical Translational Science Institute. "Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number U54GM104942. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health."

Damayanti Samanta is a member of the West Virginia Clinical Translational Science Institute. "Research reported in this publication was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number U54GM104942. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health."

References

- Morris DS, Reilly P, Rohrbach J, Telford G, Kim P, Sims CA. The influence of unit-based nurse practitioners on hospital outcomes and readmission rates for patients with trauma. *J Trauma Acute Care Surg.* 2012; 73(2):474–8. DOI: 10.1097/TA.0b013e31825882bb [PubMed: 22846959]
- "NP Fact Sheet." AANP, American Association of Nurse Practitioners. Jun 6. 2017 Retrieved from, www.aanp.org/all-about-nps/np-fact-sheet
- Rejtar M, Ransstrom L, Allcox C. Development of the 24/7 Nurse Practitioner Model on the Inpatient Pediatric General Surgery Service at a Large Tertiary Care Children's Hospital and Associated Outcomes. *J Pediatr Health Care.* 2017; 31(1):131–140. DOI: 10.1016/j.pedhc.2016.08.007 [PubMed: 27692506]
- Sise CB, Sise MJ, Kelley DM, Walker SB, Calvo RY, Shackford SR, Lome BR, Sack DI, Osler TM. Resource commitment to improve outcomes and increase value at a level I trauma center. *J Trauma.* 2011; 70(3):560–8. DOI: 10.1097/TA.0b013e31820c7b79 [PubMed: 21610343]
- Resler J, Hackworth J, Mayo E, Rouse TM. Detection of missed injuries in a pediatric trauma center with the addition of acute care pediatric nurse practitioners. *J Trauma Nurs.* 2014; 21(6):272–5. quiz 276–7. DOI: 10.1097/JTN.0000000000000080 [PubMed: 25397332]
- Collins N, Miller R, Kapu A, Martin R, Morton M, Forrester M, Atkinson S, Evans B, Wilkinson L. Outcomes of adding acute care nurse practitioners to a Level I trauma service with the goal of decreased length of stay and improved physician and nursing satisfaction. *J Trauma Acute Care Surg.* 2014; 76(2):353–7. DOI: 10.1097/TA.0000000000000097 [PubMed: 24398767]

- Dutton RP, Cooper C, Jones A, Leone S, Kramer ME, Scalea TM. Daily multidisciplinary rounds shorten length of stay for trauma patients. *J Trauma*. 2003; 55(5):913–9. [PubMed: 14608165]
- Haan JM, Dutton RP, Willis M, Leone S, Kramer ME, Scalea TM. Discharge rounds in the 80-hour workweek: importance of the trauma nurse practitioner. *J Trauma*. 2007; 63(2):339–43. [PubMed: 17693833]
- Spisso J, O'Callaghan C, McKennan M, Holcroft JW. Improved quality of care and reduction of housestaff workload using trauma nurse practitioners. *J Trauma*. 1990; 30(6):660–3. discussion 663–5. [PubMed: 2112612]
- Christmas AB, Reynolds J, Hodges S, Franklin GA, Miller FB, Richardson JD, Rodriguez JL. Physician extenders impact trauma systems. *J Trauma*. 2005; 58(5):917–20. [PubMed: 15920403]
- Kalemoglu M, Demirbas S, Akin ML, Yildirim I, Kurt Y, Uluutku H, Yildiz M. Missed injuries in military patients with major trauma: original study. *Mil Med*. 2006; 171(7):598–602. [PubMed: 16895123]
- Lawrence, RR. *Trauma Rehabilitation*. Philadelphia, PA: Lippincott Williams & Wilkins; 2006.

Key Points

- Expansion of nurse practitioner services resulted in an improvements in patient outcomes.
- Post-implementation data revealed improvements in time to placement of discharge orders, ICULOS and HLOS.
- An investment in increasing nurse practitioners presence on trauma services proved to be valuable institutional investment.

Table 1
Patient Characteristics

	Sep 2012 to Aug 2013 (N=1088)	Sep 2013 to Aug 2014 (N=1009)	Sep 2014 to Aug 2015 (N=1187)	P-Value
Age	44.60±20.62	46.16±21.48	46.49±20.95	0.079
Injury Severity Score	13.19±8.70	12.08±7.95	11.77±7.90	0.000
Glasgow Comma Scale Score at Scene	13.61±3.13	13.57±3.19	13.84±2.80	0.297
Abbreviated Injury Score				
Head/neck	2.58±1.02	2.56±0.99	2.59±1.01	0.891
Face	1.78±0.50	1.76±0.53	1.74±0.51	0.638
Chest	2.52±0.79	2.57±0.82	2.55±0.77	0.629
Abdomen/pelvic	2.47±0.80	2.43±0.73	2.31±0.80	0.054
Extremities/pelvic girdle	2.27±0.58	2.24±0.56	2.30±0.60	0.338
Comorbidities				
Pulmonary	5.5%	7.8%	2.4%	0.000
Cardiovascular	34.7%	36.1%	38.3%	0.142
Neurologic	6.1%	9.6%	8.8%	0.007
Gastrointestinal	1.5%	2.0%	1.7%	0.663
Psychiatric	56.2%	55.1%	57.4%	0.563
Renal	0.6%	1.1%	0.8%	0.503
Diabetes	12.3%	12.3%	15.3%	0.050
Hematologic	6.9%	9.0%	8.8%	0.146
Cancer	0.3%	0.3%	0.1%	0.481
Autoimmune	1.4%	1.9%	1.7%	0.656

Boldface indicates statistical significance (p < 0.05)

Table 2
Study Outcomes

	Sep 2012 to Aug 2013 (N=1088)	Sep 2013 to Aug 2014 (N=1009)	Sep 2014 to Aug 2015 (N=1187)	P-Value
Hospital length of stay <i>d</i>	6.92±8.50	6.88±9.13	5.94±8.27	0.007
ICU length of stay <i>d</i>	4.64±5.31	4.27±5.17	3.56±4.47	0.001
Time to place rehabilitation consult <i>d</i>	4.88±4.42	2.66±3.19	3.63±3.98	0.000
30-day readmission	3.8%	3.1%	2.0%	0.078
Missed injury rate	1.0%	0.4%	0.3%	0.061
Unplanned ICU admission	3.0%	3.5%	2.2%	0.346
Discharge order placed before noon	47.0%	50.3%	64.3%	0.000

Boldface indicates statistical significance (p 0.05)

Table 3
List of Complications

	Sep 2012 to Aug 2013 (N=1088)	Sep 2013 to Aug 2014 (N=1009)	Sep 2014 to Aug 2015 (N=1187)	P-Value
Pneumonia	8.6%	8.3%	5.1%	0.002
Urinary tract infection	2.5%	4.1%	2.6%	0.125
Surgical site infection/deep surgical infection	0.9%	0.7%	0.8%	0.806
Acute renal failure	4.5%	4.5%	3.0%	0.084
Deep vein thrombosis	3.4%	3.5%	1.9%	0.028
Pulmonary embolism	1.2%	0.7%	0.8%	0.454
Sepsis	0.9%	1.7%	0.8%	0.135

Boldface indicates statistical significance (p 0.05)