DISCUSSION

This technique is inexpensive and successful in preventing lead migration during trials of sacral nerve stimulation. Use of the securement device has enabled more reliable trials of sacral nerve stimulation.

References

1. National Institute for Health and Care Excellence. *Overactive Bladder in Women: Management*. Clinical Guideline CG171. London: NICE; 2015.
2. Tawadros C, Burnett KE, Betts C. Neuromodulation by Sacral Nerve Stimulation. In: Payne S, Eardley I, O'Flynn KJ, eds. *Imaging and Technology in Urology: Principles and Clinical Applications*. Dordrecht: Springer; 2012, pp307–310.
3. National Institute for Health and Care Excellence. *Faecal Incontinence in Adults: Management*. Clinical Guideline CG49. London: NICE; 2007.

Reinforcement of sutures to close surgical incisions in frail skin

CP Wakeling, A Sandiford

Trauma and Orthopaedics, St George's Hospital, London, UK

CORRESPONDENCE TO

Chris Wakeling, E: cpwakeling@gmail.com
doi 10.1308/rcsann.2017.0139

Adequate surgical incision closure protects against infection and helps produce satisfactory scars. This can be challenging with frail, 'paper-thin' older skin, unable to hold a subcuticular or skin suture without cutting out.

Using adhesive strips perpendicularly across a traumatic wound prior to suturing has been described previously.¹ Our alternative technique places 1/8" SteriStrips™ (3M Corporation, St Paul, MN, USA) longitudinally along elective incision margins, parallel to the line of incision, reinforcing the skin before suturing. A heavy nylon suture is then placed through this strip-augmented skin, permitting satisfactory

closure (Fig 1). SteriStrips and suture are removed at 14 days, the adhesive having loosened enough to avoid skin damage.

Reference

1. Nawaz SZ, Chan O, Newman K. Simple technique for closure of skin lacerations under tension. *Ann R Coll Surg Engl* 2011; **93**(8): 650.

Smartphone flashlight to aid in locating lost and dropped surgical sharps

G Aldous, J Mathers

Department of Anaesthesia, Lister Hospital, Stevenage, UK

CORRESPONDENCE TO

J Mathers, E: j.mathers@nhs.net
doi 10.1308/rcsann.2017.0182

BACKGROUND

Loss of a suture needle is a known risk with any surgical procedure that can be noted during or at the end of surgery when the final instrument counts are completed. Accounting for instruments forms a part of the World Health Organization surgical checklist.¹ Normal protocol in most UK hospitals requires the use of fluoroscopic screening of patients to confirm that no instruments have been retained within the patient, introducing surgical delay and radiation exposure. Decision algorithms have been developed to aid in decision making when a needle is lost² but apart from the use of magnets or sticking tape there have been no advances in searching for lost needles. We describe a simple, free technique to search for a needle that may have been dropped on the floor that can be used while waiting for x-ray.

TECHNIQUE

Our technique uses the widespread availability of smartphones and the flash that accompanies them. When a needle has possibly fallen on the floor, the operating theatre lights are dimmed and multiple smartphones are used in a constant sweeping motion around the theatre floor. The constant movement allows for those searching to look for a glint of reflected light, which is easy to locate. Once the needle has been located it can be retrieved using adhesive tape.

DISCUSSION

Normal hospital protocols should always be used when a lost needle or sharp is suspected. However, this simple technique can help to locate a lost needle during the period of waiting for x-ray.

References

1. Haynes AB, Weiser TG, Berry WR *et al.*, Safe Surgery Saves Lives Study Group. A surgical safety checklist to reduce morbidity and mortality in a global population. *N Engl J Med* 2009; **360**(5): 491–499.
2. Zaman S, Clarke R, Schofield A. Intraoperative loss of a surgical needle: a laparoscopic dilemma. *JSLs*; **19**(2): e2013.00401.

Figure 1 SteriStrip™ sutured wound.