Published in final edited form as:

Prev Med. 2018 June ; 111: 348–350. doi:10.1016/j.ypmed.2017.11.022.

Vascular Risk Factor Burden and New-Onset Depression in the Community

Shayna Adams, MPH^{a,b}, Sarah Conner, MPH^{b,c}, Jayandra J. Himali, PhD^{a,b,c}, Alexa Beiser, PhD^{a,b,c}, Ramachandran S. Vasan, MD^{b,d,e}, Sudha Seshadri, MD^{a,b,*}, and Matthew P. Pase, PhD^{a,b,f,*}

^aDepartment of Neurology, Boston University School of Medicine, 72 East Concord St. Boston, MA, 02118, USA

^bFramingham Heart Study, 73 Mt. Wayte Avenue, Suite 2, Framingham, MA 01702-5827 MA, USA

^cDepartment of Biostatistics, Boston University School of Public Health, 715 Albany St, Boston, MA 02118, USA

^dSections of Preventive Medicine and Epidemiology, Department of Medicine, Boston University School of Medicine, 801 Massachusetts Ave, Boston, MA, 02118, USA

^eDepartment of Epidemiology, Boston University School of Public Health, 801 Massachusetts Ave, Boston, MA, 02118, USA

^fCentre for Human Psychopharmacology, Swinburne University of Technology, Burwood Rd, Hawthorn, 3122, Australia

Abstract

Depression is associated with an increased likelihood of cardiac events and stroke. We hypothesized that the vascular risk factor burden might itself predispose to both cardiovascular events and depression. Therefore, we examined whether aggregate scores of vascular risk factor burden were associated with the new-onset of depression in the community. We studied 2023 depression- and dementia-free Framingham Heart Study (Framingham, USA) Offspring participants who attended both examination cycles 7 (1998-2001) and 8 (2005-2008). The American Heart Association Ideal Cardiovascular Health metric and the Framingham Stroke Risk Profile were calculated at exam seven. New-onset depression was adjudicated at examination cycle eight as antidepressant medication use or Centre for Epidemiologic Studies Depression Scale scores ≥ 16 , after a mean follow-up of 6.6 years (standard deviation = 0.7). Of the 2023 participants, 269 (13%) developed new-onset depression. Following adjustments for age, sex,

Corresponding Author: Dr. Matthew P. Pase, Department of Neurology, Boston University School of Medicine & Framingham Heart Study, 72 East Concord St. Boston, 02118, MA. matthewpase@gmail.com; T 1(617) 638-8064.

*Joint senior authors

Conflict of interest: All other authors report no disclosures or potential conflicts of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

education, and the time interval between baseline and follow-up, the odds of new-onset depression decreased by 10% for each one-point increase in ideal cardiovascular health scores (Odds Ratio [OR], 0.90; 95% confidence interval [CI] 0.81-0.99) and increased by 4% for each percentage point increase in the Framingham stroke risk profile (OR, 1.04; CI, 1.00-1.07). Results were not explained by interim clinical stroke or cerebral white matter injury. In conclusion, vascular risk factor burden was associated with the new onset of depression. Shared vascular risk factors may contribute to the increased risk of cardiovascular events observed in persons with depression.

Introduction

Depression and cardiovascular disease are leading causes of disability in the developed world.¹ Evidence suggests that depression and cardiovascular disease may share a bidirectional relationship. Vulnerability to both conditions may be due to shared genetic or environmental risk factors; Depression is associated with an increased likelihood of cardiac events and stroke, presumably because of its association with vascular risk factors such as smoking, obesity, physical inactivity, poor diet, and shared genetic factors.²⁻⁵ We hypothesized that vascular risk factor burden might also predispose to the risk of new-onset depression, thereby establishing common factors underlying both risks for cardiovascular disease and depression.

The ideal-cardiovascular health metric was developed as part of the American Heart Association's 2020 impact goals.⁶ The metric awards one point for meeting each recommended target for four health behaviors (non-smoking, body mass index, physical activity, and diet) and three risk factors (serum total cholesterol, blood pressure, and fasting blood glucose) with a score of seven reflecting ideal-cardiovascular health. We recently demonstrated that higher ideal-cardiovascular health scores were associated with a lower risk of incident cardiovascular disease, stroke, dementia, brain atrophy and cognitive decline.^{7,8} In our previous study, each point higher ideal-cardiovascular health scores was associated with a 20% lower chance of stroke within ten years amongst participants of the Framingham Heart Study.⁷ The Framingham stroke risk profile is a widely used algorithm to estimate the 10-year risk of stroke based on clinical information.⁹

The aim of the present study was to examine the prospective association between overall vascular risk factor burden, using the ideal-cardiovascular health and Framingham stroke risk profile metrics, and new-onset depression in participants of the community-based Framingham Heart Study.

Methods

The Framingham Heart Study Offspring cohort commenced in 1971 with the enrollment of 5124 volunteers.¹⁰ Participants have been studied across 9 examination cycles, once approximately every four years. Between 1998 and 2001, 3539 Offspring participants attended the seventh Heart Study examination cycle. Using these data as a baseline, we captured ideal-cardiovascular health and Framingham stroke risk profile scores for 2750 participants. We excluded 372 participants who were deemed to be depressed at baseline based on the use of antidepressants or a Center for Epidemiologic Studies Depression Scale

(CES-D) score ≥ 16 , and 6 participants with prevalent dementia. Of the remaining 2372 participants, our analysis sample included 2023 persons who returned for follow-up at the eighth heart study examination cycle (2005-2008) and had complete data on medication use and CES-D scores. All participants provided written informed consent, and the study was approved by the Institutional Review Board and Boston University Medical Center.

Assessment of vascular risk

Consistent with our previous work,^{7,8} the ideal-cardiovascular health score was operationalized by awarding one point for each of the following: never smoked or quit >12 months ago, body mass index $<25 \text{ kg/m}^2$, adequate physical activity (≥ 150 minutes/week moderate intensity or ≥ 75 min/week vigorous intensity or a combination of the two), a healthy diet (≥ 2 of the following components: >4.5 cups/day fruits and vegetables, $\geq 2 \times 3.5$ oz servings/week of fish, $\geq 3 \times 1$ oz servings/day of fiber-rich whole grains, $<1500 \text{ mg/d}$ of sodium, & ≤ 36 oz/week of sugar-sweetened beverages), serum total cholesterol untreated and $<200 \text{ mg/dL}$, blood pressure untreated and $<120/80 \text{ mmHg}$ (measured twice by a physician with results averaged), and fasting plasma glucose untreated $<100 \text{ mg/dL}$. Diet and physical activity were measured with the use of a validated Harvard semi-quantitative food frequency¹¹ and physical activity index questionnaire,¹² respectively, whereas all other variables were obtained from the Framingham Heart Study clinic. Smoking status was assessed in a clinical interview. Blood pressure was measured by a physician using a mercury sphygmomanometer with the participant supine and following 5 minutes rest. Fasting cholesterol was measured in 99% of our sample. We calculated the recently updated Framingham stroke risk profile according to published criteria.⁹ The algorithm combines information on age, current smoking status, prevalent cardiovascular disease, prevalent atrial fibrillation, diabetes, hypertension treatment and systolic blood pressure. This updated algorithm has been validated in 3 large community samples.⁹ Higher Framingham stroke risk profile scores indicate a higher risk of incident stroke.

Assessment of new-onset depression

New-onset depression was defined at examination cycle eight as the use of antidepressants or CES-D scores ≥ 16 . This CES-D threshold has been shown to have high sensitivity and specificity for detecting major depression in community-dwelling older adults.¹³

Statistical analysis

We used logistic regression models to relate continuous ideal-cardiovascular health and Framingham stroke risk profile scores at examination cycle seven to the new onset of depression at examination cycle eight (mean follow-up of 6.6 years, SD 0.7). Results are expressed as odds ratios (OR) and 95% confidence intervals (CI) and are adjusted for age, sex, education and the time interval between baseline and follow-up.

We performed sensitivity analysis to examine whether any associations between vascular risk and new-onset depression could be explained by clinical stroke or silent cerebral small vessel disease. Therefore, we included additional adjustments for prevalent stroke ($n=25$) at baseline and as well as prevalent stroke combined with interim stroke between baseline and follow-up ($n=51$). Next, we examined mediation by white matter hyperintensity volume,

which was quantified at a date proximal to baseline using a Siemens 1T or 1.5T field strength machine with a T2-weighted double spin-echo coronal imaging sequence in contiguous slices of 4mm.¹⁴ Values of white matter hyperintensity volume were expressed as a percentage of intracranial volume and were log transformed. Lastly, we examined the association between vascular risk and the onset of depressive symptoms, defined solely by CES-D scores ≥ 16 in participants who were not using antidepressants (participants taking antidepressants were excluded from analysis). Results were considered significant if $p < 0.05$.

Results

Cohort demographics are summarized in Table 1 with results displayed in Table 2. The age range of participants was from 37 to 88 years. Of the 2023 participants, 269 (13%) developed new-onset depression at follow-up. The odds of new-onset depression decreased by 10% for each one-point increase in ideal-cardiovascular health scores. Each percentage point increase in the Framingham stroke risk profile was associated with a 4% higher odds of new-onset depression. Effect sizes were similar when examining the association between vascular risk and the new onset of depressive symptoms (after excluding persons taking antidepressants). Effect sizes were also comparable after additional adjustment for prevalent stroke at baseline as well as both prevalent and interim stroke. White matter hyperintensity volume was not associated with new-onset depression (Odds Ratio, 1.03; 95% confidence interval, 0.88-1.22) and was not considered to be a potential mediator.

Discussion

Our large prospective population-based study demonstrates that higher ideal-cardiovascular health and lower Framingham stroke risk profile scores were associated with a lower odds of new-onset depression, as defined by new antidepressant use or the development of significant depressive symptomatology. Results were similar after adjustments for clinical stroke and were not accounted for by subclinical white matter injury. Collectively, our results suggested that vascular risk factor burden is associated with new-onset depression across the adult age range, independent of overt stroke and subclinical white matter disease.

Whereas vascular risk factors have been hypothesized to underlie the association between prevalent depression and cardiovascular events,³ our results suggest that the adherence to simple cardiovascular health targets was associated with a lower risk of developing depression in the first instance. This suggests that the association between depression and cardiovascular disease may be due to shared vascular risk factor burden. Our findings also extend our previous findings demonstrating that higher ideal-cardiovascular health scores were associated with a lower risk of incident cardiovascular disease, stroke, dementia, brain atrophy and cognitive decline.^{7,8} Further research is warranted to determine whether improving cardiovascular health lowers the risk of developing depression. This is an important topic because further promotion of heart-healthy guidelines, such as the ideal-CVH guidelines, may help lower the burden of cardiovascular disease and depression in the community.

The present study was not without limitations. The sample was comprised mostly of Caucasians, thereby limiting the generalizability of our findings to other ethnic groups. The associations between vascular risk factor burden and new-onset depression could not be explained by overt stroke or white matter hyperintensities. However, we cannot exclude the possibility that more subtle cerebrovascular disease (undetectable by our imaging methods) was responsible for the observed associations. More sensitive imaging measures of white matter integrity such as fractional anisotropy or free water diffusion will be required to explore this further. Further understanding the mechanisms linking vascular risk to new-onset depression may help clarify whether reducing cardiovascular disease risk can mitigate the risk of depression. Moreover, we did not have data on current or previous psychiatric disorders to include in our analysis. Lastly, new-onset depression was defined according to CES-D scores or medication use, rather than a physician diagnosis of clinical depression. It is possible that some participants, not captured as being depressed at baseline, may have had depression earlier in life.

In summary, vascular risk factor burden was associated with the new-onset of depression. Further research is warranted to replicate our findings and to evaluate whether reducing vascular risk factors attenuates the risk of developing depression.

Acknowledgments

Dr. Pase is funded by an Australian National Health and Medical Research Council Early Career Fellowship (APP1089698). The Framingham Heart Study is supported by the National Heart, Lung and Blood Institute (contracts N01-HC-25195 and HHSN268201500001I), the National Institute of Neurological Disorders and Stroke (NS017950, UH2 NS100605) and the National Institute on Aging (AG033193, AG033040, AG049505, AG049607, AG054076).

References

1. Kassebaum NJ, Arora M, Barber RM, et al. Global, regional, and national disability-adjusted life-years (DALYs) for 315 diseases and injuries and healthy life expectancy (HALE), 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *The Lancet*. 2016; 388(10053): 1603–1658.
2. Xian H, Scherrer JF, Franz CE, et al. Genetic vulnerability and phenotypic expression of depression and risk for ischemic heart disease in the Vietnam era twin study of aging. *Psychosomatic Medicine*. 2010; 72(4):370–375. [PubMed: 20190130]
3. Penninx BWJH. Depression and cardiovascular disease: Epidemiological evidence on their linking mechanisms. *Neuroscience & Biobehavioral Reviews*. 2017; 74, Part B:277–286. [PubMed: 27461915]
4. Dong JY, Zhang YH, Tong J, Qin LQ. Depression and Risk of Stroke: A Meta-Analysis of Prospective Studies. *Stroke*. 2012; 43(1):32–37. [PubMed: 22020036]
5. Kendler KS, Fiske A, Gardner CO, Gatz M. Delineation of Two Genetic Pathways to Major Depression. *Biological Psychiatry*. 2009; 65(9):808–811. [PubMed: 19103442]
6. Lloyd-Jones DM, Hong Y, Labarthe D, et al. Defining and Setting National Goals for Cardiovascular Health Promotion and Disease Reduction: The American Heart Association's Strategic Impact Goal Through 2020 and Beyond. *Circulation*. 2010; 121(4):586–613. [PubMed: 20089546]
7. Pase MP, Beiser A, Enserro D, et al. Association of Ideal Cardiovascular Health With Vascular Brain Injury and Incident Dementia. *Stroke*. 2016; 47(5):1201–1206. [PubMed: 27073239]

8. Xanthakis V, Enserro DM, Murabito JM, et al. Ideal cardiovascular health: associations with biomarkers and subclinical disease and impact on incidence of cardiovascular disease in the Framingham Offspring Study. *Circulation*. 2014; 130(19):1676–1683. [PubMed: 25274000]
9. Dufouil C, Beiser AS, McClure LA, et al. A Revised Framingham Stroke Risk Profile to Reflect Temporal Trends. *Circulation*. 2017; 135(12):1145–1159. [PubMed: 28159800]
10. Feinleib M, Kannel WB, Garrison RJ, McNamara PM, Castelli WP. The framingham offspring study. Design and preliminary data. *Preventive Medicine*. 1975; 4(4):518–525. [PubMed: 1208363]
11. Rimm EB, Giovannucci EL, Stampfer MJ, Colditz GA, Litin LB, Willett WC. Reproducibility and Validity of an Expanded Self-Administered Semiquantitative Food Frequency Questionnaire among Male Health Professionals. *American Journal of Epidemiology*. 1992; 135(10):1114–1126. [PubMed: 1632423]
12. Kannel WB, Sorlie P. Some health benefits of physical activity: The framingham study. *Archives of Internal Medicine*. 1979; 139(8):857–861. [PubMed: 464698]
13. Beekman ATF, Deeg DJH, Van Limbeek J, Braam AW, De Vries MZ, Van Tilburg W. Criterion validity of the Center for Epidemiologic Studies Depression scale (CES-D): Results from a community-based sample of older subjects in the Netherlands. *Psychological Medicine*. 1997; 27(1):231–235. [PubMed: 9122304]
14. DeCarli C, Massaro J, Harvey D, et al. Measures of brain morphology and infarction in the framingham heart study: establishing what is normal. *Neurobiology of Aging*. 2005; 26(4):491–510. [PubMed: 15653178]

Abbreviations

CES-D	Center for Epidemiologic Studies Depression Scale
CI	Confidence interval
OR	Odds Ratio

Highlights

- We examined whether vascular risk burden was associated with new-onset depression
- Better vascular health predicted a reduced risk of new-onset depression
- Higher stroke risk predicted an increased risk of new-onset depression

Table 1
Characteristics of the study sample at baseline

	Sample mean (SD) ^a
Age, years	61 (9)
Men, n (%)	946 (47)
Education, n (%)	
No High School	68 (3)
High-School Degree	1124 (57)
College Graduate	790 (40)
Body Mass Index, kg/m ² , median [Q1, Q3]	27 [24, 31]
Systolic blood pressure, mm Hg	126 (18)
Hypertension treatment, n (%)	638 (32)
Total cholesterol, mg/dL	201.0 (35.9)
High density lipoprotein cholesterol, mg/dL	53.8 (16.6)
Prevalent diabetes, n (%)	185 (9)
Prevalent atrial fibrillation, n (%)	65 (3)
Prevalent cardiovascular disease, n (%)	211 (10)
CES-D, score units, median [Q1, Q3]	3 [1, 5]
Ideal-cardiovascular health, score units	3 (1)
Framingham stroke risk profile, %, median [Q1, Q3]	2 [1, 5]

^a unless specified otherwise. CES-D - Center for Epidemiologic Studies Depression Scale

Table 2
Association of vascular risk factor burden and new-onset depression

Predictor	Risk for new-onset depression ^a	Risk for new-onset depressive symptoms ^b
	OR (95% CI)	OR (95% CI)
Basic model		
Ideal-Cardiovascular health	0.90 (0.81, 0.99)	0.84 (0.71, 0.99)
Framingham stroke risk profile	1.04 (1.00, 1.07)	1.06 (1.00, 1.12)
Basic model plus adjustment for prevalent stroke		
Ideal-Cardiovascular health	0.90 (0.81, 0.99)	0.84 (0.71, 0.99)
Framingham stroke risk profile	1.04 (1.00, 1.07)	1.06 (1.01, 1.12)
Basic model plus adjustment for prevalent and interim stroke		
Ideal-Cardiovascular health	0.90 (0.81, 0.99)	0.84 (0.71, 0.99)
Framingham stroke risk profile	1.03 (1.00, 1.07)	1.06 (1.00, 1.11)

All models are adjusted for age, sex, education and the time interval between baseline and follow-up.

^aDefined as CES-D scores ≥ 16 or antidepressant use. After adjustment for covariates, the sample size was 1982 with 264 events.

^bDefined as CES-D scores ≥ 16 with antidepressant users excluded from analysis. After adjustment for covariates, the sample size was 1192 with 92 events.