A Role for the Inflammasome in Spontaneous Labor at Term with Acute Histologic Chorioamnionitis

Reproductive Sciences
2017, Vol. 24(6) 934-953
© The Author(s) 2016
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/1933719116675058
journals.sagepub.com/home/rsx

Nardhy Gomez-Lopez, MSc, PhD^{1,2,3}, Roberto Romero, MD, DMedSci^{1,4,5,6},
Yi Xu, PhD^{1,2}, Olesya Plazyo, PhD^{1,2}, Ronald Unkel, BSc^{1,2},
Nandor Gabor Than, MD, PhD^{1,7,8,9}, Piya Chaemsathong, MD^{1,2},
Tinnakorn Chaiworapongsa, MD^{1,2}, Zhong Dong, PhD^{1,2},
Adi L. Tarca, PhD^{1,2}, Vikki M. Abrahams, PhD¹⁰, Lami Yeo, MD^{1,2},
and Sonia S. Hassan, MD^{1,2}

Abstract

Inflammasomes are cytosolic signaling platforms that regulate the activation of caspase (CASP)-1, which induces the maturation of interleukin (IL)-1 β and IL-18. Herein, we determined whether the chorioamniotic membranes from women in spontaneous labor at term with acute histologic chorioamnionitis express major inflammasome components and whether these changes are associated with the activation of CASP-1 and CASP-4 and the release of mature IL-1 β and IL-18. When comparing the chorioamniotic membranes from women in spontaneous labor at term with acute histologic chorioamnionitis to those without this placental lesion, we found that (1) the messenger RNA (mRNA) abundance of NLR family pyrin domain containing 3 (NLRP3), NLR family CARD domain containing 4 (NLRC4), absent in melanoma 2 (AIM2), and nucleotide binding oligomerization domain 2 (NOD2) was higher; (2) the NLRP3 and NLRC4 protein quantities were increased; (3) the mRNA and protein expressions of CASP-1 and its active forms were greater; (4) CASP-4 was increased at the mRNA level only; (5) the mRNA and protein expressions of IL-1 β and its mature form were higher; and (6) a modest increase in the total protein concentration and abundance of the mature form of IL-18 was observed. *In vitro* incubation of the chorioamniotic membranes with the CASP-1 inhibitor, VX765, decreased the release of endotoxin-induced IL-1 β and IL-18 (2-fold) but not IL-6 or tumor necrosis factor α . In conclusion, spontaneous labor at term with acute histologic chorioamnionitis is characterized by an upregulation of inflammasome components which, in turn, may participate in the activation of CASP-1 and lead to the release of mature IL-1 β by the chorioamniotic membranes. These results support a role for the inflammasome in the mechanisms responsible for spontaneous labor at term with acute histologic chorioamnionitis.

¹ Perinatology Research Branch, Program for Perinatal Research and Obstetrics, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, NICHD/NIH/DHHS, Bethesda, MD, and Detroit, MI, USA

² Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI, USA

³ Department of Immunology and Microbiology, Wayne State University School of Medicine, Detroit, MI, USA

⁴ Department of Obstetrics and Gynecology, University of Michigan, Ann Arbor, MI, USA

⁵ Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI, USA

⁶ Center for Molecular Medicine and Genetics, Wayne State University, Detroit, MI, USA

⁷ Systems Biology of Reproduction Lendulet Research Group, Institute of Enzymology, Research Centre for Natural Sciences, Hungarian Academy of Sciences, Budapest, Hungary

⁸ Maternity Private Department, Kutvolgyi Clinical Block, Semmelweis University, Budapest, Hungary

⁹ First Department of Pathology and Experimental Cancer Research, Semmelweis University, Budapest, Hungary

¹⁰ Department of Obstetrics, Gynecology and Reproductive Sciences, Yale University School of Medicine, New Haven, CT, USA

Corresponding Authors:

Roberto Romero, Perinatology Research Branch, NICHD/NIH/DHHS, Hutzel Women's Hospital 3990 John R, Box 4, Detroit, MI 48201, USA.

Email: prbchiefstaff@med.wayne.edu

Nardhy Gomez-Lopez, Perinatology Research Branch, NICHD/NIH/DHHS, Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI 48201, USA.

Email: ngomezlo@med.wayne.edu

Keywords

caspase-1, caspase-4, cytokine, endotoxin, fetal inflammatory response, funisitis, fever, intra-amniotic inflammation, intra-amniotic infection, lipopolysaccharide, microbial invasion of the amniotic cavity (MIAC), neutrophil, parturition, pregnancy

Introduction

Acute histologic chorioamnionitis is the most common placental lesion and represents the presence of intra-amniotic infection.^{1,2} Yet, acute histologic chorioamnionitis can also occur as a result of sterile intra-amniotic inflammation, which is induced by danger signals released under cellular stress, injury, or death.³⁻⁶ These signals are termed damage-associated molecular pattern molecules⁷ or alarmins.⁸ Acute histologic chorioamnionitis is strongly associated with preterm labor/birth⁹⁻¹²; however, it is frequently observed in the placentas of women who delivered after spontaneous labor at term.^{13,14} The presence of this placental lesion at term is strongly associated with intrapartum fever and elevated levels of maternal white blood cell counts, interleukin (IL)-8, and IL-6 at admission for delivery.¹⁵ Furthermore, acute histologic chorioamnionitis at term is associated with elevated concentrations of cytokines such as IL-1 α , IL-1 β , tumor necrosis factor (TNF)- α , IL-6, and IL-8 in the amniotic fluid,¹⁶ umbilical cord blood,^{15,17-20} and decidual tissues.²¹⁻²⁶ Elevated concentrations of cytokines in these compartments are linked to adverse neonatal outcomes.²⁷⁻⁴⁶ Therefore, our investigation focused on the mechanisms implicated in spontaneous labor at term with acute histologic chorioamnionitis.

Acute inflammation of the chorioamniotic membranes is characterized by the infiltration of neutrophils.^{1,47} The mechanisms implicated in the pathogenesis of this placental lesion involve chemotactic signals including those derived from IL-8, C-X-C chemokine ligand (CXCL) 6 and growth regulated oncogene (GRO) α ,¹ all of which are found in the amniotic fluid.⁴⁸⁻⁶¹ Acute histologic chorioamnionitis is also associated with elevated concentrations of cytokines, such as IL-1 β in the amniotic fluid²⁹ and chorioamniotic membranes (including the decidua).²¹ The IL-1 β cytokine is synthesized as a zymogen (or pro-form), which requires cleavage in order to be biologically active and released extracellularly.⁶²⁻⁶⁵ The cleavage of IL-1 β can be performed by intracellular protein caspase (CASP)-1⁶²⁻⁶⁸ which, in turn, could be activated by cytoplasmic complexes termed inflammasomes.⁶⁹⁻⁹⁷ Active CASP-1 can also cleave pro-IL-18 into its mature form.^{66-68,98,99} Inflammasome complexes contain (1) a pattern recognition receptor (a sensor molecule), (2) the adaptor protein ASC (an apoptosis-associated speck-like protein), and (3) pro-CASP-1.¹⁰⁰⁻¹⁰⁹ In addition to inflammasomes, active CASP-4 is also required, but not necessary, for the activation of CASP-1.^{110,111}

Inflammasomes are implicated in physiological¹¹²⁻¹¹⁴ and pathological^{113,115-124} inflammations during pregnancy. Recently, we provided evidence that supports a role for the inflammasome in the physiological inflammatory response that accompanies human spontaneous labor at term.¹¹⁴ Since amniotic fluid concentrations of CASP-1,¹¹³ IL-1 β ,¹²⁵ and IL-18¹²⁶ are greater

in women with intra-amniotic infection/inflammation (i.e. pathological inflammation) than in those without intra-amniotic infection/inflammation, we proposed that the inflammasome is also implicated in the pathological inflammatory process in acute histologic chorioamnionitis during spontaneous labor at term.

In this study, we aimed to determine (1) whether inflammasome components are expressed in the chorioamniotic membranes from women who have undergone spontaneous labor at term with histological signs of acute chorioamnionitis and (2) whether these changes are linked to the activation of CASP-1 and CASP-4 and to the release of the mature forms of IL-1 β and IL-18. In addition, we investigated whether *in vitro* incubation of term nonlabor chorioamniotic membrane explants with the CASP-1 inhibitor, VX765, decreases the release of endotoxin-induced IL-1 β and IL-18.

Materials and Methods

Human Participants, Clinical Specimens, and Definitions

This case-control study included patients who delivered at term after labor with (TIL-ACA) and without (TIL) acute histologic chorioamnionitis. Samples from the chorioamniotic membranes were collected from the Bank of Biological Specimens of the Detroit Medical Center, Wayne State University, and the Perinatology Research Branch, NICHD/NIH/DHHS (Detroit, Michigan). The review boards of these institutions approved the collection and the use of biological materials for research purposes. Participants provided written informed consent, and samples were collected within 0.5 hours after delivery. Table 1 includes the demographic and clinical characteristics of the study groups. Multiparous women or women with neonates having congenital or chromosomal abnormalities were excluded. Labor was defined by the presence of regular uterine contractions at a frequency of at least 2 contractions every 10 minutes with cervical changes resulting in delivery.¹²⁷

For the *in vitro* experiments, chorioamniotic membrane samples were collected within 30 minutes after delivery from healthy pregnant women at term without labor. Women included in this study had no medical complications and delivered neonates of appropriate weight for gestational age.¹²⁸

Placental Histopathological Examinations

Five-micrometer-thick sections of formalin-fixed, paraffin-embedded tissue specimens were cut and mounted on Superfrost Plus microscope slides (Erie Scientific LLC, Portsmouth, New Hampshire). In each case, several tissue sections of the chorioamniotic membranes, umbilical cord, and placental disc were examined. After deparaffinization, slides were rehydrated, stained with hematoxylin–eosin, and evaluated by

Table 1. Demographic and Clinical Characteristics of the Study Population.

	Spontaneous Term Labor (TIL, n = 28)	Spontaneous Term Labor with Acute Chorioamnionitis (TIL-ACA, n = 32)	P Value
Maternal age, years ^a	21.0 (18.5-26.0)	21.0 (19.0-26.5)	NS
Race ^b			
African American	24 (88.9%)	29 (90.6%)	NS
Caucasian	2 (7.4%)	1 (3.1%)	
Hispanic	0 (0.0%)	0 (0.0%)	
Other	1 (3.7%)	2 (6.3%)	
Maternal weight ^a , kg	66 (59-75)	68 (60-77)	NS
Body mass index ^a , kg/m ²	24.5 (22.1-29.1)	26.4 (22.5-28.3)	NS
Primiparity ^b	11 (39.3%)	14 (43.8%)	NS
Gestational age at delivery, weeks ^a	39.7 (38.7-40.5)	40.1 (38.6-40.6)	NS
Birth weight, g ^a	3252 (3125-3420)	3487 (3085-3765)	NS
Cesarean section ^b	35.7%	50%	NS
Acute chorioamnionitis ^b	0%	100%	<.0001

Abbreviation: NS, not significant.

^aMann-Whitney *U* test.^bFisher's exact test.

pathologists who had been blinded to the clinical outcome, according to the published criteria.^{1,129} The diagnosis of acute histologic chorioamnionitis was made when the infiltration of neutrophils into the chorionic trophoblast layer or chorioamniotic connective tissue was observed.^{1,129}

RNA Isolation, Complementary DNA Synthesis, and Quantitative Reverse Transcription Polymerase Chain Reaction Analysis

Total RNA was extracted from snap-frozen chorioamniotic membrane samples (TIL, n = 28 and TIL-ACA, n = 32) and from term nonlabor tissue explants (n = 6) using TRIzol (Invitrogen, Life Technologies Corporation, Grand Island, New York) and Qiagen RNeasy kits (Qiagen, Gaithersburg, Maryland). The purity, concentration, and integrity of the RNA were assessed by using the NanoDrop 1000 spectrophotometer (Thermo Scientific, Wilmington, Delaware), and the Bioanalyzer 2100 (Agilent Technologies, Wilmington, Delaware). Complementary DNA was synthesized using the SuperScript III First-Strand Synthesis System (Invitrogen) and oligo(dT)20 primers (Invitrogen). Gene expression profiling was performed on the Biomark System for high-throughput quantitative reverse transcription polymerase chain reaction (qRT-PCR; Fluidigm, San Francisco, California) and the ABI 7500 Fast Real-Time PCR System (Applied Biosystems, Life Technologies Corporation, Foster City, California) with TaqMan gene expression assays (Applied Biosystems) listed in Supplementary Table I.

Chorioamniotic Membrane Tissue Lysates

Tissue lysates were prepared by homogenizing snap-frozen chorioamniotic membranes (TIL and TIL-ACA; n = 10 each) in 2 mL of 1× phosphate-buffered saline (PBS; Invitrogen) containing a complete protease inhibitor cocktail (Cat. No.

11697498001; Roche Applied Science, Mannheim, Germany). Next, lysates were centrifuged at 15700g for 5 minutes at 4°C, and the supernatant was collected and stored at −80°C. Prior to enzyme-linked immunosorbent assay (ELISA) or immunoblotting, total protein determination was determined using the Quick Start Bradford Protein Assay Kit (Bio-Rad, Hercules, California).

Enzyme-Linked Immunosorbent Assays

Protein concentrations of the NLR family pyrin domain containing (NLRP)1, NLRP3, absent in melanoma 2 (AIM2), nucleotide-binding oligomerization domain (NOD)2, CASP-1, CASP-4, IL-18, pro-IL-1β, and mature IL-1β were determined in chorioamniotic membrane tissue lysates by immunoassays (NLRP1, NLRP3, and NOD2 ELISA kits from Cusabio, Wuhan, Hubei, China; AIM2, CASP-1, and CASP-4 ELISA kits from Cloud Clone, Houston, Texas; pro-IL-1β and IL-1β ELISA kits from R&D Systems, Minneapolis, Minnesota; and IL-18 ELISA kits from MBL International Corporation, Woburn, Massachusetts), as previously described.¹¹⁴ The sensitivities of the assays were <4.68 pg/mL for NLRP1, <0.039 ng/mL for NLRP3, <0.056 ng/mL for AIM2, <6.25 pg/mL for NOD2, <0.112 ng/mL for CASP-1, <0.053 ng/mL for CASP-4, 3.3 pg/mL for pro-IL-1β, <1 pg/mL for mature IL-1β, and <12.5 pg/mL for IL-18. The IL-1β ELISA kit measured about 10% of pro-IL-1β. The immunoassays for NLRP4 and NOD1 did not meet our criteria for validation; instead, immunoblotting was performed.

Concentrations of human IL-1β, IL-18, IL-6, and TNF-α in cell culture supernatants from *in vitro* experiments were measured with sensitive and specific immunoassays (Meso Scale Discovery, Gaithersburg, Maryland), according to the manufacturer's instructions. Briefly, 150 μL of 1% (w/v) Blocker B Solution were dispensed into each well of the precoated plates and incubated for 1 hour with vigorous shaking at room

temperature. Plates were then washed 3 times with $1 \times$ PBS and 0.05% Tween-20 (Sigma-Aldrich, St Louis, Missouri). Twenty-five microliters of each sample or calibrator were dispensed into separate wells of the plates and incubated for 2 hours with vigorous shaking at room temperature. The samples and calibrators were discarded, and the plates were washed 3 times with $1 \times$ PBS and 0.05% Tween-20 (Sigma-Aldrich), followed by an addition of 25 μ L of the $1 \times$ detection antibody solution into each well. Plates were incubated for 2 hours with vigorous shaking at room temperature. The detection antibody was removed and the plates were washed 3 times with $1 \times$ PBS and 0.05% Tween-20 (Sigma-Aldrich). One hundred fifty microliters of $2 \times$ Read Buffer T (Meso Scale Discovery) were added to each well, and the signals were read by the Sector 2400 Imager (Meso Scale Discovery). Standard curves were generated, and the assay values of the samples were extrapolated from the curves. The sensitivities of the assays were 0.268 pg/mL for IL-1 β , 0.216 pg/mL for IL-18, 0.206 pg/mL for IL-6, and 0.417 pg/mL for TNF- α .

Immunohistochemistry

Tissue sections (5 μ m thick) were prepared from the chorioamniotic membranes (TIL and TIL-ACA, $n = 10$ each). Immunohistochemistry for NLRP1, NLRP3, NLRC4, AIM2, NOD1, NOD2, CASP-1, CASP-4, IL-1 β , and IL-18 was performed, as previously described.¹¹⁴ Supplementary Table II includes the utilized primary antibodies and immunostaining conditions. Quantification of the intensity was performed using a Perkin-Elmer Panoramic MIDI slide scanner (PerkinElmer, Waltham, Massachusetts).

Immunoblotting

The protein quantity of NLRC4 and NOD1 and the active/mature forms of CASP-1, CASP-4, and IL-18 were determined by immunoblotting. Tissue lysates (50 μ g for NLRC4, NOD1, and IL-18; 20 μ g for CASP-1; and 40 μ g for CASP-4 per well) were subjected to 4% to 12% sodium dodecyl sulfate polyacrylamide gel electrophoresis (Invitrogen). Separated proteins were then transferred onto nitrocellulose membranes (Bio-Rad). Next, membranes were submerged in blocking solution (5% nonfat dry milk in tris-buffered saline containing 0.1% Tween-20 [Bio-Rad] or StartingBlock T20 Block Buffer [ThermoFisher Scientific, Inc, Rockford, Illinois]) and probed overnight at 4°C with the following human antibodies—mouse anti-NLRC4 antibody (BioLegend, San Diego, California), rabbit anti-NOD1 polyclonal antibody (Enzo Life Sciences, Farmingdale, New York),¹³⁰ mouse anti-CASP-1 monoclonal antibody (R&D Systems), rabbit anti-CASP-4 polyclonal antibody (Abcam, Cambridge, Massachusetts), or rabbit anti-IL-18 polyclonal antibody (Santa Cruz Biotechnology, Dallas, Texas). A horseradish peroxidase-conjugated anti-mouse or anti-rabbit IgG (Cell Signaling, Boston, Massachusetts) was used as a secondary antibody. Chemiluminescence signals were detected with ChemiGlow

West Reagents (Protein Simple, Santa Clara, California), and images were acquired using the Fujifilm LAS-4000 Imaging System (FUJIFILM North America Corporation, Valhalla, New York). Nitrocellulose membranes were then stripped with Restore Plus Western Blot Stripping Buffer (Pierce Biotechnology, ThermoFisher Scientific Inc) for 15 minutes, washed with $1 \times$ PBS, blocked, and reprobed for 1 hour at room temperature with a mouse anti-beta actin (ACTB) monoclonal antibody (Sigma-Aldrich) or a mouse anti-glyceraldehyde 3-phosphate dehydrogenase monoclonal antibody (Santa Cruz Biotechnology). Chemiluminescence signals were again detected with ChemiGlow West Reagents, and images were acquired using the Fujifilm LAS-4000 Imaging System.

In Vitro Experiments in Chorioamniotic Membrane Explants

Chorioamniotic membrane samples were collected from women undergoing elective cesarean delivery ($n = 6$) and processed within 30 minutes. Tissue samples were washed with $1 \times$ PBS and cut into 2 cm $\times$ 2 cm pieces. These tissue explants were transferred into 6-well tissue culture plates containing 2 mL of Dulbecco's Modified Eagle Medium (Invitrogen) supplemented with 10% fetal bovine serum (Invitrogen) and 1% penicillin/streptomycin (Invitrogen). Tissue explants were incubated in a humidified 5% CO₂ incubator at 37°C with or without the CASP-1 inhibitor VX765 (10 μ M in 0.1% DMSO; MedKoo Biosciences, Chapel Hill, North Carolina) for 30 minutes before an endotoxin or lipopolysaccharide (10 ng/mL; LPS, *Escherichia coli* 0111: B4, Sigma-Aldrich) was added into the culture medium. Incubations were continued for 16 hours; then tissue culture supernatants were collected for ELISA, and tissue samples were homogenized in TRIzol reagent (Invitrogen) for qRT-PCR. Triplicate supernatants were obtained from the membranes of each patient.

Statistical Analyses

The SPSS v.19.0 software (SPSS Inc, Chicago, Illinois) was used to analyze demographic and clinical data. Comparisons between the 2 groups (TIL vs TIL-ACA) were performed using Fisher's exact test for proportions as well as the Mann-Whitney *U* test for nonnormally distributed continuous variables. Gene expressions relative to *ACTB/GAPDH/RPLP0* (snap-frozen samples) or *RPLP0* (*in vitro* experiments) were calculated as $-\Delta$ Ct values, where Δ Ct (Δ Ct = Ct_{target} - Ct_{reference}) was computed for each sample after averaging the Ct values over the technical replicates. Group means of gene expression were then compared using *t* tests from an analysis of the variance linear model and the resulting *P* values were adjusted using the Benjamini-Hochberg procedure. An adjusted *P* value of $\leq .05$ was considered statistically significant.

Figure 1. Inflammasome components and NOD2 protein in the chorioamniotic membranes. **A**, Messenger RNA abundance of inflammasome components and *NOD2* in the chorioamniotic membranes from women in spontaneous labor at term with (TIL-ACA, $n = 32$) or without (TIL, $n = 28$) acute histologic chorioamnionitis. Relative gene expressions are presented as $-\Delta Ct$ values. **B**, Protein concentrations of inflammasome components and *NOD2* in chorioamniotic membrane tissue lysates ($n = 10$ each). **C**, Representative immunostainings for inflammasome components and *NOD2* in the chorioamniotic membranes ($n = 10$ each), $200\times$ magnification. NOD indicates nucleotide-binding oligomerization domain.

Results

Messenger RNA Abundance and Protein Expression of *NLRP3* and *NLRC4* Increase in the Chorioamniotic Membranes in Spontaneous Labor at Term with Acute Histologic Chorioamnionitis

The messenger RNA (mRNA) abundance of *NLRP3*, *NLRC4*, *AIM2*, and *NOD2* was greater in the chorioamniotic membranes from women who underwent spontaneous labor at term

with acute histologic chorioamnionitis than in those without this placental lesion (Figures 1A and 2A). However, no differences were observed in the mRNA abundance of *NLRP1* and *NOD1* between these groups (Figures 1A and 2D). The *NLRP3* and *NLRC4* protein concentrations were increased in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis compared to those without this placental lesion (Figures 1B and 2B). No differences were observed in the protein

Figure 2. NLRC4 and NOD1 in the chorioamnion membranes. A and D, Messenger RNA abundance of *NLRC4* and *NOD1* in the chorioamnion membranes from women in spontaneous labor at term with (TIL-ACA, n = 32) or without (TIL, n = 28) acute histologic chorioamnionitis. Relative gene expressions are presented as $-\Delta\text{Ct}$ values. B and E, Protein quantity of NLRC4 and NOD1 in chorioamnion membrane tissue lysates (n = 5 each). C and F, Intensity of the immunostainings for NLRC4 and NOD1 in the chorioamnion membranes (n = 10 each) and representative immunostainings, 200× magnifications. NOD indicates nucleotide-binding oligomerization domain.

concentrations of NLRP1, AIM2, NOD2, and NOD1 between these 2 groups of women (Figures 1B and 2E).

Immunohistochemistry revealed that all of the inflammasome components and NOD proteins were expressed by the chorioamniotic membranes from women who underwent spontaneous labor at term with and without acute histologic chorioamnionitis. These proteins were detected in the chorionic trophoblast cells, decidual stromal cells, and amniotic mesodermal and epithelial cells (Figures 1C, 2C, and 2F). However, NOD1 immunoreactivity seemed to be lower in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figure 2F).

The collective increase in the mRNA abundance and protein expression of NLRP3 and NLRC4 in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis provides evidence that the inflammasome may participate in the pathological activation of the innate immune system in the setting of intra-amniotic infection/inflammation.

Activation of CASP-1 in the Chorioamniotic Membranes in Spontaneous Labor at Term with Acute Histologic Chorioamnionitis

We then investigated whether the increased protein concentrations of NLRP3 and NLRC4 in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis were linked to the activation of CASP-1 and CASP-4. The mRNA abundance of *CASP1* and *CASP4* in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis was greater than in those without this placental lesion (Figure 3A). The protein concentrations of CASP-1 and CASP-4 were not significantly different between the 2 groups (Figure 3B). Immunohistochemistry analysis revealed that CASP-1, but not CASP-4, was greater in the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figures 3C and 3D). Also, the active forms of CASP-1 (p10 and p20) were increased in the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis compared to those without this placental lesion (Figures 3E and 3G). In contrast, the pro-form of CASP-1 was reduced in the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis (Figure 3G). The active form of CASP-4 was undetectable by immunoblotting (Figure 3E). However, the pro-form of CASP-4 was detected but no differences were observed between the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis and those without this placental lesion (Figure 3F). Altogether, these data suggest that pathological inflammation in spontaneous labor at

term with acute histologic chorioamnionitis involves the participation of NLRP3 and NLRC4 inflammasomes which, in turn, may participate in the activation of CASP-1 in the chorioamniotic membranes.

Increased mRNA Abundance and Protein Expression of IL-1 β in the Chorioamniotic Membranes in Spontaneous Labor at Term with Acute Chorioamnionitis

Active forms of CASP-1 convert inactive pro-IL-1 β into its mature form.⁶²⁻⁶⁸ We then determined whether the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis release the mature form of IL-1 β . The mRNA abundance of *IL1 β* was greater in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figure 4A). Also, the protein concentration of mature IL-1 β was greater in the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figure 4B). In contrast, the concentration of pro-IL-1 β was reduced in cases with acute histologic chorioamnionitis (Figure 4B). Immunohistochemistry analysis revealed that the immunoreactivity of IL-1 β was stronger in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion (Figure 4C). These findings suggest that pathological inflammation in spontaneous labor at term with acute histologic chorioamnionitis involves the release of mature IL-1 β , which is likely processed by the active form of CASP-1 in the chorioamniotic membranes.

A Modest Increase in the Total Protein Concentration and Abundance of the Mature Form of IL-18 in the Chorioamniotic Membranes in Spontaneous Labor at Term with Acute Histologic Chorioamnionitis

Active forms of CASP-1 also convert inactive pro-IL-18 into its mature form.⁹⁸ Therefore, we investigated whether the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis release the mature form of IL-18. No differences were observed in the mRNA abundance (Figure 5A) and immunoreactivity (Figure 5C) of IL-18 between the chorioamniotic membranes from women who had undergone spontaneous labor at term with acute histologic chorioamnionitis and those without this placental lesion. A modest but not significant increase in the total protein concentration (Figure 5B) and the abundance of the mature form (Figure 5D) of IL-18 was observed in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis compared to those without this placental lesion. These data suggest that IL-18 may be involved in the pathological inflammation of the chorioamniotic membranes

Figure 3. Inflammatory caspases in the chorioamniotic membranes. A, Messenger RNA abundance of *CASP1* and *CASP4* in the chorioamniotic membranes from women in spontaneous labor at term with (TIL-ACA, $n = 32$) or without (TIL, $n = 28$) acute histologic chorioamnionitis. Relative gene expressions are presented as $-\Delta C_t$ values. B, Protein concentrations of *CASP-1* and *CASP-4* in chorioamniotic membrane tissue lysates ($n = 10$ each). C and D, Intensity of the immunostainings for *CASP-1* and *CASP-4* in the chorioamniotic membranes ($n = 10$ each) and representative immunostainings, $200\times$ magnifications. E and F, Immunoblotting of *CASP-1*, *CASP-4*, its active forms, and *GAPDH* in the chorioamniotic membranes and their quantifications ($n = 4$ each). *CASP* indicates caspase.

Figure 4. Interleukin (IL) 1 β in the chorioamniotic membranes. A, Messenger RNA abundance of *IL1B* in the chorioamniotic membranes from women in spontaneous labor at term with (TIL-ACA, $n = 32$) or without (TIL, $n = 28$) acute histologic chorioamnionitis. Relative gene expressions are presented as $-\Delta Ct$ values. B, Protein concentrations of the pro- and mature form of IL-1 β in chorioamniotic membrane lysates ($n = 10$ each). C, Intensity of the immunostainings for IL-1 β in the chorioamniotic membranes ($n = 10$ each) and representative immunostainings, 200 $\times$ magnifications.

during spontaneous labor at term with acute histologic chorioamnionitis.

In Vitro Incubation of the Chorioamniotic Membranes with a CASP-1 Inhibitor, VX765, Reduces the LPS-Induced Release of IL-1 β and IL-18

Finally, we examined whether the CASP-1 inhibitor, VX765, would reduce the LPS-induced mRNA abundance of pro-

inflammatory cytokines in the chorioamniotic membranes. Incubation with LPS increased the mRNA abundance of *NLRP3*, *AIM2*, *NOD2*, *CASP1* (2-fold; $P = 0.058$), *CASP4*, *IL1 β* , *IL6*, and *TNF* in the chorioamniotic membranes (Figure 6A and Supplementary Table III). *In vitro* incubation with LPS also increased the release of IL-1 β , IL-18, IL-6, and TNF- α by the chorioamniotic membranes (Figure 6B). However, incubation with the CASP-1 inhibitor, VX765, reduced the LPS-induced release of inflammasome-dependent IL-1 β and IL-18 (2-fold, $P = .058$) but did not diminish the release of

Figure 5. Interleukin (IL)-8 in the chorioamniotic membranes. A, Messenger RNA abundance of IL-18 in the chorioamniotic membranes from women in spontaneous labor at term with (TIL-ACA, n = 32) or without (TIL, n = 28) acute histologic chorioamnionitis. Relative gene expressions are presented as $-\Delta\text{Ct}$ values. B, Protein concentrations of IL-18 in chorioamniotic membrane tissue lysates (n = 10 each). C, Intensity of the immunostainings for IL-18 in the chorioamniotic membranes (n = 10 each) and representative immunostainings, 200 $\times$ magnifications. D, Immunoblotting of IL-18, its mature form, and ACTB in the chorioamniotic membranes and their quantifications (n = 6-7 each).

Figure 6. *In vitro* incubation of the chorioamniotic membranes with the CASP-1 inhibitor, VX765, reduces the LPS-induced release of IL-1β and IL-18. A, Messenger RNA abundance of IL-1β, IL-6, and TNF in chorioamniotic membrane explants from women who delivered at term without labor (n = 6) that were incubated with LPS, the CASP-1 inhibitor VX765, or both. Relative gene expressions are presented as -ΔCt values. B, Protein concentrations of IL-1β, IL-18, IL-6, and TNF-α in chorioamniotic membrane tissue lysates (n = 6). Experiments were performed in triplicate. CASP-1 indicates caspase 1; IL, interleukin; LPS, lipopolysaccharide; TNF, tumor necrosis factor.

inflammasome-independent IL-6 and TNF- α (Figure 6B). These data demonstrate that the release of LPS-induced IL-1 β , and most likely IL-18, is mediated by CASP-1 in the chorioamniotic membranes.

Discussion

Principal findings of the study: (1) mRNA abundance of *NLRP3*, *NLRC4*, *AIM2*, and *NOD2* in the chorioamniotic membranes was higher in women in spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion, (2) NLRP3 and NLRC4 protein concentrations were higher in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion, (3) mRNA abundance of *CASP1* and *CASP4* was greater in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion, (4) the immunoreactivity of CASP-1 and the abundance of its active forms (p10 and p20) were increased among women who had undergone spontaneous labor at term with acute histologic chorioamnionitis compared to those without this placental lesion, (5) the mRNA abundance, protein concentration, immunoreactivity, and mature form of IL-1 β were greater in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis than in those without this placental lesion, (6) a modest increase in the total protein concentration and abundance of the mature form of IL-18 was observed in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis compared to those without this placental lesion, and (7) *in vitro* incubation of the chorioamniotic membranes with the CASP-1 inhibitor, VX765, reduced the LPS-induced release of inflammasome-dependent IL-1 β and IL-18 (2-fold) but did not diminish the release of inflammasome-independent IL-6 and TNF- α . Taken together, these findings support a role for the inflammasome in the pathological inflammation of the chorioamniotic membranes during spontaneous labor at term with acute histologic chorioamnionitis.

Previously, we proposed a role for the inflammasome in the physiological pro-inflammatory process of spontaneous labor at term.¹¹²⁻¹¹⁴ Herein, we demonstrated that the mRNA and protein expressions of NLRP3 are increased in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis. The NLRP3 inflammasome complex includes the NLRP3 protein, the adaptor molecule ASC, and pro-CASP-1.^{70,106,131,132} The activation of the NLRP3 inflammasome can be induced by microorganisms and danger signals^{75,133-161} and requires 2 steps—priming and the assembly of the inflammasome complex.^{162,163} Although the first step includes the activation of the nuclear factor kappa B (NF- κ B) pathway, which induces the upregulation of the NLR protein and IL-1 β ,

the second step permits the assembly of the inflammasome complex.¹⁶²⁻¹⁶⁴ The data presented herein provide evidence that there is priming of the NLRP3 inflammasome in the chorioamniotic membranes during spontaneous labor at term with acute histologic chorioamnionitis. Further studies are required to demonstrate that there is assembly of the NLRP3 inflammasome complex in the chorioamniotic membranes.

We also found that the NLRC4 inflammasome may be involved in the pathological pro-inflammatory process of spontaneous labor at term with acute histologic chorioamnionitis. NLRC4 is mostly expressed in myeloid cells and can also regulate CASP-1 activation and IL-1 β processing.¹⁶⁵ NLRC4 contains an N-terminal caspase activation and recruitment domain (CARD), a central NACHT (NACHT is an acronym for neuronal apoptosis inhibitor protein), C2TA (major histocompatibility complex (MHC) class 2 transcription activator), HET-E (incompatibility locus protein from *Podospora anserina*) and TP1 (telomerase-associated protein 1) domain, and C-terminal leucine-rich repeat (LRR).¹⁶⁵ The activation of the NLRC4 inflammasome is mediated by flagellins^{166,167} and requires the deletion of the LRR domain.¹⁶⁸ Comparable to NLRP3, NLRC4 activation also requires the chaperone protein, heat-shock protein 90 and the ubiquitin ligase-associated protein SGT1.¹⁶⁹ However, there are striking differences between NLRC4 and other NLR proteins. For example, activation of CASP-1 by NLRC4 primarily leads to cell death (i.e. necrosis).¹⁶⁵ However, the activation of CASP-1 by NLRP3 largely results in IL-1 β and IL-18 processing without leading to cell death (i.e. pyroptosis).¹⁶⁵ Also, activation of CASP-1 by NLRC4 does not involve potassium.¹⁷⁰ Therefore, NLRP3 and NLRC4 inflammasomes activate CASP-1 with certain specificity.¹⁶⁵ Altogether, these data suggest that the pathological pro-inflammatory process in spontaneous labor at term with acute histologic chorioamnionitis includes the participation of both NLRP3 and NLRC4, which most likely results in pyroptosis and necrosis in the chorioamniotic membranes.

Inflammasome assembly/activation leads to the processing of pro-CASP-1 into its active form.^{106,171} Herein, we found that the active forms of CASP-1 are increased in the chorioamniotic membranes from women in spontaneous labor at term with acute histologic chorioamnionitis. This finding is in line with a previous study demonstrating that amniotic fluid CASP-1 concentrations are greater in women with intra-amniotic infection/inflammation who delivered preterm than in cases without this clinical condition.¹¹³ This finding suggests that during spontaneous labor at term with acute histologic chorioamnionitis, the chorioamniotic membranes release abundant active forms of CASP-1, which is likely mediated by the inflammasome.

Activation of the noncanonical NLRP3 inflammasome pathway is induced by gram-negative bacteria and includes the participation of CASP-4 (murine homologue CASP-11) in the activation of CASP-1.^{158,172,173} Although the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis expressed high

levels of CASP-4 at the RNA level, its protein and mature form were not different between tissues with and without acute inflammation. These data do not support a role of CASP-4 in the pathological pro-inflammatory process of spontaneous labor at term with acute histologic chorioamnionitis.

The active forms of CASP-1 can convert inactive pro-IL-1 β ⁶²⁻⁶⁸ and pro-IL-18^{66-68,98,99} into their mature forms. Herein, we found that the chorioamniotic membranes from women in spontaneous labor at term with acute histologic chorioamnionitis release high amounts of mature IL-1 β . To determine whether the release of mature IL-1 β from the chorioamniotic membranes was driven by the activation of CASP-1, we used VX765 in an *in vitro* model of inflammation induced by LPS. This model is widely used *in vitro*¹⁷⁴⁻¹⁷⁷ and *in vivo*.¹⁷⁸⁻¹⁸⁴ VX765 is a potent, selective, and competitive inhibitor of CASP-1, which inhibits the CASP-1-dependent release of IL-1 β and IL-18 from peripheral blood mononuclear cells¹⁸⁵ and whole blood.¹⁸⁶ In this study, VX765 inhibited the LPS-induced release of inflammasome-dependent IL-1 β and IL-18 (2-fold) by the chorioamniotic membranes without affecting the release of inflammasome-independent IL-6 or TNF- α . The IL-1 β cytokine participates in several processes of parturition,¹⁸⁷⁻¹⁹³ and its causal role was demonstrated by the fact that injection of this cytokine in mice^{194,195} and monkeys¹⁹⁶⁻²⁰³ induces preterm labor and birth. Collectively, these data demonstrate that the chorioamniotic membranes release the mature form of IL-1 β , which is likely mediated by the inflammasome-derived active forms of CASP-1 during spontaneous labor at term with acute histologic chorioamnionitis.

We observed a modest increase in the total protein concentration and abundance of the mature form of IL-18 in the chorioamniotic membranes from women who underwent spontaneous labor at term with acute histologic chorioamnionitis compared to those without this placental lesion. The lack of statistical significance between these 2 study groups may be due to the sample size. Regardless, this finding is in line with a previous report demonstrating that amniotic fluid concentrations of IL-18 are higher in women who underwent spontaneous labor at term with microbial invasion of the amniotic cavity (MIAC) than in those who delivered at term without this clinical condition.¹²⁶ These data suggest that IL-18 may be involved in the pathological inflammation of the chorioamniotic membranes during spontaneous labor at term with acute histologic chorioamnionitis.

Conclusion

This study provides evidence supporting a role for the inflammasome in the activation of CASP-1 and the release of mature IL-1 β by the chorioamniotic membranes from women in spontaneous labor at term with acute histologic chorioamnionitis. The selective upregulation of NLRP3 and NLRC4 suggests that these inflammasomes are implicated in the mechanisms responsible for pathological inflammation (i.e. acute histologic chorioamnionitis) in term parturition.

Acknowledgments

The authors gratefully acknowledge Lorri McLuckie, Rona Wang, Yang Jiang, and Zhonghui Xu for their contributions to the execution of this study. The authors thank the physicians and nurses from the Center for Advanced Obstetrical Care and Research and the Intrapartum Unit for their help in collecting human samples. The authors also thank staff members of the PRB Clinical Laboratory and the PRB Histology/Pathology Unit for the processing and examination of the pathological sections. Finally, the authors thank Tara Mial for her critical readings of the manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported, in part, by the Perinatology Research Branch, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, US Department of Health and Human Services (NICHD/NIH/DHHS), and, in part, with federal funds from the NICHD/NIH/DHHS under Contract No. HHSN275201300006C. This research was also supported by the Wayne State University Perinatal Initiative in Maternal, Perinatal and Child Health.

Supplemental Material

The online supplementary tables are available at <http://journals.sagepub.com/doi/suppl/10.1177/1933719116675058>.

References

1. Kim CJ, Romero R, Chaemsathong P, Chaiyasit N, Yoon BH, Kim YM. Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. *Am J Obstet Gynecol*. 2015;213(4 suppl): S29-S52.
2. Romero R, Chaemsathong P, Docheva N, et al. Clinical chorioamnionitis at term VI: acute chorioamnionitis and funisitis according to the presence or absence of microorganisms and inflammation in the amniotic cavity. *J Perinat Med*. 2016; 44(1):33-51.
3. Romero R, Miranda J, Chaiworapongsa T, et al. A novel molecular microbiologic technique for the rapid diagnosis of microbial invasion of the amniotic cavity and intra-amniotic infection in preterm labor with intact membranes. *Am J Reprod Immunol*. 2014;71(4):330-358.
4. Romero R, Miranda J, Chaiworapongsa T, et al. Prevalence and clinical significance of sterile intra-amniotic inflammation in patients with preterm labor and intact membranes. *Am J Reprod Immunol*. 2014;72(5):458-474.
5. Romero R, Miranda J, Chaiworapongsa T, et al. Sterile intra-amniotic inflammation in asymptomatic patients with a sonographic short cervix: prevalence and clinical significance [published online September 24, 2014]. *J Matern Fetal Neonatal Med*:1-17.
6. Romero R, Miranda J, Kusanovic JP, et al. Clinical chorioamnionitis at term I: microbiology of the amniotic cavity using

- cultivation and molecular techniques. *J Perinat Med*. 2015; 43(1):19-36.
7. Lotze MT, Deisseroth A, Rubartelli A. Damage associated molecular pattern molecules. *Clin Immunol*. 2007;124(1):1-4.
 8. Oppenheim JJ, Yang D. Alarmins: chemotactic activators of immune responses. *Curr Opin Immunol*. 2005;17(4):359-365.
 9. Russell P. Inflammatory lesions of the human placenta: clinical significance of acute chorioamnionitis. *Am J Diagn Gynecol Obstet*. 1979;2:127-137.
 10. Guzik DS, Winn K. The association of chorioamnionitis with preterm delivery. *Obstet Gynecol*. 1985;65(1):11-16.
 11. van Hoven KH, Anyaegbunam A, Hochster H, et al. Clinical significance of increasing histologic severity of acute inflammation in the fetal membranes and umbilical cord. *Pediatr Pathol Lab Med*. 1996;16(5):731-744.
 12. Lee SM, Park JW, Kim BJ, et al. Acute histologic chorioamnionitis is a risk factor for adverse neonatal outcome in late preterm birth after preterm premature rupture of membranes. *PLoS One*. 2013;8(12): e79941.
 13. Seong HS, Lee SE, Kang JH, Romero R, Yoon BH. The frequency of microbial invasion of the amniotic cavity and histologic chorioamnionitis in women at term with intact membranes in the presence or absence of labor. *Am J Obstet Gynecol*. 2008; 199(4):375.e1-5.
 14. Park HS, Romero R, Lee SM, Park CW, Jun JK, Yoon BH. Histologic chorioamnionitis is more common after spontaneous labor than after induced labor at term. *Placenta*. 2010;31(9):792-795.
 15. Roberts DJ, Celi AC, Riley LE, et al. Acute histologic chorioamnionitis at term: nearly always noninfectious. *PLoS One*. 2012; 7(3): e31819.
 16. Halgunset J, Johnsen H, Kjollesdal AM, Qvigstad E, Espevik T, Austgulen R. Cytokine levels in amniotic fluid and inflammatory changes in the placenta from normal deliveries at term. *Eur J Obstet Gynecol Reprod Biol*. 1994;56(3):153-160.
 17. Dollner H, Vatten L, Halgunset J, Rahimipoor S, Austgulen R. Histologic chorioamnionitis and umbilical serum levels of pro-inflammatory cytokines and cytokine inhibitors. *BJOG*. 2002; 109(5):534-539.
 18. Tasci Y, Dilbaz B, Uzmez Onal B, et al. The value of cord blood interleukin-6 levels for predicting chorioamnionitis, funisitis and neonatal infection in term premature rupture of membranes. *Eur J Obstet Gynecol Reprod Biol*. 2006;128(1-2):34-39.
 19. Duncombe G, Veldhuizen RA, Gratton RJ, Han VK, Richardson BS. IL-6 and TNFalpha across the umbilical circulation in term pregnancies: relationship with labour events. *Early Hum Dev*. 2010;86(2):113-117.
 20. Chan CJ, Summers KL, Chan NG, Hardy DB, Richardson BS. Cytokines in umbilical cord blood and the impact of labor events in low-risk term pregnancies. *Early Hum Dev*. 2013;89(12): 1005-1010.
 21. Baergen R, Benirschke K, Ulich TR. Cytokine expression in the placenta. The role of interleukin 1 and interleukin 1 receptor antagonist expression in chorioamnionitis and parturition. *Arch Pathol Lab Med*. 1994;118(1):52-55.
 22. Lockwood CJ, Arcuri F, Toti P, et al. Tumor necrosis factor-alpha and interleukin-1beta regulate interleukin-8 expression in third trimester decidual cells: implications for the genesis of chorioamnionitis. *Am J Pathol*. 2006;169(4):1294-1302.
 23. Hung TH, Chen SF, Hsu JJ, Hsieh CC, Hsueh S, Hsieh TT. Tumor necrosis factor-alpha converting enzyme in human gestational tissues from pregnancies complicated by chorioamnionitis. *Placenta*. 2006;27(9-10):996-1006.
 24. Arcuri F, Toti P, Buchwalder L, et al. Mechanisms of leukocyte accumulation and activation in chorioamnionitis: interleukin 1 beta and tumor necrosis factor alpha enhance colony stimulating factor 2 expression in term decidua. *Reprod Sci*. 2009;16(5): 453-461.
 25. Lockwood CJ, Murk WK, Kayisli UA, et al. Regulation of interleukin-6 expression in human decidual cells and its potential role in chorioamnionitis. *Am J Pathol*. 2010;177(4): 1755-1764.
 26. Vanderhoeven JP, Bierle CJ, Kapur RP, et al. Group B streptococcal infection of the choriodecidua induces dysfunction of the cytokeratin network in amniotic epithelium: a pathway to membrane weakening. *PLoS Pathog*. 2014;10(3): e1003920.
 27. Hillier SL, Witkin SS, Krohn MA, Watts DH, Kiviat NB, Eschenbach DA. The relationship of amniotic fluid cytokines and preterm delivery, amniotic fluid infection, histologic chorioamnionitis, and chorioamnion infection. *Obstet Gynecol*. 1993;81(6):941-948.
 28. Saito S, Kasahara T, Kato Y, Ishihara Y, Ichijo M. Elevation of amniotic fluid interleukin 6 (IL-6), IL-8 and granulocyte colony stimulating factor (G-CSF) in term and preterm parturition. *Cytokine*. 1993;5(1):81-88.
 29. Yoon BH, Jun JK, Romero R, et al. Amniotic fluid inflammatory cytokines (interleukin-6, interleukin-1beta, and tumor necrosis factor-alpha), neonatal brain white matter lesions, and cerebral palsy. *Am J Obstet Gynecol*. 1997;177(1):19-26.
 30. Yoon BH, Romero R, Park JS, et al. Fetal exposure to an intra-amniotic inflammation and the development of cerebral palsy at the age of three years. *Am J Obstet Gynecol*. 2000;182(3): 675-681.
 31. Hitti J, Tarczy-Hornoch P, Murphy J, Hillier SL, Aura J, Eschenbach DA. Amniotic fluid infection, cytokines, and adverse outcome among infants at 34 weeks' gestation or less. *Obstet Gynecol*. 2001;98(6):1080-1088.
 32. Moon JB, Kim JC, Yoon BH, et al. Amniotic fluid matrix metalloproteinase-8 and the development of cerebral palsy. *J Perinat Med*. 2002;30(4):301-306.
 33. Combs CA, Gravett M, Garite TJ, et al; ProteoGenix/Obstetrix Collaborative Research Network. Amniotic fluid infection, inflammation, and colonization in preterm labor with intact membranes. *Am J Obstet Gynecol*. 2014;210(2):125.e121-125 e115.
 34. Kunze M, Klar M, Morfeld CA, et al. Cytokines in noninvasively obtained amniotic fluid as predictors of fetal inflammatory response syndrome. *Am J Obstet Gynecol*. 2016;215(1): 96.e1-8.
 35. Yoon BH, Romero R, Yang SH, et al. Interleukin-6 concentrations in umbilical cord plasma are elevated in neonates with white matter lesions associated with periventricular leukomalacia. *Am J Obstet Gynecol*. 1996;174(5):1433-1440.

36. Weeks JW, Reynolds L, Taylor D, Lewis J, Wan T, Gall SA. Umbilical cord blood interleukin-6 levels and neonatal morbidity. *Obstet Gynecol.* 1997;90(5):815-818.
37. Gomez R, Romero R, Ghezzi F, Yoon BH, Mazon M, Berry SM. The fetal inflammatory response syndrome. *Am J Obstet Gynecol.* 1998;179(1):194-202.
38. Berner R, Niemeyer CM, Leitis JU, et al. Plasma levels and gene expression of granulocyte colony-stimulating factor, tumor necrosis factor- α , interleukin (IL)-1 β , IL-6, IL-8, and soluble intercellular adhesion molecule-1 in neonatal early onset sepsis. *Pediatr Res.* 1998;44(4):469-477.
39. Dollner H, Vatten L, Linnebo I, Zanussi GF, Laerdal A, Austgulen R. Inflammatory mediators in umbilical plasma from neonates who develop early-onset sepsis. *Biol Neonate.* 2001;80(1):41-47.
40. Goepfert AR, Andrews WW, Carlo W, et al. Umbilical cord plasma interleukin-6 concentrations in preterm infants and risk of neonatal morbidity. *Am J Obstet Gynecol.* 2004;191(4):1375-1381.
41. An H, Nishimaki S, Ohyama M, et al. Interleukin-6, interleukin-8, and soluble tumor necrosis factor receptor-I in the cord blood as predictors of chronic lung disease in premature infants. *Am J Obstet Gynecol.* 2004;191(5):1649-1654.
42. Elsmen E, Ley D, Cilio CM, Hansen-Pupp I, Hellstrom-Westas L. Umbilical cord levels of interleukin-1 receptor antagonist and neonatal outcome. *Biol Neonate.* 2006;89(4):220-226.
43. Satar M, Turhan E, Yapicioglu H, Narli N, Ozgunen FT, Cetiner S. Cord blood cytokine levels in neonates born to mothers with prolonged premature rupture of membranes and its relationship with morbidity and mortality. *Eur Cytokine Netw.* 2008;19(1):37-41.
44. Liu J, Feng ZC. Increased umbilical cord plasma interleukin-1 β levels was correlated with adverse outcomes of neonatal hypoxic-ischemic encephalopathy. *J Trop Pediatr.* 2010;56(3):178-182.
45. Armstrong-Wells J, Donnelly M, Post MD, Manco-Johnson MJ, Winn VD, Sebire G. Inflammatory predictors of neurologic disability after preterm premature rupture of membranes. *Am J Obstet Gynecol.* 2015;212(2):212.e211-e219.
46. Cordeiro CN, Savva Y, Vaidya D, et al. Mathematical modeling of the biomarker milieu to characterize preterm birth and predict adverse neonatal outcomes. *Am J Reprod Immunol.* 2016;75(5):594-601.
47. Redline RW, Faye-Petersen O, Heller D, Qureshi F, Savell V, Vogler C; Society for Pediatric Pathology, Perinatal Section, Amniotic Fluid Infection Nosology Committee. Amniotic infection syndrome: nosology and reproducibility of placental reaction patterns. *Pediatr Dev Pathol.* 2003;6(5):435-448.
48. Romero R, Ceska M, Avila C, Mazon M, Behnke E, Lindley I. Neutrophil attractant/activating peptide-1/interleukin-8 in term and preterm parturition. *Am J Obstet Gynecol.* 1991;165(4 pt 1):813-820.
49. Cherouny PH, Pankuch GA, Romero R, et al. Neutrophil attractant/activating peptide-1/interleukin-8: association with histologic chorioamnionitis, preterm delivery, and bioactive amniotic fluid leukoattractants. *Am J Obstet Gynecol.* 1993;169(5):1299-1303.
50. Gomez R, Ghezzi F, Romero R, Munoz H, Tolosa JE, Rojas I. Premature labor and intra-amniotic infection. Clinical aspects and role of the cytokines in diagnosis and pathophysiology. *Clin Perinatol.* 1995;22(2):281-342.
51. Yoon BH, Romero R, Jun JK, et al. Amniotic fluid cytokines (interleukin-6, tumor necrosis factor- α , interleukin-1 β , and interleukin-8) and the risk for the development of bronchopulmonary dysplasia. *Am J Obstet Gynecol.* 1997;177(4):825-830.
52. Ghezzi F, Gomez R, Romero R, et al. Elevated interleukin-8 concentrations in amniotic fluid of mothers whose neonates subsequently develop bronchopulmonary dysplasia. *Eur J Obstet Gynecol Reprod Biol.* 1998;78(1):5-10.
53. Cohen J, Ghezzi F, Romero R, et al. GRO α in the fetal-maternal and amniotic fluid compartments during pregnancy and parturition. *Am J Reprod Immunol.* 1996;35(1):23-29.
54. Hsu CD, Meaddough E, Aversa K, Copel JA. The role of amniotic fluid L-selectin, GRO- α , and interleukin-8 in the pathogenesis of intraamniotic infection. *Am J Obstet Gynecol.* 1998;178(3):428-432.
55. Hsu CD, Meaddough E, Aversa K, et al. Elevated amniotic fluid levels of leukemia inhibitory factor, interleukin 6, and interleukin 8 in intra-amniotic infection. *Am J Obstet Gynecol.* 1998;179(5):1267-1270.
56. Jacobsson B, Mattsby-Baltzer I, Andersch B, et al. Microbial invasion and cytokine response in amniotic fluid in a Swedish population of women in preterm labor. *Acta Obstet Gynecol Scand.* 2003;82(2):120-128.
57. Jacobsson B, Mattsby-Baltzer I, Andersch B, et al. Microbial invasion and cytokine response in amniotic fluid in a Swedish population of women with preterm prelabor rupture of membranes. *Acta Obstet Gynecol Scand.* 2003;82(5):423-431.
58. Figueroa R, Garry D, Elimian A, Patel K, Sehgal PB, Tejani N. Evaluation of amniotic fluid cytokines in preterm labor and intact membranes. *J Matern Fetal Neonatal Med.* 2005;18(4):241-247.
59. Witt A, Berger A, Gruber CJ, Petricevic L, Apfalter P, Husslein P. IL-8 concentrations in maternal serum, amniotic fluid and cord blood in relation to different pathogens within the amniotic cavity. *J Perinat Med.* 2005;33(1):22-26.
60. Mittal P, Romero R, Kusanovic JP, et al. CXCL6 (granulocyte chemotactic protein-2): a novel chemokine involved in the innate immune response of the amniotic cavity. *Am J Reprod Immunol.* 2008;60(3):246-257.
61. Romero R, Chaemsathong P, Korzeniewski SJ, et al. Clinical chorioamnionitis at term II: the intra-amniotic inflammatory response. *J Perinat Med.* 2016;44(1):5-22.
62. Black RA, Kronheim SR, Merriam JE, March CJ, Hopp TP. A pre-aspartate-specific protease from human leukocytes that cleaves pro-interleukin-1 β . *J Biol Chem.* 1989;264(10):5323-5326.
63. Kostura MJ, Tocci MJ, Limjuco G, et al. Identification of a monocyte specific pre-interleukin 1 β convertase activity. *Proc Natl Acad Sci U S A.* 1989;86(14):5227-5231.
64. Thornberry NA, Bull HG, Calaycay JR, et al. A novel heterodimeric cysteine protease is required for interleukin-1 β processing in monocytes. *Nature.* 1992;356(6372):768-774.

65. Wilson KP, Black JA, Thomson JA, et al. Structure and mechanism of interleukin-1 beta converting enzyme. *Nature*. 1994; 370(6487):270-275.
66. Dinarello CA. Interleukin-1 beta, interleukin-18, and the interleukin-1 beta converting enzyme. *Ann N Y Acad Sci*. 1998; 856:1-11.
67. Fantuzzi G, Dinarello CA. Interleukin-18 and interleukin-1 beta: two cytokine substrates for ICE (caspase-1). *J Clin Immunol*. 1999;19(1):1-11.
68. Sansonetti PJ, Phalipon A, Arondel J, et al. Caspase-1 activation of IL-1beta and IL-18 are essential for Shigella flexneri-induced inflammation. *Immunity*. 2000;12(5):581-590.
69. Martinon F, Burns K, Tschopp J. The inflammasome: a molecular platform triggering activation of inflammatory caspases and processing of proIL-beta. *Mol Cell*. 2002;10(2):417-426.
70. Agostini L, Martinon F, Burns K, McDermott MF, Hawkins PN, Tschopp J. NALP3 forms an IL-1beta-processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder. *Immunity*. 2004;20(3):319-325.
71. Mariathasan S, Newton K, Monack DM, et al. Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature*. 2004;430(6996):213-218.
72. Petrilli V, Papin S, Tschopp J. The inflammasome. *Curr Biol*. 2005;15(15): R581.
73. Ogura Y, Sutterwala FS, Flavell RA. The inflammasome: first line of the immune response to cell stress. *Cell*. 2006;126(4): 659-662.
74. Sutterwala FS, Ogura Y, Flavell RA. The inflammasome in pathogen recognition and inflammation. *J Leukoc Biol*. 2007;82(2): 259-264.
75. Mariathasan S, Monack DM. Inflammasome adaptors and sensors: intracellular regulators of infection and inflammation. *Nat Rev Immunol*. 2007;7(1):31-40.
76. Lamkanfi M, Dixit VM. Inflammasomes: guardians of cytosolic sanctity. *Immunol Rev*. 2009;227(1):95-105.
77. Pedra JH, Cassel SL, Sutterwala FS. Sensing pathogens and danger signals by the inflammasome. *Curr Opin Immunol*. 2009; 21(1):10-16.
78. Franchi L, Eigenbrod T, Munoz-Planillo R, Nunez G. The inflammasome: a caspase-1-activation platform that regulates immune responses and disease pathogenesis. *Nat Immunol*. 2009;10(3): 241-247.
79. Netea MG, Nold-Petry CA, Nold MF, et al. Differential requirement for the activation of the inflammasome for processing and release of IL-1beta in monocytes and macrophages. *Blood*. 2009; 113(10):2324-2335.
80. Stutz A, Golenbock DT, Latz E. Inflammasomes: too big to miss. *J Clin Invest*. 2009;119(12):3502-3511.
81. Jha S, Ting JP. Inflammasome-associated nucleotide-binding domain, leucine-rich repeat proteins and inflammatory diseases. *J Immunol*. 2009;183(12):7623-7629.
82. Schroder K, Zhou R, Tschopp J. The NLRP3 inflammasome: a sensor for metabolic danger? *Science*. 2010;327(5963): 296-300.
83. Schroder K, Tschopp J. The inflammasomes. *Cell*. 2010;140(6): 821-832.
84. Franchi L, Munoz-Planillo R, Reimer T, Eigenbrod T, Nunez G. Inflammasomes as microbial sensors. *Eur J Immunol*. 2010; 40(3):611-615.
85. Khare S, Luc N, Dorfleutner A, Stehlik C. Inflammasomes and their activation. *Crit Rev Immunol*. 2010;30(5):463-487.
86. Gross O, Thomas CJ, Guarda G, Tschopp J. The inflammasome: an integrated view. *Immunol Rev*. 2011;243(1):136-151.
87. Bauernfeind F, Ablasser A, Bartok E, et al. Inflammasomes: current understanding and open questions. *Cell Mol Life Sci*. 2011;68(5):765-783.
88. Horvath GL, Schrum JE, De Nardo CM, Latz E. Intracellular sensing of microbes and danger signals by the inflammasomes. *Immunol Rev*. 2011;243(1):119-135.
89. Franchi L, Munoz-Planillo R, Nunez G. Sensing and reacting to microbes through the inflammasomes. *Nat Immunol*. 2012; 13(4):325-332.
90. Dagenais M, Skeldon A, Saleh M. The inflammasome: in memory of Dr. Jurg Tschopp. *Cell Death Differ*. 2012;19(1):5-12.
91. Ciraci C, Janczy JR, Sutterwala FS, Cassel SL. Control of innate and adaptive immunity by the inflammasome. *Microbes Infect*. 2012;14(14):1263-1270.
92. Rathinam VA, Vanaja SK, Fitzgerald KA. Regulation of inflammasome signaling. *Nat Immunol*. 2012;13(4):333-342.
93. Franchi L, Nunez G. Immunology. Orchestrating inflammasomes. *Science*. 2012;337(6100):1299-1300.
94. Henao-Mejia J, Elinav E, Strowig T, Flavell RA. Inflammasomes: far beyond inflammation. *Nat Immunol*. 2012;13(4):321-324.
95. Bauernfeind F, Hornung V. Of inflammasomes and pathogens—sensing of microbes by the inflammasome. *EMBO Mol Med*. 2013;5(6):814-826.
96. Vladimer GI, Marty-Roix R, Ghosh S, Weng D, Lien E. Inflammasomes and host defenses against bacterial infections. *Curr Opin Microbiol*. 2013;16(1):23-31.
97. Ulland TK, Ferguson PJ, Sutterwala FS. Evasion of inflammasome activation by microbial pathogens. *J Clin Invest*. 2015; 125(2):469-477.
98. Gu Y, Kuida K, Tsutsui H, et al. Activation of interferon-gamma inducing factor mediated by interleukin-1beta converting enzyme. *Science*. 1997;275(5297):206-209.
99. Ghayur T, Banerjee S, Hugunin M, et al. Caspase-1 processes IFN-gamma-inducing factor and regulates LPS-induced IFN-gamma production. *Nature*. 1997;386(6625):619-623.
100. Latz E. The inflammasomes: mechanisms of activation and function. *Curr Opin Immunol*. 2010;22(1):28-33.
101. van de Veerdonk FL, Netea MG, Dinarello CA, Joosten LA. Inflammasome activation and IL-1beta and IL-18 processing during infection. *Trends Immunol*. 2011;32(3):110-116.
102. Skeldon A, Saleh M. The inflammasomes: molecular effectors of host resistance against bacterial, viral, parasitic, and fungal infections. *Front Microbiol*. 2011;2:15.
103. Lamkanfi M, Dixit VM. Modulation of inflammasome pathways by bacterial and viral pathogens. *J Immunol*. 2011;187(2): 597-602.
104. Broz P, Monack DM. Molecular mechanisms of inflammasome activation during microbial infections. *Immunol Rev*. 2011; 243(1):174-190.

105. Koizumi Y, Toma C, Higa N, Nohara T, Nakasone N, Suzuki T. Inflammasome activation via intracellular NLRs triggered by bacterial infection. *Cell Microbiol.* 2012;14(2):149-154.
106. Latz E, Xiao TS, Stutz A. Activation and regulation of the inflammasomes. *Nat Rev Immunol.* 2013;13(6):397-411.
107. Lamkanfi M, Dixit VM. Mechanisms and functions of inflammasomes. *Cell.* 2014;157(5):1013-1022.
108. Vanaja SK, Rathinam VA, Fitzgerald KA. Mechanisms of inflammasome activation: recent advances and novel insights. *Trends Cell Biol.* 2015;25(5):308-315.
109. Guo H, Callaway JB, Ting JP. Inflammasomes: mechanism of action, role in disease, and therapeutics. *Nat Med.* 2015;21(7):677-687.
110. Wang S, Miura M, Jung YK, Zhu H, Li E, Yuan J. Murine caspase-11, an ICE-interacting protease, is essential for the activation of ICE. *Cell.* 1998;92(4):501-509.
111. Sollberger G, Strittmatter GE, Kistowska M, French LE, Beer HD. Caspase-4 is required for activation of inflammasomes. *J Immunol.* 2012;188(4):1992-2000.
112. Pineles BL, Romero R, Montenegro D, et al. "The inflammasome" in human parturition. *Reprod Sci.* 2007;14(1):59A.
113. Gotsch F, Romero R, Chaiworapongsa T, et al. Evidence of the involvement of caspase-1 under physiologic and pathologic cellular stress during human pregnancy: a link between the inflammasome and parturition. *J Matern Fetal Neonatal Med.* 2008;21(9):605-616.
114. Romero R, Xu Y, Plazyo O, et al. A role for the inflammasome in spontaneous labor at term [published online March 8, 2016]. *Am J Reprod Immunol.* doi:10.1111/aji.12440.
115. Montenegro D, Romero R, Pineles P, et al. Differential expression of the inflammasome components in the fetal inflammatory response syndrome. *Reprod Sci.* 2007;14(1):59A-60A.
116. Abrahams VM. The role of the Nod-like receptor family in trophoblast innate immune responses. *J Reprod Immunol.* 2011;88(2):112-117.
117. Mulla MJ, Myrtolli K, Potter J, et al. Uric acid induces trophoblast IL-1 β production via the inflammasome: implications for the pathogenesis of preeclampsia. *Am J Reprod Immunol.* 2011;65(6):542-548.
118. Pontillo A, Girardelli M, Agostinis C, Masat E, Bulla R, Crovella S. Bacterial LPS differently modulates inflammasome gene expression and IL-1 β secretion in trophoblast cells, decidual stromal cells, and decidual endothelial cells. *Reprod Sci.* 2013;20(5):563-566.
119. Mulla MJ, Salmon JE, Chamley LW, et al. A role for uric acid and the Nalp3 inflammasome in antiphospholipid antibody-induced IL-1 β production by human first trimester trophoblast. *PLoS One.* 2013;8(6): e65237.
120. Abrahams VM. Novel mechanisms of placenta inflammation in obstetrics antiphospholipid syndrome. *Am J Reprod Immunol.* 2014;71(suppl 1):25.
121. Hansen L, Kotla S, Mari G, Rao G. The role of NLRP3 inflammasome in preeclampsia—a translational approach. *Am J Obstet Gynecol.* 2014;210(1): S134.
122. Khan RN, Hay DP. A clear and present danger: inflammasomes DAMPING down disorders of pregnancy. *Hum Reprod Update.* 2015;21(3):388-405.
123. Pontillo A, Reis EC, Bricher PN, et al. NLRP1 L155H polymorphism is a risk factor for preeclampsia development. *Am J Reprod Immunol.* 2015;73(6):577-581.
124. Maneta E, Warren AY, Hay DP, Khan RN. Caspase-1-mediated cytokine release from gestational tissues, placental, and cord blood. *Front Physiol.* 2015;6:186.
125. Romero R, Mazor M, Brandt F, et al. Interleukin-1 α and interleukin-1 β in preterm and term human parturition. *Am J Reprod Immunol.* 1992;27(3-4):117-123.
126. Pacora P, Romero R, Maymon E, et al. Participation of the novel cytokine interleukin 18 in the host response to intra-amniotic infection. *Am J Obstet Gynecol.* 2000;183(5): 1138-1143.
127. American College of Obstetrics and Gynecology Committee on Practice Bulletins-Obstetrics. ACOG practice bulletin number 49, December 2003: dystocia and augmentation of labor. *Obstet Gynecol.* 2003;102(6):1445-1454.
128. Alexander GR, Himes JH, Kaufman RB, Mor J, Kogan M. A United States national reference for fetal growth. *Obstet Gynecol.* 1996;87(2):163-168.
129. Redline RW. Placental pathology: a systematic approach with clinical correlations. *Placenta.* 2008;29(suppl A): S86-S91.
130. Hoang M, Potter JA, Gysler SM, et al. Human fetal membranes generate distinct cytokine profiles in response to bacterial Toll-like receptor and nod-like receptor agonists. *Biol Reprod.* 2014; 90(2):39.
131. Sutterwala FS, Ogura Y, Zamboni DS, Roy CR, Flavell RA. NALP3: a key player in caspase-1 activation. *J Endotoxin Res.* 2006;12(4):251-256.
132. Chae JJ, Cho YH, Lee GS, et al. Gain-of-function pyrin mutations induce NLRP3 protein-independent interleukin-1 β activation and severe autoinflammation in mice. *Immunity.* 2011; 34(5):755-768.
133. Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature.* 2006;440(7081):237-241.
134. Kanneganti TD, Body-Malapel M, Amer A, et al. Critical role for Cryopyrin/Nalp3 in activation of caspase-1 in response to viral infection and double-stranded RNA. *J Biol Chem.* 2006; 281(48):36560-36568.
135. Mariathasan S, Weiss DS, Newton K, et al. Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature.* 2006; 440(7081):228-232.
136. Muruve DA, Petrilli V, Zaiss AK, et al. The inflammasome recognizes cytosolic microbial and host DNA and triggers an innate immune response. *Nature.* 2008;452(7183): 103-107.
137. Dostert C, Petrilli V, Van Bruggen R, Steele C, Mossman BT, Tschopp J. Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science.* 2008;320(5876): 674-677.
138. Cassel SL, Eisenbarth SC, Iyer SS, et al. The Nalp3 inflammasome is essential for the development of silicosis. *Proc Natl Acad Sci U S A.* 2008;105(26):9035-9040.
139. Eisenbarth SC, Colegio OR, O'Connor W, Sutterwala FS, Flavell RA. Crucial role for the Nalp3 inflammasome in the

- immunostimulatory properties of aluminium adjuvants. *Nature*. 2008;453(7198):1122-1126.
140. Li H, Willingham SB, Ting JP, Re F. Cutting edge: inflammasome activation by alum and alum's adjuvant effect are mediated by NLRP3. *J Immunol*. 2008;181(1):17-21.
141. Hornung V, Bauernfeind F, Halle A, et al. Silica crystals and aluminum salts activate the NALP3 inflammasome through phagosomal destabilization. *Nat Immunol*. 2008;9(8):847-856.
142. Franchi L, Nunez G. The Nlrp3 inflammasome is critical for aluminium hydroxide-mediated IL-1 β secretion but dispensable for adjuvant activity. *Eur J Immunol*. 2008;38(8):2085-2089.
143. Kool M, Petrilli V, De Smedt T, et al. Cutting edge: alum adjuvant stimulates inflammatory dendritic cells through activation of the NALP3 inflammasome. *J Immunol*. 2008;181(6):3755-3759.
144. Koo IC, Wang C, Raghavan S, Morisaki JH, Cox JS, Brown EJ. ESX-1-dependent cytolysis in lysosome secretion and inflammasome activation during mycobacterial infection. *Cell Microbiol*. 2008;10(9):1866-1878.
145. Allen IC, Scull MA, Moore CB, et al. The NLRP3 inflammasome mediates in vivo innate immunity to influenza A virus through recognition of viral RNA. *Immunity*. 2009;30(4):556-565.
146. Duncan JA, Gao X, Huang MT, et al. Neisseria gonorrhoeae activates the proteinase cathepsin B to mediate the signaling activities of the NLRP3 and ASC-containing inflammasome. *J Immunol*. 2009;182(10):6460-6469.
147. Joly S, Ma N, Sadler JJ, Soll DR, Cassel SL, Sutterwala FS. Cutting edge: *Candida albicans* hyphae formation triggers activation of the Nlrp3 inflammasome. *J Immunol*. 2009;183(6):3578-3581.
148. Munoz-Planillo R, Franchi L, Miller LS, Nunez G. A critical role for hemolysins and bacterial lipoproteins in *Staphylococcus aureus*-induced activation of the Nlrp3 inflammasome. *J Immunol*. 2009;183(6):3942-3948.
149. Yamasaki K, Muto J, Taylor KR, et al. NLRP3/cryopyrin is necessary for interleukin-1 β (IL-1 β) release in response to hyaluronan, an endogenous trigger of inflammation in response to injury. *J Biol Chem*. 2009;284(19):12762-12771.
150. Cassel SL, Joly S, Sutterwala FS. The NLRP3 inflammasome: a sensor of immune danger signals. *Semin Immunol*. 2009;21(4):194-198.
151. Demento SL, Eisenbarth SC, Foellmer HG, et al. Inflammasome-activating nanoparticles as modular systems for optimizing vaccine efficacy. *Vaccine*. 2009;27(23):3013-3021.
152. Thomas PG, Dash P, Aldridge JR Jr, et al. The intracellular sensor NLRP3 mediates key innate and healing responses to influenza A virus via the regulation of caspase-1. *Immunity*. 2009;30(4):566-575.
153. Ichinohe T, Lee HK, Ogura Y, Flavell R, Iwasaki A. Inflammasome recognition of influenza virus is essential for adaptive immune responses. *J Exp Med*. 2009;206(1):79-87.
154. Cassel SL, Sutterwala FS. Sterile inflammatory responses mediated by the NLRP3 inflammasome. *Eur J Immunol*. 2010;40(3):607-611.
155. Leemans JC, Cassel SL, Sutterwala FS. Sensing damage by the NLRP3 inflammasome. *Immunol Rev*. 2011;243(1):152-162.
156. Nakahira K, Haspel JA, Rathinam VA, et al. Autophagy proteins regulate innate immune responses by inhibiting the release of mitochondrial DNA mediated by the NALP3 inflammasome. *Nat Immunol*. 2011;12(3):222-230.
157. Menu P, Vince JE. The NLRP3 inflammasome in health and disease: the good, the bad and the ugly. *Clin Exp Immunol*. 2011;166(1):1-15.
158. Rathinam VA, Vanaja SK, Waggoner L, et al. TRIF licenses caspase-11-dependent NLRP3 inflammasome activation by gram-negative bacteria. *Cell*. 2012;150(3):606-619.
159. Iyer SS, He Q, Janczy JR, et al. Mitochondrial cardiolipin is required for Nlrp3 inflammasome activation. *Immunity*. 2013;39(2):311-323.
160. O'Neill LA. Cardiolipin and the Nlrp3 inflammasome. *Cell Metab*. 2013;18(5):610-612.
161. Clay GM, Sutterwala FS, Wilson ME. NLR proteins and parasitic disease. *Immunol Res*. 2014;59(1-3):142-152.
162. Bauernfeind FG, Horvath G, Stutz A, et al. Cutting edge: NF- κ B activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression. *J Immunol*. 2009;183(2):787-791.
163. Sutterwala FS, Haasken S, Cassel SL. Mechanism of NLRP3 inflammasome activation. *Ann N Y Acad Sci*. 2014;1319:82-95.
164. Franchi L, Eigenbrod T, Nunez G. Cutting edge: TNF- α mediates sensitization to ATP and silica via the NLRP3 inflammasome in the absence of microbial stimulation. *J Immunol*. 2009;183(2):792-796.
165. Sutterwala FS, Flavell RA. NLRC4/IPAF: a CARD carrying member of the NLR family. *Clin Immunol*. 2009;130(1):2-6.
166. Franchi L, Amer A, Body-Malapel M, et al. Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1 β in salmonella-infected macrophages. *Nat Immunol*. 2006;7(6):576-582.
167. Miao EA, Alpuche-Aranda CM, Dors M, et al. Cytoplasmic flagellin activates caspase-1 and secretion of interleukin 1 β via Ipaf. *Nat Immunol*. 2006;7(6):569-575.
168. Poyet JL, Srinivasula SM, Tnani M, Razmara M, Fernandes-Alnemri T, Alnemri ES. Identification of Ipaf, a human caspase-1-activating protein related to Apaf-1. *J Biol Chem*. 2001;276(30):28309-28313.
169. Mayor A, Martinon F, De Smedt T, Petrilli V, Tschopp J. A crucial function of SGT1 and HSP90 in inflammasome activity links mammalian and plant innate immune responses. *Nat Immunol*. 2007;8(5):497-503.
170. Petrilli V, Papin S, Dostert C, Mayor A, Martinon F, Tschopp J. Activation of the NALP3 inflammasome is triggered by low intracellular potassium concentration. *Cell Death Differ*. 2007;14(9):1583-1589.
171. Fernandes-Alnemri T, Wu J, Yu JW, et al. The pyroptosome: a supramolecular assembly of ASC dimers mediating inflammatory cell death via caspase-1 activation. *Cell Death Differ*. 2007;14(9):1590-1604.

172. Kayagaki N, Warming S, Lamkanfi M, et al. Non-canonical inflammasome activation targets caspase-11. *Nature*. 2011; 479(7371):117-121.
173. Broz P, Ruby T, Belhocine K, et al. Caspase-11 increases susceptibility to *Salmonella* infection in the absence of caspase-1. *Nature*. 2012;490(7419):288-291.
174. Romero R, Mazor M, Wu YK, Avila C, Oyarzun E, Mitchell MD. Bacterial endotoxin and tumor necrosis factor stimulate prostaglandin production by human decidua. *Prostaglandins Leukot Essent Fatty Acids*. 1989;37(3):183-186.
175. Fidel PL Jr, Romero R, Ramirez M, et al. Interleukin-1 receptor antagonist (IL-1ra) production by human amnion, chorion, and decidua. *Am J Reprod Immunol*. 1994;32(1):1-7.
176. Vega-Sanchez R, Gomez-Lopez N, Flores-Pliego A, et al. Placental blood leukocytes are functional and phenotypically different than peripheral leukocytes during human labor. *J Reprod Immunol*. 2010;84(1):100-110.
177. Khan KN, Kitajima M, Inoue T, Fujishita A, Nakashima M, Masuzaki H. 17beta-estradiol and lipopolysaccharide additively promote pelvic inflammation and growth of endometriosis. *Reprod Sci*. 2015;22(5):585-594.
178. Fidel PL Jr, Romero R, Wolf N, et al. Systemic and local cytokine profiles in endotoxin-induced preterm parturition in mice. *Am J Obstet Gynecol*. 1994;170(5 pt 1):1467-1475.
179. Fidel PL Jr, Romero R, Cutright J, et al. Treatment with the interleukin-1 receptor antagonist and soluble tumor necrosis factor receptor Fc fusion protein does not prevent endotoxin-induced preterm parturition in mice. *J Soc Gynecol Investig*. 1997;4(1):22-26.
180. Saito M, Payne MS, Miura Y, et al. Polymyxin B agonist capture therapy for intrauterine inflammation: proof-of-principle in a fetal ovine model. *Reprod Sci*. 2014;21(5):623-631.
181. Pineda-Torres M, Flores-Espinosa P, Espejel-Nunez A, et al. Evidence of an immunosuppressive effect of progesterone upon in vitro secretion of proinflammatory and prodegradative factors in a model of choriodecidual infection. *BJOG*. 2015;122(13):1798-1807.
182. Furuya H, Taguchi A, Kawana K, et al. Resveratrol protects against pathological preterm birth by suppression of macrophage-mediated inflammation. *Reprod Sci*. 2015;22(12):1561-1568.
183. Arenas-Hernandez M, Romero R, St Louis D, Hassan SS, Kaye EB, Gomez-Lopez N. An imbalance between innate and adaptive immune cells at the maternal-fetal interface occurs prior to endotoxin-induced preterm birth. *Cell Mol Immunol*. 2016; 13(4):462-473.
184. Xu Y, Romero R, Miller D, et al. An M1-like macrophage polarization in decidual tissue during spontaneous preterm labor that is attenuated by rosiglitazone treatment. *J Immunol*. 2016; 196(6):2476-2491.
185. Stack JH, Beaumont K, Larsen PD, et al. IL-converting enzyme/caspase-1 inhibitor VX-765 blocks the hypersensitive response to an inflammatory stimulus in monocytes from familial cold autoinflammatory syndrome patients. *J Immunol*. 2005;175(4): 2630-2634.
186. Wannamaker W, Davies R, Namchuk M, et al. (S)-1-((S)-2-[[1-(4-Amino-3-chloro-phenyl)-methanoyl]-amino]-3,3-dimethylbutanoyl)-pyrrolidine-2-carboxylic acid ((2R,3S)-2-ethoxy-5-oxo-tetrahydro-furan-3-yl)-amide (VX-765), an orally available selective interleukin (IL)-converting enzyme/caspase-1 inhibitor, exhibits potent anti-inflammatory activities by inhibiting the release of IL-1beta and IL-18. *J Pharmacol Exp Ther*. 2007; 321(2):509-516.
187. Romero R, Durum S, Dinarello CA, Oyarzun E, Hobbins JC, Mitchell MD. Interleukin-1 stimulates prostaglandin biosynthesis by human amnion. *Prostaglandins*. 1989;37(1): 13-22.
188. Hertelendy F, Romero R, Molnar M, Todd H, Baldassare JJ. Cytokine-initiated signal transduction in human myometrial cells. *Am J Reprod Immunol*. 1993;30(2-3):49-57.
189. Belt AR, Baldassare JJ, Molnar M, Romero R, Hertelendy F. The nuclear transcription factor NF-kappaB mediates interleukin-1beta-induced expression of cyclooxygenase-2 in human myometrial cells. *Am J Obstet Gynecol*. 1999;181(2): 359-366.
190. Watari M, Watari H, DiSanto ME, Chacko S, Shi GP, Strauss JF III. Pro-inflammatory cytokines induce expression of matrix-metabolizing enzymes in human cervical smooth muscle cells. *Am J Pathol*. 1999;154(6):1755-1762.
191. Hertelendy F, Rastogi P, Molnar M, Romero R. Interleukin-1beta-induced prostaglandin E2 production in human myometrial cells: role of a pertussis toxin-sensitive component. *Am J Reprod Immunol*. 2001;45(3):142-147.
192. Heng YJ, Liong S, Permezel M, Rice GE, Di Quinzio MK, Georgiou HM. The interplay of the interleukin 1 system in pregnancy and labor. *Reprod Sci*. 2014;21(1):122-130.
193. Stock O, Gordon L, Kapoor J, et al. Chorioamnionitis occurring in women with preterm rupture of the fetal membranes is associated with a dynamic increase in mRNAs coding cytokines in the maternal circulation. *Reprod Sci*. 2015;22(7):852-859.
194. Romero R, Mazor M, Tartakovsky B. Systemic administration of interleukin-1 induces preterm parturition in mice. *Am J Obstet Gynecol*. 1991;165(4 pt 1):969-971.
195. Romero R, Sepulveda W, Mazor M, et al. The natural interleukin-1 receptor antagonist in term and preterm parturition. *Am J Obstet Gynecol*. 1992;167(4 pt 1):863-872.
196. Gravett MG, Witkin SS, Haluska GJ, Edwards JL, Cook MJ, Novy MJ. An experimental model for intraamniotic infection and preterm labor in rhesus monkeys. *Am J Obstet Gynecol*. 1994;171(6):1660-1667.
197. Witkin SS, Gravett MG, Haluska GJ, Novy MJ. Induction of interleukin-1 receptor antagonist in rhesus monkeys after intraamniotic infection with group B streptococci or interleukin-1 infusion. *Am J Obstet Gynecol*. 1994;171(6): 1668-1672.
198. Baggia S, Gravett MG, Witkin SS, Haluska GJ, Novy MJ. Interleukin-1 beta intra-amniotic infusion induces tumor necrosis factor-alpha, prostaglandin production, and preterm contractions in pregnant rhesus monkeys. *J Soc Gynecol Investig*. 1996; 3(3):121-126.
199. Vadillo-Ortega F, Sadowsky DW, Haluska GJ, et al. Identification of matrix metalloproteinase-9 in amniotic fluid and amnio-chorion in spontaneous labor and after experimental intrauterine

- infection or interleukin-1 beta infusion in pregnant rhesus monkeys. *Am J Obstet Gynecol.* 2002;186(1):128-138.
200. Sadowsky DW, Adams KM, Gravett MG, Witkin SS, Novy MJ. Preterm labor is induced by intraamniotic infusions of interleukin-1beta and tumor necrosis factor-alpha but not by interleukin-6 or interleukin-8 in a nonhuman primate model. *Am J Obstet Gynecol.* 2006;195(6):1578-1589.
201. Aagaard K, Ganu R, Ma J, et al. Intraamniotic interleukin-1 (IL1 β) induces histological chorioamnionitis and alters the microbiome in a primate model of inflammatory preterm birth. *Am J Obstet Gynecol.* 2014;208(1): S218.
202. Prince A, Ma J, Miller L, et al. Chorioamnionitis induced by intraamniotic injection of IL1, LPS or *Ureaplasma parvum* is associated with an altered microbiome in a primate model of inflammatory preterm birth. *Am J Obstet Gynecol.* 2014;212(1): S153.
203. Presicce P, Senthamaraikannan P, Alvarez M, et al. Neutrophil recruitment and activation in decidua with intra-amniotic IL-1beta in the preterm rhesus macaque. *Biol Reprod.* 2015;92(2):56.