CORRESPONDENCE

Indications for the Surgical Management of Benign Goiter in Adults

by Prof. Dr. med. Detlef K. Bartsch, Prof. Dr. med. Markus Luster, Prof. Dr. med. Heinz J. Buhr, Prof. Dr. med. Dietmar Lorenz, Prof. Dr. med. Christoph Germer, and Prof. Dr. med. Peter E. Goretzki in issue 1–2/2018

Calcitonin Screening in Nodular Goiter—Upper Limits

In the description of calcitonin screening in nodular goiter, the specific dimension has to be considered when reporting the upper-limit values (conversion factor for calcitonin $\text{pg/mL} \times 0.28 = \text{pmol/L}$); this was corrected in issue 9. The article (1) reported a 95% probability for the presence of medullary thyroid carcinoma for limits of $>26 \text{ pg/mL}$ in women and $>60 \text{ pg/mL}$ in men—these data came from an original article by Mian et al. (2). Mian et al. used a receiver-operating characteristic curve (ROC) analysis of baseline calcitonin concentrations to test the best method for distinguishing normal persons and patients with medullary thyroid carcinoma and calculated a cut-off value of $>26 \text{ pg/mL}$ for women and $>68 \text{ pg/mL}$ for men.

When sensitivity and specificity were taken into account, the sex-specific upper limits for baseline calcitonin concentrations that led to a recommendation to operate on the suspected sporadic medullary thyroid carcinoma (MTC) were ca. 30 pg/mL in women and ca. 60 pg/mL in men. These cut-offs were measured by using the newer calcitonin assays and were also confirmed by other working groups (3, 4). This means a grey area for calcitonin of $20\text{--}30 \text{ pg/mL}$ in women and $30\text{--}60 \text{ pg/mL}$ in men, where 6–13% of smaller MTC were missed, but C-cell hyperplasia was diagnosed just as often. This uncertainty can be overcome by checking calcitonin concentrations at 3–6 month intervals. Increasing calcitonin concentrations may indicate an MTC—in which case one should operate. There is no great urgency, however, since thyroidectomy can cure MTC in almost 100% of cases where the calcitonin concentration is below 100 pg/mL . Calcitonin screening should also be undertaken in any nodule that is smaller than 1 cm. MTC of a size of $0.4\text{--}0.6 \text{ cm}$ tend to be associated with calcitonin concentrations of $20\text{--}100 \text{ pg/mL}$.

DOI: 10.3238/arztebl.2018.0221a

References

1. Bartsch DK, Luster M, Buhr HJ, Lorenz D, Germer CT, Goretzki PE on behalf of the German Society for General and Visceral Surgery quality commission: Indications for the surgical management of benign goiter in adults. *Dtsch Arztebl Int* 2018; 115: 1–7.
2. Mian C, Perrino M, Colombo C, et al.: Refining calcium test for the diagnosis of medullary thyroid cancer: cutoffs, procedures, and safety. *J Clin Endocrinol Metab* 2014; 99: 1656–64.
3. Allelein S, Ehlers M, Morneau C, et al.: Measurement of basal serum calcitonin for the diagnosis of medullary thyroid cancer. *Horm Metab Res* 2018; 50: 23–8.
4. Rosario PW, Calsolari MR: Usefulness of serum calcitonin in patients without a suspicious history of medullary thyroid carcinoma and with thyroid nodules without an indication for fine-needle aspiration or with benign cytology. *Horm Metab Res* 2016; 48: 372–6.

Prof. Dr. med. Friedhelm Raue

Prof. Dr. med. Karin Frank-Raue

Endokrinologisch-Nuklearmedizinische Gemeinschaftspraxis
Heidelberg

friedhelm.raue@raue-endokrinologie.de

In Reply:

We thank our correspondents for their contribution, which provides important additional information regarding the problem of raised serum calcitonin and the associated suspected presence of a medullary thyroid carcinoma (MTC).

Our correspondents highlight in a concise way the current problems with the available data, and we fully agree with their comments. As our article (1) was, however, subject to a definite word limit and MTC accounts for less than 10% of all thyroid cancers, we were not able to conduct a comprehensive discussion of the subject. We—as coauthors of the cited article by S Allelein et al. (2)—are fully aware of the blurred delineation between suspicious findings and confirmation of an MTC of prognostic relevance, based on the baseline calcitonin concentration. In this context, the factors of influence pointed out by our correspondents—sex and the assay used for testing—are of enormous relevance.

As MTC is, however, in any case a disorder that should conscientiously be excluded or confirmed pre-surgery, we, as surgeons and nuclear medicine specialists, recommend in cases of doubt a second opinion from an endocrinologist and to confirm the laboratory findings in a repeat test. If the suspected finding of MTC is confirmed, further investigations may be required to confirm or rule out a possible familial disorder (multiple endocrine neoplasia type 2).

DOI: 10.3238/arztebl.2018.0221b

References

1. Bartsch DK, Luster M, Buhr HJ, Lorenz D, Germer CT, Goretzki PE on behalf of the German Society for General and Visceral Surgery quality commission: Indications for the surgical management of benign goiter in adults. *Dtsch Arztebl Int* 2018; 115: 1–7.
2. Allelein S, Ehlers M, Morneau C, et al.: Measurement of basal serum calcitonin for the diagnosis of medullary thyroid cancer. *Horm Metab Res* 2018; 50: 23–8.

Prof. Dr. med. Detlef K. Bartsch

Klinik für Visceral-, Thorax- und Gefäßchirurgie
Universitätsklinikum Gießen und Marburg GmbH
Standort Marburg
bartsch@med.uni-marburg.de

Prof. Dr. med. Markus Luster

Klinik für Nuklearmedizin
Universitätsklinikum Gießen und Marburg GmbH
Standort Marburg

Prof. Dr. med. Peter E. Goretzki

Klinik für Allgemein-, Viszeral- und Endokrine Chirurgie
Lukaskrankenhaus Neuss

Conflict of interest statement

The authors of both contributions declare that no conflict of interest exists.