Published in final edited form as:

Curr Opin HIV AIDS. 2014 March ; 9(2): 168–173. doi:10.1097/COH.0000000000000030.

HIV Epidemics among Transgender Women

Tonia Poteat, PhD, MPH,

Johns Hopkins School of Public Health, 615 N. Wolfe Street, Room E7138., Baltimore, MD 21205

Sari L. Reisner, ScD, MA, and

The Fenway Institute, Fenway Health; Department of Epidemiology, Harvard School of Public Health

Anita Radix, MD, MPH

Callen-Lorde Community Health Center

Abstract

Purpose of review—Recent data on the high burden of HIV among transgender women has stimulated interest in addressing HIV in this vulnerable population. This review situates the epidemiologic data on HIV among transgender women in the context of the social determinants of health and describes opportunities for effective interventions.

Recent findings—Transgender women experience unique vulnerability to HIV that can be attributed to multi-level, intersecting factors that also influence the HIV treatment and care continuum. Stigma and discrimination, lack of social and legal recognition of their affirmed gender, and exclusion from employment and educational opportunities represent fundamental drivers of HIV risk in transgender women worldwide.

Summary—Interventions to improve engagement in HIV prevention, testing, care, and treatment among transgender women should build on community strengths and address structural factors as well as psychosocial and biologic factors that increase HIV vulnerability and prevent access to HIV services.

Keywords

transgender; HIV; social determinants

Introduction

Transgender is a term used to describe a diverse population whose gender identity or expression differs from that associated with their sex at birth. Language about and concepts of gender vary widely by region and culture and dynamically change over time. For the purposes of this review, “transgender women” refers to people who were natal males and who have a feminine gender identity or expression.

*corresponding author: TPoteat@jhsph.edu, +1-404-484-1118.

Conflicts of Interest

There are no conflicts of interest.

In 2008, Herbst and colleagues [1] published the first systematic review and meta-analysis of HIV prevalence data among transgender women in the United States. This exhaustive review used five search strategies and included 13 years of data (1990-2003), yet identified only 22 studies that included HIV prevalence for transgender women. In these studies, the average prevalence was 11.8%. The four studies that included laboratory confirmation found an average prevalence of 27.7%; and the two studies with incidence data found 3.4 and 7.8 infections per 100 person-years, respectively.

In the same year, Operario and colleagues [2] published a systematic review and meta-analysis comparing the global epidemics of HIV among transgender women sex workers to transgender women not engaged in sex work, male sex workers, and female sex workers. They identified 25 studies from 14 countries published 1980 – 2007. Overall crude HIV prevalence was 27.3% among transgender women sex workers and 14.7% among transgender women not engaged in sex work, including both self-report and laboratory-confirmed HIV status.

Since those seminal publications, attention to HIV among transgender women has grown such that a search of the literature published within the last year alone resulted in more than 30 unique citations. Using a social ecological framework [3] and focusing on the continuum of HIV prevention, care, and treatment, this review summarizes recent data on HIV burden among transgender women, describes the contexts that contribute to disproportionate vulnerability, and discusses opportunities for intervention.

HIV Burden

Baral and colleagues [4] published a meta-analysis of studies from 2000 - 2011 that assessed the burden of laboratory-confirmed HIV infection among transgender women worldwide. The pooled HIV prevalence was 19.1 % across the 15 countries with available data. Compared to all adults of reproductive age, transgender women had more than 48 times the odds of HIV infection. Data were only available from middle and high-income countries with male-predominant, concentrated epidemics. No studies were available from regions with generalized epidemics, such as sub-Saharan Africa [5]. Pooled prevalence can mask significant intra-country variability, hiding “hot spots” of extreme HIV burden. For example, while the pooled HIV prevalence among transgender women in the United States was 21.7% in the meta-analysis, a recently published study estimated HIV prevalence among transgender women in San Francisco to be a staggering 39.5% by laboratory testing [6].

HIV Vulnerability

Transgender women experience unique vulnerability to HIV that can be attributed to multi-level, intersecting factors that also influence all stages of the HIV treatment and care continuum. Transgender women who have sex with men often, but not exclusively, engage in receptive anal intercourse – an efficient route for acquisition of HIV infection [7]. High rates of sexually transmitted infections (STIs) have been reported among transgender women, including syphilis and herpes [7, 8], both ulcerative genital diseases that increase risk of HIV acquisition. Transgender women who have undergone genital reconstruction surgery may be

at additional risk for HIV through vaginal intercourse and trauma of the neovagina. However, data on HIV acquisition via the neovagina are lacking [9]. While biologic risks are important, they are not sufficient to explain the elevated risk of HIV among transgender women.

Transgender individuals around the world face high levels of stigma and discrimination, lack of social and legal recognition of their affirmed gender, and exclusion from employment and educational opportunities [7, 10, 11]. In several countries, transgender women can be arrested for “cross-dressing” or vagrancy and they may also be prosecuted under sodomy laws [12, 13]. Legal and economic marginalization increase transgender women’s vulnerability to HIV infection by limiting their employment options to sex work which in turn results in further stigmatization and disenfranchisement. For many transgender women, sex with men – including paid sex – can provide important validation of their gender identity [14, 15]. Sex work thus provides both desired gender affirmation and economic stability, often with greater financial rewards for sex without a condom [14, 16].

Experiences of gender-based discrimination, violence, victimization, rejection, and social marginalization are pervasive among transgender women worldwide [12, 17, 18]. Gender abuse in the form of verbal abuse, harassment, and/or physical abuse based on gender identity or presentation is a powerful risk factor for HIV among transgender women and often begins at an early age. A longitudinal study in New York City found that gender abuse was prospectively associated with sexual risk behavior and with incident HIV and STIs among young transgender women [19]. Lack of parental support has also been associated with HIV risk. One study among young transgender women in Los Angeles and Chicago found that sex without condoms was reported by those without parental support while consistent condom use was associated with having at least one supportive parent [20]. Brennan and colleagues [17] have shown that low self-esteem, polysubstance use, victimization related to transgender identity, and intimate partner violence were additively associated with HIV-infection and sexual risk.

The intersection of multiple stigmatized identities [21, 22] contributes to HIV vulnerability, and stigma toward people living with HIV is common in transgender and other marginalized communities [23]. Thus, transgender women living with HIV face stigma based on gender and HIV status, even within transgender communities. Intersecting stigma has important implications for engagement in care and treatment as well as for HIV prevention.

Continuum of HIV Care and Treatment

Transgender women may experience multiple barriers when attempting to access both routine and gender-related health services. Barriers include refusal of care, harassment and violence, and a lack of competent medical providers [10, 24, 25]. Faced with a lack of access to appropriate care, many transgender women utilize hormones without medical supervision [26]. Body modification may be undertaken with soft tissue fillers, usually industrial silicone. Among 335 transgender women in Peru who reported having undergone any gender enhancement procedure, 40% had injected industrial silicone[7]. Illicit hormone injections as well as injected fillers may carry HIV risk through the use of contaminated needles.

Barriers to healthcare are also associated with delayed access to urgent and preventive care, including HIV testing, care, and treatment. HIV screening rates are very low among transgender women, despite a high burden of infection [27], leaving many transgender women unaware of their HIV status. In national study of transgender people in Canada, Bauer and colleagues found that 46% of the transgender women had never been tested for HIV [28]. More than 50% of the kathoey sex workers interviewed by Nemoto and colleagues in Bangkok had never been tested [18]. Also, data suggest that few transgender women know the HIV status of their sexual partners [29]. Lack of inclusion in HIV awareness and prevention campaigns as well as competing priorities in the lives of transgender women may contribute to low rates of HIV testing [30, 31].

Once diagnosed with HIV, transgender women may continue to avoid medical care due to fear of, or actual experiences of discrimination in healthcare settings. A survey among transgender women living with HIV in the United States identified fear of disclosure of transgender identity, poor treatment by staff, and providers' lack of awareness of transgender issues, as important barriers to engagement in care [32]. Qualitative research among HIV-positive transgender women in Canada revealed similar themes [31].

Receiving antiretroviral therapy and achieving an undetectable viral load are important components of the HIV care and treatment continuum. Few studies have evaluated these components among transgender women. However, one study found that transgender women with HIV were less likely to have received antiretroviral therapy compared to non-transgender participants (59% vs. 82%) [33]. In another study, transgender women were less likely to report adherence to their antiretroviral regimen, more likely to have difficulty integrating treatment into their daily lives, and less likely to report positive interactions with HIV care providers relative to non-transgender participants [34]. Among HIV-positive women (including 28% transgender women) enrolled in a U.S.-based study, a history of recent abuse and violence was associated with a four-fold increase in likelihood of antiretroviral failure, underscoring the need to identify and address psychosocial issues, including trauma, in order to improve HIV-related outcomes [35].

In San Francisco, transgender women living with HIV have been found to have higher HIV-related mortality and a higher community viral load than non-transgender persons [36], consistent with both a lack of adequate HIV treatment and late presentation of disease. A recent retrospective analysis found similar rates for retention in care and virologic suppression for HIV-infected transgender women and nontransgender people attending 13 clinics in the USA [37]. This study included only clinics within and HIV research network, suggesting that these research-oriented clinics may have developed effective strategies for retention and adherence that could serve as models for other health facilities. No published studies were found outside the U.S. that evaluated engagement and retention in care in this population. This gap in the literature highlights the need for further research among transgender women living with HIV.

Opportunities

Scale-up of HIV interventions for transgender women must take place in the context of improved surveillance, monitoring, and evaluation so that health facilities, communities, and governments can be held accountable. It's promising to note that HIV studies in the United States and Asia have increasingly including transgender women. However, these studies often combine data from MSM and transgender women for sample size and other reasons. In order to better inform interventions for transgender women, they should be included in HIV research with sample sizes large enough for substantive analysis [9]; and results of combined studies should be disaggregated by gender-identity. Despite growing awareness of transgender identities, many current studies refer to transgender women as "transgender males," "transgender men," "males by nature but appearing as women." etc. [38] When collecting data with transgender populations, using language that is consistent with their gender identity may improve participation. Andraski et al. [39] recommend addressing barriers to research participation by transgender women through: (a) cultural competency training, (b) transgender-friendly environments, (c) true partnerships with local transgender-friendly organizations and health care providers, (d) protocols that focus on transgender-specific issues, and (e) data collection and tracking of transgender individuals.

Combination approaches that address biologic, psychosocial, and structural factors are most likely to have impact [40]. Important legal interventions include decriminalization of "cross-dressing" and "sodomy," as well as repeal of other laws used against transgender women such as anti-prostitution and vagrancy laws. Legal recognition of gender identity, prohibition of discrimination in education, employment, and housing, as well as other positive legislation will promote an enabling environment for transgender women to access HIV services.

Ensuring that transgender women have access to respectful and appropriate health care, including gender-affirming medical interventions can be accomplished both by training health care providers and integrating gender care into routine and HIV related medical care. Such interventions have the potential to improve rates of HIV testing as well as engagement and retention in care for transgender women with HIV. The HIV/AIDS Bureau of the Department of Health and Human Services in the USA has launched a new initiative to develop effective models of care for engaging and retaining transgender women of color in HIV care and treatment [41].

While data on stigma, discrimination, and violence among transgender women are growing, resilience among this population has been less studied. Resilience is a dynamic process of positive adaptation within the context of significant threat or adversity [42, 43]. Not all transgender women who have experienced adversity engage in HIV risk behaviors. For example, among 112 kathoey sex workers in Thailand, 73% consistently used condoms with clients and 65% were willing to have unprotected sex with clients for extra money[18]. Among non-transgender marginalized women with HIV, resilient coping skills have been found to mediate the relationship between HIV-stigma and depression, a factor associated with poor adherence [44]. Understanding not only risks/vulnerabilities, but also resilience among transgender women is critical to the development of effective interventions that are

relevant to and sustained by transgender communities. A strengths-based approach is especially important for transgender women, given the high levels of stigma and marginalization the community often faces. One example of such an intervention is Life Skills, a peer-led group intervention designed by and for young transgender women [45].

Empowerment has been defined as “a social process for people to gain mastery over their lives and the lives of their communities” (p. 294) [46]. It typically involves developing skills, knowledge, and confidence in their own abilities to overcome obstacles. Supporting existing transgender community structures and social networks that empower and mobilize community members is one way to support resilience and catalyze community-led structural change. In many parts of the world, transgender people have formed their own community and family structures. For example, Akhtar and colleagues [38] found that only four individuals in a sample of 306 hijras in Pakistan lived with their families of origin. The majority lived together as a clan in all-hijra communities with a guru (group leader). Similar alternative family structures have been documented in the U.S. with house mothers or fathers in house and ball communities [47].

Transgender individuals and groups are actively involved in creating websites to disseminate resources and information that target and reach other transgender people[48]. Across geographic regions and cultural contexts, transgender social networks have been shown to be efficient at engaging other transgender people [7, 28, 49]. Engaging with existing social networks may be a way to reach transgender people and deliver HIV prevention interventions that address the broader social context while being grounded in everyday lived realities.

Pre-Exposure Prophylaxis (PrEP) – the use of antiretroviral medication to reduce risk of HIV infection among HIV-negative people – can be highly effective among MSM and transgender populations [50]. A modeling study by Gomez and colleagues [51] in Peru suggests that PrEP strategies that prioritize transgender women and female sex workers can be expected to have higher impact for the same cost compared to prioritizing men who have sex with women or MSM.

The use of topical agents, microbicides, for the prevention of HIV infection has shown promise for prevention of vaginal HIV acquisition [52]. In recent years, there has also been progress in the development of microbicides for rectal use[53]. Focus groups and consultations among transgender people in both the Thailand and the United States [54, 55] suggest that, in addition to ensuring access, low cost and mitigating stigma and discrimination, education and outreach that acknowledge differences between MSM and transgender women are needed to support the effectiveness of rectal microbicide implementation.

Conclusion

The scientific literature on HIV among transgender women is growing, but large gaps remain. Stigma and discrimination remain formidable barriers to effective HIV prevention, care, and treatment. Researchers are beginning to examine the role of new biomedical

interventions, such as PrEP and microbicides, for HIV prevention in this population. They are also beginning to examine the factors that affect HIV testing and engagement in care and treatment. An increase in research and a scale-up of combination prevention programs tailored for transgender women will be necessary to effectively address the HIV epidemic among transgender women [56].

References

1. Herbst JH, Jacobs ED, Finlayson TJ, et al. Estimating HIV prevalence and risk behaviors of transgender persons in the United States: a systematic review. *AIDS Behav.* 2008 Jan; 12(1):1–17. Epub 2007/08/19. eng. [PubMed: 17694429]
2. Operario D, Soma T, Underhill K. Sex work and HIV status among transgender women: Systematic review and meta-analysis. *JAIDS Journal of Acquired Immune Deficiency Syndromes.* 2008; 48(1): 97–103. [PubMed: 18344875]
3. Baral S, Logie CH, Grosso A, et al. Modified social ecological model: a tool to guide the assessment of the risks and risk contexts of HIV epidemics. *BMC Public Health.* 2013; 13:482. Epub 2013/05/18. eng. [PubMed: 23679953]
4. Baral SD, Poteat T, Stromdahl S, et al. Worldwide burden of HIV in transgender women: a systematic review and meta-analysis. *Lancet Infect Dis.* 2013 Mar; 13(3):214–22. Epub 2012/12/25. *This provides the first global view of the burden of laboratory confirmed HIV among transgender women world-wide. It highlights the enormous disparity in HIV prevalence compared to the general population and points out the gaps in data from Africa and Eastern Europe. [PubMed: 23260128]
5. Jobson GA, Theron LB, Kaggwa JK, et al. Transgender in Africa: Invisible, inaccessible, or ignored? *SAHARA-J: Journal of Social Aspects of HIV/AIDS.* 2012; 9(3):160–3. 2012/09/01. [PubMed: 23237071]
6. Rapues J, Wilson EC, Packer T, et al. Correlates of HIV Infection Among Transfemales, San Francisco, 2010: Results From a Respondent-Driven Sampling Study. *American Journal of Public Health.* 2013; 103(8):1485–92. [PubMed: 23763398]
7. Silva-Santisteban A, Raymond HF, Salazar X, et al. Understanding the HIV/AIDS epidemic in transgender women of Lima, Peru: results from a sero-epidemiologic study using respondent driven sampling. *AIDS Behav.* 2012 May; 16(4):872–81. Epub 2011/10/11. eng. [PubMed: 21983694]
8. Guadamuz TE, Wimonasate W, Varangrat A, et al. HIV Prevalence, Risk Behavior, Hormone Use and Surgical History Among Transgender Persons in Thailand. *AIDS Behav.* 2010 Nov 20. Epub 2010/11/26 Eng.
9. van Griensven F, Na Ayutthaya PP, Wilson E. HIV surveillance and prevention in transgender women. *Lancet Infect Dis.* 2013 Mar; 13(3):185–6. Epub 2012/12/25. eng. [PubMed: 23260127]
10. Grant, JM., Mottet, LA., Tanis, J., et al. Injustice at Every Turn: A Report of the National Transgender Discrimination Survey. Washington: National Center for Transgender Equality and National Gay and Lesbian Task Force; 2011.
11. REDLACTRANS. The Night is Another country: Impunity and violence against transgender women human rights defenders in Latin America. Buenos Aires: Red Latinoamericana y del Caribe de Personas Trans, HIV/AIDS Alliance; 2012.
12. Winter, S. Lost in Transition: Transgender People, Rights and HIV Vulnerability in the Asia-Pacific Region. Bangkok: UNDP Asia Pacific Regional Center; 2012.
13. Baral, S., Beyrer, C., Poteat, T. Human Rights, the Law, and HIV among Transgender People; Working Paper prepared for the Third Meeting of the Technical Advisory Group of the Global Commission on HIV and the Law; 7-9 July 2011; 2011.
14. Melendez RM, Pinto R. 'It's really a hard life': love, gender and HIV risk among male-to-female transgender persons. *Cult Health Sex.* 2007 May-Jun; 9(3):233–45. Epub 2007/04/26. eng. [PubMed: 17457728]
15. Sevelius JM. Gender Affirmation: A Framework for Conceptualizing Risk Behavior among Transgender Women of Color. *Sex Roles.* 2013 Jun 1; 68(11-12):675–89. Epub 2013/06/05. *This

paper draws a detailed conceptual framework for HIV risk among transgender women that can be used to develop and test HIV prevention interventions. [PubMed: 23729971]

16. Sausa LA, Keatley J, Operario D. Perceived risks and benefits of sex work among transgender women of color in San Francisco. *Arch Sex Behav*. 2007 Dec; 36(6):768–77. Epub 2007/08/04. eng. [PubMed: 17674180]
17. Brennan J, Kuhns LM, Johnson AK, et al. Syndemic theory and HIV-related risk among young transgender women: the role of multiple, co-occurring health problems and social marginalization. *Am J Public Health*. 2012 Sep; 102(9):1751–7. Epub 2012/08/10. eng. [PubMed: 22873480]
18. Nemoto T, Iwamoto M, Perngparn U, et al. HIV-related risk behaviors among kathoey (male-to-female transgender) sex workers in Bangkok, Thailand. *AIDS Care*. 2012; 24(2):210–9. Epub 2011/07/26. eng. [PubMed: 21780964]
19. Nuttbrock L, Bockting W, Rosenblum A, et al. Gender abuse, depressive symptoms, and HIV and other sexually transmitted infections among male-to-female transgender persons: a three-year prospective study. *Am J Public Health*. 2013 Feb; 103(2):300–7. *This is the first longitudinal study of HIV and STIs among transgender women to appear in the published literature. It provides evidence that such studies are feasible as well as providing valuable information on predictors of HIV vulnerability. [PubMed: 22698023]
20. Wilson EC, Iverson E, Garofalo R, et al. Parental support and condom use among transgender female youth. *J Assoc Nurses AIDS Care*. 2012 Jul-Aug;23(4):306–17. Epub 2011/11/15. eng. [PubMed: 22079675]
21. Bowleg L. The problem with the phrase women and minorities: intersectionality-an important theoretical framework for public health. *Am J Public Health*. 2012 Jul; 102(7):1267–73. Epub 2012/05/19. eng. [PubMed: 22594719]
22. Logie CH, James L, Tharao W, et al. HIV, Gender, Race, Sexual Orientation, and Sex Work: A Qualitative Study of Intersectional Stigma Experienced by HIV-Positive Women in Ontario, Canada. *PLoS Med*. 2011; 8(11):e1001124. [PubMed: 22131907]
23. Ramirez-Valles J, Molina Y, Dirkes J. Stigma towards PLWHA: the role of internalized homosexual stigma in Latino gay/bisexual male and transgender communities. *AIDS Educ Prev*. 2013 Jun; 25(3):179–89. Epub 2013/05/02. eng. [PubMed: 23631713]
24. Lambda Legal. When Health Care Isn't Caring: Lambda Legal's Survey of Discrimination Against LGBT People and People with HIV. New York: Lambda Legal; 2010.
25. Obedin-Maliver J, Goldsmith ES, Stewart L, et al. Lesbian, gay, bisexual, and transgender-related content in undergraduate medical education. *JAMA*. 2011 Sep 7; 306(9):971–7. Epub 2011/09/09. eng. [PubMed: 21900137]
26. Rotondi NK, Bauer GR, Scanlon K, et al. Nonprescribed Hormone Use and Self-Performed Surgeries: "Do-It-Yourself" Transitions in Transgender Communities in Ontario, Canada. *Am J Public Health*. 2013 Oct; 103(10):1830–6. Epub 2013/08/21. Eng. [PubMed: 23948009]
27. Beattie TS, Bhattacharjee P, Suresh M, et al. Personal, interpersonal and structural challenges to accessing HIV testing, treatment and care services among female sex workers, men who have sex with men and transgenders in Karnataka state, South India. *J Epidemiol Community Health*. 2012 Oct; 66(Suppl 2):ii42–8. Epub 2012/04/13. eng. [PubMed: 22495772]
28. Bauer G, Travers R, Scanlon K, et al. High heterogeneity of HIV-related sexual risk among transgender people in Ontario, Canada: a province-wide respondent-driven sampling survey. *BMC Public Health*. 2012; 12(1):292. [PubMed: 22520027]
29. Nagaraj S, Segura ER, Peinado J, et al. A cross-sectional study of knowledge of sex partner serostatus among high-risk Peruvian men who have sex with men and transgender women: implications for HIV prevention. *BMC Public Health*. 2013; 13:181. Epub 2013/03/02. eng. [PubMed: 23448153]
30. Altaf A, Zahidie A, Agha A. Comparing risk factors of HIV among hijra sex workers in Larkana and other cities of Pakistan: an analytical cross sectional study. *BMC Public Health*. 2012; 12:279. Epub 2012/04/12. eng. [PubMed: 22490045]
31. Logie CH, James L, Tharao W, et al. "We don't exist": a qualitative study of marginalization experienced by HIV-positive lesbian, bisexual, queer and transgender women in Toronto, Canada. *J Int AIDS Soc*. 2012; 15(2):17392. Epub 2012/09/20. eng. [PubMed: 22989529]

32. Houston Area Ryan White Planning Council. Access to HIV Care among transgender and gender non-conforming people in Houston 2013. [cited 2013 October 1] Available from: <http://www.rwpchouston.org/Publications/TG%20Special%20Study%20Report%20APPROVED%2003-14-13.pdf>
33. Melendez RM, Exner TA, Ehrhardt AA, et al. Health and health care among male-to-female transgender persons who are HIV positive. *Am J Public Health*. 2006 Jun; 96(6):1034–7. Epub 2005/09/01. eng. [PubMed: 16131645]
34. Sevelius JM, Carrico A, Johnson MO. Antiretroviral therapy adherence among transgender women living with HIV. *J Assoc Nurses AIDS Care*. 2010 May-Jun;21(3):256–64. Epub 2010/03/30. eng. [PubMed: 20347342]
35. Machtinger EL, Haberer JE, Wilson TC, et al. Recent trauma is associated with antiretroviral failure and HIV transmission risk behavior among HIV-positive women and female-identified transgenders. *AIDS Behav*. 2012 Nov; 16(8):2160–70. Epub 2012/03/20. eng. [PubMed: 22426597]
36. Das M, Chu PL, Santos G-M, et al. Decreases in Community Viral Load Are Accompanied by Reductions in New HIV Infections in San Francisco. *PLoS ONE*. 2010; 5(6):e11068. *This is the first multi-site study to examine retention and virologic suppression among transgender women compared to other people living with HIV. [PubMed: 20548786]
37. Yehia BR, Fleishman JA, Moore RD, et al. Retention in Care and Health Outcomes of Transgender Persons Living With HIV. *Clin Infect Dis*. 2013 Jun 23. Epub 2013/06/01 Eng.
38. Akhtar H, Badshah Y, Akhtar S, et al. Prevalence of human immunodeficiency virus infection among transgender men in Rawalpindi (Pakistan). *Viol J*. 2012; 9:229. Epub 2012/10/09. eng. [PubMed: 23039269]
39. Andrasik MP, Yoon R, Mooney J, et al. Exploring Barriers and Facilitators to Participation of Male-to-Female Transgender Persons in Preventive HIV Vaccine Clinical Trials. *Prev Sci*. 2013 Feb 28. Epub 2013/03/01 Eng.
40. Auerbach JD, Parkhurst JO, Caceres CF. Addressing social drivers of HIV/AIDS for the long-term response: conceptual and methodological considerations. *Glob Public Health*. 2011; 6(Suppl 3):S293–309. Epub 2011/07/13. eng. [PubMed: 21745027]
41. Health Resources and Services Administration. What's Going On @ SPNS April 2013. [cited 2013 5 October 2013] Available from: <http://webcache.googleusercontent.com/search?q=cache:http://hab.hrsa.gov/about/hab/files/cyberspns042013.pdf>.
42. Luthar SS, Brown PJ. Maximizing resilience through diverse levels of inquiry: Prevailing paradigms, possibilities, and priorities for the future. *Dev Psychopathol*. 2007 Summer;19(3):931–55. Epub 2007/08/21. eng. [PubMed: 17705909]
43. Rutter M. Psychosocial resilience and protective mechanisms. *Am J Orthopsychiatry*. 1987 Jul; 57(3):316–31. Epub 1987/07/01. eng. [PubMed: 3303954]
44. Logie C, James L, Tharao W, et al. Associations between HIV-related stigma, racial discrimination, gender discrimination, and depression among HIV-positive African, Caribbean, and Black women in Ontario, Canada. *AIDS Patient Care STDS*. 2013 Feb; 27(2):114–22. Epub 2013/02/05. eng. [PubMed: 23373665]
45. Garofalo R, Johnson AK, Kuhns LM, et al. Life skills: evaluation of a theory-driven behavioral HIV prevention intervention for young transgender women. *J Urban Health*. 2012 Jun; 89(3):419–31. Epub 2012/01/10. eng. *This study provides a rare evaluation of a theory-based intervention specifically for transgender women. It can serve as a model for developing evidence-based interventions for this population. [PubMed: 22223033]
46. Glanz, K., Rimer, BK., Viswanath, K. Health behavior and health education: theory, research, and practice. John Wiley & Sons; 2008.
47. Kubicek K, Beyer WH, McNeeley M, et al. Community-engaged research to identify house parent perspectives on support and risk within the house and ball scene. *J Sex Res*. 2013; 50(2):178–89. Epub 2011/12/31. eng. [PubMed: 22206442]
48. Horvath KJ, Iantaffi A, Grey JA, et al. A Review of the Content and Format of Transgender-Related Webpages. *Health Communication*. 2012; 27(5):457–66. Epub 2011 Oct 18. [PubMed: 22007970]

49. Barrington C, Wejnert C, Guardado ME, et al. Social network characteristics and HIV vulnerability among transgender persons in San Salvador: identifying opportunities for HIV prevention strategies. *AIDS Behav.* 2012 Jan; 16(1):214–24. Epub 2011/05/04. eng. [PubMed: 21538082]
50. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med.* 2010 Dec 30; 363(27):2587–99. Epub 2010/11/26. eng. [PubMed: 21091279]
51. Gomez GB, Borquez A, Caceres CF, et al. The potential impact of pre-exposure prophylaxis for HIV prevention among men who have sex with men and transwomen in Lima, Peru: a mathematical modelling study. *PLoS Med.* 2012; 9(10):e1001323. Epub 2012/10/12. eng. [PubMed: 23055836]
52. Friend DR, Kiser PF. Assessment of topical microbicides to prevent HIV-1 transmission: Concepts, testing, lessons learned. *Antiviral Res.* 2013 Sep; 99(3):391–400. Epub 2013/07/13. eng. [PubMed: 23845918]
53. McGowan I, Dezzutti C. Rectal Microbicide Development. *Curr Top Microbiol Immunol.* 2013 Apr 24. Epub 2013/04/25 Eng.
54. Rectal Microbicides for HIV Prevention. A Transgender Update and Consultation, May 19, 2013. San Francisco: 2013.
55. Newman PA, Roungrakphon S, Tepjan S. A social ecology of rectal microbicide acceptability among young men who have sex with men and transgender women in Thailand. *J Int AIDS Soc.* 2013; 16(1):18476. Epub 2013/08/06. eng. [PubMed: 23911116]
56. Subramanian T, Ramakrishnan L, Aridoss S, et al. Increasing condom use and declining STI prevalence in high-risk MSM and TGs: evaluation of a large-scale prevention program in Tamil Nadu, India. *BMC Public Health.* 2013 Sep 17.13(1):857. Epub 2013/09/21 Eng. [PubMed: 24044766]

Key points

- Transgender women experience unique vulnerability to HIV that can be attributed to structural, psychosocial, and biologic factors that also influence the HIV treatment and care continuum.
- Stigma and discrimination, lack of social and legal recognition of their affirmed gender, and exclusion from employment and educational opportunities represent fundamental drivers of HIV risk in transgender women.
- Barriers to health care access, including stigma, discrimination, and lack of health care providers competent in gender-related care, contribute to low HIV testing rates as well as limited engagement in care.
- Interventions to improve engagement in HIV prevention, testing, care, and treatment among transgender women should build on community strengths and address structural factors as well as psychosocial and biologic factors that increase HIV vulnerability and prevent access to HIV services.