Published in final edited form as:

Am J Clin Dermatol. 2018 June ; 19(3): 293–302. doi:10.1007/s40257-017-0340-7.

The Role and Diagnosis of Allergic Contact Dermatitis in Patients with Atopic Dermatitis

Joshua L. Owen, MD, PhD^{a,*}, Paras P. Vakharia, PharmD^{a,*}, and Jonathan I. Silverberg, MD, PhD, MPH^{b,c}

^aDepartment of Dermatology, Northwestern University Feinberg School of Medicine, Chicago, IL, 60611, USA

^bDepartments of Dermatology, Preventive Medicine and Medical Social Sciences, Northwestern University Feinberg School of Medicine, Chicago, IL 60611

^cNorthwestern Medicine Multidisciplinary Eczema Center

Abstract

Patients with atopic dermatitis (AD) have increased penetration of allergens, immune dysregulation including shared cytokine pathways, frequent use of emollients and topical medications, all of which may predispose toward developing allergic contact dermatitis (ACD). Recent systematic reviews have suggested that ACD is a significant clinical problem in both children and adults with AD. While this remains controversial, ACD remains an important comorbidity and potential exacerbant of AD in clinical practice. Common relevant allergens, include lanolin, neomycin, formaldehyde, sesquiterpine lactone mix, compositae mix, and fragrances that are commonly found in AD patients' personal care products. We herein review the clinical scenarios where patch testing is indicated in AD. In addition, we review the contraindications, preferred patch testing series, pitfalls, and challenges determining the relevance of positive patch test reactions in AD patients.

1 Introduction

1.1 Atopic dermatitis (AD)

AD and allergic contact dermatitis (ACD) are both common and burdensome inflammatory skin disorders. AD is a chronic disease that is caused by a combination of genetic predisposition, skin-barrier disruption, immune factors, and environmental exposures. AD affects up to 15–20% of children and 1–10% of adults worldwide, including 13% of US children and 7.2% of US adults.[1–4] AD is a heterogeneous disorder associated with a constellation of signs and symptoms, including pruritus, skin pain [5], mental health symptoms, xerosis, oozing/weeping in acute lesions, lichenification and prurigo nodules in

Corresponding author and reprint requests: Jonathan I. Silverberg, MD, PhD, MPH, 676 N. St. Clair St, Suite 1600, Chicago, IL 60611, Phone: 312-503-4985, Fax: 312-695-0537, JonathanISilverberg@Gmail.com.

*These authors contributed equally to this manuscript.

Compliance with Ethical Standards

J Owen, P Vakharia and JI Silverberg have no relevant conflicts of interest to declare.

chronic lesions. AD also has a chronic relapsing or persistent course, with an age-related distribution of cutaneous lesions, with facial dermatitis affecting infants, extensor dermatitis in toddlers, flexural lesions in older children and adults, and more facial and hand dermatitis in adults.[6]

Given the varied presentation of AD, it is often challenging to diagnose with certainty, particularly in adults. AD is diagnosed clinically based on the signs and symptoms. There are no currently accepted tissue or blood biomarkers to diagnose AD. The original diagnostic criteria for AD are those of Hanifin and Rajka (HR),[7] published in 1980 and developed via clinical experience and expert consensus. However, various modifications of these criteria, as well as other diagnostic criteria were subsequently developed.[8–12]

1.2 ACD

ACD is caused by a delayed-type hypersensitivity response to contact allergens. The incidence of ACD is not clearly defined, but is thought to be rising.[13] A recent study found that all forms of contact dermatitis (CD), including irritant contact dermatitis (ICD) and ACD, had a claims-based prevalence of 4.17% within the US.[14] The most recent estimated annual medical costs in the US in 2013 for AD and CD is \$314,000,000 and \$1,529,000,000, respectively.[14] ACD is the second most common type of CD after irritant contact dermatitis (ICD), and may present with similar signs and symptoms to AD. The most common symptoms are pruritus, along with burning and stinging. ACD commonly presents acutely with erythematous, indurated papules and plaques, vesiculation, edema and bullae formation in severe cases. Chronic ACD can present with scaling, lichenification, and fissuring. ACD typically presents with a well-defined, exposure-dependent distribution, commonly involving the hands, face, or eyelids. However, irregular or diffuse distributions can occur due to secondary allergen transfer or systemic allergen sensitization. ACD is diagnosed via a combination of clinical signs and symptoms and patch testing, the gold standard for ACD diagnosis, where non-irritating concentrations of allergens are used to determine the presence of an allergic reaction in vivo.[15]

Importantly, due to the large symptom-burden, including the sequela of itch, pain, sleep, and mental health disturbance, AD and ACD both have a significant negative impact on quality of life (QOL).[16–20] While these two cutaneous eruptions may appear similar and often coexist,[21] the etiologies, distributions, and therapeutic options often differ. This makes differentiating the two diseases critical to the successful treatment of the dermatitis. The goal of this article is to review ACD in AD patients: when to suspect ACD and how best to test for ACD in patients with AD.

2 Mechanisms

2.1 AD

The pathogenesis of AD is multifactorial, with both epidermal barrier and immunologic defects. A subset of AD patients have filaggrin (FLG) gene null mutations that are inherited in an autosomal semi-dominant fashion. Barrier disruption may occur secondary to exogenous insults, even in those without germline FLG mutations, possibly through direct

insult to the skin-barrier and/or epigenetic alterations.[22, 23] Such factors include fragrances, pruritogens, stress, climate, pollution, among others. Thymic stromal lymphopoietin (TSLP) and other cytokines are released by damaged keratinocytes from the disrupted skin-barrier and contribute to skin inflammation, and may also be involved in gene-environment interactions in AD.[23] Due to impaired barrier function, there is an increased risk of transcutaneous allergen penetrance[24] and potentially antigen sensitization and presentation.

The AD inflammatory signature is primarily of the T-helper (Th) cell 2 type in both the acute and chronic phases, with a contribution from Th1 in the chronic phase.[25–27] Th2 cells produce interleukins (IL-) 4, 5, 13, and 31, all of which have downstream effects in AD. Notably, IL-4 and IL-13 promote skin barrier disruption. Thus, epidermal inflammation may precede and be sufficient to cause skin-barrier disruption, even in those without preceding barrier defects. Some studies have shown upregulation of IL-17 and IL-22 (secreted by Th17 and Th22 cells, respectively) in the acute phase of AD. These cytokines may induce epidermal hyperplasia and/or alter terminal differentiation proteins. Th2 cytokines also impair antimicrobial peptide (AMP) responses to pathogens, which, in conjunction with barrier disruption, allows for increased pathogen penetration.[28–35]

Recent studies have also shown a potential role for Th9 and Th17 pathways in AD. The mechanism by which IL-9, secreted by Th9 cells, contributes to AD pathogenesis is not fully known. However, IL-9 promotes mast cell activity, eosinophils, and innate immune cells. [36] IL-9 levels have been shown to be increased in both pediatric and adult AD patients and correlate with AD severity.[31, 37, 38] IL-9 also enhances the secretion of IL-13, a key cytokine in AD pathogenesis. Importantly, a significant association between IL-9 and IL-9 receptor gene polymorphisms with atopic dermatitis was found in a Korean population.[39] Th17 levels have also been found to correlate with AD severity,[32] and may be related to host-defense and skin remodeling.[40, 41] Th17 cytokines may play a greater role in intrinsic AD, i.e. AD without comorbid atopy or atopic disease.[35]

2.2 ACD

ACD is a classic type IV hypersensitivity reaction requiring two phases: sensitization and elicitation. In the sensitization phase, an allergen is captured by antigen-presenting cells (APCs), which migrate to the draining lymphoid tissue. Subsequent activation of naïve T cells occurs, leading to differentiation of memory T cells specific for that allergen. In the elicitation phase, re-exposure to the allergen or a cross-reacting allergen results in activation of memory T cells. T cytotoxic (Tc) 1 cells are activated and lead to the hallmark inflammation and adaptive immune response resulting in dermatitis.[42] The primary ACD inflammatory signature is a T cytotoxic (Tc) 1 or Th1 response. However, Th2, Th17, and Th22 responses appear to play a role in ACD, sometimes depending on the allergen.[43–45] For example, nickel was found to be a potent inducer of the innate immune Th1, Th17, and Th22 pathways, while fragrance and rubber promoted Th2 activity with less Th1 and Th17 involvement.[46]

IL-9 expression has also been found to be elevated in skin from positive patch-test reactions in ACD patients, including reactions to metals, drugs, and polymers; IL-9 also increased in

nickel-allergic patients after nickel stimulation.[47–49] Th17 cell expansion occurs upon allergen contact in individuals with ACD.[50] IL-17 secretion increases local inflammation via induction of proinflammatory cytokines, chemokines, and adhesion molecules.[51–54] The potential role of Th17 in ACD was also demonstrated by a recent experimental study showing that ACD reactions were decreased in the absence of IL-17.[55]

3 Allergic Contact Dermatitis in Patients with Atopic Dermatitis

3.1 Plausibility

Many factors are thought to affect prevalence of ACD in patients with AD. The historical perspective is that the Th2-skewed inflammatory response of AD results in less contact sensitivity.[60] For example, some studies have shown an increased elicitation threshold in patients with AD compared to controls.[60–63] Other studies have demonstrated several reasons for AD patients to have similar or even increased risk of ACD compared to those without AD. Patients with AD have skin-barrier disruption, with an approximately two-fold increase in skin absorption of irritants and contact allergens.[64–66] Irritants lead to further breakdown of the skin barrier, increased penetration of contact allergens, and eventually increased risk of contact sensitization.[67, 68] It has also been demonstrated that cutaneous responses and elicitation thresholds in ACD patients were considerably influenced by combined allergen and irritant exposure.[69–71] Additionally, the treatment of AD requires chronic topical application of emollients and anti-inflammatories, and many of these topical products have been found to be contact sensitizers.[72, 73] More recently, potential shared immune pathways have been demonstrated for subsets of AD and ACD, including Th1, Th2, Th9 and/or Th17, as reviewed above. An emerging idea is the role of bacterial colonization in AD and how, by stimulating an inflammatory environment, may lead to enhanced contact sensitization.[67, 74–76]

3.2 Evidence

A recent systematic review was performed assessing contact allergy in children with AD. The review assessed 31 studies and found that ACD was significantly greater in children without AD vs those with AD (46.6% and 41.7% sensitized to at least one allergen, respectively; I-squared=61.7%, $P<0.001$). However, the authors noted significant variability of sensitization rates, study designs and criteria that limit conclusions being drawn. The results of the available studies were conflicting with respect to whether AD patients have higher rates of ACD than the rest of the population. Nevertheless, ACD was found to be a common clinical problem in AD, with approximately one-third of children with AD that were patch-tested having at least one contact allergy.[77]

Another systematic review and meta-analysis, including 74 studies evaluating the prevalence of contact sensitization (defined as a positive patch test reaction to any allergen) in various patient populations found that AD patients had increased prevalence of contact sensitization compared to the general population.[78] However, there was an inverse association when patients with AD were compared to a patch-test referral population. The authors postulated that this latter relationship could be because AD patients in a referral population have more severe and recalcitrant disease, which has been shown to have a higher elicitation threshold

for contact sensitization.[60–62] Further, severe AD patients are often referred for patch testing to rule out contact sensitization even without clear clinical suspicion prior to initiating systemic AD therapy.[78]

3.3 Relevant allergens

Results from various studies assessing the relationship of ACD in AD patients have led to the identification of common allergens (Table 1), including nickel, cobalt, potassium dichromate, chromium, lanolin, neomycin, formaldehyde, sesquiterpine lactone mix, compositae mix, and fragrance markers (e.g. fragrance mix I, fragrance mix II, Myroxylon pereirae, and hydroxyisohexyl-3-cyclohexenecarboxaldehyde).[73, 77–90]

It has been recently demonstrated that commonly used personal care products, including those self-identified as hypoallergenic, contain potent contact allergens.[72, 91] Furthermore, AD patients with frequent emollient use were found to have increased urinary levels of such allergens, particularly parabens and phthalate metabolites, indicating that such allergens do have cutaneous penetrance.[92] Topical treatment with emollients in AD has been shown to be associated with cutaneous sensitization.[73] A retrospective Dutch study of pediatric patients found that children with AD had significantly increased reactivity to lanolin and fragrances.[87] Further, a retrospective analysis of 26,479 patients patch tested with the NACDG screening series found that patients with positive reactions to lanolin were more likely to have a history of AD.[93]

4 Clinical Assessment for Contact Dermatitis in Patients with Atopic Dermatitis

4.1 When to Consider Patch Testing in a Patient with AD

Guidelines for when to perform patch testing in AD patients are based largely upon consensus expert opinion.[94] Recommendations for when to consider patch testing include: adolescent- or adult-onset AD, as ACD can occasionally present with a flexural distribution and mimic AD. Pediatric and adult AD patients with worsening or more generalized dermatitis should also be patch tested, as there may be an allergenic trigger of their underlying AD.

Patch testing is also indicated in both children and adults when there is a lesional distribution that is atypical for AD, or one that is localized and suggestive of contact dermatitis (e.g., eyelids, head and neck, hand and foot, perioral, or periorbital). This is a particularly important consideration in adults with AD, for whom previous studies have demonstrated higher rates of lesions affecting the head and neck, or hands and feet (even in the absence of contact dermatitis).[95]

Patch testing should be considered in both children and adults if the dermatitis is recalcitrant to topical therapy and prior to initiation of systemic immunosuppressive therapy. Identification and avoidance of a relevant positive allergen on patch testing may decrease the severity of the underlying AD and abrogate the need for systemic therapy.

Patch testing should be considered in children and adults when the AD worsens with therapy, or rebounds quickly upon cessation of therapy. This may signal that the patient has developed ACD to the active ingredients or excipients in their topical therapy, e.g. corticosteroids or propylene glycol.

In addition, previous studies have shown high rates of ACD in patients with nummular eczema. Nummular lesions have been shown to occur with greater frequency in school age children with AD [96] and adult-onset AD [97]. However, widespread nummular lesions may be a sign of ACD in an AD patient.[98, 99]

4.2 When is Patch Testing Not Routinely Recommended in AD

Situations in which patch testing is less likely to be helpful include stable and well-controlled AD, AD flare and/or active dermatitis involving the back and other potential sites of application for the patch tests, current or recent use of systemic immunosuppressive medications, recent exposure to ultraviolet therapy or excessive solar radiation, and use of a limited patch testing series that do not incorporate the full-spectrum of allergens previously shown to be relevant in AD.[94]

A commonly encountered clinical situation is the patient with active, often severe, dermatitis on the back and other potential sites of application for the patch tests. This scenario should delay and may even prevent patch testing. Patch testing on actively inflamed skin may lead to both false positive and false negative reactions. The patient may also experience immense discomfort secondary to pruritus and pain from the adhesives used, increased heat and sweat and exposure to potentially irritating reagents being tested. In addition, the term “angry back syndrome” has been used to describe when patients develop positive reactions to most or all allergens tested.

Efforts should be made to first treat and resolve the active dermatitis on the back and other potential sites of application for the patch tests. Ideally, this should be done using topical therapy, e.g. corticosteroids and calcineurin inhibitors. If successful, the patient should discontinue application of topical therapy to the back for 1–2 weeks and then undergo patch testing. Systemic therapy or phototherapy may be required if the patient has an inadequate response to topical therapy or immediately experiences a flare of their dermatitis. However, such therapies may decrease the sensitivity of the patch testing process.

There is insufficient experimental data to precisely define the extent to which each immunosuppressive medication decreases the sensitivity threshold of patch testing. An expert consensus opinion from the North American Contact Dermatitis Group[100] suggested that the following medications were at high risk for leading to false negative patch test results: prednisone >10 mg/d and intramuscular triamcinolone (avoid for 4 weeks), topical corticosteroids or calcineurin inhibitors at patch test application sites (avoid for 1 week), azathioprine, cyclosporine, mycophenolate mofetil, and systemic tacrolimus. There was not enough data for the panel to make specific avoidance period recommendations for the non-corticosteroid immunosuppressants, other than to say that their effect on the results of patch testing are dose-dependent. Ultraviolet exposure to the testing site was recommended to be avoided for 1 week prior to testing. [100] The following medications

were considered generally acceptable for patients to be taking during patch testing: methotrexate, prednisone <10 mg/d, tumor necrosis factor α inhibitors, ustekinumab, and antihistamines. Another expert consensus opinion echoed these suggestions, but noted the lack of information regarding immunosuppressive agent effects on patch test reactions. [101] There is no consensus regarding avoidance of newer agents being used in the treatment of AD, including crisaborole, Janus kinase inhibitors, or dupilumab.

In the authors personal experience, many patients experience false negative reactions to patch testing up to 4 weeks (or longer) after intense ultraviolet radiation, e.g. sunny vacation, cyclosporine at a dose of >2.0 mg/kg/day, methotrexate at a dose of >0.20 mg/kg/week. If patch testing is performed in these scenarios, results should be interpreted with caution. Weak or irritant reactions should be considered as true positives. Negative patch tests should be considered as false negatives and repeat patch testing should be considered upon discontinuation and washout from such treatments.

5 Patch Testing in Patients with AD

5.1 Choosing the Right Patch Testing Series

Once the decision has been made to perform patch testing, allergen selection is critical for a satisfactory outcome and should be made on an individual patient basis. Factors to be considered during allergen selection include the region or country, occupation, hobbies and recreations, and other exposures. One option is the Thin-Layer Rapid Use Epicutaneous (T.R.U.E.) test; however, it should be noted that this test lacks multiple allergens that are commonly relevant and present on expanded patch testing series, e.g. American Contact Dermatitis Society (ACDS) or NACDG core series. Examples include cinnamic aldehyde, propylene glycol, dimethylol dimethyl hydantoin, iodopropynyl butyl carbamate, amidoamine, acrylates, tea tree oil, propolis, benzophenone-3, and sesquiterpene lactone mix.

While evidence-based guidelines are lacking for which allergens to include for patch testing in AD patients, several recommendations have been made by different authors. In summary, the majority of studies recommend expanded screening for the most commonly encountered allergens in AD patients (e.g. metals such as nickel, potassium dichromate, carba mix, formaldehyde, neomycin sulfate, balsam of Peru, fragrances, and preservatives), allergens that are common components of over-the-counter and prescription topical therapies, and allergens specific to a patient's environment (i.e. patient's personal care products or occupational exposures). For adults in North America, an expanded screening series, such as the American Contact Dermatitis Society or North American Contact Dermatitis Group core series, appear to be reasonable. Screening series may vary regionally based on the most prevalent allergens. More targeted patch testing can be considered in younger children,[73, 94, 102–105] but standardized screening series are not well-established. Patch test reads should be performed at 48 and 72 hours and preferably a delayed read between 96 and 144 hours.

All personal care products and topical medications should be inspected for possible allergens. An important limitation is the U.S. Food and Drug Administration's Cosmetic

Labeling Guide regulations. Although all ingredients, including those with less than 1% concentration, are supposed to be listed on product labels, there are numerous ways around this. “Incidental Ingredients” (ingredients present at an insignificant level and having no technical or functional effect) and/or a “Trade Secret Ingredient” (an ingredient that offers one’s business potential to obtain an advantage over those not using or knowing about it) are exempt from ingredient declaration, and the phrase “and other ingredients” may be used in place.[106] Thus, there may be additional unknown exposures to lower concentrations of allergens. Leave on products can be patch-tested as they are formulated, but may be subject to false-negative reactions. However, rinse-off products should be diluted given their high potential for irritancy.[101]

5.2 Pitfalls and Determining the Relevance of Positive Patch Test Reactions in Patients with AD

There are several potential pitfalls to be considered when patch testing patients with AD. As mentioned previously, patients with AD have a lower irritancy threshold. This may lead to higher rates of irritant or false positive reactions, the most common of which are metals, fragrance, formaldehyde, and lanolin.[67, 103, 107] Additionally, as mentioned, patch test reactions should be interpreted with caution in patients on specific immunosuppressants. Weak or irritant reactions should be considered as true positives, while negative patch tests should be considered as possible false negatives.

On the other hand, irritant reactions may also be more difficult to distinguish from true positive reactions. That is, some relevant positive reactions in AD patients may display as weaker reactions that would be mistaken as irritant reactions, i.e. a negative patch test. One reason for this is that patients with AD are less likely to exhibit the “crescendo” pattern of increasing reactivity between patch test reads seen in true positive reactions.[103] Another reason is that positive reactions may be weaker in patients with AD, especially with increasing severity of disease, as they may be less effective at acquiring sensitization.[60, 108] Delayed reads of >96 hours may be somewhat helpful to overcome this.[109–111] Some patients may benefit from an empiric trial of allergen avoidance despite only displaying an irritant or weak positive reaction. In addition, if patch testing was performed but only displayed no or irritant reactions in a patient with a compelling history and/or physical exam for ACD, then false negatives should be contemplated. In such patients, particularly those with uncontrolled dermatitis during patch testing, repeat patch testing should be considered at a later date and may successfully identify relevant positive reactions, despite false negative or weak reactions upon initial patch testing.

Finally, active AD may paradoxically result in false negative reactions on patch testing. The risk of false negatives was found to be higher with increasing severity of AD; this may be true even when the patches are applied to apparently non-lesional skin and patients are not on systemic immunosuppressive medications.[94, 101] One experimental report demonstrated that well-controlled AD (less than 10% body surface for at least one month) is associated with lower rates of false negative reactions.[60] Taken together, the results of patch testing in AD patients should be interpreted with caution.

6 Conclusion

The risk of ACD appears to be increased in patients with AD, though this association remains controversial. Regardless, ACD is an important comorbidity and potential exacerbant of AD in clinical practice. Mechanisms of ACD developing in AD patients include epidermal barrier dysfunction leading to increased allergen and irritant penetrance, repetitive exposure to allergens secondary to frequent use of topical medications and personal care products, and bacterial colonization in AD promoting inflammation and potentiating contact sensitization.

Patch testing should be considered in adolescent- or adult-onset AD, worsening or more generalized dermatitis, localized or atypical lesional distribution suggestive of contact dermatitis, refractory AD, prior to systemic immunosuppressive treatment, or when AD worsens with topical therapy. Patch testing in AD should use an expanded patch-test series, though more research is needed to determine the optimal screening series in AD patients.

Acknowledgments

This publication was made possible with support from the Agency for Healthcare Research and Quality (AHRQ), grant number K12 HS023011, and the Dermatology Foundation.

References

1. Odhiambo JA, Williams HC, Clayton TO, Robertson CF, Asher MI, Group IPTS. Global variations in prevalence of eczema symptoms in children from ISAAC Phase Three. *The Journal of allergy and clinical immunology*. 2009 Dec; 124(6):1251–8. e23. [PubMed: 20004783]
2. Garg N, Silverberg JI. Epidemiology of childhood atopic dermatitis. *Clin Dermatol*. 2015 May-Jun; 33(3):281–8. [PubMed: 25889128]
3. Silverberg JI, Garg NK, Paller AS, Fishbein AB, Zee PC. Sleep disturbances in adults with eczema are associated with impaired overall health: a US population-based study. *The Journal of investigative dermatology*. 2015 Jan; 135(1):56–66. [PubMed: 25078665]
4. Silverberg JI, Hanifin JM. Adult eczema prevalence and associations with asthma and other health and demographic factors: a US population-based study. *The Journal of allergy and clinical immunology*. 2013 Nov; 132(5):1132–8. [PubMed: 24094544]
5. Vakharia P, Chopra R, Sacotte R, Patel K, Singam V, Patel N, et al. Burden of skin pain in atopic dermatitis. *Annals of Allergy, Asthma and Immunology*. 2018 In Press.
6. Bieber T. Atopic dermatitis 2.0: from the clinical phenotype to the molecular taxonomy and stratified medicine. *Allergy*. 2012 Dec; 67(12):1475–82. [PubMed: 23106343]
7. Hanifin J, Rajka G. Diagnostic features of atopic eczema. *Acta dermato-venereologica*. 1980; 92:44–7.
8. Diepgen TL, Sauerbrei W, Fartasch M. Development and validation of diagnostic scores for atopic dermatitis incorporating criteria of data quality and practical usefulness. *Journal of clinical epidemiology*. 1996 Sep; 49(9):1031–8. [PubMed: 8780613]
9. Asher MI, Keil U, Anderson HR, Beasley R, Crane J, Martinez F, et al. International Study of Asthma and Allergies in Childhood (ISAAC): rationale and methods. *The European respiratory journal*. 1995 Mar; 8(3):483–91. [PubMed: 7789502]
10. Williams HC, Burney PG, Hay RJ, Archer CB, Shipley MJ, Hunter JJ, et al. The U.K. Working Party's Diagnostic Criteria for Atopic Dermatitis I. Derivation of a minimum set of discriminators for atopic dermatitis. *The British journal of dermatology*. 1994 Sep; 131(3):383–96. [PubMed: 7918015]
11. Kang KF, Tian RM. Criteria for atopic dermatitis in a Chinese population. *Acta Derm Venereol Suppl (Stockh)*. 1989; 144:26–7. [PubMed: 2800903]

12. Japanese Dermatological Association Criteria for the diagnosis of atopic dermatitis. *The Journal of dermatology*. 2002; 29(6):398.
13. Nguyen SH, Dang TP, MacPherson C, Maibach H, Maibach HI. Prevalence of patch test results from 1970 to 2002 in a multi-centre population in North America (NACDG). *Contact dermatitis*. 2008 Feb; 58(2):101–6. [PubMed: 18186744]
14. Lim HW, Collins SAB, Resneck JS Jr, Bologna JL, Hodge JA, Rohrer TA, et al. The burden of skin disease in the United States. *Journal of the American Academy of Dermatology*. 2017 May; 76(5):958–72. e2. [PubMed: 28259441]
15. Mowad CM, Anderson B, Scheinman P, Pootongkam S, Nedorost S, Brod B. Allergic contact dermatitis: Patient diagnosis and evaluation. *Journal of the American Academy of Dermatology*. 2016 Jun; 74(6):1029–40. [PubMed: 27185421]
16. Beattie PE, Lewis-Jones MS. A comparative study of impairment of quality of life in children with skin disease and children with other chronic childhood diseases. *The British journal of dermatology*. 2006 Jul; 155(1):145–51. [PubMed: 16792766]
17. Holm EA, Wulf HC, Stegmann H, Jemec GB. Life quality assessment among patients with atopic eczema. *The British journal of dermatology*. 2006 Apr; 154(4):719–25. [PubMed: 16536816]
18. Chamlin SL, Frieden IJ, Williams ML, Chren MM. Effects of atopic dermatitis on young American children and their families. *Pediatrics*. 2004 Sep; 114(3):607–11. [PubMed: 15342828]
19. Kadyk DL, Hall S, Belsito DV. Quality of life of patients with allergic contact dermatitis: an exploratory analysis by gender, ethnicity, age, and occupation. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2004 Sep; 15(3):117–24.
20. Brutti CS, Bonamigo RR, Cappelletti T, Martins-Costa GM, Menegat AP. Occupational and non-occupational allergic contact dermatitis and quality of life: a prospective study. *An Bras Dermatol*. 2013 Jul-Aug; 88(4):670–1. [PubMed: 24068152]
21. Czarnobilska E, Obtulowicz K, Dyga W, Spiewak R. A half of schoolchildren with 'ISAAC eczema' are ill with allergic contact dermatitis. *Journal of the European Academy of Dermatology and Venereology : JEADV*. 2011 Sep; 25(9):1104–7. [PubMed: 21134222]
22. Cork MJ, Robinson DA, Vasilopoulos Y, Ferguson A, Moustafa M, MacGowan A, et al. New perspectives on epidermal barrier dysfunction in atopic dermatitis: gene-environment interactions. *The Journal of allergy and clinical immunology*. 2006 Jul; 118(1):3–21. quiz 2–3. [PubMed: 16815133]
23. Kantor R, Silverberg JJ. Environmental risk factors and their role in the management of atopic dermatitis. *Expert Rev Clin Immunol*. 2017 Jan; 13(1):15–26. [PubMed: 27417220]
24. Kabashima K. New concept of the pathogenesis of atopic dermatitis: interplay among the barrier, allergy, and pruritus as a trinity. *J Dermatol Sci*. 2013 Apr; 70(1):3–11. [PubMed: 23473856]
25. Proksch E, Brasch J. Abnormal epidermal barrier in the pathogenesis of contact dermatitis. *Clin Dermatol*. 2012 May-Jun; 30(3):335–44. [PubMed: 22507049]
26. Guttman-Yassky E, Nogales KE, Krueger JG. Contrasting pathogenesis of atopic dermatitis and psoriasis--part II: immune cell subsets and therapeutic concepts. *The Journal of allergy and clinical immunology*. 2011 Jun; 127(6):1420–32. [PubMed: 21419481]
27. Leung DY, Boguniewicz M, Howell MD, Nomura I, Hamid QA. New insights into atopic dermatitis. *The Journal of clinical investigation*. 2004 Mar; 113(5):651–7. [PubMed: 14991059]
28. Auriemma M, Vianale G, Amerio P, Reale M. Cytokines and T cells in atopic dermatitis. *European cytokine network*. 2013 Mar; 24(1):37–44. [PubMed: 23608610]
29. Brandt EB, Sivaprasad U. Th2 Cytokines and Atopic Dermatitis. *Journal of clinical & cellular immunology*. 2011 Aug 10; 2(3)
30. Chan JL, Davis-Reed L, Kimball AB. Counter-regulatory balance: atopic dermatitis in patients undergoing infliximab infusion therapy. *Journal of drugs in dermatology : JDD*. 2004 May-Jun; 3(3):315–8. [PubMed: 15176169]
31. Gittler JK, Shemer A, Suarez-Farinas M, Fuentes-Duculan J, Gulewicz KJ, Wang CQ, et al. Progressive activation of T(H)2/T(H)22 cytokines and selective epidermal proteins characterizes acute and chronic atopic dermatitis. *The Journal of allergy and clinical immunology*. 2012 Dec; 130(6):1344–54. [PubMed: 22951056]

32. Koga C, Kabashima K, Shiraishi N, Kobayashi M, Tokura Y. Possible pathogenic role of Th17 cells for atopic dermatitis. *The Journal of investigative dermatology*. 2008 Nov; 128(11):2625–30. [PubMed: 18432274]
33. Nograles KE, Zaba LC, Shemer A, Fuentes-Duculan J, Cardinale I, Kikuchi T, et al. IL-22-producing "T22" T cells account for upregulated IL-22 in atopic dermatitis despite reduced IL-17-producing TH17 T cells. *The Journal of allergy and clinical immunology*. 2009 Jun; 123(6):1244–52. e2. [PubMed: 19439349]
34. Souwer Y, Szegedi K, Kapsenberg ML, de Jong EC. IL-17 and IL-22 in atopic allergic disease. *Current opinion in immunology*. 2010 Dec; 22(6):821–6. [PubMed: 21087848]
35. Suarez-Farinas M, Dhingra N, Gittler J, Shemer A, Cardinale I, de Guzman Strong C, et al. Intrinsic atopic dermatitis shows similar TH2 and higher TH17 immune activation compared with extrinsic atopic dermatitis. *The Journal of allergy and clinical immunology*. 2013 Aug; 132(2):361–70. [PubMed: 23777851]
36. Werfel T, Allam JP, Biedermann T, Eyerich K, Gilles S, Guttman-Yassky E, et al. Cellular and molecular immunologic mechanisms in patients with atopic dermatitis. *The Journal of allergy and clinical immunology*. 2016 Aug; 138(2):336–49. [PubMed: 27497276]
37. Ma L, Xue HB, Guan XH, Shu CM, Zhang JH, Yu J. Possible pathogenic role of T helper type 9 cells and interleukin (IL)-9 in atopic dermatitis. *Clinical and experimental immunology*. 2014 Jan; 175(1):25–31. [PubMed: 24032555]
38. Ciprandi G, De Amici M, Giunta V, Marseglia A, Marseglia G. Serum interleukin-9 levels are associated with clinical severity in children with atopic dermatitis. *Pediatric dermatology*. 2013 Mar-Apr;30(2):222–5. [PubMed: 22612419]
39. Namkung JH, Lee JE, Kim E, Park GT, Yang HS, Jang HY, et al. An association between IL-9 and IL-9 receptor gene polymorphisms and atopic dermatitis in a Korean population. *J Dermatol Sci*. 2011 Apr; 62(1):16–21. [PubMed: 21371865]
40. Al-Muhsen S, Letuve S, Vazquez-Tello A, Pureza MA, Al-Jahdali H, Bahammam AS, et al. Th17 cytokines induce pro-fibrotic cytokines release from human eosinophils. *Respiratory research*. 2013 Mar 13.14:34. [PubMed: 23496774]
41. Eyerich K, Pennino D, Scarponi C, Foerster S, Nasorri F, Behrendt H, et al. IL-17 in atopic eczema: linking allergen-specific adaptive and microbial-triggered innate immune response. *The Journal of allergy and clinical immunology*. 2009 Jan; 123(1):59–66. e4. [PubMed: 19056110]
42. Trautmann A, Akdis M, Schmid-Grendelmeier P, Disch R, Brocker EB, Blaser K, et al. Targeting keratinocyte apoptosis in the treatment of atopic dermatitis and allergic contact dermatitis. *The Journal of allergy and clinical immunology*. 2001 Nov; 108(5):839–46. [PubMed: 11692113]
43. Gober MD, Gaspari AA. Allergic contact dermatitis. *Current directions in autoimmunity*. 2008; 10:1–26. [PubMed: 18460878]
44. Kim BS, Miyagawa F, Cho YH, Bennett CL, Clausen BE, Katz SI. Keratinocytes function as accessory cells for presentation of endogenous antigen expressed in the epidermis. *The Journal of investigative dermatology*. 2009 Dec; 129(12):2805–17. [PubMed: 19554018]
45. Martin, SF, Pichler, WJ. Drug hypersensitivity. Karger, Basel: 2007. Chemical-induced contact hypersensitivity in the mouse model; p. 34-46.
46. Dhingra N, Shemer A, Correa da Rosa J, Rozenblit M, Fuentes-Duculan J, Gittler JK, et al. Molecular profiling of contact dermatitis skin identifies allergen-dependent differences in immune response. *The Journal of allergy and clinical immunology*. 2014 Aug; 134(2):362–72. [PubMed: 24768652]
47. Coulter EM, Jenkinson C, Farrell J, Lavergne SN, Pease C, White A, et al. Measurement of CD4+ and CD8+ T-lymphocyte cytokine secretion and gene expression changes in p-phenylenediamine allergic patients and tolerant individuals. *The Journal of investigative dermatology*. 2010 Jan; 130(1):161–74. [PubMed: 19657353]
48. Gutin L, Tammara A, Fischelevich R, Gaspari AA. Elevation of IL-9 in Extreme Patch Test Reactions Suggests It Is an Inflammatory Mediator in Allergic Contact Dermatitis. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2016 Jan-Feb;27(1):35–6.

49. Liu J, Harberts E, Tammaro A, Girardi N, Filler RB, Fischelevich R, et al. IL-9 regulates allergen-specific Th1 responses in allergic contact dermatitis. *The Journal of investigative dermatology*. 2014 Jul; 134(7):1903–11. [PubMed: 24487305]
50. Larsen JM, Bonefeld CM, Poulsen SS, Geisler C, Skov L. IL-23 and T(H)17-mediated inflammation in human allergic contact dermatitis. *The Journal of allergy and clinical immunology*. 2009 Feb; 123(2):486–92. [PubMed: 18986691]
51. Zhao Y, Balato A, Fischelevich R, Chapoval A, Mann DL, Gaspari AA. Th17/Tc17 infiltration and associated cytokine gene expression in elicitation phase of allergic contact dermatitis. *The British journal of dermatology*. 2009 Dec; 161(6):1301–6. [PubMed: 19785613]
52. Albanesi C, Cavani A, Girolomoni G. IL-17 is produced by nickel-specific T lymphocytes and regulates ICAM-1 expression and chemokine production in human keratinocytes: synergistic or antagonist effects with IFN-gamma and TNF-alpha. *J Immunol*. 1999 Jan 01; 162(1):494–502. [PubMed: 9886425]
53. Kebir H, Kreymborg K, Ifergan I, Dodelet-Devillers A, Cayrol R, Bernard M, et al. Human TH17 lymphocytes promote blood-brain barrier disruption and central nervous system inflammation. *Nature medicine*. 2007 Oct; 13(10):1173–5.
54. Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *The Journal of experimental medicine*. 2006 Oct 02; 203(10):2271–9. [PubMed: 16982811]
55. Nakae S, Komiyama Y, Nambu A, Sudo K, Iwase M, Homma I, et al. Antigen-specific T cell sensitization is impaired in IL-17-deficient mice, causing suppression of allergic cellular and humoral responses. *Immunity*. 2002 Sep; 17(3):375–87. [PubMed: 12354389]
56. Gimenez-Arnau A, Maurer M, De La Cuadra J, Maibach H. Immediate contact skin reactions, an update of Contact Urticaria, Contact Urticaria Syndrome and Protein Contact Dermatitis -- “A Never Ending Story”. *European journal of dermatology : EJD*. 2010 Sep-Oct;20(5):552–62. [PubMed: 20732848]
57. Winton GB, Lewis CW. Contact urticaria. *International journal of dermatology*. 1982 Dec; 21(10):573–8. [PubMed: 6761282]
58. Nosbaum A, Vocanson M, Rozieres A, Hennino A, Nicolas JF. Allergic and irritant contact dermatitis. *European journal of dermatology : EJD*. 2009 Jul-Aug;19(4):325–32. [PubMed: 19447733]
59. Slodownik D, Lee A, Nixon R. Irritant contact dermatitis: a review. *Australas J Dermatol*. 2008 Feb; 49(1):1–9. quiz 10-1. [PubMed: 18186838]
60. Uehara M, Sawai T. A longitudinal study of contact sensitivity in patients with atopic dermatitis. *Archives of dermatology*. 1989 Mar; 125(3):366–8. [PubMed: 2522294]
61. Uehara M, Ofuji S. Patch test reactions to human dander in atopic dermatitis. *Archives of dermatology*. 1976 Jul; 112(7):951–4. [PubMed: 938073]
62. Jones HE, Lewis CW, McMarlin SL. Allergic contact sensitivity in atopic dermatitis. *Archives of dermatology*. 1973 Feb; 107(2):217–22. [PubMed: 4265457]
63. Forsbeck M, Hovmark A, Skog E. Patch testing, tuberculin testing and sensitization with dinitrochlorobenzene and nitrosodimethylanilini of patients with atopic dermatitis. *Acta dermato-venereologica*. 1976; 56(2):135–8. [PubMed: 58522]
64. Thyssen JP, Kezic S. Causes of epidermal filaggrin reduction and their role in the pathogenesis of atopic dermatitis. *The Journal of allergy and clinical immunology*. 2014 Oct; 134(4):792–9. [PubMed: 25065719]
65. Jakasa I, de Jongh CM, Verberk MM, Bos JD, Kezic S. Percutaneous penetration of sodium lauryl sulphate is increased in uninvolved skin of patients with atopic dermatitis compared with control subjects. *The British journal of dermatology*. 2006 Jul; 155(1):104–9. [PubMed: 16792760]
66. Halling-Overgaard AS, Kezic S, Jakasa I, Engebretsen KA, Maibach H, Thyssen JP. Skin absorption through atopic dermatitis skin: a systematic review. *The British journal of dermatology*. 2017 Jul; 177(1):84–106. [PubMed: 27639188]
67. Thyssen JP, McFadden JP, Kimber I. The multiple factors affecting the association between atopic dermatitis and contact sensitization. *Allergy*. 2014 Jan; 69(1):28–36. [PubMed: 24372195]

68. Jakasa I, Verberk MM, Esposito M, Bos JD, Kezic S. Altered penetration of polyethylene glycols into uninvolved skin of atopic dermatitis patients. *The Journal of investigative dermatology*. 2007 Jan; 127(1):129–34. [PubMed: 17039242]
69. Agner T, Johansen JD, Overgaard L, Volund A, Basketter D, Menne T. Combined effects of irritants, allergens. Synergistic effects of nickel and sodium lauryl sulfate in nickel-sensitized individuals. *Contact dermatitis*. 2002 Jul; 47(1):21–6. [PubMed: 12225409]
70. McLelland J, Shuster S, Matthews JN. 'Irritants' increase the response to an allergen in allergic contact dermatitis. *Archives of dermatology*. 1991 Jul; 127(7):1016–9. [PubMed: 2064399]
71. Pedersen LK, Johansen JD, Held E, Agner T. Augmentation of skin response by exposure to a combination of allergens and irritants - a review. *Contact dermatitis*. 2004 May; 50(5):265–73. [PubMed: 15209808]
72. Hamann CR, Bernard S, Hamann D, Hansen R, Thyssen JP. Is there a risk using hypoallergenic cosmetic pediatric products in the United States? *The Journal of allergy and clinical immunology*. 2015 Apr; 135(4):1070–1. [PubMed: 25458001]
73. Mailhol C, Lauwers-Cances V, Rance F, Paul C, Giordano-Labadie F. Prevalence and risk factors for allergic contact dermatitis to topical treatment in atopic dermatitis: a study in 641 children. *Allergy*. 2009 May; 64(5):801–6. [PubMed: 19183418]
74. Seo KS, Park JY, Terman DS, Bohach GA. A quantitative real time PCR method to analyze T cell receptor Vbeta subgroup expansion by staphylococcal superantigens. *Journal of translational medicine*. 2010 Jan 13.8
75. Huang L, Kinbara M, Funayama H, Takada H, Sugawara S, Endo Y. The elicitation step of nickel allergy is promoted in mice by microbe-related substances, including some from oral bacteria. *International immunopharmacology*. 2011 Nov; 11(11):1916–24. [PubMed: 21839857]
76. Takahashi H, Kinbara M, Sato N, Sasaki K, Sugawara S, Endo Y. Nickel allergy-promoting effects of microbial or inflammatory substances at the sensitization step in mice. *International immunopharmacology*. 2011 Oct; 11(10):1534–40. [PubMed: 21621645]
77. Simonsen AB, Johansen JD, Deleuran M, Mortz CG, Sommerlund M. Contact allergy in children with atopic dermatitis: a systematic review. *The British journal of dermatology*. 2017 Aug; 177(2):395–405. [PubMed: 28470762]
78. Hamann CR, Hamann D, Egeberg A, Johansen JD, Silverberg J, Thyssen JP. Association between atopic dermatitis and contact sensitization: A systematic review and meta-analysis. *Journal of the American Academy of Dermatology*. 2017 Jul; 77(1):70–8. [PubMed: 28392290]
79. Paulsen E, Andersen KE. Sensitization patterns in Compositae-allergic patients with current or past atopic dermatitis. *Contact dermatitis*. 2013 May; 68(5):277–85. [PubMed: 23601063]
80. Simonsen AB, Deleuran M, Johansen JD, Sommerlund M. Contact allergy and allergic contact dermatitis in children - a review of current data. *Contact dermatitis*. 2011 Nov; 65(5):254–65. [PubMed: 21851361]
81. Schena D, Papagrigoraki A, Tessari G, Peroni A, Sabbadini C, Girolomoni G. Allergic contact dermatitis in children with and without atopic dermatitis. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2012 Nov-Dec;23(6):275–80.
82. Giordano-Labadie F, Rance F, Pellegrin F, Bazex J, Dutau G, Schwarze HP. Frequency of contact allergy in children with atopic dermatitis: results of a prospective study of 137 cases. *Contact dermatitis*. 1999 Apr; 40(4):192–5. [PubMed: 10208505]
83. Herro EM, Matiz C, Sullivan K, Hamann C, Jacob SE. Frequency of contact allergens in pediatric patients with atopic dermatitis. *The Journal of clinical and aesthetic dermatology*. 2011 Nov; 4(11):39–41.
84. Jacob SE, Yang A, Herro E, Zhang C. Contact allergens in a pediatric population: association with atopic dermatitis and comparison with other north american referral centers. *The Journal of clinical and aesthetic dermatology*. 2010 Oct; 3(10):29–35.
85. Lammintausta K, Kalimo K, Fagerlund VL. Patch test reactions in atopic patients. *Contact dermatitis*. 1992 Apr; 26(4):234–40. [PubMed: 1395560]
86. Lever R, Forsyth A. Allergic contact dermatitis in atopic dermatitis. *Acta Derm Venereol Suppl (Stockh)*. 1992; 176:95–8. [PubMed: 1476046]

87. Lubbes S, Rustemeyer T, Sillevs Smitt JH, Schuttelaar ML, Middelkamp-Hup MA. Contact sensitization in Dutch children and adolescents with and without atopic dermatitis - a retrospective analysis. *Contact dermatitis*. 2017 Mar; 76(3):151–9. [PubMed: 27861990]
88. Malajian D, Belsito DV. Cutaneous delayed-type hypersensitivity in patients with atopic dermatitis. *Journal of the American Academy of Dermatology*. 2013 Aug; 69(2):232–7. [PubMed: 23583066]
89. Netterlid E, Hindsen M, Ekqvist S, Henricson KA, Bruze M. Young individuals with atopic disease and asthma or rhinoconjunctivitis may have clinically relevant contact allergies. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2014 May-Jun;25(3):115–9.
90. Silny W, Bartoszak L, Jenerowicz D, Zukiewicz-Sobczak W, Gozdziwska M. Prevalence of contact allergy in children suffering from atopic dermatitis, seborrhoeic dermatitis and in healthy controls. *Annals of agricultural and environmental medicine : AAEM*. 2013; 20(1):55–60. [PubMed: 23540212]
91. Xu S, Kwa M, Lohman ME, Evers-Meltzer R, Silverberg JI. Consumer Preferences, Product Characteristics, and Potentially Allergenic Ingredients in Best-selling Moisturizers. *JAMA dermatology*. 2017 Nov 01; 153(11):1099–105. [PubMed: 28877310]
92. Overgaard LEK, Main KM, Frederiksen H, Stender S, Szecsi PB, Williams HC, et al. Children with atopic dermatitis and frequent emollient use have increased urinary levels of low-molecular-weight phthalate metabolites and parabens. *Allergy*. 2017 Nov; 72(11):1768–77. [PubMed: 28281298]
93. Warshaw EM, Nelsen DD, Maibach HI, Marks JG, Zug KA, Taylor JS, et al. Positive patch test reactions to lanolin: cross-sectional data from the north american contact dermatitis group, 1994 to 2006. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2009 Mar-Apr;20(2):79–88.
94. Chen JK, Jacob SE, Nedorost ST, Hanifin JM, Simpson EL, Boguniewicz M, et al. A Pragmatic Approach to Patch Testing Atopic Dermatitis Patients: Clinical Recommendations Based on Expert Consensus Opinion. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2016 Jul-Aug; 27(4):186–92.
95. Silverberg JI, Vakharia PP, Chopra R, Sacotte R, Patel N, Immaneni S, et al. Phenotypical Differences of Childhood- and Adult-Onset Atopic Dermatitis. *The journal of allergy and clinical immunology In practice*. 2017 Nov 10.
96. Julian-Gonzalez RE, Orozco-Covarrubias L, Duran-McKinster C, Palacios-Lopez C, Ruiz-Maldonado R, Saez-de-Ocariz M. Less common clinical manifestations of atopic dermatitis: prevalence by age. *Pediatric dermatology*. 2012 Sep-Oct;29(5):580–3. [PubMed: 22469300]
97. Silverberg JI, Vakharia PP, Chopra R, Sacotte R, Patel N, Immaneni S, et al. Phenotypical Differences of Childhood- and Adult-Onset Atopic Dermatitis. *The Journal of Allergy and Clinical Immunology: In Practice*.
98. Bonamonte D, Foti C, Vestita M, Ranieri LD, Angelini G. Nummular eczema and contact allergy: a retrospective study. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2012 Jul-Aug; 23(4):153–7.
99. Krupa Shankar DS, Shrestha S. Relevance of patch testing in patients with nummular dermatitis. *Indian journal of dermatology, venereology and leprology*. 2005 Nov-Dec;71(6):406–8.
100. Fowler JF Jr, Maibach HI, Zirwas M, Taylor JS, Dekoven JG, Sasseville D, et al. Effects of immunomodulatory agents on patch testing: expert opinion 2012. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2012 Nov-Dec;23(6):301–3.
101. Johansen JD, Aalto-Korte K, Agner T, Andersen KE, Bircher A, Bruze M, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing -recommendations on best practice. *Contact dermatitis*. 2015 Oct; 73(4):195–221. [PubMed: 26179009]
102. Aquino M, Fonacier L. The role of contact dermatitis in patients with atopic dermatitis. *The journal of allergy and clinical immunology In practice*. 2014 Jul-Aug;2(4):382–7. [PubMed: 25017524]

103. Brasch J, Schnuch A, Uter W. Patch-test reaction patterns in patients with a predisposition to atopic dermatitis. *Contact dermatitis*. 2003 Oct; 49(4):197–201. [PubMed: 14996068]
104. el Samahy MH, el-Kerdani T. Value of patch testing in atopic dermatitis. *American journal of contact dermatitis : official journal of the American Contact Dermatitis Society*. 1997 Sep; 8(3): 154–7. [PubMed: 9249284]
105. Nedorost ST, Babineau D. Patch testing in atopic dermatitis. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2010 Sep-Oct;21(5):251–4.
106. [Accessed November 11, 2017] Cosmetic Labeling Guide. U.S. Food and Drug Administration. Nov 05, 2017 <https://www.fda.gov/Cosmetics/Labeling/Regulations/ucm126444.htm> Updated
107. Kligman AM. The myth of lanolin allergy. *Contact dermatitis*. 1998 Sep; 39(3):103–7. [PubMed: 9771981]
108. Newell L, Polak ME, Perera J, Owen C, Boyd P, Pickard C, et al. Sensitization via healthy skin programs Th2 responses in individuals with atopic dermatitis. *The Journal of investigative dermatology*. 2013 Oct; 133(10):2372–80. [PubMed: 23528819]
109. Higgins E, Collins P. The relevance of 7-day patch test reading. *Dermatitis : contact, atopic, occupational, drug : official journal of the American Contact Dermatitis Society, North American Contact Dermatitis Group*. 2013 Sep-Oct;24(5):237–40.
110. Isaksson M, Andersen KE, Brandao FM, Bruynzeel DP, Bruze M, Camarasa JG, et al. Patch testing with corticosteroid mixes in Europe. A multicentre study of the EECDRG. *Contact dermatitis*. 2000 Jan; 42(1):27–35. [PubMed: 10644022]
111. Davis MD, Bhate K, Rohlinger AL, Farmer SA, Richardson DM, Weaver AL. Delayed patch test reading after 5 days: the Mayo Clinic experience. *Journal of the American Academy of Dermatology*. 2008 Aug; 59(2):225–33. [PubMed: 18513826]

Key points

- Patients with atopic dermatitis appear to have increased risk of developing allergic contact dermatitis.
- Patch testing should be considered in adolescent- or adult-onset atopic dermatitis, worsening or more generalized dermatitis, localized or atypical lesional distribution, refractory disease, prior to systemic immunosuppressive treatment, or when atopic dermatitis worsens with topical therapy.
- Patch testing in atopic dermatitis should use an expanded patch-test series and the results interpreted with caution.

Table 1

Common contact allergens identified in patients with atopic dermatitis.

Bacitracin
Carba mix
Chromium
Cinnamic aldehyde
Cobalt
Cocamidopropyl betaine
Colophonium
Compositae mix
Disperse blue dye 106
Epoxy resin
Formaldehyde
Fragrance markers (e.g. fragrance mix I, fragrance mix II, Myroxylon pereirae, and hydroxyisohexyl-3-cyclohexenecarboxaldehyde)
Isothiazolinones (e.g. methylisothiazolinone and methylchlorisothiazolinone)
Lanolin
Mercaptobenzothiazole and mercaptans
Myroxylon pereirae
Neomycin
Nickel
Para-tertiarybutyl-phenol (PTBP) formaldehyde resin
Paraphenylenediamine
Potassium dichromate
Quaternium-15
Rubber or rubber mixes
Sesquiterpene lactone mix
Topical antiseptics (e.g. chlorhexidine, hexamidine)

Table 2

Pitfalls in patch testing in atopic dermatitis (AD) patients.

Current or recent exposure to systemic immunosuppressive medications (Section 4.2), ultraviolet therapy or excessive solar radiation can decrease the sensitivity threshold of patch testing and lead to false negatives. Repeat patch testing should be considered upon treatment discontinuation and washout.
Patients with AD have a lower irritancy threshold, which may lead to higher rates of irritant or false positive reactions (most commonly with metals, fragrance, formaldehyde, and lanolin)
Positive reactions in AD patients may display as weaker reactions and be misdiagnosed as an irritant reaction (i.e. negative reaction)
Active or flaring AD may result in false negative reactions due to decreased contact sensitization