critical illness: a randomized controlled trial with 12 months of follow-up. *Crit Care* 2013;17:R156.

3. Moss M, Nordon-Craft A, Malone D, Van Pelt D, Frankel SK, Warner ML, *et al.* A randomized trial of an intensive physical therapy program for patients with acute respiratory failure. *Am J Respir Crit Care Med* 2016;193:1101–1110.
4. Walsh TS, Salisbury LG, Merriweather JL, Boyd JA, Griffith DM, Huby G, *et al.*; RECOVER Investigators. Increased hospital-based physical rehabilitation and information provision after intensive care unit discharge: the RECOVER randomized clinical trial. *JAMA Intern Med* 2015;175:901–910.
5. Elliott D, McKinley S, Alison J, Aitken LM, King M, Leslie GD, *et al.* Health-related quality of life and physical recovery after a critical illness: a multi-centre randomised controlled trial of a home-based physical rehabilitation program. *Crit Care* 2011;15:R142.
6. Lim WC, Black N, Lamping D, Rowan K, Mays N. Conceptualizing and measuring health-related quality of life in critical care. *J Crit Care* 2016;31:183–193.
7. Black N. Patient reported outcome measures could help transform healthcare. *BMJ* 2013;346:f167.

Copyright © 2018 by the American Thoracic Society

CAPTURE: A Screening Tool for Chronic Obstructive Pulmonary Disease or Obstructive Airway Disease?

To the Editor:

We read with great interest the article by Martinez and colleagues (1) wherein they developed a five-item questionnaire in addition to a peak expiratory flow rate (PEFR) value to screen potential cases of chronic obstructive pulmonary disease (COPD) that require further evaluation. The authors used an innovative, multimethod approach to select the best, smallest set of questionnaire items from a comprehensive list of candidate items derived from the literature, analysis of existing data, and qualitative research. Despite the merit and uniqueness of this approach, we find a major flaw in the development of the CAPTURE questionnaire. It did not have a respiratory symptoms control arm (i.e., another control group of subjects who had respiratory symptoms, but no COPD—for example, those who had asthma). When we administered the CAPTURE questionnaire to some of our subjects here, many who had asthma screened positive for COPD. This is not surprising, because some of the questions, such as “change in breathing due to seasons or air quality,” “cough around perfume, grass, or smoke,” “ever short of breath,” and “missed work/activities due to an acute respiratory condition,” will also be answered “yes” by subjects with asthma. Moreover, subjects with asthma will also have low PEFR value. As a result, CAPTURE may falsely label a large number of subjects with asthma as potential cases of COPD. We believe this is a major drawback of CAPTURE, and therefore it does not merit being labeled as a COPD screening questionnaire. Ideally, it would seem more appropriate to first detect presence of obstructive airways disease using a questionnaire and a peak flow meter and then have an additional set of questions that will help answer if the obstructive airways disease is likely to be asthma or COPD. We made an attempt using this approach with a reasonable

measure of success recently (2), although our study needs further validation.

Second, CAPTURE gives lesser importance to history of tobacco smoking than might be expected as an important question to help screen COPD, which we found interesting. Although screening for COPD in nonsmokers is indeed important, given the fact that a significantly large number of nonsmokers also develop COPD (3), we believe that omission of an explicit question about smoking history in CAPTURE reflects an inherent flaw in the study design. The authors included subjects with mild COPD in the control arm, presumably to balance the numbers in the study and control arms. But because 66% of the people in the mild COPD group were former smokers, this increased the proportion of smokers in the control group, thereby weakening the ability to use this question to find differences between cases and control subjects. We believe this was a second flaw in the study design. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Jyoti Londhe, M.S.W.
 Komalkirti Apte, M.B. B.S., P.G.D.C.R.
 Monica Barne, M.B. B.S.
 Sundeeep Salvi, M.D., D.N.B., Ph.D.
 Chest Research Foundation
 Pune, India

ORCID ID: 0000-0001-6843-0582 (J.L.).

References

1. Martinez FJ, Mannino D, Leidy NK, Malley KG, Bacci ED, Barr RG, *et al.*; High-Risk-COPD Screening Study Group. A new approach for identifying patients with undiagnosed chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2017;195:748–756.
2. Thorat Y, Salvi S, Kodgule R. Peak flow meter with a questionnaire and mini spirometer to help detect asthma and COPD in real-life clinical practice: a cross-sectional study. *NPJ Prim Care Respir Med* 2017;27:32.
3. Salvi SS, Barnes PJ. Chronic obstructive pulmonary disease in non-smokers. *Lancet* 2009;374:733–743.

Copyright © 2018 by the American Thoracic Society

Reply to Londhe *et al.*

From the Authors:

We appreciate the insightful comments of Londhe and colleagues. Their comments highlight important concepts that must be clarified. It is quite evident that patients with chronic obstructive pulmonary disease (COPD) remain underdiagnosed in primary care settings (1); these individuals suffer from impaired health status (2) and outcomes (3, 4). Importantly, therapies are available to improve lung function, reduce exacerbations, and improve health status in patients with COPD, particularly those with an FEV₁ less than 60% predicted or who are at risk for acute exacerbations (5, 6).

Originally Published in Press as DOI: 10.1164/rccm.201705-1040LE on July 19, 2017

Originally Published in Press as DOI: 10.1164/rccm.201707-1393LE on July 19, 2017

As such, an NHLBI task force suggested the identification of these patients may prove to be an ideal initial stage in systematically evaluating the potential impact of COPD case finding in primary care (7). This was the foundation behind development of the CAPTURE approach and provides insight into the populations chosen, including control subjects with no COPD or mild COPD. Our case-finding approach was not developed as a diagnostic instrument but as a simple format to identify individuals at greater likelihood of having COPD. In these individuals, a subsequent, targeted, and staged approach including lung function measurement and other diagnostic tools is warranted to ensure an accurate diagnosis. Given the clinical and therapeutic overlaps between asthma and COPD (8), it is not surprising that CAPTURE would identify these individuals and suggest to primary clinicians that further evaluation is warranted.

Similar concepts relate to the smoking query in CAPTURE. The original candidate item set included numerous items to assess smoking history (9). Importantly, an explicit smoking history question (have you ever smoked cigarettes, pipes, or cigars?) resulted in a higher error rate in the case/control comparison (16.6% vs. 15.7%). Given the strong performance of the five items without a direct smoking query, the investigative team concluded that this instrument should be taken forward for use with both smokers and nonsmokers. We believe this is particularly relevant, as many of the previously developed case-finding approaches used items from existing epidemiologic literature or expert opinion with limited or no patient input; many were heavily dependent on age and smoking history (10). As it is increasingly evident that a sizeable proportion of patients with COPD do not have a smoking history (6, 11), we believe our approach will be particularly relevant in a broader range of patient populations.

As the role of spirometry to enhance diagnostic efforts has been recently addressed by the U.S. Preventive Services Task Force report (12) and a subsequent editorial (13), the development of simple, questionnaire-based approaches should offer a practical triage or case-finding method for identifying symptomatic people in practice settings who may have unrecognized COPD and would benefit from appropriate diagnostic evaluation and treatment. The approach we have taken and that developed by Thorat and colleagues (14) offer a unique opportunity. We wholeheartedly agree that additional validation of such approaches is desperately needed. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Fernando J. Martinez, M.D., M.S.
Weill Cornell Medical College
New York, New York

MeiLan Han, M.D., M.S.
University of Michigan
Ann Arbor, Michigan

Nancy Leidy, Ph.D.
Evidera
Bethesda, Maryland

Barry Make, M.D.
National Jewish Health
Denver, Colorado

David M. Mannino, M.D.
University of Kentucky
Lexington, Kentucky

Stephen I. Rennard, M.D.
University of Nebraska Medical Center
Omaha, Nebraska
and
AstraZeneca
Cambridge, United Kingdom

Byron M. Thomashow, M.D.
Columbia University
New York, New York

Barbara P. Yawn, M.D.
University of Minnesota
Minneapolis, Minnesota

For the High-Risk-COPD Screening Study Group

References

1. Ford ES, Mannino DM, Wheaton AG, Giles WH, Presley-Cantrell L, Croft JB. Trends in the prevalence of obstructive and restrictive lung function among adults in the United States: findings from the National Health and Nutrition Examination surveys from 1988-1994 to 2007-2010. *Chest* 2013;143:1395-1406.
2. Miravittles M, Soriano JB, García-Río F, Muñoz L, Duran-Tauleria E, Sanchez G, et al. Prevalence of COPD in Spain: impact of undiagnosed COPD on quality of life and daily life activities. *Thorax* 2009;64:863-868.
3. Martinez CH, Mannino DM, Jaimes FA, Curtis JL, Han MK, Hansel NN, et al. Undiagnosed obstructive lung disease in the United States: associated factors and long-term mortality. *Ann Am Thorac Soc* 2015; 12:1788-1795.
4. Labonté LE, Tan WC, Li PZ, Mancino P, Aaron SD, Benedetti A, et al. Canadian Respiratory Research Network; CanCOLD Collaborative Research Group. Undiagnosed chronic obstructive pulmonary disease contributes to the burden of health care use: data from the CanCOLD study. *Am J Respir Crit Care Med* 2016;194:285-298.
5. Qaseem A, Snow V, Shekelle P, Sherif K, Wilt TJ, Weinberger S, et al.; Clinical Efficacy Assessment Subcommittee of the American College of Physicians. Diagnosis and management of stable chronic obstructive pulmonary disease: a clinical practice guideline from the American College of Physicians. *Ann Intern Med* 2007;147:633-638.
6. Vogelmeier CF, Criner GJ, Martinez FJ, Anzueto A, Barnes PJ, Bourbeau J, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report: GOLD executive summary. *Am J Respir Crit Care Med* 2017;195:557-582.
7. National Heart, Lung, and Blood Institute. A case-finding strategy for moderate-to-severe COPD in the United States. 2009 [accessed 2016 Jul 4]. Available from: <http://www.nhlbi.nih.gov/research/reports/2009-case-finding-exesum>.
8. Barnes PJ. Therapeutic approaches to asthma-chronic obstructive pulmonary disease overlap syndromes. *J Allergy Clin Immunol* 2015; 136:531-545.
9. Martinez FJ, Mannino D, Leidy NK, Malley KG, Bacci ED, Barr RG, et al.; High-Risk-COPD Screening Study Group. A new approach for identifying patients with undiagnosed chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2017;195:748-756.
10. Han MK, Steenrod AW, Bacci ED, Leidy NK, Mannino DM, Thomashow BM, et al. Identifying patients with undiagnosed COPD in primary care settings: insight from screening tools and epidemiologic studies. *Chronic Obstr Pulm Dis (Miami)* 2015;2:103-121.
11. Lamprecht B, McBurnie MA, Vollmer WM, Gudmundsson G, Welte T, Nizankowska-Mogilnicka E, et al.; BOLD Collaborative Research Group. COPD in never smokers: results from the population-based burden of obstructive lung disease study. *Chest* 2011;139: 752-763.

12. Guirguis-Blake JM, Senger CA, Webber EM, Mularski RA, Whitlock EP. Screening for chronic obstructive pulmonary disease: evidence report and systematic review for the US Preventive Services Task Force. *JAMA* 2016;315:1378–1393.
13. Martinez FJ, O'Connor GT. Screening, case-finding, and outcomes for adults with unrecognized COPD. *JAMA* 2016;315:1343–1344.
14. Thorat Y, Salvi S, Kodgule R. Peak flow meter with a questionnaire and mini-spirometer to help detect asthma and COPD in real-life clinical practice: a cross-sectional study. *NPJ Prim Care Respir Med* 2017;27:32.

Copyright © 2018 by the American Thoracic Society

Cigarette Smoke Is an Endothelial Cell Toxin

To the Editor:

The report by Polverino and colleagues (1) provides a wonderful example of productive interdisciplinary collaboration. The authors connect lung and kidney endothelial cell damage and cigarette smoke in patients with chronic obstructive pulmonary disease and in smoke-exposed mice. This is an important study with far-reaching implications. Mechanistically, the authors point to generation of advanced glycation end products and receptor for advanced glycation end products activation triggered by oxidant stress, yet there are likely additional mechanisms that can be considered. Because the kidneys do not inhale, it is evident that it is the “circulating cigarette smoke” that is toxic to endothelial cells. For example, one component of cigarette smoke is the highly aggressive aldehyde acrolein (2–4), which can be measured in the blood and urine of smokers. A large number of experimental studies leave no doubt that cigarette smoke extract injures cultured endothelial cells. These findings have been frequently ignored or characterized as clinically irrelevant, yet they provide mechanistic insights and allow us to consider the hypothesis that emphysema is a vascular problem (5). If the notion that cigarette smoke cannot cross the alveolar–capillary barrier is dismissed, the fact that components of cigarette smoke are circulating has been firmly established. Although cigarette smoke extract *in vitro* kills endothelial cells directly, *in vivo* cigarette smoke components could, in susceptible individuals, trigger an immune response that includes the production of antiendothelial antibodies (6). Another implication of the findings published by Polverino and colleagues is that not only are the lung and kidney microvessels affected by cigarette smoke components but similarly the microcirculation of the skeletal muscles, the myocardium, and the retina are affected (1).

Interesting correlations, not provided by the authors, are those between albuminuria and the diffusing capacity of the lung for carbon monoxide and albuminuria and urinary isoprostane concentration (7). Last, this study suggests that the vascular chronic obstructive pulmonary disease phenotype might benefit from antioxidant treatments. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Norbert F. Voelkel, M.D.
University of Colorado
Denver, Colorado

References

1. Polverino F, Lauchó-Contreras ME, Petersen H, Bijol V, Sholl LM, Choi ME, *et al*. A pilot study linking endothelial injury in lungs and kidneys in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2017;195:1464–1476.
2. Voelkel NF, Mizuno S, Yasuo M. Does drug-induced emphysema exist? *Eur Respir J* 2013;42:1464–1468.
3. Nana-Sinkam SP, Lee JD, Sotto-Santiago S, Stearman RS, Keith RL, Choudhury Q, *et al*. Prostacyclin prevents pulmonary endothelial cell apoptosis induced by cigarette smoke. *Am J Respir Crit Care Med* 2007;175:676–685.
4. Kitaguchi Y, Taraseviciene-Stewart L, Hanaoka M, Natarajan R, Kraskauskas D, Voelkel NF. Acrolein induces endoplasmic reticulum stress and causes airspace enlargement. *PLoS One* 2012;7:e38038.
5. Voelkel NF, Gomez-Arroyo J, Mizuno S. COPD/emphysema: the vascular story. *Pulm Circ* 2011;1:320–326.
6. Taraseviciene-Stewart L, Voelkel NF. Molecular pathogenesis of emphysema. *J Clin Invest* 2008;118:394–402.
7. Santini G, Mores N, Shohreh R, Valente S, Dabrowska M, Trové A, *et al*. Exhaled and non-exhaled non-invasive markers for assessment of respiratory inflammation in patients with stable COPD and healthy smokers. *J Breath Res* 2016;10:017102.

Copyright © 2018 by the American Thoracic Society

Reply to Voelkel

From the Authors:

We thank Dr. Voelkel for his complimentary comments acknowledging the importance of our study (1). As Dr. Voelkel correctly pointed out, the vascular chronic obstructive pulmonary disease (COPD) phenotype is a clinical entity that has been underrecognized. Initial studies from Dr. Voelkel's group provided the first clue that endothelial cell death is sufficient to induce airspace enlargement in rodents (2–4) and occurs in lungs of patients with COPD (5). Our studies of cigarette smoke (CS)-exposed mice and patients with COPD further support this hypothesis and demonstrate that endothelial injury also occurs in the kidneys in patients with COPD and smokers (1). Likely, CS injures endothelial cells in other tissues, including skeletal muscle and bones, and contributes to the comorbidities that are frequently detected in patients with COPD (6).

As Dr. Voelkel correctly points out, we focused on the contributions of the oxidative stress–advanced glycation end products (AGEs)–receptor for AGEs pathway in the pathogenesis of CS-induced endothelial injury. We investigated clues from studies of humans and experimental models of diabetes mellitus and hypertension (common comorbidities of COPD) that linked endothelial injury and microvascular complications of these diseases to