George A. Whitmore, Ph.D.
McGill University
Montreal, Québec, Canada

On behalf of all authors

ORCID ID: 0000-0002-4762-3542 (S.D.A.).

References

1. Sood A, Petersen H, Qualls C, Meek PM, Vazquez-Guillamet R, Celli BR, *et al.* Spirometric variability in smokers: transitions in COPD diagnosis in a five-year longitudinal study. *Respir Res* 2016;17:147.
2. Wan ES, Castaldi PJ, Cho MH, Hokanson JE, Regan EA, Make BJ, *et al.*; COPDGene Investigators. Epidemiology, genetics, and subtyping of preserved ratio impaired spirometry (PRISm) in COPDGene. *Respir Res* 2014;15:89.
3. Aaron SD, Tan WC, Bourbeau J, Sin DD, Loves RH, MacNeil J, *et al.*; Canadian Respiratory Research Network. Diagnostic instability and reversals of chronic obstructive pulmonary disease diagnosis in individuals with mild to moderate airflow obstruction. *Am J Respir Crit Care Med* 2017;196:306–314.

Copyright © 2018 by the American Thoracic Society

Developing Clinical Guidelines

To the Editor:

We read with pleasure the study by Schoenberg and colleagues (1) showing that the guideline emperor has no clothes. We have served on many a guideline committee and have watched with dismay how expert opinion has been made subservient to the false god of systematic review. This process assumes a profound lack of scholarship, suggesting we have missed crucial papers in the field that we live and breathe. As Schoenberg points out, Patient Intervention Comparison Outcome (PICO) and Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) have not been subjected to the rigorous review that they themselves require. The best exemplar is the comparison of meta-analysis with subsequent large randomized controlled trial by LeLorier and colleagues (2), where simple data aggregation was little better than flipping a coin in predicting true clinical outcome. The design of clinical trials needs careful consideration, as has repeatedly been pointed out by Sami Suissa, most recently in regard to the effect of the run-in period (3), and this is not something that is considered in GRADE.

What was particularly disappointing was the commentary by Krishnan (4), who suggests that further studies are needed. We suggest it is the methodology industry that needs to urgently provide evidence of its worth before working clinicians are robbed of timely authoritative expert opinion on the recent thinking in the field. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Alyn H. Morice, M.D., F.R.C.P.
Simon Hart, Ph.D., F.R.C.P.
University of Hull
Hull, United Kingdom

Eva Millqvist, M.D., Ph.D.
University of Gothenburg
Gothenburg, Sweden

References

1. Schoenberg NC, Barker AF, Bernardo J, Deterding RR, Ellner JJ, Hess DR, *et al.* A comparative analysis of pulmonary and critical care medicine guideline development methodologies. *Am J Respir Crit Care Med* 2017;196:621–627.
2. LeLorier J, Grégoire G, Benhaddad A, Lapierre J, Derderian F. Discrepancies between meta-analyses and subsequent large randomized, controlled trials. *N Engl J Med* 1997;337:536–542.
3. Suissa S. Run-in bias in randomised trials: the case of COPD medications. *Eur Respir J* 2017;49:1700361.
4. Krishnan JA. Developing clinical guidelines: 99% faster is not enough [editorial]. *Am J Respir Crit Care Med* 2017;196:543–544.

Copyright © 2018 by the American Thoracic Society

Reply to Morice *et al.*

From the Authors:

We thank Prof. Morice and colleagues for their interest in our study. We are pleased to learn that others share our concerns about the widespread expectation that the Institute of Medicine (IOM) standards for guideline creation will be adopted by guideline developers despite a lack of evidence that adherence to the IOM standards produces recommendations that yield better outcomes than less burdensome approaches.

Our study was designed to explore the importance of the systematic review in shaping guideline recommendations (1). We succeeded in replicating, with a high degree of concordance, those recommendations generated via the IOM standard process. However, the accompanying editorial by Dr. Krishnan argued that despite producing highly concordant recommendations, the Convergence of Opinion on Recommendations and Evidence (CORE) approach is inadequate to replace the IOM approach because the grades were not also highly concordant (2). His rationale was that concordance of the grades is important because grades “influence policies affecting clinical practice (e.g., performance measures and reimbursement by payers).” We would agree that a lack of concordance between the CORE grades and those derived using the Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) approach would be problematic if we had confidence in the validity of the latter. However, although studies demonstrate the reliability (replicability) of grades developed using the GRADE approach with proper training (3), the validity of those grades has been called into question (4).

Should we be concerned about discordance with a flawed comparator? Opinion is likely to be divided. We believe grades derived from the GRADE process and newcomer approaches such as the CORE process should face similar scrutiny and that neither should be awarded a preferred status until their validity is confirmed. We look forward to future explorations of both the GRADE approach and the CORE process to better determine the role of each in the creation of guidelines. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Noah C. Schoenberg, M.D.
Kevin C. Wilson, M.D.
Boston University School of Medicine
Boston, Massachusetts

References

1. Schoenberg NC, Barker AF, Bernardo J, Deterding RR, Ellner JJ, Hess DR, *et al*. A comparative analysis of pulmonary and critical care medicine guideline development methodologies. *Am J Respir Crit Care Med* 2017;196:621–627.
2. Krishnan JA. Developing clinical guidelines: 99% faster is not enough [editorial]. *Am J Respir Crit Care Med* 2017;196:543–544.
3. Mustafa RA, Santesso N, Brozek J, Akl EA, Walter SD, Norman G, *et al*. The GRADE approach is reproducible in assessing the quality of evidence of quantitative evidence syntheses. *J Clin Epidemiol* 2013;66:736–742.e5.
4. Gartlehner G, Dobrescu A, Evans TS, Bann C, Robinson KA, Reston J, *et al*. The predictive validity of quality of evidence grades for the stability of effect estimates was low: a meta-epidemiological study. *J Clin Epidemiol* 2016;70:52–60.

Copyright © 2018 by the American Thoracic Society

Might High-Frequency Oscillatory Ventilation Improve the Prognosis of More Severe Acute Respiratory Distress Syndrome? Not So Sure

To the Editor:

In their very insightful meta-analysis of high-frequency oscillatory ventilation (HFOV) during acute respiratory distress syndrome (ARDS), Meade and coworkers suggest that patients with more severe ARDS might benefit from HFOV in terms of prognosis (1). They use Pa_{O_2}/Fi_{O_2} as a marker of ARDS severity; however, the relationship of this ratio with prognosis is equivocal and debated. Interestingly, respiratory system compliance, another physiologic parameter that may also be considered at least as accurate as Pa_{O_2}/Fi_{O_2} to assess ARDS severity, did not modify the effect of HFOV on mortality. Meade and coworkers did not analyze the potential role of mean airway pressure. This pressure was much higher with HFOV than with conventional mechanical ventilation for ARDS, and correlated with mortality and barotrauma occurrence in recent studies (2). High mean airway pressure may aggravate prognosis as a result of hemodynamic compromise, but

also by increasing ventilator-induced lung injury (VILI) (2). VILI may result from insufficient recruitment (atelectrauma) or overdistension (volutrauma). The importance of the former may have received too much emphasis, because a moderate positive end-expiratory pressure (PEEP, or increase in mean airway pressure during HFOV) lessens or even abolishes it. Excessive belief in the role of atelectrauma may lead to overenthusiastic increases of PEEP during conventional ventilation or mean airway pressure during HFOV. Ferguson and Slutsky, proponents of HFOV, postulated that, “Because cyclic alveolar stretch is minimal, volutrauma can be avoided even when the mean airway pressure is set to higher levels than can be reasonably set with PEEP on conventional ventilation” (3). This assumption may have led to the use of very high mean airway pressures during HFOV. However, global and/or regional overdistension may well be the major culprit in VILI and can occur in both dynamic and static conditions (2). It would be very interesting to know whether mean airway pressure modified the effect of HFOV on mortality in this meta-analysis.

One may be concerned by the conclusion that HFOV is worth trying in patients with severest hypoxemia because it stems from a meta-analysis of studies in which conventional ventilator management was not optimal. As emphasized by the authors, muscle paralysis and (even more so) prone positioning were seldom used. These two therapeutic adjuncts have been shown to be life saving in randomized controlled trials during severe ARDS (4, 5). The authors are right when they point that a randomized controlled study of HFOV in patients with severest ARDS would be difficult (if not impossible) to implement. But before using HFOV as rescue therapy, physicians should first apply evidence-based strategies (4, 5), which are not used sufficiently at the present time (6). ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Didier Dreyfuss, M.D.
Stéphane Gaudry, M.D.
Hôpital Louis Mourier
Colombes, France
and
Université Paris Diderot
Paris, France

ORCID ID: 0000-0002-1105-6785 (S.G.).

References

1. Meade MO, Young D, Hanna S, Zhou Q, Bachman TE, Bollen C, *et al*. Severity of hypoxemia and effect of high-frequency oscillatory ventilation in acute respiratory distress syndrome. *Am J Respir Crit Care Med* 2017;196:727–733.
2. Dreyfuss D, Ricard JD, Gaudry S. Did studies on HFOV fail to improve ARDS survival because they did not decrease VILI? On the potential validity of a physiological concept enounced several decades ago. *Intensive Care Med* 2015;41:2076–2086.
3. Ferguson ND, Slutsky AS. Point: high-frequency ventilation is the optimal physiological approach to ventilate ARDS patients. *J Appl Physiol* (1985) 2008;104:1230–1231.
4. Guérin C, Reignier J, Richard JC, Beuret P, Gacouin A, Boulain T, *et al*; PROSEVA Study Group. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med* 2013;368:2159–2168.
5. Papazian L, Forel JM, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, *et al*; ACURASYS Study Investigators. Neuromuscular blockers

Originally Published in Press as DOI: 10.1164/rccm.201709-1927LE on November 1, 2017