questions related to WGS study design and interpretation, such as how to choose the optimal data analysis method(s) and even how to define a threshold for statistical significance (16). Replication in additional cohorts and functional validation of identified variants, both of which were undertaken by Radder and colleagues (8), are important steps in any genetic study and are clearly relevant to WGS studies going forward.

The NHLBI-sponsored Trans-Omics for Precision Medicine (TOPMed) project is currently performing WGS in tens of thousands of subjects across multiple heart, lung, blood, and sleep disorders, including asthma, COPD, sleep apnea, sarcoidosis, and pulmonary hypertension, along with quantitative measures of lung function (<https://www.nhlbiwgs.org>). These studies will surely identify novel genetic risk factors for these common diseases. Moreover, the data will be made publically available through the NCBI database of Genotypes and Phenotypes (dbGaP, <https://www.ncbi.nlm.nih.gov/gap/>), allowing investigators to perform additional studies as well as to develop and refine WGS data analysis methods. The next generation of respiratory genetics researchers needs to be prepared to handle this deluge of data to identify novel disease genes and targets for intervention. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Craig P. Hersh, M.D., M.P.H.
Channing Division of Network Medicine
and
Division of Pulmonary and Critical Care Medicine
Brigham and Women's Hospital and Harvard Medical School
Boston, Massachusetts

Anil Vachani, M.D., M.S.
Pulmonary, Allergy, and Critical Care Medicine Division
University of Pennsylvania Perelman School of Medicine
Philadelphia, Pennsylvania

ORCID ID: 0000-0002-1342-4334 (C.P.H.).

References

1. Cancer Genome Atlas Research Network. Comprehensive genomic characterization of squamous cell lung cancers. *Nature* 2012;489:519–525.
2. Campbell CD, Mohajeri K, Malig M, Hormozdiari F, Nelson B, Du G, Patterson KM, Eng C, Torgerson DG, Hu D, *et al.* Whole-genome sequencing of individuals from a founder population identifies candidate genes for asthma. *Plos One* 2014;9:e104396.
3. Qiao D, Lange C, Beaty TH, Crapo JD, Barnes KC, Bamshad M, Hersh CP, Morrow J, Pinto-Plata VM, Marchetti N, *et al.*; Lung GO; NHLBI Exome Sequencing Project; COPDGen Investigators. Exome sequencing analysis in severe, early-onset chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2016;193:1353–1363.
4. Stuart BD, Choi J, Zaidi S, Xing C, Holohan B, Chen R, Choi M, Dharwadkar P, Torres F, Girod CE, *et al.* Exome sequencing links mutations in PARN and RTEL1 with familial pulmonary fibrosis and telomere shortening. *Nat Genet* 2015;47:512–517.
5. Rhodes CJ, Im H, Cao A, Hennigs JK, Wang L, Sa S, Chen PI, Nickel NP, Miyagawa K, Hopper RK, *et al.* RNA sequencing analysis detection of a novel pathway of endothelial dysfunction in pulmonary arterial hypertension. *Am J Respir Crit Care Med* 2015;192:356–366.
6. Chong JX, Buckingham KJ, Jhangiani SN, Boehm C, Sobreira N, Smith JD, Harrell TM, McMillin MJ, Wiszniewski W, Gambin T, *et al.*; Centers for Mendelian Genomics. The genetic basis of Mendelian phenotypes: discoveries, challenges, and opportunities. *Am J Hum Genet* 2015;97:199–215.
7. Albert FW, Kruglyak L. The role of regulatory variation in complex traits and disease. *Nat Rev Genet* 2015;16:197–212.
8. Radder JE, Zhang Y, Gregory AD, Yu S, Kelly NJ, Leader JK, Kaminski N, Sciurba FC, Shapiro SD. Extreme trait whole-genome sequencing identifies *PTPRO* as a novel candidate gene in emphysema with severe airflow obstruction. *Am J Respir Crit Care Med* 2017;196:159–171.
9. Wain LV, Sayers I, Soler Artigas M, Portelli MA, Zeggini E, Obeidat M, Sin DD, Bossé Y, Nickle D, Brandsma CA, *et al.* Whole exome re-sequencing implicates CCDC38 and cilia structure and function in resistance to smoking related airflow obstruction. *Plos Genet* 2014;10:e1004314.
10. Vallath S, Hynds RE, Succony L, Janes SM, Giangreco A. Targeting EGFR signalling in chronic lung disease: therapeutic challenges and opportunities. *Eur Respir J* 2014;44:513–522.
11. Fuchsberger C, Flannick J, Teslovich TM, Mahajan A, Agarwala V, Gaulton KJ, Ma C, Fontanillas P, Moutsianas L, McCarthy DJ, *et al.* The genetic architecture of type 2 diabetes. *Nature* 2016;536:41–47.
12. Helgadottir A, Gretarsdottir S, Thorleifsson G, Hjartarson E, Sigurdsson A, Magnúsdóttir A, Jonasdóttir A, Kristjánsson H, Sulem P, Oddsson A, *et al.* Variants with large effects on blood lipids and the role of cholesterol and triglycerides in coronary disease. *Nat Genet* 2016;48:634–639.
13. Morrison AC, Huang Z, Yu B, Metcalf G, Liu X, Ballantyne C, Coresh J, Yu F, Muzny D, Feofanova E, *et al.* Practical approaches for whole-genome sequence analysis of heart- and blood-related traits. *Am J Hum Genet* 2017;100:205–215.
14. Zheng HF, Forgetta V, Hsu YH, Estrada K, Rosello-Diez A, Leo PJ, Dahia CL, Park-Min KH, Tobias JH, Kooperberg C, *et al.*; AOGC Consortium; UK10K Consortium. Whole-genome sequencing identifies EN1 as a determinant of bone density and fracture. *Nature* 2015;526:112–117.
15. McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernysky A, Garimella K, Altshuler D, Gabriel S, Daly M, *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res* 2010;20:1297–1303.
16. Goldstein DB, Allen A, Keebler J, Margulies EH, Petrou S, Petrovski S, Sunyaev S. Sequencing studies in human genetics: design and interpretation. *Nat Rev Genet* 2013;14:460–470.

Copyright © 2017 by the American Thoracic Society

Neutralizing Extracellular Histones in Acute Respiratory Distress Syndrome A New Role for an Endogenous Pathway

Since the publication of the double helical structure of DNA in 1953 (1), there has been an explosion of discovery about the genetic code that determines normal development and often dictates disease. The nucleotide bases that impart the DNA code have been

the focus of studies, but there is an emerging body of evidence that the histones, by which DNA is tightly wrapped, not only regulate transcription through post-translational modifications (2), but are also critical mediators of inflammatory and injury responses when

released in the extracellular spaces (3). Histones are an important member of a heterogeneous group of molecules termed damage-associated molecular patterns. Histones can be released from any dying cell, including inflammatory cells such as neutrophils, during the process of neutrophil extracellular trap (NET) formation (4). Histones are positively charged, which allows for close interactions with and wrapping of negatively charged DNA. However, when released in the extracellular space, histones bind strongly to other proteins and trigger activation of receptors (e.g., Toll-like receptors) and multiple signaling pathways (5). Purified histones, when administered systemically to animals, induce a septic shock-like state with widespread injury to the endothelial surfaces (3).

In conditions such as acute lung injury (ALI) and the acute respiratory distress syndrome (ARDS), investigations have shown that histones are released and are associated with lung inflammation and injury (6, 7). The cellular source of the histones is often unknown, except perhaps when derived from neutrophils, in which citrullination of histones is critical to the mechanism of chromatin decondensation and NET formation (8). Extracellular histones are now a hot topic in sepsis investigations, and a variety of approaches are being tested to neutralize their toxicity.

In this issue of the *Journal*, Wygrecka and colleagues (pp. 186–199) designed experiments to therapeutically target histone-mediated lung injury in mice (9). The authors tested an endogenous protein, C1-esterase inhibitor (C1INH), for its ability to bind and neutralize histones. C1INH is an acute phase protein and serine-protease inhibitor that targets multiple different pathways, including the complement pathway, coagulation pathway (factor XIIa), and the contact system protease kallikrein. It is probably best known for its deficiency in patients with hereditary angioedema (10). C1INH is heavily glycosylated and thus carries an overall negative charge, which allows it to interact with positively charged proteins like histones. The authors determined that C1INH can bind purified histones *in vitro* and ameliorate epithelial cell death, presumably by blocking histone interactions with key cell surface proteins (11, 12), although the mechanisms were not tested. In subjects with ARDS, extracellular histones and histone-C1INH complexes were detected in plasma and bronchoalveolar lavage samples, indicating an endogenous mechanism by which histones are potentially neutralized, and a mechanism that could be exploited therapeutically.

In mouse models of bleomycin-induced ALI and extracellular histone challenge, the therapeutic delivery of C1INH or an antihistone antibody reduced lung inflammation and injury. In compelling data, there was evidence for active binding of the exogenous C1INH to extracellular histones, which formed *in vivo* complexes with an associated decrease in free C1INH levels. The mechanism of protection by C1INH was determined to be dependent on its carbohydrate structure and not on its serpin activity. The source of the extracellular histones in the mouse models and human ARDS samples is likely a combination of cell death from lung stromal cells and the release of chromatin in the form of NETs. C1INH appears capable of binding to all extracellular histones, including citrullinated histones released during NETosis.

Where do we go from here? As stated by the authors, extracellular histones are released in many disease states associated

with excessive cell death, such as sepsis and ARDS. Although the molecular mechanisms by which histones trigger inflammatory responses are still being defined, it is certainly reasonable to develop therapeutic approaches to block the vicious cycle of histone-mediated cell injury that begets additional release of extracellular histones. C1INH has previously been shown to ameliorate endotoxin lung injury in dogs (13). Recently, a small study of patients with sepsis reported that C1INH improved inflammation and survival (14). Now that a novel mechanism of direct binding to histones has been defined, perhaps improvements could be made to enhance the binding to extracellular histones and to remove the apparently unneeded serpin activity, which could induce coagulation disturbances. The mechanism of protection of C1INH in injury and inflammation is complicated by its protean properties of protease inhibition, endotoxin scavenging, and inhibition of leukocyte rolling. Additional preclinical testing is needed, including postinjury therapeutic approaches and pathogen-induced lung injury and sepsis models. C1INH is already approved by the Food and Drug Administration for hereditary angioedema, which could accelerate clinical trials in sepsis or ARDS.

The focus of most ARDS clinical trials is on prevention or supportive care, and novel approaches such as this current study are needed to move the field into the molecular age of treatment. As a syndrome, ARDS is hampered by its heterogeneity, but perhaps cell death and the elaboration of extracellular histones, whether from neutrophils or not, is a common pathway that can be targeted. The positive charge of histones could be exploited to deliver negatively charged inhibitors that not only bind and neutralize histones, but are coupled to other candidate therapeutic agents to concentrate their activity to sites of injury and inflammation. ■

Author disclosures are available with the text of this article at www.atsjournals.org.

Emma Lefrançois, Ph.D.
Mark R. Looney, M.D.
Department of Medicine
University of California, San Francisco
San Francisco, California

References

- Watson JD, Crick FH. Molecular structure of nucleic acids; a structure for deoxyribose nucleic acid. *Nature* 1953;171:737–738.
- Berger SL. Histone modifications in transcriptional regulation. *Curr Opin Genet Dev* 2002;12:142–148.
- Xu J, Zhang X, Pelayo R, Monestier M, Ammollo CT, Semeraro F, Taylor FB, Esmon NL, Lupu F, Esmon CT. Extracellular histones are major mediators of death in sepsis. *Nat Med* 2009;15:1318–1321.
- Brinkmann V, Reichard U, Goosmann C, Fauler B, Uhlemann Y, Weiss DS, Weinrauch Y, Zychlinsky A. Neutrophil extracellular traps kill bacteria. *Science* 2004;303:1532–1535.
- Chen R, Kang R, Fan XG, Tang D. Release and activity of histone in diseases. *Cell Death Dis* 2014;5:e1370.
- Abrams ST, Zhang N, Manson J, Liu T, Dart C, Baluwa F, Wang SS, Brohi K, Kipar A, Yu W, et al. Circulating histones are mediators of trauma-associated lung injury. *Am J Respir Crit Care Med* 2013;187:160–169.
- Zhang Y, Wen Z, Guan L, Jiang P, Gu T, Zhao J, Lv X, Wen T. Extracellular histones play an inflammatory role in acid aspiration-induced acute respiratory distress syndrome. *Anesthesiology* 2015;122:127–139.
- Wang Y, Li M, Stadler S, Correll S, Li P, Wang D, Hayama R, Leonelli L, Han H, Grigoryev SA, et al. Histone hypercitrullination mediates

- chromatin decondensation and neutrophil extracellular trap formation. *J Cell Biol* 2009;184:205–213.
9. Wygrecka M, Kosanovic D, Wujak L, Reppe K, Henneke I, Frey H, Didiasova M, Kwapiszewska G, Marsh LM, Baal N, *et al.* Antihistone properties of C1 esterase inhibitor protect against lung injury. *Am J Respir Crit Care Med* 2017;196:186–199.
 10. Donaldson VH, Evans RR. A biochemical abnormality in hereditary angioneurotic edema: absence of serum inhibitor of C' 1-esterase. *Am J Med* 1963;35:37–44.
 11. Huang H, Evankovich J, Yan W, Nace G, Zhang L, Ross M, Liao X, Billiar T, Xu J, Esmon CT, *et al.* Endogenous histones function as alarmins in sterile inflammatory liver injury through toll-like receptor 9 in mice. *Hepatology* 2011;54:999–1008.
 12. Xu J, Zhang X, Monestier M, Esmon NL, Esmon CT. Extracellular histones are mediators of death through TLR2 and TLR4 in mouse fatal liver injury. *J Immunol* 2011;187:2626–2631.
 13. Guerrero R, Velasco F, Rodriguez M, Lopez A, Rojas R, Alvarez MA, Villalba R, Rubio V, Torres A, del Castillo D. Endotoxin-induced pulmonary dysfunction is prevented by C1-esterase inhibitor. *J Clin Invest* 1993;91:2754–2760.
 14. Igonin AA, Protsenko DN, Galstyan GM, Vlasenko AV, Khachatryan NN, Nekhaev IV, Shlyapnikov SA, Lazareva NB, Herscu P. C1-esterase inhibitor infusion increases survival rates for patients with sepsis. *Crit Care Med* 2012;40:770–777.

Copyright © 2017 by the American Thoracic Society

The Road Not Taken: Missed Opportunities in Managing Acute Exacerbations of Hypercapnic Respiratory Failure

What are the risk factors for recurrent exacerbations of hypercapnic respiratory failure, and how do we minimize them? Although there is frequently emphasis on controlling arterial blood gas tensions and reducing pulmonary infection, in this issue of the *Journal*, Adler and colleagues (pp. 200–207) have now shown the importance of systematic evaluation for comorbidities in both patients with chronic obstructive pulmonary disease (COPD) and patients without COPD who have experienced a severe hypercapnic exacerbation requiring intensive care unit (ICU) admission (1). Importantly, they have demonstrated that these comorbidities are not only underrecognized, despite patients being frequently hospitalized, but may be associated with poor outcome.

In a prospective, single-center study, the authors assessed pulmonary function, cardiac function using echocardiography, and the presence of sleep-disordered breathing using polysomnography in 78 consecutive chronic respiratory patients discharged from ICU after a hypercapnic exacerbation necessitating mechanical ventilation (invasive or noninvasive). Patients without COPD predominantly had obesity hypoventilation syndrome. Heart failure with preserved ejection fraction was found to be previously undiagnosed in 44% of patients, and hypertension had not been recognized before admission in 67%. The prevalence of severe obstructive sleep apnea (OSA) was surprisingly high in patients with COPD, at 51%, with an even greater prevalence in patients without COPD, at 81%.

Only a minority of patients with COPD had previously been diagnosed on spirometry, although a third of the total population had previously been hospitalized for respiratory failure. More than half of the patients studied had three or more morbidities (e.g., COPD, heart failure, obesity, and severe OSA). Multiple morbidities were associated with longer hospital admissions, and there was a trend to higher hospital-free survival in those with fewer comorbidities compared with those with a greater number of comorbidities.

How applicable are these results to general ICU practice and acute pulmonary care? As investigations for comorbidity were carried out after the index admission, only survivors could be studied, and it is not clear how representative they are of the total group admitted to ICU. Indeed, around 18% of the 197 patients screened for eligibility were not admitted to ICU, as the outcome was

judged likely to be poor or futile (life expectancy estimated at <3 months), so coexisting problems are likely to be even more prevalent in this group with acute hypercapnic decompensation. Median pH on admission for the total patient cohort was 7.29 (interquartile range, 7.23–7.34), and in some centers, patients with a pH >7.26 would be managed with noninvasive ventilation (NIV) on acute pulmonary wards or high-dependency units, and not admitted to ICU (2, 3). There was no difference in outcome in patients treated with NIV or invasive ventilation, but only a minority required the latter.

Although pulmonary function tests were performed in all patients, 41 of 78 patients were unable to attend a sleep laboratory for polysomnography 3 months after discharge. The members of this group were older than those who did undergo polysomnography, which may have caused an underestimate of degree of sleep-disordered breathing. In a small subset of patients, simpler respiratory polygraphy was successfully performed, and as this may be sufficient for diagnosis of OSA, polygraphy rather than polysomnography may be a more practical and cost-effective solution to maximize the diagnosis of OSA. It is notable that only 29% of patients had previously been diagnosed with OSA, and less than a third with such a finding were being treated with continuous positive airway pressure or NIV before admission.

There was a preexisting diagnosis of heart failure in 21% of patients. In the 57/78 patients who underwent echocardiography postdischarge, the most notable finding was of heart failure with preserved ejection fraction in 44% (95% confidence interval, 31–58%). Right ventricular dilation was highly prevalent, at 42% in both COPD and non-COPD groups.

A limitation of the study is that the non-COPD group primarily consisted of obese patients, and other causes of hypercapnic respiratory failure such as neuromuscular disease were excluded. However, the prevalence of obesity in the Western world is increasing rapidly, and in many countries, obesity hypoventilation syndrome is now the prime indication for long-term NIV (4–6). The authors have extensive experience in home NIV and report that 67% of patients with multiple comorbidities were discharged with home NIV or continuous positive airway pressure, and a significant number of those with fewer comorbidities also received home ventilatory