Table 2. Differentially Expressed SNPs between Patients with Fibrosing Mediastinitis and Control Subjects

SNP	Gene	A1	A2	MAF Cases	MAF Controls (BioVU)	MAF Controls (1000 Genomes)	P (BioVU)	P (1000 Genomes)
rs143105143 (missense)	MRPL44	T	G	7.27%	0%	0.04%	9.35×10^{-18}	1.23×10^{-10}
7:45122058 (missense)	NACAD	T	G	9%	0.09%	0.13%	9.29×10^{-15}	5.06×10^{-11}
rs7728764 (synonymous)	NKX2-5	T	C	5.56%	0%	0.69%	1.6×10^{-13}	2.87×10^{-4}
rs146674681 (missense)	IRX4	A	G	5.56%	0%	0.22%	1.61×10^{-13}	2.66×10^{-4}
rs57550068 (missense)	NOC3L	T	C	5.08%	0.01%	1.77%	7.53×10^{-12}	2.43×10^{-2}
rs35381228 (splice region)	ABL1	T	C	5.08%	0.03%	1.95%	5.57×10^{-11}	3.49×10^{-2}
rs78841685 (missense)	CD97	T	C	5.08%	0.03%	1.04%	5.61×10^{-11}	2.63×10^{-3}
8:143781950 (missense)	LY6K	C	T	23.64%	0.28%	—	5.82×10^{-39}	—
4:77069485 (synonymous)	NUP54	A	C	10.91%	0%	—	2.25×10^{-26}	—
11:118917318 (missense)	HYOU1	T	C	10.19%	0%	—	2.64×10^{-24}	—
16:70897062 (synonymous)	HYDIN	T	C	11.21%	0.37%	—	3.68×10^{-15}	—
16:70942706 (synonymous)	HYDIN	T	A	6.78%	0.18%	—	1.93×10^{-10}	—
rs34292278 (missense)	SMTN	T	C	6.78%	0.20%	—	4.12×10^{-10}	—
19:44377930 (synonymous)	ZNF404	T	G	7.76%	0.32%	—	5.03×10^{-10}	—
16:70917997 (missense)	HYDIN	A	C	5.08%	0.10%	—	9.61×10^{-9}	—
rs74128855 (missense)	SYT15	A	G	5.08%	0.12%	—	2.46×10^{-8}	—

Definition of abbreviation: MAF = minor allele frequency.

For the BioVU cohort, see Reference 8. For the 1000 Genomes Project, see Reference 10.

Lung Gene Expression Analysis web portal (11), we found that all the genes in Table 2 except *CD97* were expressed in the human lung, but only *MRPL44* (mitochondrial ribosomal protein L44), *NOC3L* (NOC3-like DNA replication regulator), and *ABL1* (ABL proto-oncogene 1, nonreceptor tyrosine kinase) were expressed by lung immune cells.

The small sample size of our case cohort and the lack of a replication cohort created significant limitations for this study and reflect the rarity of this disease. These data should be viewed as exploratory, pending replication in an independent cohort. However, our results suggest that FM may have a genetic basis and open the door for future investigations into the genetic underpinnings of this enigmatic disease. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Bradley W. Richmond, M.D., Ph.D.
Yan Guo, Ph.D.
Chang Yu, Ph.D.
Peggy L. Kendall, M.D.
Wendi R. Mason, M.S.N.
Joy D. Cogan, Ph.D.
James E. Loyd, M.D.
Vanderbilt University Medical Center
Nashville, Tennessee

References

- Edwards PQ, Billings EL. Worldwide pattern of skin sensitivity to histoplasmin. *Am J Trop Med Hyg* 1971;20:288–319.
- Edwards LB, Acquaviva FA, Livesay VT, Cross FW, Palmer CE. An atlas of sensitivity to tuberculin, PPD-B, and histoplasmin in the United States. *Am Rev Respir Dis* 1969;99:1–132.
- Goodwin RA, Loyd JE, Des Prez RM. Histoplasmosis in normal hosts. *Medicine (Baltimore)* 1981;60:231–266.
- Goodwin RA, Nickell JA, Des Prez RM. Mediastinal fibrosis complicating healed primary histoplasmosis and tuberculosis. *Medicine (Baltimore)* 1972;51:227–246.
- Peikert T, Shrestha B, Aubry MC, Colby TV, Ryu JH, Sekiguchi H, et al. Histopathologic overlap between fibrosing mediastinitis and IgG4-related disease. *Int J Rheumatol* 2012;2012:207056.

- Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, et al; 1000 Genomes Project Consortium. A global reference for human genetic variation. *Nature* 2015;526:68–74.
- Sherrick AD, Brown LR, Harms GF, Myers JL. The radiographic findings of fibrosing mediastinitis. *Chest* 1994;106:484–489.
- Roden DM, Pulley JM, Basford MA, Bernard GR, Clayton EW, Balser JR, et al. Development of a large-scale de-identified DNA biobank to enable personalized medicine. *Clin Pharmacol Ther* 2008;84:362–369.
- Richmond BW, Guo Y, Yu C, Mason WR, Cogan J, Loyd JE. A genome-wide association study of 58 patients with fibrosing mediastinitis [abstract]. *Am J Respir Crit Care Med* 2017;195:A3237.
- The 1000 Genomes Project Consortium. A map of human genome variation from population-scale sequencing. *Nature* 2010;467:1061–1073.
- Du Y, Kitzmiller JA, Sridharan A, Perl AK, Bridges JP, Misra RS, et al. Lung Gene Expression Analysis (LGEA): an integrative web portal for comprehensive gene expression data analysis in lung development. *Thorax* 2017;72:481–484.

Copyright © 2018 by the American Thoracic Society

Accuracy of Models to Identify Lung Nodule Cancer Risk in the National Lung Screening Trial

To the Editor:

Clinical models to estimate the probability of cancer in pulmonary nodules detected incidentally on imaging tests are used both in the clinic setting and for research purposes (1). Recently, the Brock model was developed for nodules detected by low-dose computed tomography screening (2). Patients with incidentally detected nodules and those with screening-detected nodules have different risk profiles because those screened for lung cancer must have a number of risk factors for lung cancer, whereas the patients with incidentally detected nodules do not.

Author Contributions: Conception and design: V.S.N., V.S., and M.K.G.; analysis and interpretation: V.S.N., V.S., M.D., and M.K.G.; and drafting the manuscript for important intellectual content: V.S.N., V.S., M.D., and M.K.G.

Originally Published in Press as DOI: 10.1164/rccm.201708-1632LE on October 24, 2017

Existing cancer prediction models for patients with incidentally or screening-detected nodules have not been compared in a large group of screened patients, and which ones should be used in this population has not been determined. We performed a secondary analysis of data from the National Lung Screening Trial (NLST) to compare the accuracy and calibration of the Brock model, developed for patients with nodules detected by screening, with two other models developed for patients with incidentally detected nodules (2–4). Some of the results of these studies have been previously reported in the form of an abstract (5).

Methods

We examined patients with a positive baseline screening test result in the low-dose computed tomography group of the NLST ($n = 26,722$) and compared the discrimination/accuracy of existing models for identifying cancerous nodules (6). We limited the analysis to nodules detected on baseline screening because NLST data for subsequent rounds of annual screening did not distinguish between a new nodule or an existing nodule seen previously on the baseline scan. We assumed that cancers detected before the next round of annual screening (T1) after the baseline low-dose computed tomography scan (T0) were attributed to the largest nodule present on the baseline screening exam.

We excluded all patients diagnosed with cancer after T1 ($n = 341$), patients with masses larger than 3 cm in maximum diameter ($n = 108$), and patients with incomplete data ($n = 169$). For patients with

multiple nodules on the baseline screening exam, we used the dimensions of the largest nodule diameter size when calculating the probability of lung cancer. Relevant variables were extracted from the participant and spiral computed tomography abnormalities data files of the NLST data set. Patients were excluded from the Brock model cohort if data were not available for a complete calculation (Table 1).

We calculated the accuracy of the full and parsimonious Brock models that used spiculation as a covariate, the Mayo Clinic (Mayo) model and the Veterans Affairs (VA) model, to identify cancer for patients with nodules 8 mm or larger (primary analysis) and nodules 4 mm or larger (secondary analysis), applying the published β coefficients for each model to each patient in the sample (1, 2). We chose the 8 mm or larger threshold because probability assessment is critically important in the evaluation of patients with nodules of this size. To assess model discrimination (accuracy), we calculated the area under the receiver operating characteristic curve (AUC) for each model (SAS v9.4). Differences in AUCs were compared using the nonparametric approach of DeLong and colleagues (7). To assess calibration, we plotted the predicted and observed frequencies of cancer (MS Excel). All tests were two-sided with an α level of 0.05.

Results

Among patients with nodules at least 8 mm on baseline screening, those diagnosed with cancer after T0 and before T1 were significantly older, were more likely to have a history of cancer, and had nodules that were larger and more likely to be solid or

Table 1. Characteristics of the Analytic Cohorts by Cancer Diagnosis

Variable for Modeling	≥8 mm			≥4 mm		
	All Patients	Benign Nodules	Cancer	All Patients	Benign Nodules	Cancer
<i>n</i>	2,196	2,003	193	6,568	6,348	220
Age, yr* ^{†‡§}	62.6 (5.2)	62.5 (5.2)	64.0 (5.6)	62.0 (5.2)	61.8 (5.1)	63.7 (5.7)
Sex, male, <i>n</i> (%) [§]	1,307 (59.5)	1,202 (60.0)	105 (54.4)	3,811 (58)	3,690 (58)	121 (55)
Smoking history, <i>n</i> (%) ^{†‡}	2,196 (100)	2,003 (100)	193 (100)	6,845 (100)	6,348 (100)	220 (100)
Former	1,166 (53.1)	1,074 (53.6)	92 (47.7)	3,337 (51)	3,236 (51)	101 (46)
Current	1,030 (46.9)	929 (46.4)	101 (52.3)	3,231 (49)	3,112 (49)	119 (54)
Years quit (if former smoker) [†]	7.2 (4.7)	7.3 (4.8)	6.6 (4.5)	7.2 (4.7)	7.2 (4.7)	6.4 (4.5)
History of cancer, <i>n</i> (%) [†]	103 (4.7)	88 (4.4)	15 (7.8)	244 (3.7)	228 (3.6)	16 (7.3)
Family history of lung cancer, <i>n</i> (%) [*]	518 (23.6)	467 (23.3)	51 (26.4)	1,462 (22.3)	1,402 (22.1)	60 (27.3)
Emphysema, <i>n</i> (%) [§]	247 (11.3)	226 (11.3)	21 (10.9)	575 (8.8)	553 (8.7)	22 (10.0)
Nodule size, mm* ^{†‡}	12.1 (4.9)	11.6 (4.5)	17.2 (6.1)	7.6 (4.3)	7.4 (3.9)	15.9 (6.8)
Nodule density, <i>n</i> (%) [*]						
Nonsolid	458 (20.9)	443 (22.1)	15 (7.8)	1,012 (15.4)	993 (15.6)	19 (8.6)
Semisolid/part solid	176 (8.0)	156 (7.8)	20 (10.4)	338 (5.2)	318 (5.0)	20 (9.1)
Solid	1,500 (68.3)	1,348 (67.3)	152 (78.8)	4,879 (74.3)	4,706 (74.1)	173 (78.7)
Upper lobe location, <i>n</i> (%) ^{†‡}	935 (42.6)	819 (40.9)	116 (60.1)	2,546 (38.8)	2,414 (38.0)	132 (60.0)
Spiculation, <i>n</i> (%) ^{†‡§}	487 (22.2)	373 (18.6)	114 (59.1)	748 (11.4)	630 (9.9)	118 (53.6)
Nodule count [*]	2.6 (1.8)	2.6 (1.8)	2.2 (1.4)	2.4 (1.6)	2.4 (1.6)	2.1 (1.4)

Age reported is age at randomization. Continuous variables are reported as mean (SD). The years quit variable is missing 5 and 13 observations in ≥8 mm and ≥4 mm cohorts, respectively. The family history (defined as parents, siblings, or child having lung cancer) variable is missing 36 and 105 observations in ≥8 mm and ≥4 mm cohorts, respectively. The nodule density variable is missing 62 and 354 observations in ≥8 mm and ≥4 mm cohorts, respectively. Differences of distributions in patients with and without lung cancer for categorical variables were assessed by chi-squared tests or Fisher's exact tests, as appropriate. Student's *t* test was used to compare distributions for continuous variables.

*Full Brock model variables.

[†]Mayo model variables.

[‡]Veterans Affairs model variables.

[§]Parsimonious Brock model variables.

^{||} $P < 0.01$ between groups.

Table 2. Model Accuracy by Probability Threshold for Identifying Lung Cancer in the First Year after Baseline Screen

Model and Probability (%)	Sp	Sn	PPV	NPV	–LR	+LR	Total <i>n</i>	Cancer [<i>n</i> (%)]	AUC (95% CI)
Nodules ≥ 8 mm									
Brock (full)							2,100	187 (8.9)	0.84 (0.81–0.86)
1	0.03	1.0	0.09	1.00	0.00	1.03	59		
5	0.47	0.95	0.15	0.99	0.11	1.79	907		
10	0.68	0.81	0.20	0.97	0.28	2.53	1,335		
15	0.79	0.71	0.25	0.97	0.37	3.38	1,567		
20	0.85	0.61	0.28	0.96	0.46	4.07	1,696		
65	1.0	0.02	0.39	0.91	0.98	6.67	2,091		
Mayo							2,196	193 (8.8)	0.82 (0.79–0.85)
5	0.01	0.99	0.09	0.91	1.00	1.00	24		
10	0.33	0.95	0.12	0.99	0.15	1.42	661		
15	0.51	0.90	0.15	0.98	0.20	1.84	1,048		
20	0.65	0.83	0.19	0.98	0.26	2.37	1,340		
65	0.96	0.24	0.37	0.93	0.79	6.00	2,079		
VA							2,196	193 (8.8)	0.77 (0.73–0.80)
12	0	1.0	0.09	ND	ND	1.00	1		
15	0.01	1.0	0.09	1.00	0.00	1.01	21		
20	0.05	0.98	0.09	0.96	0.40	1.03	95		
65	0.92	0.37	0.31	0.94	0.68	4.63	1,959		
Brock (parsimonious)							2,196	193 (8.8)	0.83 (0.80–0.86)
5	0.43	0.95	0.14	0.99	0.12	1.67	862		
10	0.66	0.83	0.19	0.98	0.26	2.44	1,347		
15	0.78	0.70	0.24	0.96	0.38	3.18	1,625		
20	0.84	0.60	0.27	0.96	0.48	3.75	1,769		
65	1.0	0.02	0.39	0.91	0.98	6.67	2,186		
Nodules ≥ 4 mm									
Brock (full)							6,127	211 (3.4)	0.90 (0.87–0.92)
1	0.55	0.94	0.07	1.00	0.11	2.08	3,259		
5	0.82	0.84	0.14	0.99	0.19	4.76	4,900		
10	0.90	0.72	0.20	0.99	0.31	7.20	5,362		
15	0.93	0.63	0.24	0.99	0.40	9.32	5,594		
20	0.95	0.54	0.28	0.98	0.48	10.8	5,723		
65	1.0	0.02	1.00	0.97	0.98	ND	6,118		
Mayo							6,568	220 (3.3)	0.88 (0.85–0.90)
3	0.02	1.0	0.03	1.00	0.00	1.02	106		
5	0.27	0.98	0.04	1.00	0.07	1.34	1,731		
10	0.64	0.88	0.08	0.99	0.19	2.44	4,069		
15	0.80	0.81	0.12	0.99	0.24	4.05	5,103		
20	0.86	0.75	0.16	0.99	0.29	5.36	5,510		
65	0.99	0.21	0.42	0.97	0.80	21.0	6,451		
VA							6,568	220 (3.3)	0.85 (0.82–0.87)
7	0.01	1.0	0.03	1.00	0.00	1.01	1		
10	0.01	1.0	0.03	1.00	0.00	1.01	55		
15	0.07	0.99	0.04	1.00	0.14	1.06	466		
20	0.24	0.97	0.04	1.00	0.13	1.28	1,560		
65	0.97	0.32	0.27	0.98	0.70	10.7	6,331		
Brock (parsimonious)							6,568	220 (3.3)	0.89 (0.87–0.91)
1	0.56	0.93	0.07	1.00	0.13	2.11	3,597		
5	0.81	0.84	0.13	0.99	0.20	4.42	5,198		
10	0.89	0.73	0.19	0.99	0.30	6.63	5,719		
15	0.93	0.61	0.23	0.99	0.42	8.71	5,997		
20	0.95	0.53	0.27	0.98	0.49	10.6	6,141		
65	1.0	0.02	1.00	0.97	0.98	ND	6,558		

Definition of abbreviations: AUC = area under the receiver operating characteristic curve; CI = confidence interval; LR = likelihood ratio; ND = not defined (undefined) based on calculation; NPV = negative predictive value = $[\text{specificity} \times (1 - \text{prevalence})] / [(1 - \text{sensitivity}) \times \text{prevalence} + \text{specificity} \times (1 - \text{prevalence})]$; PPV = positive predictive value = $(\text{sensitivity} \times \text{prevalence}) / [(\text{sensitivity} \times \text{prevalence}) + (1 - \text{specificity}) \times (1 - \text{prevalence})]$; Sn = sensitivity; Sp = specificity; VA = Veterans Affairs.

Number in total *n* column represents all nodules up to the risk threshold specified.

part-solid, spiculated, and located in the upper lobe compared with patients who were never diagnosed with cancer (Table 1).

Among 2,196 patients with nodules at least 8 mm, there were 2,003 benign nodules and 193 cancers (8.8% prevalence); sufficient data were available to evaluate the full Brock model in a slightly smaller sample of 2,100 patients: 1,913 with benign nodules and 187 with cancer (Table 2). By unpaired AUC analysis, the full Brock (AUC, 0.84; 95% confidence interval [95% CI], 0.81–0.86) and Mayo (AUC, 0.82; 95% CI, 0.79–0.85) models had similar accuracy ($P = 0.41$), as did the full and parsimonious (AUC, 0.83; 95% CI, 0.80–0.86) Brock models ($P = 0.79$), but the VA model (AUC, 0.77; 95% CI, 0.73–0.80) was less accurate than the full Brock model ($P = 0.003$).

Restricting our analysis to only solid nodules at least 8 mm in size showed that Brock ($n = 1,480$; AUC, 0.84; 95% CI, 0.81–0.87) and Mayo ($n = 1,500$; AUC, 0.83; 95% CI, 0.80–0.87) had similar accuracy, whereas the VA model ($n = 1,500$; AUC, 0.78, 95% CI, 0.73–0.82) was less accurate ($P = 0.02$).

Among 6,568 patients with nodules of at least 4 mm, there were 6,348 benign nodules and 220 cancers (3.3% prevalence); sufficient data were available to evaluate the full Brock model in a smaller sample of 6,127 patients: 5,916 with benign nodules and 211 with cancer (Table 2). Once again, the full Brock and Mayo models had similar accuracy ($P = 0.21$), as did the full and parsimonious Brock models ($P = 0.68$), but the VA model was less accurate ($P = 0.006$).

Across a wide range of probability thresholds (1–20%), all the models had excellent negative predictive values, reflecting the low prevalence of cancer in the sample, but positive predictive values were poor (Table 2). When comparing expected with observed frequencies of cancer across deciles of risk, the full Brock model had similar numbers of predicted and observed cancers, whereas the Mayo and VA models both overestimated the risk for cancer (figures available on request).

Discussion

Among patients with a nodule size 8 mm or larger or nodule size 4 mm or larger on baseline screen in the NLST, we found that the Brock model was most accurate for identifying cancerous nodules, and that the Brock model was reasonably well calibrated in this external validation of its performance. The overestimation of probabilities by the Mayo and VA models likely is a result of their development in study populations with a much higher prevalence of cancer compared with the screening population.

For the purpose of clinical decision-making in the setting of lung cancer screening, we suggest that using either of the Brock models is preferred over other models, as it is well calibrated and can be more generally applied to small and nonsolid nodules. Because the trade-off between sensitivity and specificity varies in distinct ways for each model according to the threshold used to define a positive result, researchers should be cognizant of the underlying prevalence of cancer in their study population when using additional biomarkers to “rule-out” or “rule-in” disease. Our findings have implications for the evaluation of novel molecular or imaging biomarkers, as it is important to demonstrate that the marker provides incremental value above and beyond the best available clinical tools (8). ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Viswam S. Nair, M.D., M.S.
University of South Florida Morsani College of Medicine
Tampa, Florida

and
Moffitt Cancer Center and Research Institute
Tampa, Florida

Vandana Sundaram, M.P.H.
Manisha Desai, Ph.D.
Stanford University School of Medicine
Stanford, California

Michael K. Gould, M.D., M.S.
Kaiser Permanente Research
Pasadena, California

References

- Schultz EM, Sanders GD, Trotter PR, Patz EF Jr, Silvestri GA, Owens DK, et al. Validation of two models to estimate the probability of malignancy in patients with solitary pulmonary nodules. *Thorax* 2008;63:335–341.
- McWilliams A, Tammemagi MC, Mayo JR, Roberts H, Liu G, Soghrati K, et al. Probability of cancer in pulmonary nodules detected on first screening CT. *N Engl J Med* 2013;369:910–919.
- Swensen SJ, Silverstein MD, Ilstrup DM, Schleck CD, Edell ES. The probability of malignancy in solitary pulmonary nodules: application to small radiologically indeterminate nodules. *Arch Intern Med* 1997;157:849–855.
- Gould MK, Ananth L, Barnett PG; Veterans Affairs SNAP Cooperative Study Group. A clinical model to estimate the pretest probability of lung cancer in patients with solitary pulmonary nodules. *Chest* 2007;131:383–388.
- Nair VS, Sundaram V, Desai M, Gould MK. Lung nodule cancer risk in the national lung screening trial [abstract]. *Am J Respir Crit Care Med* 2017;95:A7054.
- Aberle DR, Adams AM, Berg CD, Black WC, Clapp JD, Fagerstrom RM, et al.; National Lung Screening Trial Research Team. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* 2011;365:395–409.
- DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;44:837–845.
- Steyerberg EW, Vickers AJ, Cook NR, Gerds T, Gonen M, Obuchowski N, et al. Assessing the performance of prediction models: a framework for traditional and novel measures. *Epidemiology* 2010;21:128–138.

Copyright © 2018 by the American Thoracic Society

Dual Bronchodilation Response by Exacerbation History and Eosinophilia in the FLAME Study

To the Editor:

The Global Initiative for Chronic Obstructive Lung Disease (GOLD) 2017 strategy notes that exacerbation history is the best predictor of future exacerbations in patients with chronic obstructive pulmonary disease (COPD) (1). In addition, blood eosinophil levels

Supported by Novartis Pharma AG.

Author Contributions: All authors contributed to the design and conduct of this *post hoc* analysis. The authors take full responsibility for the scope, direction, and content of the manuscript, and editorial decisions relating to it; were involved at all stages of development; and have approved the submitted letter to the editor.

Originally Published in Press as DOI: 10.1164/rccm.201709-1822LE on November 3, 2017