Published in final edited form as:

Arch Sex Behav. 2018 October ; 47(7): 1959–1968. doi:10.1007/s10508-017-1061-y.

Depression, loneliness, and sexual risk-taking among HIV-negative/unknown men who have sex with men in China

Xiaoyou Su, PhD¹, A. Ning Zhou, MD², Jianjun Li, MPH³, Ling-en Shi, MPH³, Xiping Huan, MPH³, Hongjing Yan, MD³, and Chongyi Wei, DrPH⁴

¹School of Public Health, Peking Union Medical College, Chinese Academy of Medical Sciences, Beijing, China

²Department of Psychiatry, University of California – San Francisco, CA, USA

³Jiangsu Provincial Center for Disease Control and Prevention, Nanjing, Jiangsu, China

⁴Department of Epidemiology and Biostatistics & Global Health Sciences, University of California – San Francisco, CA, USA

Abstract

Background—While research conducted among men who have sex with men (MSM) in high-income countries has demonstrated that negative mental health is one of the significant drivers of HIV infection, few studies have examined the status of mental health among MSM in China. We sought to describe depression and loneliness and identify their correlates among Chinese MSM.

Methods—A cross-sectional study was conducted among HIV-negative or unknown status MSM in 2014. Time-location sampling and online convenience sampling methods were employed. Depression was measured via a short version of CES-D (CES-D 10). Loneliness was measured from a single item in CES-D 10. Multivariable logistic regressions were conducted to identify independent correlates of depression and loneliness.

Results—A total of 507 individuals participated in the study. Of them, 26.8% and 35.5% reported moderate-to-severe symptoms of depression and feeling lonely, respectively. Depressed participants were more likely to have a sense of hopelessness for the future (AOR = 3.2, 95% CI: 1.9, 5.2) and report higher levels of internalized homophobia (AOR = 2.3, 95% CI: 1.47, 3.67). Participants who reported feeling lonely were more likely to have had condomless receptive anal intercourse in the past six months (AOR = 1.7, 95% CI: 1.1, 2.6) and feel hopeless for the future (AOR = 2.4, 95% CI: 1.6, 3.7).

Conclusions—MSM in China have significant rates of depression and loneliness. HIV-prevention efforts should address the mental health needs of Chinese MSM such as providing safe environments for social support and role models.

Correspondence: Chongyi Wei, 50 Beale Street, Suite 1300, San Francisco, CA 94105, US. Phone: 1-415-597-9182; Fax: 1-415-597-9213; Chongyi.Wei@ucsf.edu.

There are no conflicts of interest

Keywords

MSM; HIV; depression; loneliness; mental health

INTRODUCTION

Men who have sex with men (MSM) are one of the most affected groups by the HIV/AIDS epidemic across the world (Neme et al., 2015). Globally, the MSM population represents 78% of all men who have been newly diagnosed with HIV (Centers for Disease Control and Prevention, 2015). In the United States, MSM comprise more than half of all newly infected people with HIV annually and continue to be the largest group of individuals infected with HIV (Hall et al., 2008). In China, the prevalence of HIV among MSM has also increased substantially in recent years, and 29.4% of new HIV infections in 2011 were through male-to-male contact (Wu et al., 2015). While numerous behavioral interventions focused on reducing sexual risk of MSM in China have been adopted in the past two decades, effective, long-term and sustainable behavioral change will be hard to obtain without considering psychosocial and mental health factors.

Psychosocial and mental health issues such as depression contribute significantly to HIV risk in the MSM population. For example, gender role-related stress, homophobia, gay-related stigma and discrimination, poor social support, and high rates of drug & alcohol use lead to riskier sexual behavior and an increased prevalence of HIV among this population (Safren et al., 2010). Moreover, MSM suffer from a range of mental health issues, most notably depression. Clinical depression is characterized by low mood, loss of interest or pleasure in most activities, feelings of worthlessness or inappropriate guilt, changes in appetite, difficulty sleeping, loss of energy, difficulty concentrating, appearing restless or slowed down, and thoughts of suicide (Ophir et al., 2017). The prevalence of moderate-to-high depressive symptoms in MSM varies in different countries, ranging from 11% in India (Tomori et al., 2016), to 24.6% in the United States (Dyer et al., 2015), to 44.5% in Peru (Ferro et al., 2015), to 46.3% in Tanzania (Ahaneku et al., 2016). In China, the prevalence of depression among MSM is 46.1% (Yan et al., 2014), and among the general population is 3% (Chen et al., 2007). Furthermore, depression has been shown to affect sexual functioning and increase the risk of condomless sex and substance use among MSM (Ahaneku et al., 2016; Watkins et al., 2016). Finally, depression can be a barrier to accessing HIV prevention services and care (Du Bois and McKirnan, 2012; Wei et al., 2016).

Loneliness is also common among MSM, many of whom do not disclose their sexual orientation to family and friends for fear of rejection. Loneliness is an unpleasant, subjective experience that results from the lack of quantity or quality of social relationships (Peplau and Perlman, 1982). Loneliness may stem from minority stress (Kuyper and Fokkema, 2010), perceived and enacted stigma, lack of social support (Hubach et al., 2015), and loss of friends and intimate partners due to HIV/AIDS (Dispenza et al., 2015). Moreover, loneliness has been shown to be a barrier to receiving HIV testing and counseling among MSM (Neme et al., 2015). Similar to depression, feeling lonely is associated with higher rates of condomless sex and increased risk for HIV infection (Doskoch, 2011); in fact,

condomless sex was reported to be a coping mechanism for lonely and isolated MSM to decrease their feelings of loneliness (Martin and Knox, 1997; Shernoff, 2006). When not directly using sexual behavior as a coping strategy, some lonely MSM may engage in activities that indirectly lead to unsafe sexual behaviors; for example, lonely MSM may turn to substances to escape feelings of loneliness, leading to lowered sexual inhibitions, resulting in increased risky sexual behavior (Kurtz, 2005).

Not only do psychosocial and mental health factors disproportionately affect MSM, they interact to lead to elevated HIV-risk behavior among MSM. In fact, the more psychosocial and mental health issues an individual endorses, the greater his risk for both participating in risky sexual behaviors (unprotected anal intercourse, multiple sex partners) and becoming infected with HIV. These co-occurring psychosocial health problems may actually be driving the HIV epidemic among MSM (Mustanski et al., 2007; Stall et al., 2003). Moreover, a few theoretical models have been developed to explain the connection between negative affective states and HIV risk behaviors, such as the AIDS Risk Reduction Model (ARRM) (Catania et al., 1990), the Information-Motivation-Behavioral Skill Model of HIV prevention behavior (Fisher et al., 2009), the Integrative Model of Behavioral Prediction (Fishbein, 2009), and the Loneliness and Sexual Risk Model (LSRM) (Torress and Gore-Felton, 2007). As an example, the LSRM has shown that loneliness is associated with increased risky sexual behavior and that this relationship is mediated by the influence of substance use and compulsive behavior (Torress and Gore-Felton, 2007). Additionally, a modified version of the LSRM suggests that internalized homophobia mediates the relationship between loneliness and sexual behavior among MSM (DeLonga et al., 2011).

While the impact of depression and loneliness on sexual risk-taking among MSM has been well documented in the mental health and HIV-prevention literature in the past decade from developed countries, their association with risky sexual behaviors is poorly understood among MSM in Asian countries such as China. One study in China examining social support and depressive symptoms found the prevalence of depression among general MSM of various serostatuses was 46.1% (Yan et al., 2014). Another study discovered the prevalence of depression among a group of HIV-positive MSM to be 36.0%, and that the level of depressive symptoms had an impact on whether participants initiated antiretroviral therapy (Tao et al., 2017). However, there are still no studies addressing the relationship between depression, loneliness and risky sexual behavior among MSM in China. Moreover, many of the existing studies were conducted among MSM who are HIV-positive, so it is unclear whether these findings apply to HIV-negative MSM. Exploring the connection between depression, loneliness, and risky sexual behavior among HIV-negative and unknown status MSM in China may shed additional light on how to best reduce HIV transmission in this vulnerable population.

In this study, we sought to describe the prevalence of depression and loneliness among a diverse sample of HIV-negative/unknown MSM in China, and examined their associations with demographic factors, psychosocial conditions, and HIV-related risk behaviors.

METHODS

Study Design and Recruitment

We conducted a cross-sectional survey study of MSM in Jiangsu Province, China between November and January 2014. This survey was the baseline data collection for a longitudinal cohort study designed to assess the effects of an intervention to increase HIV testing uptake among MSM in the province. To be eligible for the study, participants had to meet the following criteria: 1) be biologically male; 2) be at least 18 years of age; 3) be living in Jiangsu province; 4) have had oral or anal sex with another male in the past year; and 5) self-report being HIV-negative or having unknown HIV status. In addition, since this was part of a cohort study, we asked eligible participants to provide their contact information during the informed consent process. We employed two sampling methods to recruit participants: time-location sampling (TLS) and online convenience sampling.

Time-location sampling (TLS): TLS is used to sample MSM populations through the creation of a sampling frame that comprises a universe of venues, days, and time periods where and when the population can be found to congregate. It has been described in detail elsewhere (Zhao et al., 2015). Briefly, during the formative phase, we constructed an up-to-date universe or sampling frame of venues frequented by MSM, including bars, dance clubs, bathhouses, and cruising areas, as well as the days and time periods of attendance in the Chinese city of Nanjing. From the roster of all possible venue-day-time (VDT) periods, a random sample of VDT periods were drawn. At the randomly selected VDT, all potentially eligible subjects were counted. Men entering/exiting the venue or crossing a predetermined line were intercepted, assessed for eligibility, and invited to participate.

During the assessment, recruiters briefly described the study to the men and asked if they were willing to participate. Men who had not previously participated were referred to an interviewer who screened for eligibility. Eligibility screening occurred in a private area of the venue or in a designated interviewing space near the venue. In addition to the above eligibility criteria, participants recruited through TLS also had to be consecutively approached by the staff at the randomly selected VDT (i.e., they could not approach staff on their own or at a later time). Once participants were determined eligible, staff discussed informed consent using a tablet computer and addressed questions. To consent to the study, participants had to click the “Agree” button on the electronic informed consent form. Staff then oriented participants to the tablet computer-assisted interview. Once participants were familiar with the operation of the tablet computer, they completed a self-administered survey.

Participants were recruited during 23 randomly selected VDT periods from November to December 2013. Staff enumerated 777 men from 10 venues, consecutively approached 478, screened 342 (71.5% eligibility determination), found 290 to be eligible (84.8% eligibility), and 261 (90.0% participation) consented to the study and completed the survey.

Online convenience sampling: We posted our study advertisement on the most popular provincial gay-oriented website to invite online participation in our study. By clicking on the

advertisement, interested MSM were taken to an Internet-based survey webpage. The eligibility screening and informed consent procedures implemented for the online survey component were the same as the afore-mentioned TLS component. To prevent duplicate participation, a feature within the survey software program was enabled which would not allow potential participants to access the survey more than once from the same IP address. Furthermore, our staff double-checked the contact information provided by participants before giving out incentives for completing the survey. From mid-November 2013 to mid-January 2014, 985 potential participants clicked on our online survey advertisement or link. Of these, 941 entered the eligibility screening page and 823 answered all eligibility questions (87.5% eligibility determination), 592 met the eligibility criteria (71.9% eligibility), and 271 (45.8% participation) consented to the study and completed the survey.

Participants who completed the survey received an incentive in the form of a 50RMB ($\approx$ 8USD) pre-paid cell-phone card. The study was approved by the University of California – San Francisco’s Committee on Human Research and Jiangsu Provincial Center for Disease Control and Prevention’s Institutional Review Board.

Measures

Outcome variables

Depression (CES-D10): The Chinese version of 10-item CES-D depression scale was used to screen depressive symptomatology within the past week. It has been validated among a clinically depressed population as well as a non-clinical, general population in China (Yu et al., 2013; Kohout et al., 1993). Scores of the CES-D 10 range from 0 to 30, with higher scores indicating more depressive symptoms (Cronbach’s $\alpha = 0.811$). Those who had scores above 20 were classified as having moderate-to-severe depressive symptoms.

Loneliness: Loneliness was measured from a single item (“I felt lonely”) in the CES-D 10 (Kohout et al., 1993). Participants who reported feeling lonely “sometimes” or “most or all of the time” were categorized as experiencing loneliness. Although the utility of a single-item loneliness measure may be called into question, due to the stigma associated with being identified as lonely, it has been widely used in previous studies in China and other countries (Zhong et al., 2016; Luo and Waite, 2014; Savikko et al., 2005) and has been proven to have good face and predictive validity (Tilvis et al., 2000). In addition, this single-question measure was the original standard against which multi-item instruments were validated. It has been shown to be highly correlated with the two most widely used loneliness assessment tools: the UCLA Loneliness Scale (Russell, 1996) and the de Jong Gierveld Loneliness Scale (de Jong Gierveld and van Tilburg, 1999). Single-item surveys have also been used in other studies to measure mood status and general health, and are considered to have the advantage of simplicity while preserving validity (DeSalvo et al., 2006; Luria, 1975).

Independent variables

Sociodemographics: Participants were asked about their age, educational level, legal marital status, whether or not they were cohabiting with a man or woman, employment status, monthly gross income, and sexual orientation.

Sexual behaviors and HIV testing: Participants reported the number of male partners they engaged in anal intercourse with in the past six months, and whether or not they consistently used condoms when engaging in insertive and/or receptive anal intercourse. Participants were also asked if they were tested for HIV in the past 12 months.

Internalized homophobia: Five out of the seven items from the Reactions to Homosexuality scale were used to measure internalized homonegativity (“I feel comfortable in gay bars,” “I feel comfortable discussing homosexuality in a public situation,” “I feel comfortable being a homosexual man,” “Homosexuality is morally acceptable to me,” “Even if I could change my sexual orientation, I wouldn’t”)(Berg et al., 2013). The face validity of the five items were confirmed by a panel including Chinese professionals and members from the MSM community. Scores of the 5-item scale ranged from 0–4, with higher cores indicating fewer experiences of homophobia. (*Cronbach’s α = 0.655*).

Future expectations: Five items from the Chinese version of the Beck Hopelessness Scale were used to measure participants’ negative expectancies for the future (e.g., “It is very unlikely that I will get any real satisfaction in the future,” “the future seems uncertain to me”) (Shek, 1993; Beck and Weissman, 1974). The individual item scores ranged from 0 to 5 (*Cronbach α = 0.732*), and the total “hopelessness score” was the sum of the scores on the individual items, with higher scores indicating more hopelessness.

ANALYSIS

Potential duplication between TLS and online participants was examined by comparing the contact information provided by participants; seven duplicated records were excluded from analysis. The analytical sample included 507 MSM participants who completed the CES-D 10. We conducted all analyses in STATA/SE 11 (College Station, Texas) with level of significance determined at a 0.05 *p-value*.

Descriptive analyses were conducted to describe characteristics of the sample, including socio-demographics, HIV-related sexual or health behavior, and psychosocial conditions. The internal consistency of the scales was assessed using Cronbach’s alpha coefficients. Bivariate analyses, using Chi-square tests, of depression and loneliness began with the full set of demographic, sexual behavior, HIV testing, hopelessness, and internalized homophobia variables. Variables that were associated with depression and loneliness at $p < 0.10$ in bivariate analysis were entered into multivariable logistic regression models.

RESULTS

We enrolled a total of 507 participants. Of these participants, 30% were between the ages of 18–25, 41.6% had college degree or above, 19.8% of the participants were married, 20.3% were cohabitating with another man, and 67.9% were self-identified as gay. 78.5% reported having prior HIV-testing at the time of interview. The percentage of condomless insertive anal intercourse (CIAI) and condomless receptive anal intercourse (CRAI) were 38.7% and 31.2%, respectively (Table 1). Of the participants, 26.8% experienced moderate-to-severe symptoms of depression (hereafter “depressed”) and 35.5% experienced loneliness in the

past week. Meanwhile, 41.2% of the participants felt hopeless and 46.2% had high levels of internalized homophobia (Table 1).

Table 2 presents the bivariate correlates of moderate-to-severe depression and loneliness among the study participants. Compared to those who were not depressed (i.e., non-to-mild), participants who had experienced moderate-to-severe depression were significantly more likely to be 18–25 years old ($p = 0.002$), have single marital status ($p = 0.079$), and be cohabiting with men ($p = 0.003$). They were significantly less likely to have been HIV tested in the past 12 months ($p = 0.058$), and more likely to feel hopeless ($p < 0.001$) and have higher levels of internalized homophobia ($p = 0.001$). Meanwhile, compared to those who had not experienced loneliness, participants who experienced loneliness were significantly more likely to be aged between 26–35 years old ($p = 0.026$) and have no main male partner ($p = 0.04$). They were significantly less likely to have been HIV tested in the past 12 months ($p = 0.04$), more likely to have CRAI ($p = 0.04$) and feel hopeless about the future ($p < 0.001$).

Table 3 presents the multivariable correlates of moderate-to-severe depression and loneliness of the study participants. Participants who were cohabiting with men were 3.37 times more likely to be depressed (AOR = 3.37, 95% CI: 1.31, 8.64) compared to those who were cohabiting with women. Participants who felt hopelessness about the future (AOR = 3.20, 95% CI: 1.90, 5.20) and reported higher level of internalized homophobia (AOR = 2.32, 95% CI: 1.47, 3.67) were more likely to be depressed. Meanwhile, participants who were aged 26–35 years old had greater odds of experiencing loneliness compared to those aged between 18–25 years old (AOR = 1.55, 95% CI: 0.95, 2.52). Participants who had CRAI had greater odds of experiencing loneliness compared to those who did not have CRAI in the past six months (AOR = 1.67, 95% CI: 1.08, 2.58). Finally, participants who had a sense of hopelessness were 2.4 times more likely to experience loneliness compared to those who felt hopeful for the future (AOR = 2.40, 95% CI: 1.60, 3.70).

DISCUSSION

In this study, we found that about one-fourth of the HIV-negative or unknown status MSM participants in China suffered from moderate-to-severe depression and one-third suffered from loneliness. The prevalence of moderate-to-severe depression in our sample, 26.8%, was lower than the rates reported in previous studies of Chinese MSM of various serostatuses, which was 46.1% (Yan et al., 2014), and Chinese MSM who were HIV-positive, which was 36.0% (Tao et al., 2017). The rate was within the range reported by other developing countries, with 11% to 46.3% experiencing depressive symptoms among general MSM (Ahaneku et al., 2016; Tomori et al., 2016). The results of our study also revealed a high prevalence of other negative psychological states such as hopelessness and internalized homophobia, which were significantly correlated with depression and loneliness among our sample of MSM participants. In general, MSM are disproportionately impacted by mental health issues, such as depression and suicidality (Dyer et al., 2015; Li et al., 2016). These mental health issues may be consequences of stigma and discrimination and may result in adverse behavioral and health outcomes, including HIV infection and poorer health outcomes due to lack of adherence in testing and treatment (Dyer et al., 2015).

A global online survey reported that internalized homophobia was negatively associated with having ever tested for HIV among Chinese MSM (Pyun et al., 2014). As one of the social barriers of HIV care and treatment, internalized homophobia, i.e., negative feelings about one's own sexual orientation resulting from the internalization of anti-gay stigma and prejudice (Herek, 2004), may cause negative self-worth, increased self-judgment and the belief that something is wrong with oneself; this may further contribute to minority stress and feelings of depression, and ultimately affect health-seeking behaviors such as HIV testing. Therefore, campaigns that focus on destigmatizing homosexuality and promoting acceptance of one's sexuality may be helpful in reducing internalized homophobia and improving HIV-prevention efforts among MSM.

Previous studies reported that the MSM population in China was vulnerable to poor social support and increased depression and loneliness, which are associated with higher rates of suicidal ideation (Li et al., 2016; Yan et al., 2014). Depression and loneliness may arise due to genetic predisposition, personality development (lack of self-esteem and self-worth) and early or later life experiences. Lonely and depressed people are more likely to experience cognitive distortions, such as seeing neutral stimuli as negative and threatening, which then perpetuate the cycle of psychological distress (Blatt et al., 2005). Moreover, many MSM in China do not disclose their same-sex behaviors to others, including health care workers, for fear of being rejected and discriminated against (Wei et al., 2014; Choi et al., 2006). This might discourage them from accessing HIV prevention and other services provided by the Chinese government. Furthermore, decreased communication between MSM and their family and friends about their lifestyle and constantly concealing their sexual orientation may further exacerbate their already existing depressive symptoms and loneliness. To alleviate their psychological distress, they may be prone to pursue high-risk sexual behaviors (Martin and Knox, 1997; Shernoff, 2006). Therefore, development of a nuanced HIV-prevention program considering psychological, social and personality issues should be adopted among MSM in China.

An interesting finding in this study is that participants who are cohabiting with other men have greater odds of being depressed. A possible explanation may be that those MSM who are cohabitating with men are doing so out of necessity because they have been rejected by their family and peers. The rejection from family and peers may contribute to higher levels of isolation, shame, loneliness, and depression, which may then lead to increased risky sexual behavior. Another possibility is that MSM who cohabit with other men may not feel they can do so openly, but rather feel the need to conceal their living arrangements. The additional stress of hiding their home situation may contribute to increased feelings of stress and depression. To corroborate these hypotheses, a previous study among Chinese MSM showed that cohabiting with a male partner was associated with having increased rates of condomless anal sex (Zhang et al., 2015). This finding supports the idea that MSM who cohabit with men have higher rates of loneliness and depression, which then increases their risk of having condomless receptive anal intercourse.

People feel lonely when they perceive their interpersonal relationships to be deficient, situationally or chronically. A variety of coping strategies, from constructive to highly destructive, have been adopted by people who feel lonely (Woodward and Kalyanmasih,

1990). Loneliness was shown to be associated with instability of self-esteem, avoidance behaviors, and lower social support among MSM (Martin and Knox, 1997). Loneliness may contribute to difficulties with safer sex negotiation and condom use (Venable et al., 2006). A recent study showed that a one-unit increase in the UCLA loneliness score was met with a 10% decrease in the odds of condom usage among a sample of HIV-positive MSM (Hubach et al., 2015). In our study, lonely participants were significantly more likely to have condomless receptive anal intercourse, which is consistent with data from previous studies (Hubach et al., 2015; Martin and Knox, 1997). Lonely MSM may engage in condomless sexual behaviors to nurture committed relationships or seek casual sexual encounters to ward off feelings of loneliness and foster a sense of connection with sex partners (Pope et al., 2007; de Voux et al., 2016). Therefore, when examining barriers to condom use, psychological factors such as loneliness and depression should be strong considerations in addition to the usual access barriers (e.g., obtaining condoms) and usage barriers (e.g., decreased sexual sensation) (Mimiaga et al., 2013). A better understanding of the social context of the MSM community will benefit HIV-prevention programs.

Mental health of MSM is influenced by a collection of environmental, social and individual factors, especially the contexts they are living in. While previous studies have shown that HIV behavioral interventions are effective in reducing the odds of having unprotected anal sex, deeply rooted social problems and inequities, such as poverty, homelessness, racism, stigma and homophobia and the resulting mental health problems, may affect HIV risk and can impact the effective delivery of prevention programs. Thus, maximizing the effectiveness of these interventions through continued assessment of mental health issues is critical for a sustainable impact on the HIV epidemic (Herbst et al., 2007). Current HIV programming for MSM in China is heavily focused on promoting condom use and HIV testing, and there are no evidence-based interventions addressing the social and mental health contexts of HIV-related sexual behavior in this population. Measures to prevent HIV in China may have limited efficacy if mental and behavioral health are not considered together. Although the attitude of the general public toward MSM populations has improved dramatically in the past decade, MSM- and HIV-related stigma and discrimination are still highly prevalent in the community. Being attuned to psychological barriers to HIV-prevention behaviors that stem from depression and loneliness will be crucial to more effectively combat the HIV epidemic in China. To have more impact, HIV prevention programs should not be limited to screening for mental health and psychosocial issues, but also provide a supportive community for MSM via peer-led education and health communication. A comprehensive intervention program addressing the sexual health and overall wellbeing of MSM at both the individual and the community level should be developed and scaled up nationally. Such interventions should focus on the interaction of psychological and behavioral mechanisms for HIV risk, and integrate with treatment of mental health problems found among Chinese MSM. Meanwhile, improving competency of screening and identify early mental disorders of patients among health professionals in HIV organizations will strengthen the current health system in coping with HIV epidemic across China and the world.

This study had several limitations. First, participants were recruited from MSM venues and a gay-oriented website in one Chinese province. Thus, our findings may not be generalizable

to MSM who do not frequent these venues, do not visit the website, or live in other parts of China. Second, data was collected via self-report, which is subject to social desirability bias. This bias was decreased by having participants self-administer our survey. In addition, participants were asked to recall information from the past six months, which allows for the potential of recall bias. Finally, loneliness was measured from a single-item on the CES-D 10, so may be conflated with our measure for depression. Although the single question has face-validity, future studies may consider other fully validated short scales measuring loneliness, such as the short version of the UCLA Loneliness Scale, which contains three items measuring three aspects of loneliness (Hughes et al., 2004).

In summary, our study suggests that MSM in China have significant rates of depression and loneliness. Loneliness may lead to higher rates of sexual behavior that place MSM at risk for becoming infected with HIV. Further exploration of the pathway through which loneliness leads to increased risky sexual behavior is warranted among Chinese MSM. Prevention programs should integrate mental health and behavioral assessments and interventions with HIV and STI screening in order to be maximally effective. Providing a supportive community to address psychosocial and mental health factors may be the critical step in more efficiently reducing HIV-transmission among MSM in China.

Acknowledgments

Acknowledgement/Support: NIH/NIMH (Grant number: R00MH093201)

This study was funded by a grant from the U.S. National Institute of Mental Health (grant number: R00MH093201).

References

- Ahaneku H, Ross MW, Nyoni JE, Selwyn B, Troisi C, Mbwambo J, Adeboye A, McCurdy S. Depression and HIV risk among men who have sex with men in Tanzania. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2016; 28:140–147.
- Beck A, Weissman A. The Measurement of Pessimism: The Hopelessness Scale. *Journal of Consulting and Clinical Psychology*. 1974; 42(6):861–865. [PubMed: 4436473]
- Berg RC, Ross MW, Weatherburn P, Schmidt AJ. Structural and environmental factors are associated with internalised homonegativity in men who have sex with men: Findings from the European MSM Internet Survey (EMIS) in 38 countries. *Social Science & Medicine*. 2013; 78:61–69. [PubMed: 23261257]
- Blatt SJ, Luyten P, Corveleyn J. Toward a dynamic interaction model of depression and its treatment. *Psyche-Zeitschrift Fur Psychoanalyse Und Ihre Anwendungen*. 2005; 59(9–10):864–891.
- Catania JA, Kegeles SM, Coates TJ. TOWARDS AN UNDERSTANDING OF RISK BEHAVIOR - AN AIDS RISK REDUCTION MODEL (ARRM). *Health Education Quarterly*. 1990; 17(1):53–72. [PubMed: 2318652]
- Centers for Disease Control and Prevention, C. [Accessed 24 Mar 2015] HIV among gay/bisexual men. 2015. <http://www.cdc.gov/hiv/risk/gender/msm/facts/index.html>
- Chen Y, Bennett D, Clarke R, Guo Y, Yu C, Bian Z, Ma L, Huang Y, Sun Q, Zhang N, Zheng X, Chen J, Peto R, Kendler KS, Li L, Chen Z. Patterns and correlates of major depression in Chinese adults: a cross-sectional study of 0.5 million men and women. *Psychological Medicine*. 2007; 47(5):958–970.
- Choi KH, Lui H, Guo YQ, Han L, Mandel JS. Lack of HIV testing and awareness of HIV infection among men who have sex with men, Beijing, China. *Aids Education and Prevention*. 2006; 18(1): 33–43. [PubMed: 16539574]

- de Jong Gierveld J, van Tilburg T. Living arrangements of older adults in the Netherlands and Italy: Coresidence values and behavior and their consequences for loneliness. *Journal of Crosscultural Gerontology*. 1999; 14:1–24.
- de Voux A, Baral SD, Bekker LG, Beyrer C, Phaswana-Mafuya N, Siegler AJ, Sullivan PS, Winskell K, Stephenson R. A social network typology and sexual risk-taking among men who have sex with men in Cape Town and Port Elizabeth, South Africa. *Culture Health & Sexuality*. 2016; 18(5): 509–523.
- DeLonga K, Torres HL, Kamen C, Evans SN, Lee S, Koopman C, Gore-Felton C. Loneliness, internalized homophobia, and compulsive internet use: Factors associated with sexual risk behavior among a sample of adolescent males seeking services at a community LGBT center. *Sexual Addiction & Compulsivity*. 2011; 18(2):61–74.
- DeSalvo KB, Bloser N, Reynolds K, He J, Muntner P. Mortality prediction with a single general self-rated health question. *Journal of General Internal Medicine*. 2006; 21(3):267–275. [PubMed: 16336622]
- Dispenza F, Dew BJ, Tatum AK, Wolf ECM. Sexual Health Risk Behaviors Among Older Men Who Have Sex With Men: Implications for Interventions. *Adultspan Journal*. 2015; 14(1):35–48.
- Doskoch P. Drug Use and Loneliness Are Linked to Unprotected Sex in Older Adults with HIV. *Perspectives on Sexual and Reproductive Health*. 2011; 43(1):70–70.
- Du Bois SN, McKirnan DJ. A longitudinal analysis of HIV treatment adherence among men who have sex with men: A cognitive escape perspective. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2012; 24(11):1425–1431.
- Dyer TP, Regan R, Pacek LR, Acheampong A, Khan MR. Psychosocial Vulnerability and HIV-Related Sexual Risk Among Men who have Sex with Men and Women in the United States. *Archives of Sexual Behavior*. 2015; 44(2):429–441. [PubMed: 25183549]
- Ferro EG, Weikum D, Vagenas P, Copenhaver MM, Gonzales P, Peinado J, Cabello R, Lama JR, Sanchez J, Altice FL. Alcohol use disorders negatively influence antiretroviral medication adherence among men who have sex with men in Peru. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2015; 27(1):93–104.
- Fishbein M. An Integrative Model for Behavioral Prediction and its application to health promotion. In: DiClemente RJ, Crosby RA, Kegler MC, editors *Emerging theories in health promotion practice and research*. 2. San Francisco, CA: Jossey-Bass; 2009.
- Fisher JD, Fisher WA, Shuper PA. The Informational-Motivational-Behavioral Skills Model of HIV Prevention Behavior. In: DiClemente RJ, Crosby RA, Kegler MC, editors *Emerging theories in health promotion practice and research*. 2. San Francisco, CA: Jossey-Bass; 2009.
- Hall HI, Song RG, Rhodes P, Prejean J, An Q, Lee LM, Karon J, Brookmeyer R, Kaplan EH, McKenna MT, Janssen RS. Estimation of HIV incidence in the United States. *Jama-Journal of the American Medical Association*. 2008; 300(5):520–529.
- Herbst JH, Beeker C, Mathew A, McNally T, Passin WF, Kay LS, Crepaz N, Lyles CM, Briss P, Chattopadhyay S, Johnson RL. The effectiveness of individual-, group-, and community-level HIV behavioral risk-reduction interventions for adult men who have sex with men: a systematic review. *American Journal of Preventive Medicine*. 2007; 32(4):S38–S67. [PubMed: 17386336]
- Herek G. Beyond homophobia: thinking about sexual prejudice and stigma in the twenty-first century. *Sex Res Soc Policy*. 2004; 1:6–24.
- Hubach RD, Dodge B, Li MJ, Schick V, Herbenick D, Ramos WD, Cola T, Reece M. LONELINESS, HIV-RELATED STIGMA, AND CONDOM USE AMONG A PREDOMINANTLY RURAL SAMPLE OF HIV-POSITIVE MEN WHO HAVE SEX WITH MEN (MSM). *Aids Education and Prevention*. 2015; 27(1):72–83. [PubMed: 25646731]
- Hughes ME, Waite LJ, Hawkey LC, Cacioppo JT. A short scale for measuring loneliness in large surveys - Results from two population-based studies. *Research on Aging*. 2004; 26(6):655–672. [PubMed: 18504506]
- Kohout F, Berkman L, DAE, et al. Tow shorter forms of the CES-D depression symptoms index. *J Aging Health*. 1993; 5:179–193. [PubMed: 10125443]
- Kurtz SP. Post-circuit blues: Motivations and consequences of crystal meth use among gay men in Miami. *Aids and Behavior*. 2005; 9(1):63–72. [PubMed: 15812614]

- Kuyper L, Fokkema T. Loneliness Among Older Lesbian, Gay, and Bisexual Adults: The Role of Minority Stress. *Archives of Sexual Behavior*. 2010; 39(5):1171–1180. [PubMed: 19629672]
- Li R, Cai Y, Wang Y, Sun Z, Zhu CD, Tian YL, Jiang XQ, Gan F. Psychosocial Syndemic Associated With Increased Suicidal Ideation Among Men Who Have Sex With Men in Shanghai, China. *Health Psychology*. 2016; 35(2):148–156. [PubMed: 26462059]
- Luo Y, Waite LJ. Loneliness and Mortality Among Older Adults in China. *Journals of Gerontology Series B-Psychological Sciences and Social Sciences*. 2014; 69(4):633–645.
- Luria RE. VALIDITY AND RELIABILITY OF VISUAL ANALOGUE MOOD SCALE. *Journal of Psychiatric Research*. 1975; 12(1):51–57. [PubMed: 806683]
- Martin JI, Knox J. Loneliness and sexual risk behavior in gay men. *Psychological Reports*. 1997; 81(3):815–825. [PubMed: 9400073]
- Mimiaga MJ, Reisner SL, Closson EF, Perry N, Perkovich B, Nguyen T, Trang NNN, Lan HX, Thien DD, Mayer KH, Colby D. Self-perceived HIV risk and the use of risk reduction strategies among men who engage in transactional sex with other men in Ho Chi Minh City, Vietnam. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2013; 25(8):1039–1044.
- Mustanski B, Garofalo R, Herrick A, Donenberg G. Psychosocial health problems increase risk for HIV among urban young men who have sex with men: preliminary evidence of a syndemic in need of attention. *Annals of Behavioral Medicine*. 2007; 34(1):37–45. [PubMed: 17688395]
- Neme S, Goldenberg T, Stekler JD, Sullivan PS, Stephenson R. Attitudes towards couples HIV testing and counseling among Latino men who have sex with men in the Seattle area. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2015; 27(10):1354–1359.
- Ophir Y, Asterhan CSC, Schwarz BB. Unfolding the notes from the walls: Adolescents' depression manifestations on Facebook. *Computers in Human Behavior*. 2017; 72:96–107.
- Peplau LA, Perlman D. Perspectives on loneliness. In: Peplau LA, Perlman D, editors *Loneliness: A Sourcebook of Current Theory, Research and Therapy*. New York: Wiley; 1982. 1–8.
- Pope M, Wierzalis EA, Barrett B, Rankins M. Sexual and intimacy issues for aging gaymen. *Adultspan Journal*. 2007; 6:68–82.
- Pyun T, Santos GM, Arreola S, Do T, Hebert P, Beck J, Makofane K, Wilson PA, Ayala G. Internalized Homophobia and Reduced HIV Testing Among Men Who Have Sex With Men in China. *Asia-Pacific Journal of Public Health*. 2014; 26(2):118–125. [PubMed: 24554493]
- Russell DW. UCLA Loneliness Scale (Version 3): Reliability, validity, and factor structure. *Journal of Personality Assessment*. 1996; 66(1):20–40. [PubMed: 8576833]
- Safren SA, Reisner SL, Herrick A, Mimiaga MI, Stall RD. Mental Health and HIV Risk in Men Who Have Sex With Men. *J AIDS-Journal of Acquired Immune Deficiency Syndromes*. 2010; 55:S74–S77.
- Savikko N, Routasalo P, Tilvis RS, Strandberg TE, Pitkala KH. Predictors and subjective causes of loneliness in an aged population. *Archives of Gerontology and Geriatrics*. 2005; 41(3):223–233. [PubMed: 15908025]
- Shek DTL. MEASUREMENT OF PESSIMISM IN CHINESE ADOLESCENTS - THE CHINESE HOPELESSNESS SCALE. *Social Behavior and Personality*. 1993; 21(2):107–119.
- Shermoff M. Condomless sex: Gay men, barebacking, and harm reduction. *Social Work*. 2006; 51(2):106–113. [PubMed: 16858916]
- Stall R, Mills TC, Williamson J, Hart T, Greenwood G, Paul J, Pollack L, Binson D, Osmond D, Catania JA. Association of co-occurring psychosocial health problems and increased vulnerability to HIV/AIDS among urban men who have sex with men. *American Journal of Public Health*. 2003; 93(6):939–942. [PubMed: 12773359]
- Tao J, Vermund SH, Lu HY, Ruan YH, Shepherd BE, Kipp AM, Amico KR, Zhang XJ, Shao YM, Qian HZ. Impact of Depression and Anxiety on Initiation of Antiretroviral Therapy Among Men Who Have Sex with Men with Newly Diagnosed HIV Infections in China. *Aids Patient Care and Stds*. 2017; 31(2):96–104. [PubMed: 28170305]
- Tilvis RS, Pitkala KH, Jolkkonen J, Strandberg TE. Social networks and dementia. *Lancet*. 2000; 356(9223):77–78. [PubMed: 10892794]
- Tomori C, McFall AM, Srikrishnan AK, Mehta SH, Solomon SS, Anand S, Vasudevan CK, Solomon S, Celentano DD. Diverse Rates of Depression Among Men Who Have Sex with Men (MSM)

- Across India: Insights from a Multi-site Mixed Method Study. *Aids and Behavior*. 2016; 20(2): 304–316.
- Torress HL, Gore-Felton C. Compulsivity, substance use, and loneliness: the loneliness and sexual risk model (LSRM). *Sexual Addiction & Compulsivity*. 2007; 14(1):63–75.
- Vanable PA, Carey MP, Blair DC, Littlewood RA. Impact of HIV-related stigma on health behaviors and psychological adjustment among HIV-positive men and women. *Aids and Behavior*. 2006; 10(5):473–482. [PubMed: 16604295]
- Watkins TL, Simpson C, Cofield SS, Davies S, Kohler C, Usdan S. The Relationship of Religiosity, Spirituality, Substance Abuse, and Depression Among Black Men Who Have Sex with Men (MSM). *Journal of Religion & Health*. 2016; 55(1):255–268. [PubMed: 26286843]
- Wei CY, Yan HJ, Yang CK, Raymond HF, Li JJ, Yang HT, Zhao JK, Huan XP, Stall R. Accessing HIV testing and treatment among men who have sex with men in China: A qualitative study. *Aids Care-Psychological and Socio-Medical Aspects of Aids/Hiv*. 2014; 26(3):372–378.
- Wei CY, Cheung DH, Yan HJ, Li JJ, Shi LE, Raymond HF. The Impact of Homophobia and HIV Stigma on HIV Testing Uptake Among Chinese Men Who Have Sex With Men: a Mediation Analysis. *J AIDS-Journal of Acquired Immune Deficiency Syndromes*. 2016; 71(1):87–93.
- Woodward JC, Kalyanmasih V. LONELINESS, COPING STRATEGIES AND COGNITIVE STYLES OF THE GIFTED RURAL ADOLESCENT. *Adolescence*. 1990; 25(100):977–988. [PubMed: 2275450]
- Wu YL, Yang HY, Wang J, Yao H, Zhao X, Chen J, Ding XX, Zhang HB, Bi P, Sun YH. Prevalence of suicidal ideation and associated factors among HIV-positive MSM in Anhui, China. *International Journal of Std & Aids*. 2015; 26(7):496–503. [PubMed: 25060699]
- Yan HM, Wong FY, Zheng T, Ning Z, Ding YY, Nehl EJ, Lin L, He N. Social support and depressive symptoms among ‘money’ boys and general men who have sex with men in Shanghai, China. *Sexual Health*. 2014; 11(3):285–287. [PubMed: 24919661]
- Yu SC, Lin YH, Hsu WH. Applying structural equation modeling to report psychometric properties of Chinese version 10-item CES-D depression scale. *Quality & Quantity*. 2013; 47(3):1511–1518.
- Zhang H, Lu HY, Pan SW, Xia DY, Zhao YJ, Xiao Y, He X, Yue H, Sun ZY, Xu YA, Ruan YH, Shao YM. Correlates of Unprotected Anal Intercourse: The Influence of Anal Sex Position Among Men Who Have Sex with Men in Beijing, China. *Archives of Sexual Behavior*. 2015; 44(2):375–387. [PubMed: 25548064]
- Zhao J, Cai R, Chen L, Cai WD, Yang ZR, Richardus JH, de Vlas SJ. A Comparison Between Respondent-Driven Sampling and Time-Location Sampling Among Men Who Have Sex with Men in Shenzhen, China. *Archives of Sexual Behavior*. 2015; 44(7):2055–2065. [PubMed: 25239658]
- Zhong BL, Xu YM, Jin D, Zou XW, Liu TB. Prevalence and correlates of loneliness among Chinese service industry migrant workers A cross-sectional survey. *Medicine*. 2016; 95(24)

Table 1

Prevalence of negative mental health issues and sample characteristics among HIV-negative/unknown men who have sex with men, Jiangsu, China, 2013

Characteristics		N	%
Depression	No	371	73.2
	Yes	136	26.8
Loneliness	No	327	64.5
	Yes	180	35.5
Age	18–25	151	30.0
	26–35	186	36.9
	36+	167	33.1
Education	Middle school or less	74	14.6
	High school or technical	222	43.8
	College or above	211	41.6
Marital status	Single	292	57.7
	Married	100	19.8
	Other	114	22.5
Cohabiting partners	With women	93	18.3
	With men	103	20.3
	Other	311	61.3
Employment	Fulltime	418	82.4
	Other	89	17.6
Income	2,999	173	34.1
	3,000–4,999	195	38.5
	5,000	139	27.4
Sexual orientation	Gay	344	67.9
	Bisexual/straight/unsure	163	32.1
Out to anyone	No	230	45.4
	Yes	277	54.6
Age of 1 st anal intercourse	18	72	16
	19–25	260	57.6
	26	119	26.4
Condom use in 1 st anal intercourse	No	163	36.1
	Yes	288	63.9
Have a main male partner	No	276	54.4
	Yes	231	45.6
Number of male anal sex partners	1	217	42.8
	2–5	217	42.8
	6	73	14.4
CIAI	No	277	61.3
	Yes	175	38.7
CRAI	No	311	68.8

Characteristics		N	%
HIV testing	Yes	141	31.2
	No	109	21.5
Hopelessness	Yes	398	78.5
	Hopeful	298	58.8
Internalized homophobia	Hopeless	209	41.2
	Low	254	53.8
	High	218	46.2

Note. CIAI=condomless insertive anal intercourse, CRAI=condomless receptive anal intercourse.

Bivariate correlates of moderate-to-severe depressive symptoms and loneliness among HIV-negative/unknown men who have sex with men, Jiangsu, China, 2013.

Table 2

Characteristics	Depression			Loneliness		
	yes	N	%	yes	N	%
Age						
18–25	47	151	31.13	54	151	35.76
26–35	59	186	31.72	78	186	41.94
36+	28	167	16.77	47	167	28.14
Education						
Middle school or less	16	74	21.62	23	74	31.08
High school or technical	64	222	28.83	82	222	36.94
College or above	56	211	26.54	75	211	35.55
Marital status						
Single	86	292	29.45	107	292	36.64
Married	18	100	18.00	31	100	31.00
Other	32	114	28.07	42	114	36.84
Cohabiting partners						
With women	12	93	12.90	28	93	30.11
With men	34	103	33.01	31	103	30.10
Other	90	311	28.94	121	311	38.91
Employment						
Fulltime	115	418	27.51	155	418	37.08
Other	21	89	23.60	25	89	28.09
Income						
2,999	43	173	24.86	60	173	34.68
3,000–4,999	56	195	28.72	68	195	34.87
5,000	37	139	26.62	52	139	37.41
Sexual orientation						
Gay	95	344	27.62	122	344	35.47
Bisexual/straight/unsure	41	163	25.15	58	163	35.58
Out to anyone						
No	67	230	29.13	82	230	35.65
Yes	69	277	24.91	98	277	35.38
Age of 1 st anal intercourse						
18	20	72	27.78	24	72	33.33
19–25	80	260	30.77	102	260	39.23
26	26	119	21.85	37	119	31.09
Condom use in 1 st anal intercourse						
No	49	163	30.06	61	163	37.42
Yes	77	288	26.74	102	288	35.42

Characteristics	Depression					Loneliness				
	yes	N	%	p		yes	N	%	p	
Have a main male partner										
	81	276	29.35	0.161	No	109	276	39.49	0.040	
	55	231	23.81		Yes	71	231	30.74		
Number of male anal sex partners					1	60	217	27.65	0.758	0.427
	59	217	27.19		2—5	79	217	36.41		
	17	73	23.29		6	21	73	28.77		
CIAI					No	81	277	29.24	0.415	0.460
	45	175	25.71		Yes	66	175	37.71		
CRAI					No	81	311	26.05	0.197	0.022
	45	141	31.91		Yes	61	141	43.26		
HIV testing					No	37	109	33.94	0.058	0.036
	99	398	24.87		Yes	132	398	33.17		
Hopelessness					Hopeful	29	209	13.88	0.000	0.000
	107	298	35.91		Hopeless	130	298	43.62		
Internalized homophobia					Low	52	254	20.47	0.001	0.455
	74	218	33.94		High	81	218	37.16		

Note: CIAI=condomless insertive anal intercourse, CRAI=condomless receptive anal intercourse.

Multivariable correlates of moderate-to-severe depressive symptoms and loneliness among HIV-negative/unknown men who have sex with men, Jiangsu, China, 2013.

Table 3

Characteristics	Depression			Loneliness		
	AOR	95%CI	p	AOR	95%CI	p
Age						
18–25	1		0.191	1		0.043
26–35	1.41	(0.84,2.73)		1.55	(0.95,2.52)	
36+	0.67	(0.34,1.31)		0.86	(0.50,1.45)	
Marital status						
Single	1		0.964	---	---	---
Married	0.98	(0.46,2.12)		---	---	
Other	1.07	(0.59,1.94)		---	---	
Cohabiting partners						
With women	1		0.042	---	---	
With men	3.37	(1.31,8.64)		---	---	
Other	2.28	(1.01,5.00)		---	---	
Have a main male partner						
No	1		0.580	1		0.054
Yes	0.87	(0.54,1.41)		0.66	(0.44,1.00)	
CRAI						
No	---	---	---	1		0.021
Yes	---	---	---	1.67	(1.08,2.58)	
HIV testing						
No	1		0.315	1		0.445
Yes	0.77	(0.45,1.29)		0.83	(0.51,1.34)	
Hopelessness						
Hopeful	1		0.000	1		0.000
Hopeless	3.20	(1.90,5.20)		2.40	(1.60, 3.70)	
Internalized homophobia						
Low	1		0.000	---	---	---
High	2.32	(1.47,3.67)		---	---	

Note: ---not included in the final model; CRAI=condomless receptive anal intercourse.