Bowel obstruction caused by broad ligament hernia successfully repaired by laparoscopy

K Toolabi^{1,2}, Abbas Zamanian¹, R Parsaei¹

¹Department of Surgery, Erfan Hospital, Tehran, Iran

²Department of Surgery, Tehran University of Medical Sciences, Tehran, Iran

ABSTRACT

Internal hernias are rare bowel obstructions. We present a case of small bowel obstruction in a 37-year-old woman caused by internal herniation through a defect in broad ligament, which was managed by laparoscopic surgery.

KEYWORDS

Small intestine – Obstruction – Broad ligament – Hernia – Laparoscopy

Accepted 8 January 2018

CORRESPONDENCE TO

Reza Parsaei, E: rezparsaei@gmail.com

Introduction

Internal hernias are reported in 1% of mechanical bowel obstructions.¹ It occurs when small bowel herniates through a recess or defect within abdominal cavity. Herniation through a defect in broad ligament is quite rare and accounts for 4% of internal herniations.² We present a case of small bowel obstruction in a 37-year-old woman caused by internal herniation through a defect in broad ligament, which was managed by laparoscopic surgery.

Case history

A 37-year-old woman presented to our hospital with a three-day history of vomiting and abdominal pain. Her symptoms had worsened gradually until she could not tolerate any oral intake. She was quite well before the onset of symptoms and had no prior abdominal complaints. Her last bowel movement was the day before admission. She had an umbilical hernia repair surgery as a child. On examination, she had no fever or tachycardia. Her abdomen was distended but no local tenderness was detected. Laboratory findings showed a slight rise in creatinine level and elevated amylase and lipase levels. Plasma electrolytes and blood cell counts were normal. Plain abdominal radiography showed distended small bowel loops (Fig 1). Abdominopelvic computed tomography showed distended stomach and small bowel loops. Collapsed small bowel also noted, suggestive of mechanical small bowel obstruction (Fig 2).

On laparoscopy, it was found that a loop of small bowel had herniated through a defect in the left broad ligament (Fig 3). Bowel was pulled out of the defect and proved to be healthy and viable. The defect was repaired using continuous PDS 2-0 sutures (Fig 4). The patient's postoperative

course was uneventful and oral intake started the day after surgery and she was discharged two days after surgery.

Discussion

Small bowel obstruction is a relatively common surgical emergency. Most obstructions of small intestines are caused by adhesions developed after abdominal surgeries. Internal hernias are the result of protrusion of a bowel loop through a defect within the abdominal cavity. Instances include para-duodenal hernia, pericecal hernia, hernia through the foramen of Winslow and transmesenteric hernia.² Hernia through a defect in the broad ligament is quite rare and accounts for 4% of internal hernias.² Hunt has classified a broad ligament defect into: 1) fenestra type, the most common, which is indicated by the presence of a defect in the two peritoneal layers; 2) pouch type, which is indicated by the presence of a defect in only one of the two layers, whereby the visceral structures would be trapped in the parametrial tissue; and 3) hernia sac type, whereby a double layer of attenuated peritoneum lines the herniated bowel, forming a true internal hernia.³ Cilley has classified broad ligament defects based on the anatomical location: type 1, defect caudal to the round ligament; type 2, defect above the broad ligament; and type 3, defect between the round ligament and remainder of the broad ligament, through the mesoligamentum teres.⁴

Prior surgery, pelvic inflammatory disease and trauma during delivery or vaginal manipulations are the probable aetiologies of a broad ligament defect.⁵ However, as in our patient, none of these potential causes is found in some cases. These defects are believed to be congenital.

Patients present with symptoms of small bowel obstruction. Internal herniations are included in the differential

Figure 1 Plain abdominal radiographs showing distended small bowel loops with air and fluid levels

Figure 2 Computed tomography showing dilated bowel loops (red arrow) and collapsed loops (blue arrow) suggesting a transition point in left pelvis

diagnosis of small bowel obstructions in patients without prior surgery. A diagnosis of broad ligament hernia is not necessary before surgery, since evidence of complete bowel obstruction or closed loop bowel obstruction mandates surgical intervention. Computed tomography is most helpful in the diagnosis of small bowel obstruction and finding its aetiology. A transition zone located in the pelvis, a cluster of dilated small bowel loops herniated laterally to the uterus in the pelvic cavity and enlargement of the distance between the uterus and one of the ovaries, which are deviated in opposite directions, are present in computed tomography of broad ligament herniation.⁶ We did not reach a diagnosis of broad ligament herniation before surgery and, based on

Figure 3 Defect in the left broad ligament

Figure 4 Closure of the defect in the left broad ligament

imaging, we only guessed a pelvic origin for the small bowel obstruction.

Most surgeons perform laparotomy to address small bowel obstruction.⁷ Limited work space due to distended bowel loops and fear of iatrogenic injury to intestine is main reason that many surgeons are reluctant to adopt laparoscopy in small bowel obstruction, although its use is becoming more common. Behman *et al.* have reported an increase from 4.3% to 14.3% in laparoscopic adhesion lysis from 2005 to 2014.⁷ They studied 8584 patients and found that bowel injury was higher in patients undergoing laparoscopy.⁷

Many studies have reported that laparoscopy can be performed safely with comparable morbidity and earlier recovery.^{8,9} However, in these studies, most patients still underwent laparotomy and, in those for whom laparoscopy

was selected, the conversion rate was over 30%.^{8,9} It should be noted that most patients in these studies had small bowel obstruction caused by adhesions due to previous surgery. Laparoscopy in internal herniation is technically easier since abdominal adhesions pose more challenge to laparoscopy in comparison with internal herniation.

Based on current evidence, we believe that laparoscopy can be safely attempted to diagnose and repair internal herniation but the decision should be individual, based on the patient's condition and the surgeon's expertise.

References

1. Varela GG, López-Loredo A, García León JF. Broad ligament hernia-associated bowel obstruction. *JSLS* 2007; **11**(1): 127–130.
2. Matsunami M, Kusanagi H, Hayashi K *et al.* Broad ligament hernia successfully treated by laparoscopy: case report and review of literature. *Asian J Endosc Surg* 2014; **7**(4): 327–329.
3. Hunt AB. Fenestra and pouches in the broad ligament as an actual and potential cause of strangulated intra-abdominal hernia. *Surg Gynecol Obstet* 1934; **58**: 906–913.
4. Cilley R, Poterack K, Lemmer J *et al.* Defects of the broad ligament of the uterus. *Am J Gastroenterol* 1986; **81**: 389–391.
5. Ishihara H, Terahara M, Kigawa J *et al.* Strangulated herniation through a defect of the broad ligament of the uterus. *Gynecol Obstet Invest* 1993; **35**: 187–189.
6. Barbier Brion B, Daragon C, Idelcadi O *et al.* Small bowel obstruction due to broad ligament hernia: Computed tomography findings. *Hernia* 2011; **15**: 353–355.
7. Behman R, Nathens AB, Byrne JP *et al.* Laparoscopic surgery for adhesive small bowel obstruction is associated with a higher risk of bowel injury: a population-based analysis of 8584 patients. *Ann Surg* 2017; **266**(3): 489–498.
8. Byrne J, Saleh F, Ambrosini L *et al.* Laparoscopic versus open surgical management of adhesive small bowel obstruction: a comparison of outcomes. *Surg Endosc* 2015; **29**(9): 2,525–2,532.
9. Khaikin M, Schneider N, Cera S *et al.* Laparoscopic vs. open surgery for acute adhesive small-bowel obstruction: patients' outcome and cost-effectiveness. *Surg Endosc* 2007; **21**(5): 742–746.