Published in final edited form as:

Curr Clin Microbiol Rep. 2017 December ; 4(4): 208–217. doi:10.1007/s40588-017-0076-4.

Examining the potential for South American arboviruses to spread beyond the New World

Víctor Hugo Peña-García¹, Michael K. McCracken², and Rebecca C. Christofferson^{3,*}

¹Department of Biology and Microbiology, Faculty of Science and Engineering, University of Boyacá, Tunja, Boyacá, Colombia

²Viral Diseases Branch, Walter Reed Army Institute of Research, Silver Spring, Maryland 20910, USA

³Department of Pathobiological Sciences, Louisiana State University School of Veterinary Medicine, Baton Rouge, Louisiana 70803, USA

Abstract

Zika and chikungunya viruses emerged as public health emergencies in the Western Hemisphere where previously they had not been reported on a large scale. Millions were infected as the viruses met with virtually no herd immunity upon emergence.

Purpose of the Review—We explore the histories of these two recent arbovirus experiences in South America. We then review similarly three endemic South American viruses: yellow fever, Oropouche, and Mayaro viruses.

Recent Findings—We discuss the commonalities of the transmission systems and the possibility of an atypical emergence, that of New World virus to the Old World.

Summary—We discuss the avenues for research that would increase preparedness and efficiency of response should a South American arbovirus emerge in the Eastern Hemisphere.

Keywords

emerging viruses; Zika; chikungunya; yellow fever; Mayaro; Oropouche

Introduction

Arthropod-borne viruses (arboviruses) have worldwide distributions. The arboviruses of greatest human and/or animal health importance are grouped into six taxonomic families: *Bunyaviridae*, *Flaviviridae*, *Rhabdoviridae*, *Togaviridae*, *Reoviridae*, and *Orthomyxoviridae*.

*corresponding author: rcarri1@lsu.edu.

Disclaimer: The opinions or assertions contained herein are the private views of the authors and are not to be construed as reflecting the official views of the US Army, NAMRU6, the US Navy, or the US Department of Defense.

Compliance with Ethics Guidelines

Conflict of Interest

The authors declare that they have no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

However, when considering the most recent impacts to human public health, the arboviruses of most concern belong to the *Flaviviridae*, *Bunyaviridae*, or *Togaviridae* families [1]. The greatest number of pathogenic arboviruses originate from Africa and South America, where the relatively high level of biodiversity offers a wide range of both host and vector species [1].

In the Western Hemisphere, emergent arboviruses like dengue (DENV), chikungunya (CHIKV), West Nile (WNV), and, more recently, Zika (ZIKV) viruses have led to outbreaks of considerable scale, which resulted in high rates of morbidity and mortality [2]. It is interesting to note that the most significant intercontinental spread of arboviruses, particularly those listed above, appears to originate beyond the Americas rather than within. WNV was introduced via a traveler, likely from Israel; DENV is believed to have originally crossed to South America from Southeast Asia; and evidence suggests that ZIKV and CHIKV were introduced to the Americas from the Pacific region [3]. Presumably, the incursion of an American virus into the urban populations of the Eastern Hemisphere would encounter a similar lack of widespread herd immunity, and yet no such jump has been reported despite increasing global travel and trade [4].

In this review we explore the experiences of the most recent incursions of ZIKV and CHIKV into the Americas, as well as the history and most recent outbreaks of yellow fever virus (YFV), an Old World virus already established in the New World. We also review the history and transmission systems of two South American viruses with the potential to establish within global, urban transmission cycles: Mayaro (MAYV) and Oropouche (OROV) viruses.

Chikungunya (Alphavirus)

CHIKV is a member of the *Alphavirus* genus in the *Togaviridae* family. Infection with this virus usually presents as a nondescript febrile disease, but may also be accompanied by rash and severe, debilitating and often chronic arthralgia, which are its main discriminating characteristics. Historical data regarding CHIKV outbreaks may be confounded by its similarities in clinical presentation and overlapping transmission ecology to DENV, and the lack of molecular diagnostic tools that we have today. Still, some authors have suggested that many previous “dengue” records were actually misdiagnosed cases of CHIKV (reviewed in [5]). The first officially recognized outbreak of CHIKV occurred in 1952 in the coastal plateaus of Rondo, Makonde, and Mawia of modern day Tanzania. The etiological agent of this outbreak was isolated and identified as a virus subsequently called chikungunya virus. “Chikungunya” is a Tanzanian word that describes the distinctive posture caused by severe arthralgia in infected people [5]. Shortly after this episode, another outbreak was recorded in Uganda with CHIKV isolated from patients [6] as well as mosquitoes (*Aedes africanus*) [7].

In 1958, the first confirmed outbreak of CHIKV outside of Africa was reported in Bangkok, Thailand. Importantly, it was also the first report of CHIKV isolation from *Ae. aegypti* and of its co-circulation with DENV, which was also isolated from *Ae. aegypti* during this episode [8]. Reviews of historical reports suggested that CHIKV was already circulating in those areas at this time, and that CHIKV has circulated in Asia since the 18th century [9, 10]. Accordingly, phylogenetic analyses indicate three main genotypes: West African, East/

Central/South African (ECSA), and Asian, with the West African genotype considered to be the most ancestral [11, 12].

Evidence of CHIKV circulation has continued to present in Africa and sporadically in Asia. In 2005, a major outbreak of CHIKV took place among the islands of the Indian Ocean, particularly on La Reunion Island where almost a third of the population was infected [13]. From this outbreak, strains were isolated, sequenced, and demonstrated a consistent amino acid change from alanine to valine at position 226 of the E1 (envelope) glycoprotein. This mutation conferred some fitness advantage in the secondary vector, *Ae. albopictus*, while showing little to no advantage in *Ae. aegypti* [14, 15]. In addition, this outbreak was associated with the first death attributed to CHIKV and the broader nature of the *Ae. albopictus* habitat may be credited with the explosive expansion of CHIKV during this period [16]. This particular adaptation is specific to a lineage of the ECSA genotype of CHIKV whereas different amino acid shifts in strains of the Asian genotype have also conferred fitness advantages in *Ae. albopictus* [17]. Studies have shown that the epidemic lineage (often called Indian Ocean Lineage [IOL]) is closest to the ECSA lineage, supporting the theory of reintroduction to Asia from Africa [11, 18].

In 2013, CHIKV was detected and isolated for the first time in the western hemisphere on the island of St. Martin. Soon after the virus spread to 45 countries and territories in the Caribbean, and North, Central, and South America [19]. It was determined that this initial epidemic was caused by strains of the Asian lineage and not by ECSA-IOL lineage [20, 21]. Although some have postulated that molecular elements could play a role in the American epidemic (e.g., a duplication in the 3' UTR end that led to increased titers of these strains in mosquito cells [12]), it is important to note that ecological, immunological, and environmental factors likely play the dominant role in this explosive epidemic, as the major vector implicated in the Americas was *Ae. aegypti*, one for which there are no consistent fitness differences among genotypes/lineages [22, 23].

Zika (Flavivirus)

ZIKV is a member of the *Flaviviridae* family belonging to genus *Flavivirus*, which also includes DENV, WNV, Japanese encephalitis virus, and YFV. ZIKV was first identified in a sentinel rhesus monkey caged in the canopy of the Zika forest in 1947 as part of a sylvatic YFV surveillance network in Uganda [24]. A second isolation was achieved in 1948 from *Ae. africanus* mosquitoes in the same forest [24]. Additionally, while there was evidence of the presence of neutralizing antibodies against ZIKV in humans, the virus was not isolated from a human patient until 1952 in Nigeria [25, 26]. The first isolation in Asia occurred in Malaysia in 1966 from one pool of mosquitoes (*Ae. aegypti*) collected in Bentong in west-central Malaysia [27]. This isolation is particularly important because it was the first demonstration that ZIKV could be transmitted by this urban mosquito species, especially after a failed attempt to experimentally transmit the virus using *Ae. aegypti* fed on an exposed human [28]. Remarkably, it took eleven years before the first case of human ZIKV disease was reported in Asia when seven patients in Indonesia (1977–78) showed a significant rise in ZIKV antibody titers following presentation with symptoms clinically consistent with ZIKV infection [29]. Phylogenetic analyses have shown that there are two

major lineages of ZIKV, the Asian and the African lineages, though some refer to a third lineage (African II); when full genomes are considered, these isolates usually group with the major African lineage [30–32].

The emergence of Zika as a major public health problem began in 2007 on the island of Yap, Federated States of Micronesia. It was estimated that 73% of residents over the age of 3 years were infected by ZIKV, and the main vector identified was *Ae. hensilli* [33]. This outbreak was the first time ZIKV had been reported outside Africa and mainland Asia. In 2013, ZIKV was reported in French Polynesia where more than 5800 cases were recorded and a total 19,000 suspected [34]. Subsequently, the virus spread to New Caledonia, Cook Islands, and Easter Island [35]. In 2013–2014, the virus arrived in the Americas, particularly in Brazil, where numerous world sport championships were held including the FIFA Confederations Cup 2013, the International Va'a Federation's world sprints 2014, and the 2014 FIFA World Cup. There is some disagreement on exactly when the virus arrived, with some arguing 2013 [36] while others propose 2014 [37, 38] due to its first detection and rapid spread in this year. ZIKV spread to other states of Brazil in 2014 [39, 40] and, in 2015, the circulation of ZIKV was reported in 12 countries of South America, Central America and the Caribbean (reviewed in [41]). Previously, ZIKV infection was characterized by a flu-like illness, often asymptomatic or manifested as a mild clinical syndrome that may include fever, rash, arthralgia, conjunctivitis, pruritus, muscle pain, headache, and malaise. In these recent outbreaks, ZIKV was newly associated with severe manifestations, most notably Guillain-Barré syndrome, microcephaly, and ocular scarring [42–45]. Characterization of these recent outbreaks revealed that the circulating strains were from the Asian lineage [34, 36].

Yellow fever (Flavivirus)

YFV infections can be mild, but severe and often-fatal disease includes chills, fatigue, nausea, vomiting, and jaundice. The first New World YFV epidemics are believed to have occurred between 1647 and 1649 in Barbados, Cuba, Guadeloupe, and Mexico before spreading to the United States [46]. YFV continued to infect people in the tropical and subtropical areas of Americas and Africa, with additional outbreaks reported in Europe [47]. Josiah Nott was the first to present the idea that mosquitoes may transmit both yellow fever and malaria, and Carlos Finlay of Cuba formally described the theory of transmission in 1881, proposing, but not directly demonstrating, that a *Culex* mosquito was the vector [46, 48]. At the end of the 19th century, Walter Reed demonstrated that *Ae. aegypti* was the likely vector [49]. Work in the late 19th – early 20th centuries led to successful YFV reduction campaigns in Havana and Panamá [50, 51], but it wasn't until post-WWI that researchers expanded characterization of YFV to Africa with the intention of connecting it to disease in the Americas and Africa [52]. Stemming from these efforts, Max Theiler eventually attenuated the virus and thereby created the strain 17D, which is still the basis of the yellow fever vaccine [53]. Despite the existence of an effective vaccine, outbreaks have still been the cause of significant morbidity and mortality, especially in populations where vaccination levels are low [54, 55].

Phylogenetic analyses have demonstrated that YFV originated in Africa and was brought to America by slave trade hundreds of years ago [56]. One of the first comprehensive phylogenetic studies on YFV showed four main genotypes that included two genotypes from Africa and two from America [57]. Later, the existence of five genotypes in Africa was demonstrated: west African genotypes I and II, and the east African, east and central African, and Angola genotypes [58]. In America, two genotypes have been described: genotype I (grouping viruses from Brazil, Panamá, Venezuela, Colombia, Trinidad and Ecuador) and genotype II (grouping viruses mainly from Perú and some from Trinidad) [56, 57].

Discussion I: Learning from Experience

The recent emergence of ZIKV and CHIKV first in the Pacific regions and then in the Americas caused millions of infections and prompted a public health emergency [19, 59]. During these experiences, the body of knowledge regarding these viruses grew exponentially but also uncovered needed, unexplored avenues for better understanding of transmission and ultimately control. While the explosive nature of the epidemics was, in part, due to a lack of herd immunity, additional questions regarding transmission remain and the answers have significant impact on public health response and decision-making. First, which are the vector species most likely to play a role in transmission? Even with the adaptation of CHIKV to *Ae. albopictus*, the traditionally more-competent vector *Ae. aegypti* was implicated in its emergence in Brazil. Additionally, *Ae. albopictus*, which was shown to be highly competent for ZIKV in Singapore [60], is only moderately competent for ZIKV in the Americas [61]. Still, the mosquito most implicated in this ZIKV outbreak is *Ae. aegypti* [62, 63]. This suggests that other elements of the human-*aegypti*-virus interaction are equally or more important to understanding emergence potential than just vector competence in tropical regions [64]. Second, how similar must two ecologies be to support the maintenance of a newly emerged virus in an enzootic cycle? The long-term success of YFV is, in part, driven by the establishment of a robust enzootic cycle involving the sylvatic *Haemagogus spp.* mosquito vectors and a different set of non-human primates from its assumed African origins [65]. Thus, is there a similar consideration for S. American viruses that may establish sylvatic cycles in similar, but geographically distinct, regions? Lastly, there is a need for proactive, specific and consistent diagnostics for detection of new viruses and discrimination from related, endemic viruses. Because many members of the genus *Flavivirus* have historically circulated in similar ecologies and the geographic regions of these viruses overlap with the zone of emergence of ZIKV, there has been some difficulty in diagnostic development. Serological and immunological diagnostics like hemagglutination and neutralization assays have significant cross reactivity with other Flaviviruses, specifically DENV [66]. Additionally, there is antibody cross-reactivity among alphaviruses of the Semliki forest group, suggesting a similar problem may occur when differentiating CHIKV from MAYV [67]. This lack of diagnostic specificity confounded efforts to quickly and definitively diagnose ZIKV patients in the Americas, and may lead to future issues in distinguishing disease etiologies.

Mayaro (Alphavirus)

Mayaro virus (MAYV) was isolated for the first time in 1954 in Trinidad from five patients [68] and identified as closely related to Semliki Forest serogroup of viruses [69]. Like CHIKV, MAYV is a member of the genus *Alphavirus*. MAYV has been recorded from many countries in Latin America (Bolivia, Brazil, Colombia, Costa Rica, Ecuador, French Guiana, Guatemala, Guyana, Honduras, Mexico, Panamá, Perú, Suriname, Trinidad and Tobago, and Venezuela (reviewed by Mota and colleagues, 2015 [70]). Clinical presentation of MAYV is non-descript and “flu-like,” similar to other arboviruses such as dengue or chikungunya [71]. Due to the clinical similarities, some have estimated that ~1% of dengue-like illness cases in northern South America may be caused by MAYV, but are misdiagnosed [72]. However, there have been three major epidemics documented to be caused by MAYV. First, in the mid-20th century, an outbreak of febrile illness occurred 120 miles from Belem, Brazil with subsequent isolation of MAYV from six samples and further evidence of MAYV seroconversion in 18.9% of those sampled [73]. Second, an epidemic occurred in 1955 in Bolivia and killed fifteen visiting Japanese pioneers. The etiologic virus was originally named Uruma, but phylogenetic analysis revealed this to be a strain of MAYV [74, 75]. Lastly, in 1977–78, another outbreak in Belterra, Brazil was reported with approximately 20% of residents infected and more than 43 isolates collected [71, 76]. It was during this final outbreak that *Haemagogus janthinomys* was described as a possible vector and that marmosets (*Calithrix argentata*) were described as the potential enzootic reservoir [77]. Importantly, experimental transmission of MAYV by *Ae. aegypti* has been demonstrated, raising the possibility of urban transmission of MAYV in a system similar to DENV, CHIKV, and ZIKV [78]

MAYV is not as well characterized as dengue, CHIKV, or ZIKV. However, historically, two phylogenetic groups have been described: genotypes D and L. Most strains group into genotype D, and are geographically clustered with isolates from Trinidad, Brazil, French Guiana, Surinam, Perú, and Bolivia. Genotype L, on the other hand, is comprised mostly of isolates from the north central region of Brazil [75, 79, 80]. Additionally, Auguste and co-workers suggest the existence of a third lineage called N (New) and further phylogenetic analysis from Llagonne-Barets supports this assertion [81, 82].

Oropouche (Bunyavirus)

Oropouche virus (OROV) is a member of genus *Orthobunyavirus*, which belongs to family *Bunyaviridae*, and its geographic distribution is restricted to the Americas. OROV was isolated for the first time in 1955 from a febrile human in Vega de Oropouche, Trinidad and again from a pool of mosquitoes (*Coquillettidia venezuelensis*) [83]. In the same study, antibodies against OROV were detected in forest workers and non-human primates [83]. In 1960, OROV was isolated in Brazil from the blood of a sloth (*Bradypus tridactylus*) in the proximities of Belem [84]. In 1961, the first known outbreak of OROV was recorded in the Pará state of Brazil, where an estimated 11,000 people were affected by the flu-like illness [84]. At least six other outbreaks of OROV were recorded in Pará over the next decades with *Culex paraensis* implicated as the main vector and biting midges (Culicoides) have been indicated as competent vectors [85–88]. After 1981, additional outbreaks were reported outside of Pará, including the Brazilian states of Amazonas, Amapá, Acre, Rondônia,

Maranhao and Tocantins [89–93]. In 1989, the virus was detected in Panamá during an outbreak in Bejuco, located approximately 60 km west of Panamá City [94]. In 1992, OROV was diagnosed in five patients from Iquitos, Perú who were previously suspected of having DENV [95]. A subsequent serosurvey showed 35% antibody prevalence in urban residents and confirmed infection in a further 28% of residents of rural communities and 2% of urban and forest communities [96]. Since then, there have been consistent reports of OROV infections in Brazil, Panamá, and Perú, and there is evidence of OROV circulation in Ecuador, Bolivia, and Argentina [72, 97, 98].

OROV has a tri-segment, negative-sense RNA genome consisting of the large (L), medium (M) and small (S) segments [99]. Initial phylogenetic analyses of OROV have identified at least three lineages named I, II and III. Lineage I grouped isolates from Trinidad with most of the Brazilian strains. Lineage II contained Peruvian strains and some from Brazil. Finally, lineage III comprised Panamanian strains [100, 101]. Later, with the three initial lineages conserved, a fourth lineage (IV) was introduced consisting of Brazilian strains isolated in Amazonas [102, 103]. In addition, a reassortant of OROV (Madre de Dios Virus) was isolated in Venezuela [104].

Discussion II: Needed Research for Enhanced Preparedness

MAYV and OROV have been thus far restricted to the Americas and responsible for sporadic epidemics nowhere near the scale of ZIKV or CHIKV, likely due to at least a modest level of herd immunity. However, YFV, which is maintained in a sylvatic cycle in South America, has caused smaller outbreaks, including a significant outbreak in 2016–17 [105, 106]. This outbreak was successful even in the face of an effective vaccine and likely at least some pre-existing immunity of the population. Thus, we must conclude that either there are factors that preclude these viruses from establishing in regions where they've not been recorded, or that there is a real possibility that these viruses will pose public health threats in the future. The critical information needed to accurately predict and plan responses to possible emergences of these viruses mirror those questions brought up during the ZIKV and CHIKV experience.

1. Has there been undetected circulation of these viruses and are current diagnostics capable of differentiating related viruses? The combination of often non-specific symptoms, potential for high rates of asymptomatic infections, and a lack of infrastructure for definitive diagnosis may confound detection of these viruses. Thus, while the detection of South American viruses outside the Americas has not been widespread, it does not necessarily indicate the absence of their spread. In Africa, for example, etiologies have been misdiagnosed with the assumption that febrile illnesses are most likely to be malaria [107], and reports have shown cross-reactivity between MAYV and CHIKV, at least. Human-lead movement of OROV from Brazil to Peru has been hypothesized via Amazon River traffic, which subsequently seeded a series of self-contained outbreaks [72]. The MAYV outbreak that occurred in 1955 involved 200 foreign workers was in a time where travel was not as fast. Today, travel from South America to Japan is a matter of hours, well within the incubation time for this and most other viruses.

2. What are the vector species most likely to be involved in emergence and what factors make these vectors likely? The presence of a competent species is of course necessary but it may not be sufficient. For example, while DENV has been endemic to SE Asia, YFV has not spread to that region, even though the two share a common urban vector, *Ae. aegypti*. Perhaps this is because to establish a robust transmission cycle, other factors are necessary. Identification and assessment of the relative importance of elements driving regional transmission must be investigated, such as biting behavior, contact rates with human/zoonotic hosts, and the variability in vector competence among strains of the same species across geographic distances.
3. What are the sufficient and necessary elements of an urban and/or sylvatic ecology that would encourage maintenance of these viruses? For example, when a virus is introduced into new regions or ecosystems, new species of mosquitoes may be involved in transmission, resulting in different transmission patterns [108]. Certainly the presence of a robust sylvatic maintenance system has contributed to the long-term success of YFV in South America. Is it a lack of the necessary elements of a robust sylvatic transmission system that hinders YFV spread to Asia, and does a similar hurdle preclude spread of MAYV and OROV across the ocean? The role of different ecologies, potential vector species, and human urban migration patterns should be the focus of predictions for the movement of viruses across the globe and their ultimate transmission trajectories. Understanding of the existing urban and ecological factors of endemic viral success as well as investigations into the alterations of transmission cycles that promote successful expansions are needed to predict and then respond to emerging arboviral pathogens for global public health security.

Conclusions

As CHIKV and ZIKV indicate, emergent viruses are a serious and ongoing threat to global public health. Table 1 shows the reports of exportation of the viruses herein as reported to ProMed Mail (<https://www.promedmail.org>). Currently, understanding the factors of international arbovirus emergence is paramount for reasons such as vector expansion corresponding to climate change, global trade and travel, and increased urbanization and land-use changes such as deforestation [109]. More than likely, these viruses may not have emigrated from the Americas, meaning that populations outside of the Western Hemisphere likely exhibit widespread susceptibility. For that reason, MAYV and OROV have the potential to cause large-scale epidemics in these susceptible populations on the scale of ZIKV and CHIKV. Critically, not only is the community unprepared from a detection standpoint, but also the ecological determinants of emergence success are not understood. Studies of these neglected South American diseases are needed so that the community is prepared for a data-informed response in case of emergence.

Acknowledgments

Funding: This work was funded by NAMRU-6 (contract #N44852-16-P-0016) and by NIH/NIGMS R01GM12207.

References

1. Weaver SC, Reisen WK. Present and future arboviral threats. *Antiviral Res.* 2010; 85(2):328–45. [PubMed: 19857523]
2. Liang G, Gao X, Gould EA. Factors responsible for the emergence of arboviruses; strategies, challenges and limitations for their control. *Emerg Microbes Infect.* 2015; 4(3):e18. These two papers review the recent emergence of arboviruses threats and the potential environmental and anthropologic factors contributing to the seeming increase in emergence events. Marcondes takes a vector-centric view while Liang discusses anthropologic history and present issues contributing to arbovirus emergence. [PubMed: 26038768]
3. Cao-Lormeau VM, Musso D. Emerging arboviruses in the Pacific. *Lancet.* 2014; 384(9954):1571–2. [PubMed: 25443481]
4. Morens DM, Fauci AS. Emerging infectious diseases: threats to human health and global stability. *PLoS Pathog.* 2013; 9(7):e1003467. This reference highlights the overarching problem of a lack of preparedness to tackle emerging diseases of public health concern, regardless of the pathogen. [PubMed: 23853589]
5. Kuno G. A Re-Examination of the History of Etiologic Confusion between Dengue and Chikungunya. *PLoS Negl Trop Dis.* 2015; 9(11):e0004101. [PubMed: 26562299]
6. Weinbren MP. The occurrence of Chikungunya virus in Uganda. II. In man on the Entebbe peninsula. *Trans R Soc Trop Med Hyg.* 1958; 52(3):258–9. [PubMed: 13556870]
7. Weinbren MP, Haddow AJ, Williams MC. The occurrence of Chikungunya virus in Uganda. I. Isolation from mosquitoes. *Trans R Soc Trop Med Hyg.* 1958; 52(3):253–7. [PubMed: 13556869]
8. Hammon WM, Rudnick A, Sather GE. Viruses associated with epidemic hemorrhagic fevers of the Philippines and Thailand. *Science.* 1960; 131(3407):1102–3. [PubMed: 14399343]
9. Carey DE. Chikungunya and dengue: a case of mistaken identity? *J Hist Med Allied Sci.* 1971; 26(3):243–62. [PubMed: 4938938]
10. Weaver SC, Forrester NL. Chikungunya: Evolutionary history and recent epidemic spread. *Antiviral Res.* 2015; 120:32–9. [PubMed: 25979669]
11. Volk SM, Chen R, Tssetsarkin KA, Adams AP, Garcia TI, Sall AA, Nasar F, Schuh AJ, Holmes EC, Higgs S, Maharaj PD, Brault AC, Weaver SC. Genome-scale phylogenetic analyses of chikungunya virus reveal independent emergences of recent epidemics and various evolutionary rates. *J Virol.* 2010; 84(13):6497–504. [PubMed: 20410280]
12. Stapleford KA, Moratorio G, Henningsson R, Chen R, Matheus S, Enfissi A, Weissglas-Volkov D, Isakov O, Blanc H, Mounce BC, Dupont-Rouzeyrol M, Shomron N, Weaver S, Fontes M, Rousset D, Vignuzzi M. Whole-Genome Sequencing Analysis from the Chikungunya Virus Caribbean Outbreak Reveals Novel Evolutionary Genomic Elements. *PLoS Negl Trop Dis.* 2016; 10(1):e0004402. [PubMed: 26807575]
13. Renault P, Solet JL, Sissoko D, Balleydier E, Larrieu S, Filleul L, Lassalle C, Thiria J, Rachou E, de Valk H, Illef D, Ledrans M, Quatresous I, Quenel P, Pierre V. A major epidemic of chikungunya virus infection on Reunion Island, France, 2005–2006. *Am J Trop Med Hyg.* 2007; 77(4):727–31. [PubMed: 17978079]
14. Tssetsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS Pathog.* 2007; 3(12):e201. [PubMed: 18069894]
15. Vazeille M, Moutailler S, Coudrier D, Rousseaux C, Khun H, Huerre M, Thiria J, Dehecq JS, Fontenille D, Schuffenecker I, Despres P, Failloux AB. Two Chikungunya isolates from the outbreak of La Reunion (Indian Ocean) exhibit different patterns of infection in the mosquito, *Aedes albopictus*. *PLoS One.* 2007; 2(11):e1168. [PubMed: 18000540]
16. Renault P, Jossesan L, Pierre V. Chikungunya-related fatality rates, Mauritius, India, and Reunion Island. *Emerg Infect Dis.* 2008; 14(8):1327.
17. Tssetsarkin KA, Chen R, Leal G, Forrester N, Higgs S, Huang J, Weaver SC. Chikungunya virus emergence is constrained in Asia by lineage-specific adaptive landscapes. *Proc Natl Acad Sci U S A.* 2011; 108(19):7872–7. [PubMed: 21518887]

18. Duong V, Andries AC, Ngan C, Sok T, Richner B, Asgari-Jirhandeh N, Bjorge S, Huy R, Ly S, Laurent D, Hok B, Roces MC, Ong S, Char MC, Deubel V, Tarantola A, Buchy P. Reemergence of Chikungunya virus in Cambodia. *Emerg Infect Dis.* 2012; 18(12):2066–9. [PubMed: 23171736]
19. Yactayo S, Staples JE, Millot V, Cibrelus L, Ramon-Pardo P. Epidemiology of Chikungunya in the Americas. *J Infect Dis.* 2016; 214(suppl 5):S441–S445. [PubMed: 27920170]
20. Lanciotti RS, Lambert AJ. Phylogenetic Analysis of Chikungunya Virus Strains Circulating in the Western Hemisphere. *Am J Trop Med Hyg.* 2016; 94(4):800–3. [PubMed: 26856917]
21. Lanciotti RS, Valadere AM. Transcontinental movement of Asian genotype chikungunya virus. *Emerg Infect Dis.* 2014; 20(8):1400–2. [PubMed: 25076384]
22. Christofferson RC, Chisenhall DM, Wearing HJ, Mores CN. Chikungunya viral fitness measures within the vector and subsequent transmission potential. *PLoS One.* 2014; 9(10):e110538. [PubMed: 25310016]
23. Vega-Rua A, Lourenco-de-Oliveira R, Mousson L, Vazeille M, Fuchs S, Yebakima A, Gustave J, Girod R, Dusfour I, Leparc-Goffart I, Vanlandingham DL, Huang YJ, Lounibos LP, Mohamed Ali S, Nougaiere A, de Lamballerie X, Failloux AB. Chikungunya virus transmission potential by local Aedes mosquitoes in the Americas and Europe. *PLoS Negl Trop Dis.* 2015; 9(5):e0003780. [PubMed: 25993633]
24. Dick GW, Kitchen SF, Haddow AJ. Zika virus. I. Isolations and serological specificity. *Trans R Soc Trop Med Hyg.* 1952; 46(5):509–20. [PubMed: 12995440]
25. Dick GW. Zika virus. II. Pathogenicity and physical properties. *Trans R Soc Trop Med Hyg.* 1952; 46(5):521–34. [PubMed: 12995441]
26. Macnamara FN. Zika virus: a report on three cases of human infection during an epidemic of jaundice in Nigeria. *Trans R Soc Trop Med Hyg.* 1954; 48(2):139–45. [PubMed: 13157159]
27. Marchette NJ, Garcia R, Rudnick A. Isolation of Zika virus from Aedes aegypti mosquitoes in Malaysia. *Am J Trop Med Hyg.* 1969; 18(3):411–5. [PubMed: 4976739]
28. Bearcroft WG. Zika virus infection experimentally induced in a human volunteer. *Trans R Soc Trop Med Hyg.* 1956; 50(5):442–8. [PubMed: 13380987]
29. Olson JG, Ksiazek TG, Triwibowo Suhandiman. Zika virus, a cause of fever in Central Java, Indonesia. *Trans R Soc Trop Med Hyg.* 1981; 75(3):389–93. [PubMed: 6275577]
30. Ye Q, Liu ZY, Han JF, Jiang T, Li XF, Qin CF. Genomic characterization and phylogenetic analysis of Zika virus circulating in the Americas. *Infect Genet Evol.* 2016; 43:43–9. [PubMed: 27156653]
31. Angeletti S, Lo Presti A, Giovanetti M, Grifoni A, Amicosante M, Ciotti M, Alcantara LJ, Cella E, Ciccozzi M. Phylogenesis and homology modeling in Zika virus epidemic: food for thought. *Pathog Glob Health.* 2016; 110(7–8):269–274. [PubMed: 27670692]
32. Shen S, Shi J, Wang J, Tang S, Wang H, Hu Z, Deng F. Phylogenetic analysis revealed the central roles of two African countries in the evolution and worldwide spread of Zika virus. *Virol Sin.* 2016; 31(2):118–30. [PubMed: 27129451]
33. Duffy MR, Chen TH, Hancock WT, Powers AM, Kool JL, Lanciotti RS, Pretrick M, Marfel M, Holzbauer S, Dubray C, Guillaumot L, Griggs A, Bel M, Lambert AJ, Laven J, Kosoy O, Panella A, Biggerstaff BJ, Fischer M, Hayes EB. Zika virus outbreak on Yap Island, Federated States of Micronesia. *N Engl J Med.* 2009; 360(24):2536–43. [PubMed: 19516034]
34. Cao-Lormeau VM, Roche C, Teissier A, Robin E, Berry AL, Mallet HP, Sall AA, Musso D. Zika virus, French polynesia, South pacific, 2013. *Emerg Infect Dis.* 2014; 20(6):1085–6. [PubMed: 24856001]
35. Musso D, Nilles EJ, Cao-Lormeau VM. Rapid spread of emerging Zika virus in the Pacific area. *Clin Microbiol Infect.* 2014; 20(10):O595–6. [PubMed: 24909208]
36. Faria NR, Azevedo RoS, Kraemer MU, Souza R, Cunha MS, Hill SC, Thézé J, Bonsall MB, Bowden TA, Rissanen I, Rocco IM, Nogueira JS, Maeda AY, Vasami FG, Macedo FL, Suzuki A, Rodrigues SG, Cruz AC, Nunes BT, Medeiros DB, Rodrigues DS, Nunes Queiroz AL, da Silva EV, Henriques DF, Travassos da Rosa ES, de Oliveira CS, Martins LC, Vasconcelos HB, Casseb LM, De Simith B, Messina JP, Abade L, Lourenço J, Carlos Junior Alcantara L, de Lima MM, Giovanetti M, Hay SI, de Oliveira RS, Lemos PaS, de Oliveira LF, de Lima CP, da Silva SP, de Vasconcelos JM, Franco L, Cardoso JF, Vianez-Júnior JL, Mir D, Bello G, Delatorre E, Khan K, Creatore M, Coelho GE, de Oliveira WK, Tesh R, Pybus OG, Nunes MR, Vasconcelos PF. Zika

- virus in the Americas: Early epidemiological and genetic findings. *Science*. 2016; 352(6283):345–9. [PubMed: 27013429]
37. Weaver SC, Costa F, Garcia-Blanco MA, Ko AI, Ribeiro GS, Saade G, Shi PY, Vasilakis N. Zika virus: History, emergence, biology, and prospects for control. *Antiviral Res*. 2016; 130:69–80. [PubMed: 26996139]
 38. Musso D. Zika Virus Transmission from French Polynesia to Brazil. *Emerg Infect Dis*. 2015; 21(10):1887.
 39. Cardoso CW I, Paploski A, Kikuti M, Rodrigues MS, Silva MM, Campos GS, Sardi SI, Kitron U, Reis MG, Ribeiro GS. Outbreak of Exanthematous Illness Associated with Zika, Chikungunya, and Dengue Viruses, Salvador, Brazil. *Emerg Infect Dis*. 2015; 21(12):2274–6. [PubMed: 26584464]
 40. Brito CA, Brito CC, Oliveira AC, Rocha M, Atanásio C, Asfora C, Matos JD, Lima AS, Albuquerque MF. Zika in Pernambuco: rewriting the first outbreak. *Rev Soc Bras Med Trop*. 2016; 49(5):553–558. [PubMed: 27812648]
 41. Musso D, Gubler DJ. Zika Virus. *Clin Microbiol Rev*. 2016; 29(3):487–524. [PubMed: 27029595]
 42. Besnard M, Lastere S, Teissier A, Cao-Lormeau V, Musso D. Evidence of perinatal transmission of Zika virus, French Polynesia, December 2013 and February 2014. *Euro Surveill*. 2014; 19(13)
 43. de Paula Freitas B, de Oliveira Dias JR, Prazeres J, Sacramento GA, Ko AI, Maia M, Belfort R Jr. Ocular Findings in Infants With Microcephaly Associated With Presumed Zika Virus Congenital Infection in Salvador, Brazil. *JAMA Ophthalmol*. 2016
 44. Heukelbach J, Werneck GL. Surveillance of Zika virus infection and microcephaly in Brazil. *Lancet*. 2016; 388(10047):846–7. [PubMed: 27372396]
 45. Oehler E, Watrin L, Larre P, Leparc-Goffart I, Lastere S, Valour F, Baudouin L, Mallet H, Musso D, Ghawche F. Zika virus infection complicated by Guillain-Barre syndrome--case report, French Polynesia, December 2013. *Euro Surveill*. 2014; 19(9)
 46. Tomori O. Impact of yellow fever on the developing world. *Adv Virus Res*. 1999; 53:5–34. [PubMed: 10582093]
 47. Augustin, G. S.a.P. Ltd. History of yellow fever. New Orleans, LA, USA: 1909. p. 1230
 48. Finlay, Carlos J. (1833–1915) student of yellow fever. *JAMA*. 1966; 198(11):1210–1. [PubMed: 5332541]
 49. Reed W, Carroll J, Agramonte A. The etiology of yellow fever: an additional note. 1901. *Mil Med*. 2001; 166(9 Suppl):44–53.
 50. Reed W, Carroll J, Agramonte A. The etiology of yellow fever: an additional note. *J Amer Med Assoc*. 1901; 431–440:431–440.
 51. Reed W, Carroll J. The Prevention of Yellow Fever. *Public Health Pap Rep*. 1901; 27:113–29. [PubMed: 19600973]
 52. Stokes A, Bauer JH, Hudson NP. The transmission of yellow fever to *Macacus rhesus*: Preliminary note The transmission of yellow fever to *Macacus rhesus*: Preliminary note. *J Amer Med Assoc*. 1928; 90(4):253–254.
 53. Theiler M. SUSCEPTIBILITY OF WHITE MICE TO THE VIRUS OF YELLOW FEVER. *Science*. 1930; 71(1840):367.
 54. Nasidi A, Monath TP, DeCock K, Tomori O, Cordellier R, Olaleye OD, Harry TO, Adeniyi JA, Sorungbe AO, Ajose-Coker AO. Urban yellow fever epidemic in western Nigeria, 1987. *Trans R Soc Trop Med Hyg*. 1989; 83(3):401–6. [PubMed: 2617590]
 55. Romano AP, Costa ZG, Ramos DG, Andrade MA, Jayme VeS, Almeida MA, Vettorello KC, Mascheretti M, Flannery B. Yellow Fever outbreaks in unvaccinated populations, Brazil, 2008–2009. *PLoS Negl Trop Dis*. 2014; 8(3):e2740. [PubMed: 24625634]
 56. Bryant JE, Holmes EC, Barrett AD. Out of Africa: a molecular perspective on the introduction of yellow fever virus into the Americas. *PLoS Pathog*. 2007; 3(5):e75. [PubMed: 17511518]
 57. Wang E, Weaver SC, Shope RE, Tesh RB, Watts DM, Barrett AD. Genetic variation in yellow fever virus: duplication in the 3' noncoding region of strains from Africa. *Virology*. 1996; 225(2): 274–81. [PubMed: 8918913]

58. Mutebi JP, Wang H, Li L, Bryant JE, Barrett AD. Phylogenetic and evolutionary relationships among yellow fever virus isolates in Africa. *J Virol.* 2001; 75(15):6999–7008. [PubMed: 11435580]
59. Anderson KB, Thomas SJ, Endy TP. The Emergence of Zika Virus: A Narrative Review. *Ann Intern Med.* 2016; 165(3):175–83. [PubMed: 27135717]
60. Wong PS, Li MZ, Chong CS, Ng LC, Tan CH. *Aedes (Stegomyia) albopictus* (Skuse): a potential vector of Zika virus in Singapore. *PLoS Negl Trop Dis.* 2013; 7(8):e2348. [PubMed: 23936579]
61. Chouin-Carneiro T, Vega-Rua A, Vazeille M, Yebakima A, Girod R, Goindin D, Dupont-Rouzeyrol M, Lourenço-de-Oliveira R, Failloux AB. Differential Susceptibilities of *Aedes aegypti* and *Aedes albopictus* from the Americas to Zika Virus. *PLoS Negl Trop Dis.* 2016; 10(3):e0004543. [PubMed: 26938868]
62. Guerbois M, Fernandez-Salas I, Azar SR, Danis-Lozano R, Alpuche-Aranda CM, Leal G, Garcia-Malo IR, Diaz-Gonzalez EE, Casas-Martinez M, Rossi SL, Del Rio-Galvan SL, Sanchez-Casas RM, Roundy CM, Wood TG, Widen SG, Vasilakis N, Weaver SC. Outbreak of Zika Virus Infection, Chiapas State, Mexico, 2015, and First Confirmed Transmission by *Aedes aegypti* Mosquitoes in the Americas. *J Infect Dis.* 2016; 214(9):1349–1356. [PubMed: 27436433]
63. Fernandez-Salas I, Diaz-Gonzalez EE, Lopez-Gatell H, Alpuche-Aranda C. Chikungunya and zika virus dissemination in the Americas: different arboviruses reflecting the same spreading routes and poor vector-control policies. *Curr Opin Infect Dis.* 2016; 29(5):467–75. [PubMed: 27472289]
64. Kramer LD, Ciota AT. Dissecting vectorial capacity for mosquito-borne viruses. *Curr Opin Virol.* 2015; 15:112–8. [PubMed: 26569343]
65. Gardner CL, Ryman KD. Yellow fever: a reemerging threat. *Clin Lab Med.* 2010; 30(1):237–60. [PubMed: 20513550]
66. Felix AC, Souza NCS, Figueiredo WM, Costa AA, Inenami M, da Silva RMG, Levi JE, Pannuti CS, Romano CM. Cross reactivity of commercial anti-dengue immunoassays in patients with acute Zika virus infection. *J Med Virol.* 2017; 89(8):1477–1479. [PubMed: 28229481]
67. Hassing RJ, Leparac-Goffart I, Tolou H, van Doornum G, van Genderen PJ. Cross-reactivity of antibodies to viruses belonging to the Semliki forest serocomplex. *Euro Surveill.* 2010; 15(23)
68. Anderson CR, Downs WG, Wattlely GH, Ahin NW, Reese AA. Mayaro virus: a new human disease agent. II. Isolation from blood of patients in Trinidad, B.W.I. *Am J Trop Med Hyg.* 1957; 6(6):1012–6. [PubMed: 13487973]
69. Casals J, Whitman L. Mayaro virus: a new human disease agent. I. Relationship to other arbor viruses. *Am J Trop Med Hyg.* 1957; 6(6):1004–11. [PubMed: 13487972]
70. de Oliveira Mota MT, Ribeiro MR, Vedovello D, Nogueira ML. Mayaro virus: a neglected arbovirus of the Americas. *Future virology.* 2015; 10(9):73–86.
71. Pinheiro FP, Freitas RB, Travassos da Rosa JF, Gabbay YB, Mello WA, LeDuc JW. An outbreak of Mayaro virus disease in Belterra, Brazil. I. Clinical and virological findings. *Am J Trop Med Hyg.* 1981; 30(3):674–81. [PubMed: 6266263]
72. Forshey BM, Guevara C, Laguna-Torres VA, Cespedes M, Vargas J, Gianella A, Vallejo E, Madrid C, Aguayo N, Gotuzzo E, Suarez V, Morales AM, Beingolea L, Reyes N, Perez J, Negrete M, Rocha C, Morrison AC, Russell KL, Blair PJ, Olson JG, Kochel TJ, Group NFSW. Arboviral etiologies of acute febrile illnesses in Western South America, 2000–2007. *PLoS Negl Trop Dis.* 2010; 4(8):e787. [PubMed: 20706628]
73. Causey OR, Maroja OM. Mayaro virus: a new human disease agent. III. Investigation of an epidemic of acute febrile illness on the river Guama in Pará, Brazil, and isolation of Mayaro virus as causative agent. *Am J Trop Med Hyg.* 1957; 6(6):1017–23. [PubMed: 13487974]
74. Schaeffer M, Gajdusek DC, Lema AB, Eichenwald H. Epidemic jungle fevers among Okinawan colonists in the Bolivian rain forest. I. Epidemiology. *Am J Trop Med Hyg.* 1959; 8(3):372–96. [PubMed: 13661542]
75. Powers AM, Aguilar PV, Chandler LJ, Brault AC, Meakins TA, Watts D, Russell KL, Olson J, Vasconcelos PF, Da Rosa AT, Weaver SC, Tesh RB. Genetic relationships among Mayaro and Una viruses suggest distinct patterns of transmission. *Am J Trop Med Hyg.* 2006; 75(3):461–9. [PubMed: 16968922]

76. LeDuc JW, Pinheiro FP, Travassos da Rosa AP. An outbreak of Mayaro virus disease in Belterra, Brazil. II. Epidemiology. *Am J Trop Med Hyg.* 1981; 30(3):682–8. [PubMed: 6266264]
77. Hoch AL, Peterson NE, LeDuc JW, Pinheiro FP. An outbreak of Mayaro virus disease in Belterra, Brazil. III. Entomological and ecological studies. *Am J Trop Med Hyg.* 1981; 30(3):689–98. [PubMed: 6266265]
78. Long KC, Ziegler SA, Thangamani S, Hausser NL, Kochel TJ, Higgs S, Tesh RB. Experimental transmission of Mayaro virus by *Aedes aegypti*. *Am J Trop Med Hyg.* 2011; 85(4):750–7. [PubMed: 21976583]
79. Estofolete CF, Mota MT, Vedovello D, Góngora DV, Maia IL, Nogueira ML. Mayaro fever in an HIV-infected patient suspected of having Chikungunya fever. *Rev Soc Bras Med Trop.* 2016; 49(5):648–652. [PubMed: 27812665]
80. Serra OP, Cardoso BF, Ribeiro AL, Santos FA, Silhessarenko RD. Mayaro virus and dengue virus 1 and 4 natural infection in culicids from Cuiabá, state of Mato Grosso, Brazil. *Mem Inst Oswaldo Cruz.* 2016; 111(1):20–9. [PubMed: 26784852]
81. Auguste AJ, Liria J, Forrester NL, Giambalvo D, Moncada M, Long KC, Morón D, de Manzione N, Tesh RB, Halsey ES, Kochel TJ, Hernandez R, Navarro JC, Weaver SC. Evolutionary and Ecological Characterization of Mayaro Virus Strains Isolated during an Outbreak, Venezuela, 2010. *Emerg Infect Dis.* 2015; 21(10):1742–50. [PubMed: 26401714]
82. Llagonne-Barets M, Icard V, Leparç-Goffart I, Prat C, Perpoint T, André P, Ramière C. A case of Mayaro virus infection imported from French Guiana. *J Clin Virol.* 2016; 77:66–8. [PubMed: 26921736]
83. Anderson CR, Spence L, Downs WG, Aitken TH. Oropouche virus: a new human disease agent from Trinidad, West Indies. *Am J Trop Med Hyg.* 1961; 10:574–8. [PubMed: 13683183]
84. Pinheiro FP, Pinheiro M, Bensabath G, Causey OR, Shope R. Epidemia de vírus Oropouche em Belém. *Rev Serv Esp Saúde públ.* 1962; 12(1):15–23.
85. Pinheiro FP, Travassos da Rosa AP, Gomes ML, LeDuc JW, Hoch AL. Transmission of Oropouche virus from man to hamster by the midge *Culicoides paraensis*. *Science.* 1982; 215(4537):1251–3. [PubMed: 6800036]
86. Pinheiro FP, Travassos da Rosa AP, Travassos da Rosa JF, Ishak R, Freitas RB, Gomes ML, LeDuc JW, Oliva OF. Oropouche virus. I. A review of clinical, epidemiological, and ecological findings. *Am J Trop Med Hyg.* 1981; 30(1):149–60. [PubMed: 6782898]
87. Pinheiro FP, Hoch AL, Gomes ML, Roberts DR. Oropouche virus. IV. Laboratory transmission by *Culicoides paraensis*. *Am J Trop Med Hyg.* 1981; 30(1):172–6. [PubMed: 7212164]
88. Mercer DR, Castillo-Pizango MJ. Changes in relative species compositions of biting midges (Diptera: Ceratopogonidae) and an outbreak of Oropouche virus in Iquitos, Peru. *J Med Entomol.* 2005; 42(4):554–8. [PubMed: 16119543]
89. Borborema CA, Pinheiro FP, Albuquerque BC, da Rosa AP, da Rosa JF, Dourado HV. 1st occurrence of outbreaks caused by Oropouche virus in the State of Amazonas. *Rev Inst Med Trop Sao Paulo.* 1982; 24(3):132–9. [PubMed: 6818662]
90. Rosa AP, Rodrigues SG, Nunes MR, Magalhães MT, Rosa JF, Vasconcelos PF. Outbreak of oropouche virus fever in Serra Pelada, municipality of Curionópolis, Pará, 1994. *Rev Soc Bras Med Trop.* 1996; 29(6):537–41. [PubMed: 9011877]
91. Vasconcelos PF, Travassos Da Rosa JF, Guerreiro SC, Dégallier N, Travassos Da Rosa ES, Travassos Da Rosa AP. 1st register of an epidemic caused by Oropouche virus in the states of Maranhão and Goiás, Brazil. *Rev Inst Med Trop Sao Paulo.* 1989; 31(4):271–8. [PubMed: 2516642]
92. LeDuc JW, Hoch AL, Pinheiro FP, da Rosa AP. Epidemic Oropouche virus disease in northern Brazil. *Bull Pan Am Health Organ.* 1981; 15(2):97–103. [PubMed: 7326505]
93. Vasconcelos HB, Azevedo RS, Casseb SM, Nunes-Neto JP, Chiang JO, Cantuária PC, Segura MN, Martins LC, Monteiro HA, Rodrigues SG, Nunes MR, Vasconcelos PF. Oropouche fever epidemic in Northern Brazil: epidemiology and molecular characterization of isolates. *J Clin Virol.* 2009; 44(2):129–33. [PubMed: 19117799]
94. Pinheiro, FP., Travassos da Rosa, A., Vasconcelos, P. Oropouche fever. In: Feigin, RD., editor. *Textbook of pediatric infectious diseases*. 5. Editora Saunders; Philadelphia: 2004. p. 2418-2423.

95. Chavez R, Colan E, Phillips I. Fiebre de Oropuche en Iquitos: Reporte preliminar de 5 casos. *Rev farmacol terap (Lima)*. 1992; 2(1):12–14.
96. Tesh RB. The emerging epidemiology of Venezuelan hemorrhagic fever and Oropouche fever in tropical South America. *Ann N Y Acad Sci*. 1994; 740:129–37. [PubMed: 7840443]
97. Manock SR, Jacobsen KH, de Bravo NB, Russell KL, Negrete M, Olson JG, Sanchez JL, Blair PJ, Smalligan RD, Quist BK, Espín JF, Espinoza WR, MacCormick F, Fleming LC, Kochel T. Etiology of acute undifferentiated febrile illness in the Amazon basin of Ecuador. *Am J Trop Med Hyg*. 2009; 81(1):146–51. [PubMed: 19556580]
98. Nunes, M., Vasconcelos, H., Medeiros, D., Rodrigues, S., Azevedo, R., Chiang, J., Martins, L., Vasconcelos, P. Oropouche fever: an overview of the epidemiological and molecular aspects in the brazilian Amazon region. Vol. 15. *Cad Saúde Colet*; Rio de Janeiro: 2007. p. 303-318.
99. Elliott RM. Orthobunyaviruses: recent genetic and structural insights. *Nat Rev Microbiol*. 2014; 12(10):673–85. [PubMed: 25198140]
100. Saeed MF, Wang H, Nunes M, Vasconcelos PF, Weaver SC, Shope RE, Watts DM, Tesh RB, Barrett AD. Nucleotide sequences and phylogeny of the nucleocapsid gene of Oropouche virus. *J Gen Virol*. 2000; 81(Pt 3):743–8. [PubMed: 10675412]
101. Azevedo RS, Nunes MR, Chiang JO, Bensabath G, Vasconcelos HB, Pinto AY, Martins LC, Monteiro HA, Rodrigues SG, Vasconcelos PF. Reemergence of Oropouche fever, northern Brazil. *Emerg Infect Dis*. 2007; 13(6):912–5. [PubMed: 17553235]
102. Vasconcelos HB, Nunes MR, Casseb LM, Carvalho VL, Pinto da Silva EV, Silva M, Casseb SM, Vasconcelos PF. Molecular epidemiology of Oropouche virus, Brazil. *Emerg Infect Dis*. 2011; 17(5):800–6. [PubMed: 21529387]
103. Travassos da Rosa JF, de Souza WM, Pinheiro FP, Figueiredo ML, Cardoso JF, Acrani GO, Nunes MRT. Oropouche Virus: Clinical, Epidemiological, and Molecular Aspects of a Neglected Orthobunyavirus. *Am J Trop Med Hyg*. 2017; 96(5):1019–1030. [PubMed: 28167595]
104. Navarro JC, Giambalvo D, Hernandez R, Auguste AJ, Tesh RB, Weaver SC, Montañez H, Liria J, Lima A, Travassos da Rosa JF, da Silva SP, Vasconcelos JM, Oliveira R, Vianez JL, Nunes MR. Isolation of Madre de Dios Virus (Orthobunyavirus; Bunyaviridae), an Oropouche Virus Species Reassortant, from a Monkey in Venezuela. *Am J Trop Med Hyg*. 2016; 95(2):328–38. [PubMed: 27215299]
105. Saad LD, Barata RB. Yellow fever outbreaks in Sao Paulo State, Brazil, 2000–2010. *Epidemiol Serv Saude*. 2016; 25(3):531–540. [PubMed: 27869924]
106. Ortiz-Martinez Y, Patino-Barbosa AM, Rodriguez-Morales AJ. Yellow fever in the Americas: the growing concern about new epidemics. *F1000Res*. 2017; 6:398. [PubMed: 28529708]
107. Amexo M, Tolhurst R, Barnish G, Bates I. Malaria misdiagnosis: effects on the poor and vulnerable. *Lancet*. 2004; 364(9448):1896–8. [PubMed: 15555670]
108. Higgs S. Zika Virus: Emergence and Emergency. *Vector Borne Zoonotic Dis*. 2016; 16(2):75–6. [PubMed: 26824625]
109. Romero-Alvarez D, Escobar LE. Vegetation loss and the 2016 Oropouche fever outbreak in Peru. *Mem Inst Oswaldo Cruz*. 2017; 112(4):292–298. [PubMed: 28327792]

Table 1

Exports of Zika, chikungunya, yellow fever, and Mayaro viruses due to returning travelers as reported in PromedMail. There have been no reports of Oropouche virus exports.

Virus	Imported To	Exported From	link
Zika	Australia	Cook Islands	https://www.promedmail.org/post/2378034
Zika	Canada	Thailand	https://www.promedmail.org/post/1744108
Zika	Chile (Easter Island)	Tahiti	https://www.promedmail.org/post/2804821
Zika	Germany	Malaysia	https://www.promedmail.org/post/3367412
Zika	Germany	Thailand	https://www.promedmail.org/post/2139786
Zika	Mexico	Columbia	https://www.promedmail.org/post/3818813
Zika	Japan	French Polynesia	https://www.promedmail.org/post/2126046
Zika	Japan	Thailand	https://www.promedmail.org/post/2716731
Zika	Micronesia: Guam	Yap	https://www.promedmail.org/post/15805
Zika	Netherlands	Suriname	https://www.promedmail.org/post/3858300
Zika	New Caledonia	French Polynesia	https://www.promedmail.org/post/2082225
Zika	New Zealand	Cook Islands	https://www.promedmail.org/post/2762121
chikungunya	New Caledonia	Tonga	https://www.promedmail.org/post/2409660
chikungunya	Portugal	Angola	https://www.promedmail.org/post/2338670
chikungunya	Australia	Papua New Guinea	https://www.promedmail.org/post/1974825
chikungunya	Japan	Cambodia	https://www.promedmail.org/post/1909917
chikungunya	Australia	Indonesia	https://www.promedmail.org/post/1798136
chikungunya	Australia	Papua New Guinea	https://www.promedmail.org/post/1798136
chikungunya	Australia	Papua New Guinea	https://www.promedmail.org/post/1675486
chikungunya	New Guinea, Australia	Indonesia (Bali)	https://www.promedmail.org/post/1594512
chikungunya	Germany	Indonesia (Bali)	https://www.promedmail.org/post/1562598
chikungunya	Malaysia (SL)	Malaysia (PK)	https://www.promedmail.org/post/1040110
chikungunya	New Caledonia	Indonesia	https://www.promedmail.org/post/663016
chikungunya	France	Benin	https://www.promedmail.org/post/631448
chikungunya	Brazil	India & Indonesia	https://www.promedmail.org/post/601534
chikungunya	France	Reunion	https://www.promedmail.org/post/423855
chikungunya	Taiwan	Indonesia	https://www.promedmail.org/post/390011
chikungunya	France	Maldives	https://www.promedmail.org/post/388657
chikungunya	Germany	Maldives	https://www.promedmail.org/post/297273
chikungunya	Belgium	Thailand, India	https://www.promedmail.org/post/212872
chikungunya	Germany	S. E. Asia	https://www.promedmail.org/post/170276
chikungunya	Japan	Malaysia	https://www.promedmail.org/post/153351
chikungunya	Australia	Malaysia (Johor)	https://www.promedmail.org/post/141130
chikungunya	Singapore, Italy	Sri Lanka	https://www.promedmail.org/post/102398
chikungunya	Singapore	Indonesia, India (Kerala)	https://www.promedmail.org/post/78513
chikungunya	Taiwan	Indonesia	https://www.promedmail.org/post/75879
chikungunya	Hong Kong	Sri Lanka	https://www.promedmail.org/post/54478
chikungunya	Taiwan	Indonesia	https://www.promedmail.org/post/47575

Virus	Imported To	Exported From	link
chikungunya	France	Madagascar	https://www.promedmail.org/post/8163
chikungunya	Japan	Sri Lanka	https://www.promedmail.org/post/2209612
chikungunya	Taipei	Singapore	https://www.promedmail.org/post/4792
chikungunya	USA	Asia, Africa	https://www.promedmail.org/post/4759
chikungunya	France (Bourdeau)	Senegal	https://www.promedmail.org/post/4622
chikungunya	French Guiana	Madagascar	https://www.promedmail.org/post/2209424
yellow fever	China	Angola	https://www.promedmail.org/post/4179477
yellow fever	China	Angola	https://www.promedmail.org/post/4136699
yellow fever	Mauritania	Angola	https://www.promedmail.org/post/4119691
yellow fever	China	Angola	https://www.promedmail.org/post/4119422
yellow fever	China	Angola	https://www.promedmail.org/post/4109684
yellow fever	China	Angola	https://www.promedmail.org/post/4106312
yellow fever	China	Angola	https://www.promedmail.org/post/4105105
yellow fever	Kenya	Angola	https://www.promedmail.org/post/4104819
yellow fever	Kenya	Angola	https://www.promedmail.org/post/4097945
yellow fever	China	Angola	https://www.promedmail.org/post/4089857
yellow fever	Senegal	Gambia	https://www.promedmail.org/post/545254
yellow fever	Brazil (MG)	Brazil (RS)	https://www.promedmail.org/post/152217
yellow fever	Paraguay	Argentina (MN)	https://www.promedmail.org/post/146084
yellow fever	Belgium	Gambia	https://www.promedmail.org/post/2201007
yellow fever	Belgium	Gambia	https://www.promedmail.org/post/2200994
yellow fever	Belgium	Gambia	https://www.promedmail.org/post/2200990
yellow fever	Brazil (Brasilia)	Brazil (W. Central)	https://www.promedmail.org/post/2197496
yellow fever	Netherlands	Suriname	https://www.promedmail.org/post/2197331
yellow fever	Netherlands	Suriname	https://www.promedmail.org/post/2197328
yellow fever	Netherlands	Suriname	https://www.promedmail.org/post/2197319
yellow fever	Brazil (Brasilia)	Brazil (W. Central)	https://www.promedmail.org/post/2197244
yellow fever	Brazil (Brasilia)	Brazil (W. Central)	https://www.promedmail.org/post/2197216
yellow fever	USA	Venezuela	https://www.promedmail.org/post/2196966
yellow fever	USA	Venezuela	https://www.promedmail.org/post/2196947
yellow fever	Germany	Cote D'ivoire	https://www.promedmail.org/post/2196220
Mayaro	France	Brazil	https://www.promedmail.org/post/444435
Mayaro	France	Brazil	https://www.promedmail.org/post/423267