Published in final edited form as:

Parasitology. 2017 September ; 144(10): 1288–1301. doi:10.1017/S0031182017000488.

The role of rare innate immune cells in Type 2 immune activation against parasitic helminths

Lauren M. Webb¹ and Elia D. Tait Wojno¹

¹Baker Institute for Animal Health, Department of Microbiology and Immunology, Cornell University College of Veterinary Medicine, 235 Hungerford Hill Road, Ithaca, NY 14853, USA

Summary

The complexity of helminth macroparasites is reflected in the intricate network of host cell types that participate in the Type 2 immune response needed to battle these organisms. In this context, adaptive T helper 2 cells and the Type 2 cytokines interleukin (IL)-4, IL-5, IL-9 and IL-13 have been the focus of research for years, but recent work has demonstrated that the innate immune system plays an essential role. Some innate immune cells that promote Type 2 immunity are relatively abundant, such as macrophages and eosinophils. However, we now appreciate that more rare cell types including group 2 innate lymphoid cells, basophils, mast cells and dendritic cells make significant contributions to these responses. These cells are found at low frequency but they are specialized to their roles – located at sites such as the skin, lung and gut, where the host combats helminth parasites. These cells respond rapidly and robustly to worm antigens and worm-induced damage to produce essential cytokines, chemokines, eicosanoids and histamine to activate damaged epithelium and to recruit other effectors. Thus, a greater understanding of how these cells operate is essential to understanding how the host protects itself during helminth infection.

Keywords

innate; helminth; Type 2; basophil; ILC2; dendritic; mast cell; mucosal

Introduction

Helminths are complex macroparasites with large genomes and a highly evolved capacity to infect mammalian hosts, sometimes persisting in a single host for decades. Parasitic helminths infect over two billion of the world's population, many of whom suffer significant levels of morbidity and disability as a result. Parasitic helminths are also important pathogens of domestic livestock, inflicting significant economic cost on the agricultural industry. Type 2 immune activation is the primary protective immune response against these multicellular macroparasites. This response can lead to expulsion of the parasite and resolution of infection, but this is not always the case. In many situations, Type 2 immunity functions essentially as a wound repair mechanism that protects the host from worm-induced damage. Both helminth expulsion and wound repair mechanisms require host production of the Type 2 cytokines IL-4, IL-5, IL-9 and IL-13 (Allen and Maizels, 2011; Gause *et al.*,

To whom correspondence should be addressed: Elia D. TaitWojno. elia.taitwojno@cornell.edu.

2013). Up until recently the overarching view was that the primary mediator of this immune response was the adaptive T helper 2 (Th2) CD4⁺ T cell population, which is a rich source of Type 2 cytokines following helminth-induced activation (Van Dyken *et al.*, 2016). However, more recently we have come to appreciate the complex interplay of the innate and adaptive immune systems during helminth infection. Work in the last decade has revealed how cells of the innate immune system contribute to Type 2 immune responses, both as early instigators of Type 2 immune activation – as key sources of Type 2 cytokines – and as important downstream effectors that orchestrate modulation of the tissue microenvironment (Fig. 1).

The innate immune system is comprised of many different cell types including some that are found in relative abundance during inflammation such as macrophages, monocytes, eosinophils and neutrophils. Other innate immune cell types are much more rare, making up less than 5% of the circulating leukocyte population. These include group 2 innate lymphoid cells (ILC2s), basophils, mast cells (MCs) and dendritic cells (DCs). Many other reviews have focused on the importance of the more abundant cell types, such as macrophages and eosinophils (Murray and Wynn, 2011; Gause *et al.*, 2013; Huang and Appleton, 2016). Thus, here we will focus on the importance of rare innate immune cell populations, which make up for what they lack in abundance with a potent capacity to modulate the immune microenvironment. The main focus of this review will be recent important findings highlighting the role of ILC2s, basophils, MCs and DCs in Type 2 inflammation against helminths.

Rare innate immune cells – Right time, right place?

ILC2s, basophils, MCs and DCs are highly specialized for their specific roles, be that activation in response to IgE or antigen, or a capacity to rapidly release large quantities of immune chemical messengers such as cytokines, chemokines, histamine and prostaglandins. Thus, while these innate immune cells may be rare, they are poised to perform their specific role robustly. Further, while these cells can be found throughout the body in lymphoid and mucosal tissues, they are particularly enriched in sites that are subject to invasion by helminth parasites and other pathogens (Rivera *et al.*, 2016) such as the skin, lung and gut. Since helminth infections often affect more than one mucosal or barrier tissue, the conserved presence of these cells at multiple sites ensures a coordinated response on many fronts. The importance of these cells at mucosal surfaces is clear, but their influence on immune responses in lymphoid tissues should not be underestimated. We know that DC migration to the lymph node is essential for DC function in priming naïve T cells. ILC2s and granulocytes can also be found in lymphoid tissues and may influence the formation of adaptive immune activation by their potent production of inflammatory mediators (Pulendran and Artis, 2012). In addition, these rare innate immune cells are uniquely specialized to instigate immune activation promptly. ILC2s have been shown to employ genomic adaptations that maintain important genes in open states that allow rapid transcription of effector molecules in response to stimulation (Shih *et al.*, 2016), and future studies will address if this is also the case in other innate cell types. Basophils and MCs also maintain stores of premade mediators, such as cytokines, eicosanoids and histamine, and are ready to rapidly release high concentrations of these chemicals at sites of inflammation

(Voehringer, 2013). Finally, rare innate cell function is highly tunable by effector molecules found in the microenvironment, and these cells are responsive to multiple pathways that control molecular mechanisms of activation and gene expression, such as worm-induced damage signals from the epithelium, Notch signaling and eicasonoids (Rivera *et al.*, 2016; Zhang *et al.*, 2017). Hence, innate cell function can be rapidly modulated by tissue factors, making rare innate cells highly responsive to changes in the immune microenvironment and key players in Type 2 immune activation.

ILC2s – Unique cytokine power houses of the innate immune system

ILC2s are the most recently described rare innate immune cell that we will discuss. The existence of a novel innate cell population that could influence Type 2 immunity was first reported in the early 2000s (Fallon *et al.*, 2006). ILC2s were fully described in 2010 by three independent laboratories as a non-B, non-T cell population that drove Type 2 immunity in mice (Moro *et al.*, 2010; Neill *et al.*, 2010; Price *et al.*, 2010). They are characterized by their lack of expression of markers for common cell lineages and their expression of c-kit, CD90 (Thy1) and CD127 (IL-7R) (Moro *et al.*, 2010; Neill *et al.*, 2010; Price *et al.*, 2010). ILC2s arise from a common lymphoid progenitor found in the bone marrow. Their development and function is dependent on a range of transcription factors (TFs), including ROR α , Id2, GATA-3 and ETS1 (Cherrier *et al.*, 2012; Hoyler *et al.*, 2012; Zook *et al.*, 2016; Klose and Artis, 2016).

ILC2s are found in the skin (Kim *et al.*, 2013) and at mucosal surfaces, as well as other distal sites and lymphoid tissues. At these sites, they are poised to respond to a range of signals, including epithelial-derived damage signals elicited by worm feeding or migration, through their expression of receptors for the cytokines IL-25 (IL-17RB), IL-33 (ST2), and thymic stromal lymphopoietin (TSLP) (Tait Wojno and Artis, 2016). In response to IL-25, IL-33, and TSLP, ILC2s potentially produce large quantities of IL-5 and IL-13 (Moro *et al.*, 2010; Neill *et al.*, 2010; Price *et al.*, 2010), which is likely important for inducing eosinophil recruitment, epithelial cell hyperplasia, tuft cell differentiation and goblet cell activity – all mechanisms involved in worm expulsion (Cliffe *et al.*, 2005; Hammad and Lambrecht, 2015; Gerbe *et al.*, 2016; Moltke *et al.*, 2016). In addition, epithelial cell-derived IL-33 and TSLP induce IL-9 release from ILC2s that acts in an autocrine manner to potentiate production of IL-5, IL-13 and the growth factor amphiregulin during infection with *Nippostrongylus brasiliensis* (Turner *et al.*, 2013; Mohapatra *et al.*, 2015), a mouse-adapted helminth used as a model for intestinal hookworm infection (Camberis *et al.*, 2003). Recent studies have identified a key role for tuft cells, a population of specialized epithelial cells that expand upon worm infection and respond to IL-4 by producing large amounts of IL-25, in inducing IL-13 production by ILC2s (Gerbe *et al.*, 2016; Moltke *et al.*, 2016; Howitt *et al.*, 2016). Thus, ILC2s are particularly tuned to the state of the epithelium and respond to damage signals with rapid and robust production of Type 2 cytokines.

Notably, the ability of ILC2s to respond to signals from the damaged epithelium also promotes their interactions with other rare innate immune cell types to coordinate Type 2 immunity. ILC2s respond to IL-33 that has been cleaved by proteases released by MCs, leading to the potent expansion of the ILC2 population, demonstrating cooperation between

these two cell types that enhances the function of ILC2s (Lefrançois *et al.*, 2014). ILC2s have also been shown to act on DCs, with ILC2-derived IL-13 inducing migration of DCs into the draining lymph node (dLN), where DCs prime Th2 cells. ILC2-derived IL-13 is also required for CCL-17 production by DCs, which recruits activated Th2 cells into the tissue during Type 2 inflammation (Halim *et al.*, 2014; 2015), further demonstrating how innate cells work in a complex network to drive Type 2 inflammatory responses.

In addition to their production of IL-5 and IL-13, ILC2s promote the development of Th2 activation by interacting with adaptive CD4⁺ T cells. In this context, ILC2s are an essential source of IL-4 for the effective development of lamina propria (LP) Th2 responses against the gastrointestinal helminth *Heligmosomoides polygyrus* (Pelly *et al.*, 2016), which causes a persistent natural rodent infection in which larvae burrow into the small intestine submucosa and emerge as adult worms into the lumen (Reynolds *et al.*, 2012). While this study did not address where the interaction between ILC2s and T cells may occur – in the tissue site (LP) or in the dLN – the authors showed that ILC2s could drive the differentiation of Th2 cells without direct contact (Pelly *et al.*, 2016). In addition, IL-13-producing ILC2s can also cooperate with CD4⁺ T cells during *N. brasiliensis* infection to induce larval killing following secondary infection (Bouchery *et al.*, 2015) as the larvae transit through the lung before becoming resident in the small intestine (Camberis *et al.*, 2003), and these IL-13⁺ ILC2s are dependent on epithelial cell production of Surfactant Protein D (Thawer *et al.*, 2016). Notably, whether ILC2s can initiate priming of Th2 cells via direct contact with naïve T cells is somewhat contentious. Though a number of studies have suggested that ILC2s can directly prime Th2 responses via expression of MHC II (Mirchandani *et al.*, 2014; Oliphant *et al.*, 2014), other work suggests that these cells do not act as antigen presenting cells (Maizels and Withers, 2014).

While early studies suggested that ILC2s were a relatively homogenous population, recent work has identified that not all ILC2s are created equal, at least in the mouse. Paul and colleagues showed that there are “natural” ILC2s resident in the lung tissue that express high levels of ST2, while upon helminth-induced inflammation, there is an influx of ST2^{lo} KLRG1^{hi} inflammatory ILC2s that potently respond to IL-25 (Huang *et al.*, 2014). Exactly what the function of KLRG1 is in this context is not currently known, though it can be inhibitory in other settings. These inflammatory ILC2s appear to differentiate into natural ILC2s and are important in the expulsion of *N. brasiliensis* from the intestine (Huang *et al.*, 2014). Inflammatory ILC2s were also active in IL-17-associated infection with *Candida albicans* (Huang *et al.*, 2014), with the Notch pathway acting as a key regulator of these cells' ability to demonstrate functional plasticity (Zhang *et al.*, 2017).

In addition, another subset of ILC2s has recently been identified, a long-lived persistent population that was maintained after amelioration of Type 2 antigen exposure in mouse models (Martinez-Gonzalez *et al.*, 2016). How these “memory-like” ILC2s function in secondary worm infection has not yet been addressed, but IL-13-producing ILC2s and IL-13⁺ T cells must be recruited to the lung to activate alternatively-activated macrophages to kill larvae in *N. brasiliensis* reinfection (Bouchery *et al.*, 2015), suggesting one possible functional pathway. Notably, these long-lived memory-like ILC2s cells appear to resemble

memory CD8⁺ T cells genetically (Martinez-Gonzalez *et al.*, 2016), suggesting that they may share some characteristics with adaptive cells in this context.

Complementing these data, recent studies have shown that some subsets of ILC2s can be key mediators of tissue repair in mouse models through their production of Type 2 cytokines, including IL-9 (Turner *et al.*, 2013). In addition, ILC2s produce amphiregulin, a growth factor that can facilitate wound repair after pathogen insult (Monticelli *et al.*, 2011; 2015). However, IL-25 activation of IL-13⁺ ILC2s can also induce a pathological wound healing response (Hams *et al.*, 2014). This pathology is characterized by collagen deposition and fibrosis in the murine lung in response to eggs of the parasitic trematode, *Schistosoma mansoni*, a major human parasite that lives in the hepatic blood system, depositing eggs that induce a striking Type 2 response (Pearce and MacDonald, 2002). Thus, ILC2s can act as important regulators of inflammation and wound repair, and in some cases, their primary role may be to promote wound healing, but this response can become pathological if left unchecked.

These studies show that there is a level of sophistication to the form and function of ILC2s that is only now becoming appreciable. Recent work has also identified numerous novel molecular and genetic checkpoints that control the recruitment and diverse functions of these cells. For instance, recruitment of ILC2s into the murine lung during *N. brasiliensis* infection is dependent on the eicosanoid prostaglandin D₂ and ILC2 expression of its receptor, CRTH2, and this pathway mediated persistence of Type 2 inflammation in the lung after worm clearance (Tait Wojno *et al.*, 2015). Further, expression of the enzyme Arginase 1 is required for optimal proliferative capacity and cytokine production by these cells (Monticelli *et al.*, 2016), and metabolically, ILC2s depend on fatty acid oxidation during helminth infection (Wilhelm *et al.*, 2016). In addition, a key danger signal released by damaged epithelium, adenosine, induces IL-33 through the A2B receptor to drive ILC2 activation (Patel *et al.*, 2014). Finally, a key metabolic enzyme, AMPK α 1, must be expressed by myeloid cells in the lung to ensure IL-13⁺ ILC2s are recruited to this tissue (Nieves *et al.*, 2016). Together, these studies show that a variety of pathways regulate the myriad functions of ILC2s and key interactions between innate cell types.

Alongside these functional studies, with the development of new genomic tools, we are now able to appreciate how ILC2s are genetically adapted for their function. Recent work has demonstrated that even at the chromatin level, naïve ILC2s are programmed differently than their adaptive counterparts, naïve T cells. In the absence of Type 2 inflammation, naïve ILC2s maintain key effector genes, such as *Il13*, in an open state, allowing these genes to be readily transcribed upon pathogen insult (Shih *et al.*, 2016; Van Dyken *et al.*, 2016). In contrast, T cells only show this level of differentiation following priming towards a Th2 cell fate. However, ILC2s and Th2 cells are surprisingly similar in their genetic makeup, both at the chromatin level and in their gene expression (Shih *et al.*, 2016; Van Dyken *et al.*, 2016). Thus, it seems fair to assume that ILC2s act as an innate counterpart to their adaptive cousins, perhaps responding readily and robustly in the tissue site upon pathogen challenge, prior to the development of the adaptive response. Notably, these studies were performed in mouse models of disease, so it remains to be determined whether this finding holds true for human ILC2s.

Most of the studies discussed here have highlighted the importance of ILC2s to the development of Type 2 inflammation in mouse models (Moro *et al.*, 2010; Neill *et al.*, 2010; Price *et al.*, 2010), but some studies have been performed evaluating these cells' function in humans. Surprisingly, recent work has suggested that ILCs are not required for the proper functioning of the immune system in humans (Vély *et al.*, 2016). This work demonstrated that humans suffering from severe combined immunodeficiency that underwent a hematopoietic stem cell transplant did not efficiently reconstitute ILC populations, yet did not suffer any ill effects from the absence of ILC2s, as long as their T and B cell compartment was intact (Vély *et al.*, 2016). However, there are clear constraints to this study, specifically given that it did not investigate the incidence of Type 2 inflammation in these patients, and given that this work was done in the Western world, where there is little incidence of Type 2-inducing parasites. In studies performed in regions where humans are commonly affected by helminth parasites, children endemically infected with the trematode fluke *S. haematobium* had reduced ILC2 numbers in the peripheral blood, perhaps suggesting that peripheral ILC2s have been recruited into the tissue, but more work is required to verify this (Nausch *et al.*, 2015). In contrast, in human filarial infection, IL-13-producing ILCs were expanded in the blood of filarial-infected individuals compared to uninfected subjects, and these ILCs were found to express MHC II and a range of cytokine receptors, rendering them responsive to antigen and damage signals alike (Boyd *et al.*, 2014). Ultimately, while mouse models of helminth infection have advantages, allowing for the manipulation and fine dissection of pathways and cell types in various tissues, these models do not accurately represent the complexities of human infection, which features varied and repeated dosing and high incidence of coinfection. Thus, further studies in healthy humans and in patients that selectively lack ILCs in natural Type 2 immune settings will be required to clarify the importance of ILC2s to human immunity, and to verify findings from the mouse.

BASOPHILS – Rare but essential for parasite clearance

Basophils are perhaps the most rare of the innate immune cells that we will discuss here, making up less than 1% of circulating leukocytes. The rarity of these cells can make them difficult to study, particularly in tissues where purifying cells is difficult, such as the gastrointestinal tract. However, the development of mouse models such as basophil-deficient mice in which the mast cell protease 8 promoter drives expression of the high-affinity diphtheria toxin receptor, allowing for toxin-mediated basophil deletion, and a basophil-specific cre-lox system have made these studies easier to perform (Voehringer, 2013). Basophils are known perhaps most significantly for their capacity to respond to IgE cross-linking due to their expression of the high-affinity IgE receptor, FcεR1 (Mitre *et al.*, 2004; Schramm *et al.*, 2006). However, basophils also respond to cytokines, such as IL-3, IL-18, IL-33 and TSLP (Yoshimoto *et al.*, 1999; Shen *et al.*, 2008; Pecaric-Petkovic *et al.*, 2009; Kroeger *et al.*, 2009; Siracusa *et al.*, 2011; Schwartz *et al.*, 2014c), as well as helminth-derived products (Phillips *et al.*, 2003; Tawill *et al.*, 2004; Reese *et al.*, 2007).

Basophils are potent producers of IL-4 following stimulation (Min *et al.*, 2004; Gessner *et al.*, 2005; van Panhuys *et al.*, 2011), a response that has been shown to be dependent on the TF GATA-1 (Nei *et al.*, 2013). They also degranulate in response to activation, releasing

large quantities of pre-stored mediators, including histamine, chemokines and IL-6. Perhaps surprisingly, mouse basophils release much more IL-6 than IL-4 during Type 2 immune inflammation, but the role of this cytokine in this setting is largely under-appreciated. Previous studies have tested whether basophil-derived IL-6 may drive worm-specific Th17 responses, but it does not seem to play a role in this facet of immune activation (Sharma *et al.*, 2007; 2015). Finally, studies in mice show that one of the key mechanisms of action of basophils is their capacity to recruit other cell types to the site of inflammation, interacting with and attracting innate immune effectors (Perrigoue *et al.*, 2009). This capacity has primarily been demonstrated in allergic models, where basophils recruit ILC2s, alternatively-activated macrophages and eosinophils through production of chemokines and cytokines (Egawa *et al.*, 2013; Noti *et al.*, 2013; Kim *et al.*, 2014; Schwartz *et al.*, 2014c; Cheng *et al.*, 2015). It seems likely that this process is also active in helminth infection, though this has yet to be shown.

In addition, some work suggests that basophils are players in the mobilization of adaptive Th2 responses. Basophils may act as the essential IL-4 producers that instigate Type 2 inflammation by activating naïve T cells towards a Th2 cell fate (Perrigoue *et al.*, 2009). However, basophils are found at very low levels in the dLN, though at greater numbers in the spleen, and thus it is unclear whether they can truly act as the essential source of IL-4 for Th2 priming in the dLN. Indeed, they are dispensable for the onset of primary immunity against some helminth species, including *H. polygyrus* and *N. brasiliensis* (Ohnmacht *et al.*, 2010; Smith *et al.*, 2012), suggesting that their role in Th2 priming in primary infection is limited in some cases. Previous studies have also suggested that basophils may act directly as an antigen-presenting cell (APC) population in mice, as they express some level of MHC II and can prime IL-4 production from T cells and Ag-specific T cell proliferation *in vitro* (Perrigoue *et al.*, 2009), as well as have some brief interaction with T cells in the dLN (Sokol *et al.*, 2009). However, basophils lack the specific machinery to take up and process antigen (Eckl-Dorna *et al.*, 2012), and they are not required for Th2 priming against egg products of the helminth parasite *S. mansoni* and during infection with *H. polygyrus* and *N. brasiliensis* (Phythian-Adams *et al.*, 2010; Smith *et al.*, 2012). In fact, two photon microscopy studies have demonstrated that T cell-basophil interactions in the dLN are rare. However, basophils do participate in prolonged interactions with activated T cells in the tissue (Sullivan *et al.*, 2011), suggesting that they interact primarily with activated effector T cells, presenting antigen to this population in order to maintain T cell cytokine production.

Despite the ambiguous nature of basophil-T cell interactions, basophils expand and can act as an important early source of IL-4 in some infection settings, including against the gastrointestinal helminthes *N. brasiliensis*, *Trichuris muris*, *Trichinella spiralis*, and *H. polygyrus* in murine models (Siracusa *et al.*, 2011; Giacomini *et al.*, 2012); (Herbst *et al.*, 2012). During *N. brasiliensis* infection, a non-T cell, non-eosinophil source of IL-4 was found to be important at tissue sites (Voehringer *et al.*, 2006). During infection with *T. muris*, a murine whipworm that infects mice orally and resides in the epithelium of the cecum, basophils expand significantly from naïve levels, which promotes the effective clearance of the parasite (Siracusa *et al.*, 2011). In this setting, expansion of basophils is dependent on the cytokine TSLP (Siracusa *et al.*, 2011), which also drives extramedullary hematopoiesis in the spleen and has both basophils and MCs as its product (Siracusa *et al.*,

2013a). Further, basophil development in response to murine *T. spiralis* infection, an infection in which adult worms in the small intestine produce live larvae that burrow through the intestine and encyst in muscle, depends on TSLP, and these TSLP-induced basophils were required for optimal Type 2 cytokine responses during infection (Giacomin *et al.*, 2012). Early during *H. polygyrus* infection, IL-3 production by effector T cells or direct antibody binding induced basophil expansion in the bone marrow and spleen (Herbst *et al.*, 2012), though basophils were not required for primary clearance of this parasite (Smith *et al.*, 2012) or for Type 2 inflammation and infection outcome in *S. mansoni* infected mice (Schwartz *et al.*, 2014a). During *S. mansoni* infection, basophils were also dispensable for granuloma formation and protection from fatal disease (Schwartz *et al.*, 2014b), suggesting that basophils are not required for wound healing responses in this setting. Further studies will be required to determine how early basophil expansion and cytokine production during primary infection affects the course of disease and wound healing responses in multiple models of murine helminth infection.

In addition to their roles during early primary infection, basophils are also an essential component of the cellular milieu that facilitates clearance of the parasite during secondary infection in mice, responsible for IgE-elicited IL-4 production (Schwartz *et al.*, 2014c). They are also required for protection in secondary infection with *N. brasiliensis* (Ohnmacht and Voehringer, 2010; Ohnmacht *et al.*, 2010), where they are thought to co-operate with eosinophils to trap larvae in the skin (Obata-Ninomiya *et al.*, 2013). Any parasites that make it past the onslaught of alternatively-activated macrophages in the lung will be confronted by a further increased number of basophils compared to primary infection, where again IgE-mediated activation results in IL-4 and IL-13 production that confers expulsion of worms (Schwartz *et al.*, 2014c). However, basophils are not required for vaccine-induced clearance of *H. polygyrus* following injection of excretory/secretory products (Hewitson *et al.*, 2015), suggesting that while some helminth-associated products can directly activate basophils, this is not the case in all settings.

As concerns human helminth infection, basophils act as an important source of IL-4 in filarial infection (Min *et al.*, 2004). In an experimental infection of human subjects with a gastrointestinal parasite, basophilia was a reliable biomarker of early infection prior to the upregulation of IgE (Falcone *et al.*, 2009). In contrast, basophil responsiveness was abrogated in chronic human hookworm infection, suggesting that the parasite may modulate basophil responses to maintain chronicity (Larson *et al.*, 2012a; Pinot de Moira *et al.*, 2014). Work in mice suggests that downregulation of basophil responsiveness to IgE stimulation may be due to chronic production of IL-10, which could be induced by secretory/excretory products from *Litomosoides sigmodontis*, a rodent parasite used as a model of lymphatic filariasis (Larson *et al.*, 2012b). However, this murine model does not recapitulate all aspects of human filarial infection and it remains unclear whether this mechanism is at play in humans during infection with filarial parasites or hookworms. Thus, while basophils are key mediators of Type 2 inflammation during helminth infection, acting as part of a complex network of innate immune cells within the tissue site, further studies in mouse models and patients will be needed to tease out the role of these cells in worm expulsion in primary and secondary infection with various helminths in humans.

MAST CELLS – Long-lived producers of inflammatory mediators

Like other rare innate cells, MCs are found at barrier sites such as the skin and mucosal tissues and will accumulate at inflamed sites (Hepworth *et al.*, 2012b). Unlike basophils, which have a lifespan of just a matter of hours, MCs are long-lived, surviving for weeks or months, at least in mice, producing cytokines, chemokines, and releasing pre-stored mediators (Voehringer, 2013). While the developmental pathways that govern MC differentiation and accumulation in tissues during inflammation are not fully described, recent work from Siracusa and colleagues show that MC development depends on the expression of carbonic anhydrase enzymes in MC progenitors (Henry *et al.*, 2016). Expression of these enzymes was required for infection-induced mastocytosis during *T. spiralis* infection, and absence of these enzymes led to a reduction in MC-induced inflammation in the host during murine infection. The cytokines TSLP and IL-33 are also associated with MC population expansion (Hepworth *et al.*, 2012a), with TSLP inducing MC hematopoiesis in the spleen (Siracusa *et al.*, 2013b). The Notch pathway has also been implicated in MC development and distribution during infection with the gastrointestinal murine parasite *Strongyloides venezuelensis* (Sakata-Yanagimoto *et al.*, 2008; 2011). MC accumulation in the skin during *L. sigmodontis* infection was controlled by CCL17 production by DCs, preventing larval entry by reducing MC-dependent vascular permeability (Specht *et al.*, 2011).

A number of studies have suggested that MCs are necessary for helminth expulsion in mice, employing *Kit*-deficient mouse models that lack MCs. While these studies should be viewed with caution as these mice have other defects, including defective intestinal peristalsis (Voehringer, 2013), this work suggests that immune clearance of worms during *N. brasiliensis*, *T. muris* and *T. spiralis* infection are all delayed in the absence of MCs (Voehringer, 2013). During *S. mansoni* infection, mice deficient in IgE responses have smaller granulomas and have a higher worm burden, demonstrating that IgE-mediated activation of MCs and basophils contributes to host immunity and potentially wound healing mechanisms during infection with this parasite (King *et al.*, 1997). These antibody-mediated responses have also been shown to be protective in humans suffering from *S. haematobium* infection and with the filarial parasite *Brugia malayi* (Lobos *et al.*, 2003; Mitchell *et al.*, 2012). The mechanism of action in this context is unclear, although it is likely that IgE induced immune attack of the larvae by MCs and other Fc receptor-bearing cells, inhibiting establishment of infection.

Whether MCs directly contribute to worm expulsion or not, during Type 2 inflammation they rapidly release large quantities of a multitude of inflammatory mediators, including histamine, proteases and cytokines. For instance, during murine infection with *T. spiralis*, MCs contribute to an increase in intestinal permeability via their production of proteases, particularly MCP1 (Lawrence *et al.*, 2004), which can in this case contribute to enhanced expulsion of the parasite (McDermott *et al.*, 2003). In the later stages of worm infection, MCs respond to helminth-specific IgE crosslinking of the surface receptor FcεR1 by degranulating and releasing large quantities of inflammatory mediators, including IL-4 and IL-6 (Gessner *et al.*, 2005; Liu *et al.*, 2013). Another MC-derived mediator is prostaglandin D₂ (Nakamura *et al.*, 2015), an eicosanoid that recruits ILC2s to the site of Type 2

inflammation via their expression of the CCR2 receptor (Xue *et al.*, 2014; Tait Wojno *et al.*, 2015). Thus, it is likely that MCs are an important source of this molecule to mediate further innate cell recruitment during helminth infection. Finally, MC cytokine production is critical in the Type 2 inflammatory response in tissues, as mice deficient in MC IL-4 or TNF α responses displayed delayed clearance of *T. spiralis*, suggesting that MC production of these cytokines is required for anti-helminthic responses (Ierna *et al.*, 2008).

An important role for MCs is to recruit and support the activation of other innate cell types. In addition to their ability to produce prostaglandins that recruit innate immune cells (Nakamura *et al.*, 2015), MCs produce TSLP, a mediator of innate and adaptive cell recruitment, at least at the mRNA level (Dewas *et al.*, 2015). MCs are also active at early stages of helminth infection in mice, inducing production of protective tissue factors such as IL-25, IL-33 and TSLP from the epithelium that activate and recruit other innate immune cell types in response to *H. polygyrus* and *T. muris* (Hepworth *et al.*, 2012a). MC-derived proteases may elicit production of these tissue factors by activating protease-activated receptor 2, which is required for optimal Type 2 immunity against *T. spiralis* infection (Park *et al.*, 2011). Further, proteases secreted by MCs are actively involved in the cleavage of IL-33, enhancing its potency and leading to the activation of ILC2s (Lefrançois *et al.*, 2014). It is also thought that MC degranulation and histamine release may have adjuvant activity during allergic inflammation, enhancing adaptive Th2 responses by inducing emigration of DCs to the dLN (Dudeck *et al.*, 2011), but it is unknown if this mechanism is at play during helminth infection. Myeloid-derived suppressor cells induce optimal secretion of cytokines from MC and are thought to help mediate clearance of *N. brasiliensis* and *T. spiralis* in this regard (Morales *et al.*, 2014), highlighting that MCs may interact with APCs and myeloid cells in numerous contexts. Finally, ILC2s in the skin have also been shown to recruit and interact with activated MCs in contact dermatitis models in mice, though whether this interaction is active in helminth infection has not yet been addressed (Roediger *et al.*, 2013). These results underscore the importance of MCs' ability to communicate with a wide variety of cell types in the tissue.

Together, these data show that MCs are key mediators of immune activation at the early stages of infection and also function at later stages, mediating worm clearance and participating in protective immune responses in secondary infection. Conversely, one recent study has suggested that MCs may play a role in immune hyporesponsiveness in the tissue site following repeated infection with *S. mansoni* in a rodent model of reinfection, as deletion of MCs was associated with a reduction in IL-10 and an increase in MHC II⁺ infiltrating cells (Prendergast *et al.*, 2016). Further studies will be needed to dissect the molecular mechanisms by which MCs promote and limit Type 2 inflammation and whether these cells mediate wound healing mechanisms, both in mouse models and particularly in helminth-infected humans. However, MCs seemingly have a multifaceted role to play in helminth infection, with a wide array of functions that differ based on the tissue location and stimulation.

DENDRITIC CELLS – An essential bridge between innate and adaptive immunity

DCs come in many flavors, but their essential role in priming adaptive Type 2 immune activation is a feature in a variety of infection settings (Phythian-Adams *et al.*, 2010; Smith *et al.*, 2012; Connor *et al.*, 2014). DC subsets can be defined, broadly, by the TFs they express. Conventional DCs (cDCs) are dependent on the TF Zbtb46 (Meredith *et al.*, 2012; Satpathy *et al.*, 2012a), and cDC subsets can be further subdivided by the TFs they depend on (Satpathy *et al.*, 2012b), while plasmacytoid DCs (pDCs) depend on E2-2 (Cisse *et al.*, 2008). Recently it has become clear that these various DC subsets have different roles in Type 2 immunity against helminths. Specifically, IRF4-dependent CD11b⁺ DCs (cDC2s) are required for Th2 immune activation (Tussiwand *et al.*, 2015), while IRF8-dependent CD103⁺/CD8⁺ DCs (cDC1s) instigate Th1 immune activation during worm infection (Everts *et al.*, 2016). While the role of pDCs in Type 2 immunity has been contentious, with some suggestion that they modulate Type 2 activation (de Heer *et al.*, 2004), it now seems that they are not essential for Th2 responses in the tissue site (Lundie *et al.*, 2016). However, they can elicit Type 2 cytokine production from activated T cells (Lundie *et al.*, 2016) and may have a role to play in promoting Th2 responses the dLN (unpublished findings).

Very recent studies have revealed that DCs can perform other functions that support their direct induction of Th2 responses during helminth infection in mice, dependent upon the subset and the immune microenvironment. In this context, DCs adopt a Type 1 interferon (IFN-I) immune signature in response to stimulation with Type 2-associated antigen from *S. mansoni* and *N. brasiliensis* in murine models (Connor *et al.*, 2016). This immune signature was previously described in DCs following stimulation with *S. mansoni* egg products (Trottein *et al.*, 2004), but it has now been shown to emerge in response to cytokine release from activated DCs (unpublished findings). Notably, this IFN-I activation and responsiveness was required for DCs to prime Th2 immune activation effectively (Connor *et al.*, 2016). Thus, for the first time, a specific cytokine signature has been identified that typifies DC phenotype and function following exposure to Type 2 inflammation-associated antigens.

This is significant, as a specific suite of surface molecules expressed by murine DCs that initiate a Th2 program from naïve T cells has not been identified to date. Th2-inducing DCs express CD301b and PDL2 (Gao *et al.*, 2013; Kumamoto *et al.*, 2013) and the Notch ligand Jagged 1 (Okamoto *et al.*, 2009). However, these markers are not unique to DCs that will induce polarization of Th2 and only Th2 cells. Thus, researchers have looked further afield to characterize Type 2-specific signals from DCs. Specifically, gene expression changes specific for Th2-inducing DCs have been identified (Connor *et al.*, 2016). Further, epigenetic control of a Th2-inducing profile has been described, with methyl-CpG-binding protein 2 (Mbd2) isolated as an epigenetic regulator of a range of genes involved in Th2 immune activation by DCs, including *Irf4*, *Cd40*, *Cd80*, *Cd86* and *Retnla* (Cook *et al.*, 2015), all molecules associated with Type 2 activation by DCs (MacDonald and Maizels, 2008; Cook *et al.*, 2012; Bouchery *et al.*, 2014). In addition, DC responsiveness to Type 2-associated antigen may depend on the TF STAT5, at least in TSLP-induced responses (Bell *et al.*, 2013). Finally, specific patterns of chemokine responsiveness may be a hallmark of Th2-inducing DCs. For instance, expression of CCR7 was critical for the migration of DCs in a

Type 2 environment (unpublished findings), and Th2-inducing DCs are directed by their expression of CXCR5 to locate at the fringe of B cell follicles (León *et al.*, 2012), a site which may favor the development of T follicular helper cells, key adaptive producers of IL-4 in the dLN (King and Mohrs, 2009; Fairfax *et al.*, 2015). However, the search continues for a set of cell surface markers that typify all Th2-inducing DCs and could be used to identify and characterize these cells *ex vivo* in mice and humans.

Regardless, significant evidence exists that DC function in guiding Type 2 inflammatory responses is directly dependent on environmental signals that occur in the context of helminth infection. DCs, similar to other innate effectors, respond to innate cytokines produced by damaged epithelium, including TSLP (Ito *et al.*, 2005; Wang *et al.*, 2006). In the absence of TSLP, Th2 immune activation is reduced in *T. muris* infection, with a consequent increase in IL-12, IFN γ and IL-17 (Taylor *et al.*, 2009), suggesting that TSLP is an important polarizing signal for Th2 induction in mice. The epithelium- and macrophage-derived factor IL-33 may also shape DC responses, as IL-33 can activate DCs (Besnard *et al.*, 2011; de Kleer *et al.*, 2016) and can induce polarization of Th2 immunity from naïve T cells (Besnard *et al.*, 2011), though whether IL-33 acts on DCs directly in helminth infection has not been investigated. Further, it has been suggested that DCs themselves must produce IL-33 to polarize Th2 immune activation (Williams *et al.*, 2013), and inflammatory DCs and respiratory DCs may upregulate IL-33 in certain circumstances (Wills-Karp *et al.*, 2012; Tjota *et al.*, 2013), though IL-33 is not essential for Th2 induction (unpublished findings).

In addition to their Th2 priming capabilities, DCs also influence Type 2 inflammation by interacting with adaptive and innate cells in other ways. DCs are important recruiters of Th2 cells, via their production of CCL17, dependent on ILC2-derived IL-13 (Halim *et al.*, 2015). Further, migration of DCs to the dLN during Type 2 inflammation may well also depend on eosinophil activity (Chu *et al.*, 2014), another example of an innate-innate cell interaction that is key to Type 2 immunity. As previously discussed, it is possible that basophils act as an early source of IL-4 for polarizing Th2 cells in the presence of activating signals from DCs, while it is also possible that ILC2s are an important early source of IL-4 in the dLN (Pelly *et al.*, 2016). IL-4 has also been shown to act directly on DCs to induce Relm α expression, which was required for the induction of optimal IL-10 and IL-13 production from T cells (Cook *et al.*, 2012). Importantly, these data suggest that DCs can also play a critical role in wound healing responses during helminth infection, since Relm α and other factors, including Relm β and Arginase1, are actively involved in wound healing mechanisms and can also limit Type 2 inflammation, perhaps through the induction of T cell IL-10 (Pesce *et al.*, 2009a; b; Nair *et al.*, 2009; Cook *et al.*, 2012; Esser-von Bieren *et al.*, 2013; Knipper *et al.*, 2015).

Taken together, these studies demonstrate that DCs exist within a complex immune microenvironment that includes many players, both innate and adaptive. Clearly, more work is required to address how DCs from humans reflect the complex interplay that has been identified in mice (de Jong *et al.*, 2002). One study suggests that a mechanism of immune hyporesponsiveness during chronic helminth infection may reflect the functional impairment of DCs in humans chronically infected with *S. haematobium* (Everts *et al.*, 2010), but human DC subsets that act in infected tissues and prime human Th2 responses remain poorly

understood. Thus, while DCs act as an essential bridge between innate and adaptive immune states, controlling the greater immune state of the host, future work will be needed to fully dissect the form and function of these cells during helminth infection in both murine models and in humans.

Conclusions

Innate immune activation – A collaborative effort

One over-arching message from our discussion of rare innate immune cell types during helminth infection is that Type 2 immune activation is a collaborative effort that requires a host of cells (Fig. 1). For example, here we have described basophils recruiting ILC2s and eosinophils, while ILC2s are responsible for effective recruitment of DCs, and MCs produce mediators that recruit ILC2s. Clearly, rare innate cells, and more numerous alternatively-activated macrophages and eosinophils, act in a complex interconnected network that mediates immune activation against helminth parasites. Importantly, these cells are uniquely adapted to their function, employing a variety of cellular and molecular mechanisms to perform the right task at the right time and place. While we do not yet fully understand these mechanisms, ongoing and future efforts will unravel novel pathways that control these highly regulated and powerful cell types.

It is worth emphasizing that different cell types are required for protective immunity and wound repair mechanisms against different species of helminths – there is not one unified response that can be described as protective for all helminth infections. Likely, the route of infection, the tissues affected, the parasite lifecycle, and the parasite's arsenal of excretory/secretory products dictate which cell types are active against a particular species of helminth. Many helminth pathogens are highly immunomodulatory and may act to inhibit or exacerbate certain responses, which could also shape the host immune response and influence the function of rare innate immune cell types. Thus, innate cells and helminths participate in a complex interplay that is unique to every parasite. In light of this, conclusions from studies in murine models of infection must be evaluated in studies of human helminth infection to reveal the nature of the interplay between rare innate immune cells and helminth parasites in humans.

Importantly, the studies discussed here and such future work will also shed light on potential novel pathways and cell types that could be targeted to enhance anti-helminth immune responses. Given that anti-helminth therapies are rarely protective and that resistance to current chemotherapeutics is now becoming a reality, it is important to focus efforts towards the development of therapies that can induce protective Type 2 immunity and effective wound repair pathways. Further, the increasing scourge of allergic Type 2 immune activation in the Western world means that mechanisms to inhibit Type 2 immunity are also required. With a greater understanding of the role of rare innate immune cells we can move one step closer to tailored therapies that could alleviate the suffering of millions of people around the world. However, we must take into account the fact that these cells do not function in isolation, rather acting in a complex, highly adapted network that orchestrates Type 2 immunity.

Acknowledgments

We thank members of the Tait Wojno laboratory for assistance in preparing the manuscript. Research in the Tait Wojno laboratory is supported by Cornell University, the Cornell University College of Veterinary Medicine Research in Animal Health Award (to E.D.T.W.), and the National Institutes of Health/National Institute of Allergy and Infectious Diseases (K22 AI116729 to E.D.T.W.).

References

- Allen JE, Maizels RM. Diversity and dialogue in immunity to helminths. *Nature Reviews Immunology*. 2011; 11:375–388. DOI: 10.1038/nri2992
- Bell BD, Kitajima M, Larson RP, Stoklasek TA, Dang K, Sakamoto K, Wagner KU, Reizis B, Hennighausen L, Ziegler SF. The transcription factor STAT5 is critical in dendritic cells for the development of TH2 but not TH1 responses. *Nature Immunology*. 2013; 14:364–371. DOI: 10.1038/ni.2541 [PubMed: 23435120]
- Besnard AG, Togbe D, Guillou N, Erard F, Quesniaux V, Ryffel B. IL-33-activated dendritic cells are critical for allergic airway inflammation. *European journal of immunology*. 2011; 41:1675–1686. DOI: 10.1002/eji.201041033 [PubMed: 21469105]
- Bouchery T, Kyle R, Camberis M, Shepherd A, Filbey K, Smith A, Harvie M, Painter G, Johnston K, Ferguson P, Jain R, Roediger B, Delahunt B, Weninger W, Forbes-Blom E, Le Gros G. ILC2s and T cells cooperate to ensure maintenance of M2 macrophages for lung immunity against hookworms. *Nature communications*. 2015; 6:6970.doi: 10.1038/ncomms7970
- Bouchery T, Kyle R, Ronchese F, Le Gros G. The Differentiation of CD4(+) T-Helper Cell Subsets in the Context of Helminth Parasite Infection. *Frontiers in immunology*. 2014; 5:487.doi: 10.3389/fimmu.2014.00487 [PubMed: 25360134]
- Boyd A, Ribeiro JMC, Nutman TB. Human CD117 (cKit)+ Innate Lymphoid Cells Have a Discrete Transcriptional Profile at Homeostasis and Are Expanded during Filarial Infection. *PLoS ONE*. 2014; 9:e108649–11. DOI: 10.1371/journal.pone.0108649 [PubMed: 25255226]
- Camberis, M., Le Gros, G., Urban, J. Animal model of *Nippostrongylus brasiliensis* and *Heligmosomoides polygyrus*. In: Coligan, John E., et al., editors. *Current protocols in immunology*. Vol. Chapter 19. 2003. p. 12.19-12.27.
- Cheng LE, Sullivan BM, Retana LE, Allen CDC, Liang HE, Locksley RM. IgE-activated basophils regulate eosinophil tissue entry by modulating endothelial function. *The Journal of experimental medicine*. 2015; 212:513–524. DOI: 10.1084/jem.20141671 [PubMed: 25779634]
- Cherrier M, Sawa S, Eberl G. Notch, Id2, and ROR γ t sequentially orchestrate the fetal development of lymphoid tissue inducer cells. *Journal of Experimental Medicine*. 2012; 209:729–740. DOI: 10.1084/jem.20111594 [PubMed: 22430492]
- Chu DK, Jimenez-Saiz R, Verschoor CP, Walker TD, Goncharova S, Llop-Guevara A, Shen P, Gordon ME, Barra NG, Bassett JD, Kong J, Fattouh R, McCoy KD, Bowdish DM, Erjefalt JS, Pabst O, Humbles AA, Kolbeck R, Wasserman S, Jordana M. Indigenous enteric eosinophils control DCs to initiate a primary Th2 immune response in vivo. *Journal of Experimental Medicine*. 2014; 211:1657–1672. DOI: 10.1084/jem.20020656 [PubMed: 25071163]
- Cisse B, Caton ML, Lehner M, Maeda T, Scheu S, Locksley R, Holmberg D, Zweier C, Hollander den NS, Kant SG, Holter W, Rauch A, Zhuang Y, Reizis B. Transcription Factor E2-2 Is an Essential and Specific Regulator of Plasmacytoid Dendritic Cell Development. *Cell*. 2008; 135:37–48. DOI: 10.1016/j.cell.2008.09.016 [PubMed: 18854153]
- Cliffe LJ, Humphreys NE, Lane TE, Potten CS, Booth C, Grecis RK. Accelerated intestinal epithelial cell turnover: a new mechanism of parasite expulsion. *Science*. 2005; 308:1463–1465. DOI: 10.1126/science.1108661 [PubMed: 15933199]
- Connor LM, Tang SC, Camberis M, Le Gros G, Ronchese F. Helminth-Conditioned Dendritic Cells Prime CD4+ T Cells to IL-4 Production In Vivo. *The Journal of Immunology*. 2014; 193:2709–2717. DOI: 10.4049/jimmunol.1400374 [PubMed: 25108019]
- Connor LM, Tang SC, Cognard E, Ochiai S, Hilligan KL, Old SI, Pellefigues C, White RF, Patel D, Smith AAT, Eccles DA, Lamiable O, McConnell MJ, Ronchese F. Th2 responses are primed by

- skin dendritic cells with distinct transcriptional profiles. *Journal of Experimental Medicine*. 2016; jem.20160470-18. doi: 10.1084/jem.20160470
- Cook PC, Jones LH, Jenkins SJ, Wynn TA, Allen JE, MacDonald AS. Alternatively activated dendritic cells regulate CD4⁺ T-cell polarization in vitro and in vivo. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; doi: 10.1073/pnas.1121231109
- Cook PC, Owen H, Deaton AM, Borger JG, Brown SL, Clouaire T, Jones GR, Jones LH, Lundie RJ, Marley AK, Morrison VL, Phythian-Adams AT, Wachter E, Webb LM, Sutherland TE, Thomas GD, Grainger JR, Selfridge J, McKenzie ANJ, Allen JE, Fagerholm SC, Maizels RM, Ivens AC, Bird A, MacDonald AS. A dominant role for the methyl-CpG-binding protein Mbd2 in controlling Th2 induction by dendritic cells. *Nature communications*. 2015; 6:6920.doi: 10.1038/ncomms7920
- de Heer HJ, Hammad H, Soullié T, Hijdra D, Vos N, Willart MAM, Hoogsteden HC, Lambrecht BN. Essential role of lung plasmacytoid dendritic cells in preventing asthmatic reactions to harmless inhaled antigen. *The Journal of experimental medicine*. 2004; 200:89–98. DOI: 10.1084/jem.20040035 [PubMed: 15238608]
- de Jong EC, Vieira PL, Kalinski P, Schuitemaker JHN, Tanaka Y, Wierenga EA, Yazdanbakhsh M, Kapsenberg ML. Microbial compounds selectively induce Th1 cell-promoting or Th2 cell-promoting dendritic cells in vitro with diverse th cell-polarizing signals. *Journal of immunology* (Baltimore, Md : 1950). 2002; 168:1704–1709.
- de Kleer IM, Kool M, de Bruijn MJW, Willart M, van Moorleghem J, Schuijs MJ, Plantinga M, Beyaert R, Hams E, Fallon PG, Hammad H, Hendriks RW, Lambrecht BN. Perinatal Activation of the Interleukin-33 Pathway Promotes Type 2 Immunity in the Developing Lung. *Immunity*. 2016; 45:1285–1298. DOI: 10.1016/j.immuni.2016.10.031 [PubMed: 27939673]
- Dewas C, Chen X, Honda T, Junttila I, Linton J, Udey MC, Porcella SF, Sturdevant DE, Feigenbaum L, Koo L, Williams J, Paul WE. TSLP expression: analysis with a ZsGreen TSLP reporter mouse. *The Journal of Immunology*. 2015; 194:1372–1380. DOI: 10.4049/jimmunol.1400519 [PubMed: 25539812]
- Dudeck A, Dudeck J, Scholten J, Petzold A, Surianarayanan S, Köhler A, Peschke K, Vöhringer D, Waskow C, Krieg T, Müller W, Waisman A, Hartmann K, Gunzer M, Roers A. Mast Cells Are Key Promoters of Contact Allergy that Mediate the Adjuvant Effects of Haptens. *Immunity*. 2011; 34:973–984. DOI: 10.1016/j.immuni.2011.03.028 [PubMed: 21703544]
- Eckl-Dorna J, Ellinger A, Blatt K, Ghanim V, Steiner I, Pavelka M, Valent P, Valenta R, Niederberger V. Basophils are not the key antigen-presenting cells in allergic patients. *Allergy*. 2012; 67:601–608. DOI: 10.1111/j.1398-9995.2012.02792.x [PubMed: 22335568]
- Egawa M, Mukai K, Yoshikawa S, Iki M, Mukaida N, Kawano Y, Minegishi Y, Karasuyama H. Inflammatory Monocytes Recruited to Allergic Skin Acquire an Anti-inflammatory M2 Phenotype via Basophil-Derived Interleukin-4. *Immunity*. 2013; 38:570–580. DOI: 10.1016/j.immuni.2012.11.014 [PubMed: 23434060]
- Esser-von Bieren J, Mosconi I, Guet R, Piersgilli A, Volpe B, Chen F, Gause WC, Seitz A, Verbeek JS, Harris NL. Antibodies Trap Tissue Migrating Helminth Larvae and Prevent Tissue Damage by Driving IL-4R α -Independent Alternative Differentiation of Macrophages. *PLoS pathogens*. 2013; 9:e1003771–15. DOI: 10.1371/journal.ppat.1003771 [PubMed: 24244174]
- Everts B, Adegnika AA, Kruize YCM, Smits HH, Kremsner PG, Yazdanbakhsh M. Functional impairment of human myeloid dendritic cells during *Schistosoma haematobium* infection. *PLoS neglected tropical diseases*. 2010; 4:e667.doi: 10.1371/journal.pntd.0000667 [PubMed: 20422029]
- Everts B, Tussiwand R, Dreesen L, Fairfax KC, Huang SCC, Smith AM, O'Neill CM, Lam WY, Edelson BT, Urban JF, Murphy KM, Pearce EJ. Migratory CD103⁺ dendritic cells suppress helminth-driven type 2 immunity through constitutive expression of IL-12. *Journal of Experimental Medicine*. 2016; 213:35–51. DOI: 10.1084/jem.20150235 [PubMed: 26712805]
- Fairfax KC, Everts B, Amiel E, Smith AM, Schramm G, Haas H, Randolph GJ, Taylor JJ, Pearce EJ. IL-4-Secreting Secondary T Follicular Helper (T_{fh}) Cells Arise from Memory T Cells, Not Persisting T_{fh} Cells, through a B Cell-Dependent Mechanism. *The Journal of Immunology*. 2015; doi: 10.4049/jimmunol.1401225
- Falcone FH, Telford G, Hooi D, Brown AP, Seabra R, Feary J, Venn A, Britton J, Pritchard DI. Antigen-driven basophil activation is indicative of early *Necator americanus* infection in IgE-

seronegative patients. *The Journal of Allergy and Clinical Immunology*. 2009; 124:1343–50.e7. DOI: 10.1016/j.jaci.2009.07.039 [PubMed: 19800679]

- Fallon PG, Ballantyne SJ, Mangan NE, Barlow JL, Dasvarma A, Hewett DR, McIlgorm A, Jolin HE, McKenzie ANJ. Identification of an interleukin (IL)-25-dependent cell population that provides IL-4, IL-5, and IL-13 at the onset of helminth expulsion. *The Journal of experimental medicine*. 2006; 203:1105–1116. DOI: 10.1084/jem.20051615 [PubMed: 16606668]
- Gao Y, Nish SA, Jiang R, Hou L, Licona-Limón P, Weinstein JS, Zhao H, Medzhitov R. Control of T Helper 2 Responses by Transcription Factor IRF4-Dependent Dendritic Cells. *Immunity*. 2013; 39:722–732. DOI: 10.1016/j.immuni.2013.08.028 [PubMed: 24076050]
- Gause WC, Wynn TA, Allen JE. Type 2 immunity and wound healing: evolutionary refinement of adaptive immunity by helminths. *Nature Reviews Immunology*. 2013; doi: 10.1038/nri3476
- Gerbe F, Sidot E, Smyth DJ, Ohmoto M, Matsumoto I, Dardalhon V, Cesses P, Garnier L, Pouzolles M, Brulin B, Bruschi M, Hargus Y, Zimmermann VS, Taylor N, Maizels RM, Jay P. Intestinal epithelial tuft cells initiate type 2 mucosal immunity to helminth parasites. *Nature*. 2016; 529:226–230. DOI: 10.1038/nature16527 [PubMed: 26762460]
- Gessner A, Mohrs K, Mohrs M. Mast cells, basophils, and eosinophils acquire constitutive IL-4 and IL-13 transcripts during lineage differentiation that are sufficient for rapid cytokine production. *Journal of immunology* (Baltimore, Md : 1950). 2005; 174:1063–1072.
- Giacomin PR, Siracusa MC, Walsh KP, Grencis RK, Kubo M, Comeau MR, Artis D. Thymic stromal lymphopoietin-dependent basophils promote Th2 cytokine responses following intestinal helminth infection. *The Journal of Immunology*. 2012; 189:4371–4378. DOI: 10.4049/jimmunol.1200691 [PubMed: 23024277]
- Halim TYF, Hwang YY, Scanlon ST, Zaghouni H, Garbi N, Fallon PG, McKenzie ANJ. Group 2 innate lymphoid cells license dendritic cells to potentiate memory T. *Nature Immunology*. 2015; : 1–10. DOI: 10.1038/ni.3294 [PubMed: 25521670]
- Halim TYF, Steer CA, Mathä L, Gold MJ, Martinez-Gonzalez I, McNagny KM, McKenzie ANJ, Takei F. Group 2 innate lymphoid cells are critical for the initiation of adaptive T helper 2 cell-mediated allergic lung inflammation. *Immunity*. 2014; 40:425–435. DOI: 10.1016/j.immuni.2014.01.011 [PubMed: 24613091]
- Hammad H, Lambrecht BN. Barrier Epithelial Cells and the Control of Type 2 Immunity. *Immunity*. 2015; 43:29–40. DOI: 10.1016/j.immuni.2015.07.007 [PubMed: 26200011]
- Hams E, Armstrong ME, Barlow JL, Saunders SP, Schwartz C, Cooke G, Fahy RJ, Crotty TB, Hirani N, Flynn RJ, Voehringer D, McKenzie ANJ, Donnelly SC, Fallon PG. IL-25 and type 2 innate lymphoid cells induce pulmonary fibrosis. *Proceedings of the National Academy of Sciences of the United States of America*. 2014; 111:367–372. DOI: 10.1073/pnas.1315854111 [PubMed: 24344271]
- Henry EK, Sy CB, Inclan-Rico JM, Espinosa V, Ghanny SS, Dwyer DF, Soteropoulos P, Rivera A, Siracusa MC. Carbonic anhydrase enzymes regulate mast cell-mediated inflammation. *Journal of Experimental Medicine*. 2016; 151 jem.20151739-11. doi: 10.1084/jem.20151739
- Hepworth MR, Daniłowicz-Luebert E, Rausch S, Metz M, Klotz C, Maurer M, Hartmann S. Mast cells orchestrate type 2 immunity to helminths through regulation of tissue-derived cytokines. *Proceedings of the National Academy of Sciences of the United States of America*. 2012a; 109:6644–6649. DOI: 10.1073/pnas.1112268109 [PubMed: 22493240]
- Hepworth MR, Maurer M, Hartmann S. Regulation of type 2 immunity to helminths by mast cells. *Gut Microbes*. 2012b; 3:476–481. DOI: 10.4161/gmic.21507 [PubMed: 22892692]
- Herbst T, Esser J, Prati M, Kulagin M, Stettler R, Zaiss MM, Hewitson JP, Merky P, Verbeek JS, Bourquin C, Camberis M, Prout M, Maizels RM, Le Gros G, Harris NL. Antibodies and IL-3 support helminth-induced basophil expansion. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109:14954–14959. DOI: 10.1073/pnas.1117584109 [PubMed: 22930820]
- Hewitson JP, Filbey KJ, Esser-von Bieren J, Camberis M, Schwartz C, Murray J, Reynolds LA, Blair N, Robertson E, Hargus Y, Boon L, Huang SCC, Yang L, Tu Y, Miller MJ, Voehringer D, Le Gros G, Harris N, Maizels RM. Concerted activity of IgG1 antibodies and IL-4/IL-25-dependent effector cells trap helminth larvae in the tissues following vaccination with defined secreted

- antigens, providing sterile immunity to challenge infection. *PLoS pathogens*. 2015; 11:e1004676.doi: 10.1371/journal.ppat.1004676 [PubMed: 25816012]
- Howitt MR, Lavoie S, Michaud M, Blum AM, Tran SV, Weinstock JV, Gallini CA, Redding K, Margolskee RF, Osborne LC, Artis D, Garrett WS. Tuft cells, taste-chemosensory cells, orchestrate parasite type 2 immunity in the gut. *Science*. 2016; 351:1329–1333. DOI: 10.1126/science.aaf1648 [PubMed: 26847546]
- Hoyler T, Klose CSN, Souabni A, Turqueti-Neves A, Pfeifer D, Rawlins EL, Voehringer D, Busslinger M, Diefenbach A. The transcription factor GATA-3 controls cell fate and maintenance of type 2 innate lymphoid cells. *Immunity*. 2012; 37:634–648. DOI: 10.1016/j.immuni.2012.06.020 [PubMed: 23063333]
- Huang L, Appleton JA. Eosinophils in Helminth Infection: Defenders and Dupes. *Trends in parasitology*. 2016; 32:798–807. DOI: 10.1016/j.pt.2016.05.004 [PubMed: 27262918]
- Huang Y, Guo L, Qiu J, Chen X, Hu-Li J, Siebenlist U, Williamson PR, Urban JF, Paul WE. IL-25-responsive, lineage-negative KLRG1hi cells are multipotential “inflammatory” type 2 innate lymphoid cells. *Nature Immunology*. 2014; 16:161–169. DOI: 10.1038/ni.3078 [PubMed: 25531830]
- Ierna MX, Scales HE, Saunders KL, Lawrence CE. Mast cell production of IL-4 and TNF may be required for protective and pathological responses in gastrointestinal helminth infection. *Mucosal Immunology*. 2008; 1:147–155. DOI: 10.1038/mi.2007.16 [PubMed: 19079172]
- Ito T, Wang YH, Duramad O, Hori T, Delespesse GJ, Watanabe N, Qin FFX, Yao Z, Cao W, Liu YJ. TSLP-activated dendritic cells induce an inflammatory T helper type 2 cell response through OX40 ligand. *The Journal of experimental medicine*. 2005; 202:1213–1223. DOI: 10.1084/jem.20051135 [PubMed: 16275760]
- Kim BS, Siracusa MC, Saenz SA, Noti M, Monticelli LA, Sonnenberg GF, Hepworth MR, Van Voorhees AS, Comeau MR, Artis D. TSLP elicits IL-33-independent innate lymphoid cell responses to promote skin inflammation. *Science translational medicine*. 2013; 5:170ra16.doi: 10.1126/scitranslmed.3005374
- Kim BS, Wang K, Siracusa MC, Saenz SA, Brestoff JR, Monticelli LA, Noti M, Tait Wojno ED, Fung TC, Kubo M, Artis D. Basophils promote innate lymphoid cell responses in inflamed skin. *The Journal of Immunology*. 2014; 193:3717–3725. DOI: 10.4049/jimmunol.1401307 [PubMed: 25156365]
- King CL, Xianli J, Malhotra I, Liu S, Mahmoud AA, Oettgen HC. Mice with a targeted deletion of the IgE gene have increased worm burdens and reduced granulomatous inflammation following primary infection with *Schistosoma mansoni*. *Journal of immunology (Baltimore, Md : 1950)*. 1997; 158:294–300.
- King IL, Mohrs M. IL-4-producing CD4+ T cells in reactive lymph nodes during helminth infection are T follicular helper cells. *Journal of Experimental Medicine*. 2009; 206:1001–1007. DOI: 10.1084/jem.20090313 [PubMed: 19380638]
- Klose CSN, Artis D. Innate lymphoid cells as regulators of immunity, inflammation and tissue homeostasis. *Nature Immunology*. 2016; 17:765–774. DOI: 10.1038/ni.3489 [PubMed: 27328006]
- Knipper JA, Willenborg S, Brinckmann J, Bloch W, Maaß T, Wagener R, Krieg T, Sutherland T, Munitz A, Rothenberg ME, Niehoff A, Richardson R, Hammerschmidt M, Allen JE, Eming SA. Interleukin-4 Receptor a Signaling in Myeloid Cells Controls Collagen Fibril Assembly in Skin Repair. *Immunity*. 2015; 43:803–816. DOI: 10.1016/j.immuni.2015.09.005 [PubMed: 26474656]
- Kroeger KM, Sullivan BM, Locksley RM. IL-18 and IL-33 elicit Th2 cytokines from basophils via a MyD88- and p38alpha-dependent pathway. *Journal of Leukocyte Biology*. 2009; 86:769–778. DOI: 10.1189/jlb.0708452 [PubMed: 19451398]
- Kumamoto Y, Linehan M, Weinstein JS, Laidlaw BJ, Craft JE, Iwasaki A. CD301b+ Dermal Dendritic Cells Drive T Helper 2 Cell-Mediated Immunity. *Immunity*. 2013; 39:733–743. DOI: 10.1016/j.immuni.2013.08.029 [PubMed: 24076051]
- Larson D, Cooper PJ, Hübner MP, Reyes J, Vaca M, Chico M, Kong HH, Mitre E. Helminth infection is associated with decreased basophil responsiveness in human beings. *The Journal of Allergy and Clinical Immunology*. 2012a; 130:270–272. DOI: 10.1016/j.jaci.2012.04.017 [PubMed: 22608572]

- Larson D, Hubner MP, Torrero MN, Morris CP, Brankin A, Swierczewski BE, Davies SJ, Vonakis BM, Mitre E. Chronic Helminth Infection Reduces Basophil Responsiveness in an IL-10-Dependent Manner. *Journal of immunology* (Baltimore, Md : 1950). 2012b; 188:4188–4199. DOI: 10.4049/jimmunol.1101859
- Lawrence CE, Paterson YYW, Wright SH, Knight PA, Miller HRP. Mouse mast cell protease-1 is required for the enteropathy induced by gastrointestinal helminth infection in the mouse. *Gastroenterology*. 2004; 127:155–165. DOI: 10.1053/j.gastro.2004.04.004 [PubMed: 15236182]
- Lefrançois E, Duval A, Mirey E, Roga S, Espinosa E, Cayrol C, Girard JP. Central domain of IL-33 is cleaved by mast cell proteases for potent activation of group-2 innate lymphoid cells. *Proceedings of the National Academy of Sciences of the United States of America*. 2014; 111:15502–15507. DOI: 10.1073/pnas.1410700111 [PubMed: 25313073]
- León B, Ballesteros-Tato A, Browning JL, Dunn R, Randall TD, Lund FE. Regulation of TH2 development by CXCR5+ dendritic cells and lymphotoxin-expressing B cells. *Nature Immunology*. 2012; 13:681–690. DOI: 10.1038/ni.2309 [PubMed: 22634865]
- Liu AY, Dwyer DF, Jones TG, Bankova LG, Shen S, Katz HR, Austen KF, Gurish MF. Mast cells recruited to mesenteric lymph nodes during helminth infection remain hypogranular and produce IL-4 and IL-6. *The Journal of Immunology*. 2013; 190:1758–1766. DOI: 10.4049/jimmunol.1202567 [PubMed: 23319739]
- Lobos E, Nutman TB, Hothersall JS, Moncada S. Elevated Immunoglobulin E against Recombinant *Brugia malayi* -Glutamyl Transpeptidase in Patients with Bancroftian Filariasis: Association with Tropical Pulmonary Eosinophilia or Putative Immunity. *Infection and Immunity*. 2003; 71:747–753. DOI: 10.1128/IAI.71.2.747-753.2003 [PubMed: 12540554]
- Lundie RJ, Webb LM, Marley AK, Phythian-Adams AT, Cook PC, Jackson-Jones LH, Brown S, Maizels RM, Boon L, O'Keeffe M, MacDonald AS. A central role for hepatic conventional dendritic cells in supporting Th2 responses during helminth infection. *Immunology and Cell Biology*. 2016; 94:400–410. DOI: 10.1038/icb.2015.114 [PubMed: 26657145]
- MacDonald AS, Maizels RM. Alarming dendritic cells for Th2 induction. *J Exp Med*. 2008; 205:13–17. DOI: 10.1084/jem.20040662 [PubMed: 18195077]
- Maizels RM, Withers DR. MHC-II: A Mutual Support System for ILCs and T Cells? *Immunity*. 2014; 41:174–176. DOI: 10.1016/j.immuni.2014.07.006 [PubMed: 25148019]
- Martinez-Gonzalez I, Mathä L, Steer CA, Ghaedi M, Poon GFT, Takei F. Allergen-Experienced Group 2 Innate Lymphoid Cells Acquire Memory-like Properties and Enhance Allergic Lung Inflammation. *Immunity*. 2016; 45:198–208. DOI: 10.1016/j.immuni.2016.06.017 [PubMed: 27421705]
- McDermott JR, Bartram RE, Knight PA, Miller HRP, Garrod DR, Grencis RK. Mast cells disrupt epithelial barrier function during enteric nematode infection. *Proceedings of the National Academy of Sciences*. 2003; 100:7761–7766. DOI: 10.1073/pnas.1231488100
- Meredith MM, Liu K, Darrasse-Jeze G, Kamphorst AO, Schreiber HA, Guermontprez P, Idoyaga J, Cheong C, Yao KH, Niec RE, Nussenzweig MC. Expression of the zinc finger transcription factor zDC (Zbtb46, Btbd4) defines the classical dendritic cell lineage. *Journal of Experimental Medicine*. 2012; 209:1153–1165. DOI: 10.1084/jem.20112675 [PubMed: 22615130]
- Min B, Prout M, Hu-Li J, Zhu J, Jankovic D, Morgan ES, Urban JF, Dvorak AM, Finkelman FD, LeGros G, Paul WE. Basophils produce IL-4 and accumulate in tissues after infection with a Th2-inducing parasite. *The Journal of experimental medicine*. 2004; 200:507–517. DOI: 10.1084/jem.20040590 [PubMed: 15314076]
- Mirchandani AS, Besnard AG, Yip E, Scott C, Bain CC, Cerovic V, Salmond RJ, Liew FY. Type 2 innate lymphoid cells drive CD4+ Th2 cell responses. *The Journal of Immunology*. 2014; 192:2442–2448. DOI: 10.4049/jimmunol.1300974 [PubMed: 24470502]
- Mitchell KM, Mutapi F, Savill NJ, Woolhouse MEJ. Protective immunity to *Schistosoma haematobium* infection is primarily an anti-fecundity response stimulated by the death of adult worms. *Proceedings of the National Academy of Sciences of the United States of America*. 2012; 109:13347–13352. DOI: 10.1073/pnas.1121051109 [PubMed: 22847410]
- Mitre E, Taylor RT, Kubofcik J, Nutman TB. Parasite Antigen-Driven Basophils Are a Major Source of IL-4 in Human Filarial Infections. *Journal of immunology* (Baltimore, Md : 1950). 2004; 172:2439–2445. DOI: 10.4049/jimmunol.172.4.2439

- Mohapatra A, Van Dyken SJ, Schneider C, Nussbaum JC, Liang HE, Locksley RM. Group 2 innate lymphoid cells utilize the IRF4-IL-9 module to coordinate epithelial cell maintenance of lung homeostasis. 2015; :1–12. DOI: 10.1038/mi.2015.59
- von Moltke J, Ji M, Liang HE, Locksley RM. Tuft-cell-derived IL-25 regulates an intestinal ILC2-epithelial response circuit. *Nature*. 2016; 529:221–225. DOI: 10.1038/nature16161 [PubMed: 26675736]
- Monticelli LA, Buck MD, Flamar AL, Saenz SA, Wojno EDT, Yudanin NA, Osborne LC, Hepworth MR, Tran SV, Rodewald HR, Shah H, Cross JR, Diamond JM, Cantu E, Christie JD, Pearce EL, Artis D. Arginase 1 is an innate lymphoid-cell-intrinsic metabolic checkpoint controlling type 2 inflammation. *Nature Immunology*. 2016; :1–12. DOI: 10.1038/ni.3421
- Monticelli LA, Osborne LC, Noti M, Tran SV, Zaiss DMW, Artis D. IL-33 promotes an innate immune pathway of intestinal tissue protection dependent on amphiregulin–EGFR interactions. *Proceedings of the National Academy of Sciences*. 2015; 112:10762–10767. DOI: 10.1073/pnas.1509070112
- Monticelli LA, Sonnenberg GF, Abt MC, Alenghat T, Ziegler CGK, Doering TA, Angelosanto JM, Laidlaw BJ, Yang CY, Sathaliyawala T, Kubota M, Turner D, Diamond JM, Goldrath AW, Farber DL, Collman RG, Wherry EJ, Artis D. Innate lymphoid cells promote lung-tissue homeostasis after infection with influenza virus. *Nature Immunology*. 2011; 12:1045–1054. DOI: 10.1038/ni.2131 [PubMed: 21946417]
- Morales JK, Saleem SJ, Martin RK, Saunders BL, Barnstein BO, Faber TW, Pullen NA, Kolawole EM, Brooks KB, Norton SK, Sturgill J, Graham L, Bear HD, Urban JF, Lantz CS, Conrad DH, Ryan JJ. Myeloid-derived suppressor cells enhance IgE-mediated mast cell responses. *Journal of Leukocyte Biology*. 2014; 95:643–650. DOI: 10.1189/jlb.0913510 [PubMed: 24338630]
- Moro K, Yamada T, Tanabe M, Takeuchi T, Ikawa T, Kawamoto H, Furusawa JI, Ohtani M, Fujii H, Koyasu S. Innate production of T(H)2 cytokines by adipose tissue-associated c-Kit(+)Sca-1(+) lymphoid cells. *Nature*. 2010; 463:540–544. DOI: 10.1038/nature08636 [PubMed: 20023630]
- Murray PJ, Wynn TA. Protective and pathogenic functions of macrophage subsets. *Nature Reviews Immunology*. 2011; 11:723–737. DOI: 10.1038/nri3073
- Nair MG, Du Y, Perrigoue JG, Zaph C, Taylor JJ, Goldschmidt M, Swain GP, Yancopoulos GD, Valenzuela DM, Murphy A, Karow M, Stevens S, Pearce EJ, Artis D. Alternatively activated macrophage-derived RELM- α is a negative regulator of type 2 inflammation in the lung. *Journal of Experimental Medicine*. 2009; 206:937–952. DOI: 10.1084/jem.20082048 [PubMed: 19349464]
- Nakamura T, Maeda S, Horiguchi K, Maehara T, Aritake K, Choi BI, Iwakura Y, Urade Y, Murata T. PGD2 deficiency exacerbates food antigen-induced mast cell hyperplasia. *Nature communications*. 2015; 6:7514.doi: 10.1038/ncomms8514
- Nausch N, Appleby LJ, Sparks AM, Midzi N, Mduluzi T, Mutapi F. Group 2 Innate Lymphoid Cell Proportions Are Diminished in Young Helminth Infected Children and Restored by Curative Anti-helminthic Treatment. *PLoS neglected tropical diseases*. 2015; 9:e0003627–16. DOI: 10.1371/journal.pntd.0003627 [PubMed: 25799270]
- Nei Y, Obata-Ninomiya K, Tsutsui H, Ishiwata K, Miyasaka M, Matsumoto K, Nakae S, Kanuka H, Inase N, Karasuyama H. GATA-1 regulates the generation and function of basophils. *Proceedings of the National Academy of Sciences of the United States of America*. 2013; 110:18620–18625. DOI: 10.1073/pnas.1311668110 [PubMed: 24167252]
- Neill DR, Wong SH, Bellosi A, Flynn RJ, Daly M, Langford TKA, Bucks C, Kane CM, Fallon PG, Pannell R, Jolin HE, McKenzie ANJ. Nuocytes represent a new innate effector leukocyte that mediates type-2 immunity. *Nature*. 2010; doi: 10.1038/nature08900
- Nieves W, Hung LY, Oniskey TK, Boon L, Foretz M, Viollet B, Herbert DR. Myeloid-Restricted AMPK α 1 Promotes Host Immunity and Protects against IL-12/23p40–Dependent Lung Injury during Hookworm Infection. *Journal of immunology (Baltimore, Md : 1950)*. 2016; 196:4632–4640. DOI: 10.4049/jimmunol.1502218
- Noti M, Wojno EDT, Kim BS, Siracusa MC, Giacomini PR, Nair MG, Benitez AJ, Ruymann KR, Muir AB, Hill DA, Chikwava KR, Moghaddam AE, Sattentau QJ, Alex A, Zhou C, Yearley JH, Menard-Katcher P, Kubo M, Obata-Ninomiya K, Karasuyama H, Comeau MR, Brown-Whitehorn T, de Waal Malefyt R, Sleiman PM, Hakonarson H, Cianferoni A, Falk GW, Wang ML, Spergel

- JM, Artis D. Thymic stromal lymphopoietin–elicited basophil responses promote eosinophilic esophagitis. *Nature Medicine*. 2013; 19:1005–1013. DOI: 10.1038/nm.3281
- Obata-Ninomiya K, Ishiwata K, Tsutsui H, Nei Y, Yoshikawa S, Kawano Y, Minegishi Y, Ohta N, Watanabe N, Kanuka H, Karasuyama H. The skin is an important bulwark of acquired immunity against intestinal helminths. *Journal of Experimental Medicine*. 2013; 210:2583–2595. DOI: 10.1084/jem.20130761 [PubMed: 24166714]
- Ohnmacht C, Voehringer D. Basophils protect against reinfection with hookworms independently of mast cells and memory Th2 cells. *The Journal of Immunology*. 2010; 184:344–350. DOI: 10.4049/jimmunol.0901841 [PubMed: 19955520]
- Ohnmacht C, Schwartz C, Panzer M, Schiedewitz I, Naumann R, Voehringer D. Basophils orchestrate chronic allergic dermatitis and protective immunity against helminths. *Immunity*. 2010; 33:364–374. DOI: 10.1016/j.immuni.2010.08.011 [PubMed: 20817571]
- Okamoto M, Matsuda H, Joetham A, Lucas JJ, Domenico J, Yasutomo K, Takeda K, Gelfand EW. Jagged1 on dendritic cells and Notch on CD4+ T cells initiate lung allergic responsiveness by inducing IL-4 production. *The Journal of Immunology*. 2009; 183:2995–3003. DOI: 10.4049/jimmunol.0900692 [PubMed: 19667086]
- Oliphant CJ, Hwang YY, Walker JA, Salimi M, Wong SH, Brewer JM, Englezakis A, Barlow JL, Hams E, Scanlon ST, Ogg GS, Fallon PG, McKenzie ANJ. MHCII-Mediated Dialog between Group 2 Innate Lymphoid Cells and CD4. *Immunity*. 2014; :1–13. DOI: 10.1016/j.immuni.2014.06.016
- Park MK, Cho MK, Kang SA, Park HK, Kim YS, Kim KU, Ahn SC, Kim DH, Yu HS. Protease-Activated Receptor 2 Is Involved in Th2 Responses against *Trichinella spiralis* Infection. *The Korean Journal of Parasitology*. 2011; 49:235–10. DOI: 10.3347/kjp.2011.49.3.235 [PubMed: 22072823]
- Patel N, Wu W, Mishra PK, Chen F, Millman A, Csóka B, Koscsó B, Eltzschig HK, Haskó G, Gause WC. A2B Adenosine Receptor Induces Protective Antihelminth Type 2 Immune Responses. *CHOM*. 2014; 15:339–350. DOI: 10.1016/j.chom.2014.02.001
- Pearce EJ, MacDonald AS. The immunobiology of schistosomiasis. *Nature Reviews Immunology*. 2002; 2:499–511. DOI: 10.1038/nri843
- Pecaric-Petkovic T, Didichenko SA, Kaempfer S, Spiegl N, Dahinden CA. Human basophils and eosinophils are the direct target leukocytes of the novel IL-1 family member IL-33. *Blood*. 2009; 113:1526–1534. DOI: 10.1182/blood-2008-05-157818 [PubMed: 18955562]
- Pelly VS, Kannan Y, Coomes SM, Entwistle LJ, Ruckerl DRU, Seddon B, MacDonald AS, McKenzie A, Wilson MS. IL-4-producing ILC2s are required for the differentiation of TH2 cells following *Heligmosomoides polygyrus* infection. *Mucosal Immunology*. 2016; :1–11. DOI: 10.1038/mi.2016.4 [PubMed: 25872482]
- Perrigoue JG, Saenz SA, Siracusa MC, Allenspach EJ, Taylor BC, Giacomini PR, Nair MG, Du Y, Zaph C, van Rooijen N, Comeau MR, Pearce EJ, Laufer TM, Artis D. MHC class II-dependent basophil-CD4+ T cell interactions promote T(H)2 cytokine-dependent immunity. *Nature Immunology*. 2009; 10:697–705. DOI: 10.1038/ni.1740 [PubMed: 19465906]
- Pesce JT, Ramalingam TR, Mentink-Kane MM, Wilson MS, Kasmi EI KC, Smith AM, Thompson RW, Cheever AW, Murray PJ, Wynn TA. Arginase-1-expressing macrophages suppress Th2 cytokine-driven inflammation and fibrosis. *PLoS pathogens*. 2009a; 5:e1000371.doi: 10.1371/journal.ppat.1000371 [PubMed: 19360123]
- Pesce JT, Ramalingam TR, Wilson MS, Mentink-Kane MM, Thompson RW, Cheever AW, Urban JF, Wynn TA. Retnla (relmalph/fizz1) suppresses helminth-induced Th2-type immunity. *PLoS pathogens*. 2009b; 5:e1000393.doi: 10.1371/journal.ppat.1000393 [PubMed: 19381262]
- Phillips C, Coward WR, Pritchard DI, Hewitt CRA. Basophils express a type 2 cytokine profile on exposure to proteases from helminths and house dust mites. *Journal of Leukocyte Biology*. 2003; 73:165–171. [PubMed: 12525574]
- Phythian-Adams AT, Cook PC, Lundie RJ, Jones LH, Smith KA, Barr TA, Hochweller K, Anderton SM, Hammerling GJ, Maizels RM, MacDonald AS. CD11c depletion severely disrupts Th2 induction and development in vivo. *J Exp Med*. 2010; 207:2089–2096. DOI: 10.1038/ni.1737 [PubMed: 20819926]

- Pinot de Moira A, Fitzsimmons CM, Jones FM, Wilson S, Cahen P, Tukahebwa E, Mpairwe H, Mwatha JK, Bethony JM, Skov PS, Kabatereine NB, Dunne DW. Suppression of basophil histamine release and other IgE-dependent responses in childhood *Schistosoma mansoni*/hookworm coinfection. *The Journal of Infectious Diseases*. 2014; 210:1198–1206. DOI: 10.1093/infdis/jiu234 [PubMed: 24782451]
- Prendergast CT, Sanin DE, Mountford AP. CD4 T-cell hyporesponsiveness induced by schistosome larvae is not dependent upon eosinophils but may involve connective tissue mast cells. *Parasite immunology*. 2016; 38:81–92. DOI: 10.1111/pim.12300 [PubMed: 26679416]
- Price AE, Liang HE, Sullivan BM, Reinhardt RL, Eisley CJ, Erle DJ, Locksley RM. Systemically dispersed innate IL-13-expressing cells in type 2 immunity. *Proceedings of the National Academy of Sciences of the United States of America*. 2010; 107:11489–11494. DOI: 10.1073/pnas.1003988107 [PubMed: 20534524]
- Pulendran B, Artis D. New Paradigms in Type 2 Immunity. *Science*. 2012; 337:431–435. DOI: 10.1126/science.1221064 [PubMed: 22837519]
- Reese TA, Liang HE, Tager AM, Luster AD, van Rooijen N, Voehringer D, Locksley RM. Chitin induces accumulation in tissue of innate immune cells associated with allergy. *Nature*. 2007; 447:92–96. DOI: 10.1038/nature05746 [PubMed: 17450126]
- Reynolds LA, Filbey KJ, Maizels RM. Immunity to the model intestinal helminth parasite *Heligmosomoides polygyrus*. *Seminars in Immunopathology*. 2012; doi: 10.1007/s00281-012-0347-3
- Rivera A, Siracusa MC, Yap GS, Gause WC. Innate cell communication kick-starts pathogen-specific immunity. *Nature Immunology*. 2016; 17:356–363. DOI: 10.1038/ni.3375 [PubMed: 27002843]
- Roediger B, Kyle R, Yip KH, Sumaria N, Guy TV, Kim BS, Mitchell AJ, Tay SS, Jain R, Forbes-Blom E, Chen X, Tong PL, Bolton HA, Artis D, Paul WE, de St Groth BF, Grimbaldston MA, Le Gros G, Weninger W. Cutaneous immunosurveillance and regulation of inflammation by group 2 innate lymphoid cells. *Nature Immunology*. 2013; 14:564–573. DOI: 10.1038/ni.2584 [PubMed: 23603794]
- Sakata-Yanagimoto M, Nakagami-Yamaguchi E, Saito T, Kumano K, Yasutomo K, Ogawa S, Kurokawa M, Chiba S. Coordinated regulation of transcription factors through Notch2 is an important mediator of mast cell fate. *Proceedings of the National Academy of Sciences*. 2008; 105:7839–7844. DOI: 10.1073/pnas.0801074105
- Sakata-Yanagimoto M, Sakai T, Miyake Y, Saito TI, Maruyama H, Morishita Y, Nakagami-Yamaguchi E, Kumano K, Yagita H, Fukayama M, Ogawa S, Kurokawa M, Yasutomo K, Chiba S. Notch2 signaling is required for proper mast cell distribution and mucosal immunity in the intestine. *Blood*. 2011; 117:128–134. DOI: 10.1182/blood-2010-07-289611 [PubMed: 20971948]
- Satpathy AT, KC W, Albring JC, Edelson BT, Kretzer NM, Bhattacharya D, Murphy TL, Murphy KM. Zbtb46 expression distinguishes classical dendritic cells and their committed progenitors from other immune lineages. *Journal of Experimental Medicine*. 2012a; 209:1135–1152. DOI: 10.1084/jem.20120030 [PubMed: 22615127]
- Satpathy AT, Wu X, Albring JC, Murphy KM. Re(de)fining the dendritic cell lineage. *Nature Immunology*. 2012b; 13:1145–1154. DOI: 10.1038/ni.2467 [PubMed: 23160217]
- Schramm G, Gronow A, Knobloch J, Wippersteg V, Grevelding CG, Galle J, Fuller H, Stanley RG, Chiodini PL, Haas H, Doenhoff MJ. IPSE/alpha-1: a major immunogenic component secreted from *Schistosoma mansoni* eggs. *Molecular and biochemical parasitology*. 2006; 147:9–19. DOI: 10.1016/j.molbiopara.2006.01.003 [PubMed: 16480783]
- Schwartz C, Oeser K, Prazeres da Costa C, Layland LE, Voehringer D. T Cell-Derived IL-4/IL-13 Protects Mice against Fatal *Schistosoma mansoni* Infection Independently of Basophils. *Journal of immunology* (Baltimore, Md : 1950). 2014a; 193:3590–3599. DOI: 10.4049/jimmunol.1401155
- Schwartz C, Oeser K, Prazeres da Costa C, Layland LE, Voehringer D. T Cell-Derived IL-4/IL-13 Protects Mice against Fatal *Schistosoma mansoni* Infection Independently of Basophils. *The Journal of Immunology*. 2014b; doi: 10.4049/jimmunol.1401155
- Schwartz C, Turqueti-Neves A, Hartmann S, Yu P, Nimmerjahn F, Voehringer D. Basophil-mediated protection against gastrointestinal helminths requires IgE-induced cytokine secretion.

- Proceedings of the National Academy of Sciences. 2014c; 111:E5169–E5177. DOI: 10.1073/pnas.1412663111
- Sharma M, Stephen-Victor E, Poncet P, Kaveri SV, Bayry J. Basophils are inept at promoting human Th17 responses. *Human immunology*. 2015; 76:176–180. DOI: 10.1016/j.humimm.2014.12.015 [PubMed: 25526920]
- Sharma R, Zheng L, Deshmukh US, Jarjour WN, Sung SSJ, Fu SM, Ju ST. A regulatory T cell-dependent novel function of CD25 (IL-2Ralpha) controlling memory CD8(+) T cell homeostasis. *Journal of immunology* (Baltimore, Md : 1950). 2007; 178:1251–1255.
- Shen T, Kim S, Do JS, Wang L, Lantz C, Urban JF, Le Gros G, Min B. T cell-derived IL-3 plays key role in parasite infection-induced basophil production but is dispensable for in vivo basophil survival. *International Immunology*. 2008; 20:1201–1209. DOI: 10.1093/intimm/dxn077 [PubMed: 18632726]
- Shih HY, Sciumè G, Mikami Y, Guo L, Sun HW, Brooks SR, Urban JF Jr, Davis FP, Kanno Y, O'Shea JJ. Developmental Acquisition of Regulomes Underlies Innate Lymphoid Cell Functionality. *Cell*. 2016; :1–24. DOI: 10.1016/j.cell.2016.04.029
- Siracusa MC, Saenz SA, Hill DA, Kim BS, Headley MB, Doering TA, Wherry EJ, Jessup HK, Siegel LA, Kambayashi T, Dudek EC, Kubo M, Cianferoni A, Spergel JM, Ziegler SF, Comeau MR, Artis D. TSLP promotes interleukin-3-independent basophil haematopoiesis and type 2 inflammation. *Nature*. 2011; 477:229–233. DOI: 10.1038/nature10329 [PubMed: 21841801]
- Siracusa MC, Saenz SA, Wojno EDT, Kim BS, Osborne LC, Ziegler CG, Benitez AJ, Ruymann KR, Farber DL, Sleiman PM, Hakonarson H, Cianferoni A, Wang ML, Spergel JM, Comeau MR, Artis D. Thymic stromal lymphopoietin-mediated extramedullary hematopoiesis promotes allergic inflammation. *Immunity*. 2013a; 39:1158–1170. DOI: 10.1016/j.immuni.2013.09.016 [PubMed: 24332033]
- Siracusa MC, Saenz SA, Wojno EDT, Kim BS, Osborne LC, Ziegler CG, Benitez AJ, Ruymann KR, Farber DL, Sleiman PM, Hakonarson H, Cianferoni A, Wang ML, Spergel JM, Comeau MR, Artis D. Thymic Stromal Lymphopoietin-Mediated Extramedullary Hematopoiesis Promotes Allergic Inflammation. *Immunity*. 2013b; 39:1158–1170. DOI: 10.1016/j.immuni.2013.09.016 [PubMed: 24332033]
- Smith KA, Harcus Y, Garbi N, Hämmerling GJ, MacDonald AS, Maizels RM. Type 2 innate immunity in helminth infection is induced redundantly and acts autonomously following CD11c(+) cell depletion. *Infection and Immunity*. 2012; 80:3481–3489. DOI: 10.1128/IAI.00436-12 [PubMed: 22851746]
- Sokol CL, Chu NQ, Yu S, Nish SA, Laufer TM, Medzhitov R. Basophils function as antigen-presenting cells for an allergen-induced T helper type 2 response. *Nature Immunology*. 2009; 10:713–720. DOI: 10.1038/ni.1738 [PubMed: 19465907]
- Specht S, Frank JK, Alferink J, Dubben B, Layland LE, Denece G, Bain O, Förster I, Kirschning CJ, Martin C, Hoerauf A. CCL17 controls mast cells for the defense against filarial larval entry. *The Journal of Immunology*. 2011; 186:4845–4852. DOI: 10.4049/jimmunol.1000612 [PubMed: 21398605]
- Sullivan BM, Liang HE, Bando JK, Wu D, Cheng LE, McKerrow JK, Allen CDC, Locksley RM. Genetic analysis of basophil function in vivo. *Nature Immunology*. 2011; 12:527–535. DOI: 10.1038/ni.2036 [PubMed: 21552267]
- Tait Wojno ED, Artis D. Emerging concepts and future challenges in innate lymphoid cell biology. *The Journal of experimental medicine*. 2016; 213:2229–2248. DOI: 10.1084/jem.20160525 [PubMed: 27811053]
- Tait Wojno ED, Monticelli LA, Tran SV, Alenghat T, Osborne LC, Thome JJ, Willis C, Budelsky A, Farber DL, Artis D. The prostaglandin D2 receptor CRTH2 regulates accumulation of group 2 innate lymphoid cells in the inflamed lung. *Mucosal Immunology*. 2015; doi: 10.1038/mi.2015.21
- Tawill S, Le Goff L, Ali F, Blaxter M, Allen JE. Both free-living and parasitic nematodes induce a characteristic Th2 response that is dependent on the presence of intact glycans. *Infection and Immunity*. 2004; 72:398–407. DOI: 10.1128/IAI.72.1.398-407.2004 [PubMed: 14688121]

- Taylor BC, Zaph C, Troy AE, Du Y, Guild KJ, Comeau MR, Artis D. TSLP regulates intestinal immunity and inflammation in mouse models of helminth infection and colitis. *The Journal of experimental medicine*. 2009; 206:655–667. DOI: 10.1084/jem.20081499 [PubMed: 19273626]
- Thawer S, Auret J, Schnoeller C, Chetty A, Smith K, Darby M, Roberts L, Mackay RM, Whitwell HJ, Timms JF, Madsen J, Selkirk ME, Brombacher F, Clark HW, Horsnell WGC. Surfactant Protein-D Is Essential for Immunity to Helminth Infection. *PLoS pathogens*. 2016; 12:e1005461–18. DOI: 10.1371/journal.ppat.1005461 [PubMed: 26900854]
- Tjota MY, Williams JW, Lu T, Clay BS, Byrd T, Hrusch CL, Decker DC, de Araujo CA, Bryce PJ, Sperling AI. IL-33-dependent induction of allergic lung inflammation by Fc γ RIII signaling. *The Journal of clinical investigation*. 2013; 123:2287–2297. DOI: 10.1172/JCI63802DS1 [PubMed: 23585480]
- Trottein F, Pavelka N, Vizzardelli C, Angeli V, Zouain CS, Pelizzola M, Capozzoli M, Urbano M, Capron M, Belardelli F, Granucci F, Ricciardi-Castagnoli P. A type I IFN-dependent pathway induced by *Schistosoma mansoni* eggs in mouse myeloid dendritic cells generates an inflammatory signature. *Journal of immunology* (Baltimore, Md : 1950). 2004; 172:3011–3017.
- Turner JE, Morrison PJ, Wilhelm C, Wilson M, Ahlfors H, Renauld JC, Panzer U, Helmby H, Stockinger B. IL-9-mediated survival of type 2 innate lymphoid cells promotes damage control in helminth-induced lung inflammation. *The Journal of experimental medicine*. 2013; 210:2951–2965. DOI: 10.1084/jem.20130071 [PubMed: 24249111]
- Tussiwand R, Everts B, Grajales-Reyes GE, Kretzer NM, Iwata A, Bagaitkar J, Wu X, Wong R, Anderson DA, Murphy TL, Pearce EJ, Murphy KM. Klf4 Expression in Conventional Dendritic Cells Is Required for T Helper 2 Cell Responses. *Immunity*. 2015; 42:916–928. DOI: 10.1016/j.immuni.2015.04.017 [PubMed: 25992862]
- Van Dyken SJ, Nussbaum JC, Lee J, Molofsky AB, Liang HE, Pollack JL, Gate RE, Haliburton GE, Ye CJ, Marson A, Erle DJ, Locksley RM. A tissue checkpoint regulates type 2 immunity. *Nature Immunology*. 2016; :1–9. DOI: 10.1038/ni.3582
- van Panhuys N, Prout M, Forbes E, Min B, Paul WE, Le Gros G. Basophils are the major producers of IL-4 during primary helminth infection. *The Journal of Immunology*. 2011; 186:2719–2728. DOI: 10.4049/jimmunol.1000940 [PubMed: 21270410]
- Vély F, Barlogis V, Vallentin B, Neven B, Piperoglou C, Ebbo M, Perchet T, Petit M, Yessaad N, Touzot F, Bruneau J, Mahlaoui N, Zucchini N, Farnarier C, Michel G, Moshous D, Blanche S, Dujardin A, Spits H, Distler JHW, Ramming A, Picard C, Golub R, Fischer A, Vivier E. Evidence of innate lymphoid cell redundancy in humans. *Nature Immunology*. 2016; 17:1291–1299. DOI: 10.1038/ni.3553 [PubMed: 27618553]
- Voehringer D. Protective and pathological roles of mast cells and basophils. *Nature Reviews Immunology*. 2013; 13:362–375. DOI: 10.1038/nri3427
- Voehringer D, Reese TA, Huang X, Shinkai K, Locksley RM. Type 2 immunity is controlled by IL-4/IL-13 expression in hematopoietic non-eosinophil cells of the innate immune system. *The Journal of experimental medicine*. 2006; 203:1435–1446. DOI: 10.1084/jem.20052448 [PubMed: 16702603]
- Wang YH, Ito T, Wang YH, Homey B, Watanabe N, Martin R, Barnes CJ, McIntyre BW, Gilliet M, Kumar R, Yao Z, Liu YJ. Maintenance and polarization of human TH2 central memory T cells by thymic stromal lymphopoietin-activated dendritic cells. *Immunity*. 2006; 24:827–838. DOI: 10.1016/j.immuni.2006.03.019 [PubMed: 16782037]
- Wilhelm C, Harrison OJ, Schmitt V, Pelletier M, Spencer SP, Urban JF Jr, Ploch M, Ramalingam TR, Siegel RM, Belkaid Y. Critical role of fatty acid metabolism in ILC2-mediated barrier protection during malnutrition and helminth infection. *The Journal of experimental medicine*. 2016; 213:1409–1418. DOI: 10.1084/jem.20151448 [PubMed: 27432938]
- Williams JW, Tjota MY, Clay BS, Vander Lugt B, Bandukwala HS, Hrusch CL, Decker DC, Blaine KM, Fixsen BR, Singh H, Sciammas R, Sperling AI. Transcription factor IRF4 drives dendritic cells to promote Th2 differentiation. *Nature communications*. 2013; 4:2990.doi: 10.1038/ncomms3990
- Wills-Karp M, Rani R, Dienger K, Lewkowich I, Fox JG, Perkins C, Lewis L, Finkelman FD, Smith DE, Bryce PJ, Kurt-Jones EA, Wang TC, Sivaprasad U, Hershey GK, Herbert DR. Trefoil factor 2 rapidly induces interleukin 33 to promote type 2 immunity during allergic asthma and

- hookworm infection. *The Journal of experimental medicine*. 2012; 209:607–622. DOI: 10.1084/jem.20110079 [PubMed: 22329990]
- Xue L, Salimi M, Panse I, Mjösberg JM, McKenzie ANJ, Spits H, Klenerman P, Ogg G. Prostaglandin D2 activates group 2 innate lymphoid cells through chemoattractant receptor-homologous molecule expressed on TH2 cells. *The Journal of Allergy and Clinical Immunology*. 2014; 133:1184–1194. DOI: 10.1016/j.jaci.2013.10.056 [PubMed: 24388011]
- Yoshimoto T, Tsutsui H, Tominaga K, Hoshino K, Okamura H, Akira S, Paul WE, Nakanishi K. IL-18, although antiallergic when administered with IL-12, stimulates IL-4 and histamine release by basophils. *Proceedings of the National Academy of Sciences*. 1999; 96:13962–13966.
- Zhang K, Xu X, Asghar Pasha M, Siebel CW, Costello A, Haczku A, MacNamara K, Liang T, Zhu J, Bhandoola A, Maillard I, Yang Q. Cutting Edge: Notch Signaling Promotes the Plasticity of Group-2 Innate Lymphoid Cells. *Journal of immunology (Baltimore, Md : 1950)*. 2017; : 1601421–9. DOI: 10.4049/jimmunol.1601421
- Zook EC, Ramirez K, Guo X, van der Voort G, Sigvardsson M, Svensson EC, Fu YX, Kee BL. The ETS1 transcription factor is required for the development and cytokine-induced expansion of ILC2. *The Journal of experimental medicine*. 2016; 213:687–696. DOI: 10.1084/jem.20150851 [PubMed: 27069114]

Key Findings

- Innate cells make up a complex network that fights infection with helminths
- Innate cells are specifically located at mucosal and barrier sites to combat helminths
- Innate cells are highly adapted to respond rapidly and robustly to helminth infection
- Innate cells are producers of a variety of mediators, including cytokines, histamine and eicosanoids

Figure 1. Innate immune cells participate in a network that facilitates anti-helminth Type 2 immunity

Worms cause significant damage to the epithelium as they migrate and complete their lifecycle. This causes release of damage signals such as interleukin (IL)-25, IL-33 and thymic stromal lymphopoietin (TSLP) that promote the activation of innate cells present in the tissue. Activated DCs sense antigen (Ag) and migrate to the draining lymph node (dLN), where they release pro-Th2 mediators such as Type I interferon (IFN-I), CCL17 and Relma and induce polarization of naïve T cells to a T helper 2 (Th2) cell fate. Tissue-released damage signals also activate basophils, ILC2s and MCs that play distinct roles in the tissue. Basophils produce various inflammatory mediators such as chemokines and histamine and may interact with ILC2s. ILC2s are rich sources of Type 2 cytokines and wound repair-related molecules (not shown). Activated MCs release inflammatory mediators including prostaglandin D₂ (PGD₂), which activates and recruits ILC2s. Critically, activated basophils, ILC2s and MCs produce significant amounts of IL-4, IL-5 and IL-13 that can contribute to the polarization of Th2 cells in the dLN and may potentiate the activity of Th2 cells within the tissue. Finally, Type 2 cytokines from these innate cells and activated Th2 cells in the tissue recruit eosinophils, induce the alternative activation of macrophages (AAMΦ) that produce anti-helminth and pro-wound healing mediators such as Relma, and activate anti-helminth responses from the epithelium, such as goblet cell hyperplasia, mucus production, tuft cell activation and epithelial cell turnover.