CORRESPONDENCE

Spondylodiscitis: Diagnosis and Treatment Options

A Systematic Review

by Dr. med. Christian Herren, PD Dr. med. Dipl. chem. Norma Jung, Dr. med. Miguel Pishnamaz, Dr. med. Marianne Breuninger, PD Dr. med. Jan Siewe, and Prof. Dr. med. Rolf Sobottke in issue 51–52/2017

Questions upon Questions

As a primary care internist and a trauma surgery layperson, I still am left with numerous clarifying questions.

The authors mentioned (on page 876, at the bottom left) results from Sobottke et al., that spondylodiscitis occurs in 22.2% of patients after conservative treatment, and in 50.4% after invasive treatments, such as catheterization (urinary?) or fine-needle aspiration puncture (where was the puncture made?). However, if correct, this would signify that the absolute number of disease cases is exorbitantly high and not 30 per 250 000 per year, as initially mentioned in the summary (1).

These figures would be meaningful if the text should be inverted: that is, if 22.2% of patients with spondylodiscitis previously underwent conservative treatment, and 50.4%, invasive treatments (such as catheterization or puncture). Would this still be considered hematogenous, which according to the introduction should be the most common pathway of infection? Indeed, how do exogenous, endogenous, and hematogenous infections differ? Take as an example a case of spondylodiscitis following thyroid gland puncture. As a layperson, I would say that it is an exogenous form, as germs are introduced from the outside „exogenously“; however, as the germs are carried through the bloodstream, is it therefore „hematogenic“?

Furthermore, I would have liked a more detailed antibiotic recommendation with dosages, which is after all relevant to everyday life, especially in cases of intolerance or allergy to a previous medication and the antibiotic used as an alternative. DOI: 10.3238/arztebl.2018.0210a

References

1. Herren C, Jung N, Pishnamaz M, Breuninger M, Siewe J, Sobottke R: Spondylodiscitis: diagnosis and treatment options—a systematic review. *Dtsch Arztebl Int* 2017; 114: 875–82.

Dr. med. Cornel Certain

Konstanz, dr.cornel-certain@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank Dr. Certain for the critical discussion of our article (1). As Dr. Certain correctly stated, our wording was ambiguous, and it should mean that, according to

Sobottke et al., 22.2% of spondylodiscitis cases are preceded by spinal surgery, and more than half of cases of spondylodiscitis are associated with punctures or similar procedures (2).

Several ways in which pathogens can transmit progressive infections to the spine have been described. Hematogenous dissemination describes the way through the arterial system or after flow reversal in the Batson venous plexus, which increases pressure on the venous system. Here, the cause can be endogenous, such as an infection associated with endocarditis (and distant from the vertebrae), or it can be due to exogenous deposition, such as following surgery or puncture/injection. Thus, the thyroid puncture example mentioned by Dr. Certain would be considered as a hematogenous deposition. Lymphogenous spread and infection per continuitatem are rarer causes of spinal infection.

We agree with the objection of Dr. Certain: our article only presents an overview of antibiotic therapy. However, we would like to point out that, due to the space limitations, we deliberately provided an overview of the most common pathogens and their antibiotic therapies (see *Table 4* of the article). To address further questions and alternative therapy recommendations, we recommend referring to the current IDSA guideline (3). DOI: 10.3238/arztebl.2018.0210b

References

1. Herren C, Jung N, Pishnamaz M, Breuninger M, Siewe J, Sobottke R: Spondylodiscitis: diagnosis and treatment options—a systematic review. *Dtsch Arztebl Int* 2017; 114: 875–82.
2. Sobottke R, Rollingshoff M, Zarghooni K, et al.: Spondylodiscitis in the elderly patient: clinical mid-term results and quality of life. *Arch Orthop Trauma Surg* 2010; 130: 1083–91.
3. Berbari EF, Kanj SS, Kowalski TJ, et al.: Infectious Diseases Society of America (IDSA): clinical practice for the diagnosis and therapy of native vertebral osteomyelitis in adults. *Clin Infect Dis* 2015; 61: 26–46.

On behalf of the authors

Dr. med. Christian Herren

Department for Trauma and Reconstructive Surgery,
University Hospital RWTH Aachen, Germany
cherren@ukaachen.de

Conflict of interest statement

The author declares that no conflict of interest exists.