STUDY FUNDING

No targeted funding reported.

DISCLOSURES

The author reports no disclosures. Full disclosure form information provided by the author is available with the **full text of this article at Neurology.org/cp**

COMMENT: Bumping up Kernohan notch phenomenon

In 1929, Drs. Kernohan and Woltman described a phenomenon of hemiparesis that paradoxically localized to an ipsilateral brain tumor.¹ The mechanism involved uncus herniation from tumoral mass effect leading to brainstem displacement and compression of the contralateral cerebral peduncle against the tentorium cerebelli.

In this issue of *Neurology*[®] Clinical Practice, Dr. Chang² reports a case similar to the Kernohan notch phenomenon (KNP), but with a mechanism that is distinct from previously reported cases. Whereas classic KNP implied that the etiology is a supratentorial intraparenchymal mass or subdural hematoma leading to uncus herniation, in this case, the supratentorial injury is not exerting mass effect and a different mechanism is evoked. In this contusional KNP, it seems that the kinetic displacement of the brainstem against the tentorium directly led to midbrain edge compression injury. MRI reveals the presence of linear restricted diffusion in the contralateral midbrain. Hemorrhages in the ipsilateral splenium suggest a second site of crush injury that occurred from the kinetic central brain shift against the falx cerebri. Another less likely mechanism would be a transient uncus herniation from the supratentorial contusion that resolved by the time of imaging. More likely, uncus herniation is not a sine qua non for the diagnosis of KNP. Indeed, transtentorial herniation with a downward-lateral shift can produce the same type of injury.³ Similarly, in the absence of uncus herniation, ipsilateral mydriasis may not occur, and in fact, further brainstem shift can lead to compression of the contralateral third cranial nerve, causing contralateral mydriasis.³ Finally, ipsilateral hemiparesis similar to KNP can also occur in the absence of any type of herniation or contusion, e.g., due to cerebellar edema with brainstem displacement.⁴

KNP can thus be bumped up a notch in classification: A classic KNP, as initially described; a contusional KNP, with velocity-related recoil injury without evidence of herniation; and a modified KNP, with contralateral mydriasis.

1. Kernohan J. Incisura of the crus due to contralateral brain tumour. *Arch Neurol Psychiatry* 1929; 21:274.
2. Chang GY. Contusional Kernohan notch phenomenon. *Neurol Clin Pract* 2016;6:553–554.
3. Chung KH, Chandran KN. Paradoxical fixed dilatation of the contralateral pupil as a false-localizing sign in intraparenchymal frontal hemorrhage. *Clin Neurol Neurosurg* 2007;109: 455–457.
4. Kanis KB, Ropper AH, Adelman LS. Homolateral hemiparesis as an early sign of cerebellar mass effect. *Neurology* 1994;44:2194–2197.

Heustein Sy, MD and Gregory Kapinos, MD, MS

Departments of Neurosurgery and Neurology, Hofstra Northwell School of Medicine, Manhasset, NY.

Corresponding Author: Gregory Kapinos, MD, MS, North Shore University Hospital, Dept of Neurosurgery, 300 Community Drive, Tower, 9th Floor, Manhasset, NY 11030, Ph 516-562-3020, Fax: 516-562-3631, kapigreg@gmail.com

Author Contributions: H Sy: drafting/revising the manuscript. G Kapinos: drafting/revising the manuscript.

Study funding: No targeted funding reported.

Disclosures: H.L. Sy reports no disclosures. G. Kapinos is copyright owner of a patent for Advanced Neurologic Life Support training program, course, and manual. Go to Neurology.org/cp for full disclosures.