Published in final edited form as:

Glob Heart. 2017 June ; 12(2): 81–90. doi:10.1016/j.ghheart.2017.01.007.

Regional and sex differences in the prevalence and awareness of hypertension across six sites in sub-Saharan Africa: an H3Africa AWI-Gen study

F. Xavier Gómez-Olivé^{1,11,14}, Stuart A. Ali², Felix Made², Catherine Kyobutungi³, Engelbert Nonterah⁴, Lisa Micklesfield⁵, Marianne Alberts⁶, Romuald Boua⁸, Scott Hazelhurst^{2,9}, Cornelius Debpuur⁴, Felisatas Mashinya⁶, Sekgothe Dikotope⁶, Hermann Sorgho⁸, Ian Cook⁷, Stella Muthuri³, Cassandra Soo², Freedom Mukomana², Godfred Agongo⁴, Christopher Wandabwa³, Sulaimon Afolabi¹, Abraham Oduro⁴, Halidou Tinto⁸, Ryan G. Wagner¹, Tilahun Haregu³, Alisha Wade¹, Kathleen Kahn¹, Shane A. Norris⁵, Nigel J. Crowther¹⁰, Stephen Tollman¹, Osman Sankoh^{11,12,13}, Michèle Ramsay^{2,15,*}, and as members of AWI-Gen and the H3Africa Consortium

¹MRC/Wits Rural Public Health and Health Transitions Research Unit (Agincourt), School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

²Sydney Brenner Institute for Molecular Bioscience, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

³African Population and Health Research Center, Nairobi, Kenya

⁴Navrongo Health Research Centre, Navrongo, Ghana

⁵MRC/Wits Developmental Pathways for Health Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

⁶Department of Pathology and Medical Sciences, School of Health Care Sciences, Faculty of Health Sciences, University of Limpopo, Polokwane, South Africa

⁷Physical Activity Epidemiology Laboratory (EDST), School of Education, Faculty of Humanities, University of Limpopo, Polokwane, South Africa

⁸Institut de Recherche en Sciences de la Sante, Clinical Research Unit of Nanoro, Burkina Faso

⁹School of Electrical & Information Engineering, University of the Witwatersrand, Johannesburg, South Africa

*Corresponding Author: Michèle Ramsay, Sydney Brenner Institute for Molecular Bioscience, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, 9 Jubilee Road, Parktown, 2050, South Africa; michele.ramsay@wits.ac.za.

Conflict of interest statements for authors

The authors state that no conflicts of interest exist.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

¹⁰Department of Chemical Pathology, National Health Laboratory Service, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

¹¹INDEPTH Network, Accra, Ghana

¹²School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

¹³Njala University, Njala, Sierra Leone

¹⁴Harvard Center for Population and Development Studies, Harvard T.H. Chan School of Public Health, Harvard University, Cambridge, MA

¹⁵Division of Human Genetics, National Health Laboratory Service and School of Pathology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

Abstract

Background—There is a high prevalence of hypertension and related cardiovascular diseases in sub-Saharan Africa yet few large studies exploring hypertension in Africa are available. The actual burden of disease is poorly understood and awareness and treatment to control it is often suboptimal.

Objectives—To report the prevalence of measured hypertension and to assess awareness and control of blood pressure among older adults in rural and urban settings in six sites located in west, east and southern Africa. In addition, we examined regional, sex, and age differences related to hypertension.

Methods—A population-based cross-sectional study was performed at six sites in four African countries - Burkina Faso (Nanoro), Ghana (Navrongo), Kenya (Nairobi) and South Africa (Agincourt, Dikgale, Soweto). Blood pressure measurements were taken using standardized procedures on 10,696 adults aged 40 to 60 years. Hypertension was defined as SBP ≥ 140mm Hg and/or DBP ≥ 90 mm Hg or taking anti-hypertensive medication.

Results—The mean prevalence of hypertension ranged from 15.1% in Nanoro to 54.1% in Soweto. All three of the South African sites had a mean prevalence of hypertension of over 40.0%, significantly higher than Nairobi (25.6%) and Navrongo (24.5%). Prevalence increased with age in both sexes and at all sites. A significantly higher prevalence of hypertension was observed in women in Agincourt, Dikgale and Nairobi, while in Nanoro this trend was reversed. Within the hypertensive group the average proportion of participants who were aware of their blood pressure status was only 39.4% for men and 53.8% for women, and varied widely across sites.

Conclusions—Our study demonstrates that the prevalence of hypertension and the level of disease awareness differ not only between but also within sub-Saharan African countries. Each nation must tailor their regional hypertension awareness and screening programs to match the characteristics of their local populations.

Keywords

Hypertension; awareness; controlled; AWI-Gen; H3Africa; cardiovascular disease; cardiometabolic disease

INTRODUCTION

Hypertension is a rising global health problem with an estimated 1 billion people living with hypertension, and 9.4 million related annual deaths worldwide (1, 2). Low and Middle Income Countries (LMIC) suffer two thirds of the global burden of cardiovascular disease (CVD), which is associated with inadequate treatment of hypertension (3). In sub-Saharan Africa (SSA) hypertension has become a major public health problem (4, 5). At present, cardiovascular diseases are considered to be a significant contributor to premature death in this region, with a high prevalence of hypertension found in many SSA settings. The World Health Organization Study on Global Ageing and Adult Health (WHO-SAGE) (6), which compared nationally representative populations over 50 years of age from China, Russia, India, Ghana, South Africa and Mexico, reported that the prevalence of hypertension in South Africa was amongst the highest at over 77%. At the same time, the study reported low levels of awareness and blood pressure control in all of these countries (6). In the absence an effective intervention strategy to control this epidemic, it is expected that hypertension-associated CVD will continue to rise, with a concomitant increase in mortality and disability (7, 8).

Despite global concerns over hypertension, there seems to be a dearth of information for most African countries on its prevalence, the levels of awareness by individuals on their blood pressure status, and how effective treatment is (9). This is especially so in the population age group 40 to 60 years, which is the age at which individuals suffer the highest levels of blood-pressure related disease (10, 11).

The principal aims of the current study were therefore to measure the prevalence, level of awareness and control of hypertension in a large population of 40–60 year-old participants of both sexes in six research sites across four SSA countries.

METHODS

This study draws upon the Africa Wits-INDEPTH (University of the Witwatersrand, Johannesburg (Wits), and the International Network for the Demographic Evaluation of Populations and Their Health (INDEPTH)) (12) partnership for Genomic Studies (AWI-Gen), which is an NIH funded Collaborative Centre of the Human Heredity and Health in Africa (H3Africa) Consortium (13, 14). The primary aim of AWI-Gen is to study genetic and environmental factors that contribute to cardiometabolic disease in African populations. Six study sites in four SSA countries are involved in AWI-Gen: in South Africa the MRC/Wits Agincourt Health and Demographic Surveillance System Site (HDSS) (15), Dikgale HDSS (16) and the Soweto cohort located within the MRC/Wits Developmental Pathways for Health Research Unit (DPHRU) (17), in Kenya the Nairobi HDSS (18), in Ghana the Navrongo HDSS (19) and in Burkina Faso the Nanoro HDSS (20).

Study Design and Participant Recruitment

This was a cross-sectional population study to serve as a baseline time point for adults aged between 40 and 60 years, who were recruited into the study between August 2013 and August 2016.

Participants who were recruited into the AWI-Gen study (21) were residents at six sites that span the African continent. In West Africa, these were the rural areas of Nanoro and Navrongo; in East Africa an urban informal settlement in Nairobi; and in South Africa, the rural areas of Agincourt in the province of Mpumalanga and Dikgale in the province of Limpopo, and the suburban township of Soweto that lies on the outskirts of Johannesburg.

Pregnant women, first degree relatives of existing participants, and individuals with physical impairments preventing measurement of blood pressure, were excluded from the study. A resident was defined according to the HDSS's criteria (staying within the area for at least three months and expecting to stay more than three months after the census).

Informed Consent, and Anonymization of Participant Identity

All participants provided written informed consent before any study procedures were conducted. Each participant was assigned a random AWI-Gen study participant identity code which was used on study documentation, and the key linking participant identity and their identity code locked securely away at the research site, and not entered onto any further study documentation.

Measurement of Blood Pressure

Blood pressure was measured using a sphygmomanometer (Omron M6, Omron, Kyoto, Japan), according to the following standardized procedure: participants were seated upright in a chair, with feet resting firmly on the floor and not crossed. To ensure that the antecubital fossa was at the level of the heart participants were seated with the left arm resting on a desk or arm rest. The cuff was placed on the arm about 2–3 cm above the antecubital fossa, and not restricted by clothing. The participant was allowed to relax for 3–5 minutes before blood pressure was measured. Three measurements were taken, at two minute intervals. During this period, participants remained quiet and still. The mean of the last two measurements was used to calculate the final systolic blood pressure (SBP) and diastolic blood pressure (DBP).

Diagnosis of Hypertension

Blood pressure measurements were classified according to the JNC7 guidelines of the Joint National Committee of Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (22). Hypertension was defined as presenting one or more of the following conditions: mean SBP $\geq$ 140mm Hg, mean DBP $\geq$ 90 mm Hg, or the participant reported that they were currently taking medication for hypertension.

Data Collection and Quality Control

A paper questionnaire was administered to participants as part of a wider survey by AWI-Gen. This questionnaire asked for demographic information about the participant, as well as historical information about their blood pressure, previous diagnosis of hypertension, and how it was treated. Blood pressure data were also recorded on this questionnaire. These data were then entered into a REDCap (Research Electronic Data Capture) computerized database (23, 24). For quality control purposes, 10% of all entries were checked for consistency between the paper form and the electronic versions. Anonymized data were then

transferred through a secure File Transfer Protocol (FTP) connection to the central REDCap server at the University of the Witwatersrand, where a second process searched for outliers and missing data. Sites were requested to refer back to the paper forms for reconciliation.

Calculation of the Awareness of Hypertension and Statistical Analyses

Awareness of hypertension was calculated using the question “Have you ever been told by a doctor, nurse, or other healthcare worker that you have hypertension (high blood pressure)?”. The response to this question was used to quantify self-reported hypertension and these data in conjunction with the gold standard of measured blood pressure were used to calculate sensitivity, specificity, positive predictive value (PPV) and negative predictive value (NPV) of the self-reported disease status.

Normally distributed continuous variables were described using means and standard deviations, and non-normally distributed variables were described using medians and interquartile ranges. Proportions were used to describe categorical data. Descriptive analyses were performed using proportions, means, and confidence intervals and data were compared across populations using the χ^2 test. Statistical analyses were performed using the statistics program STATA, version 13 (StataCorp, College Station, Texas, USA).

Research Ethics Board Approval

This study was approved by the Human Research Ethics Committee (Medical) of the University of the Witwatersrand (Protocol Number: M121029). In addition, each study site obtained ethics approval from their respective institutional and national ethics committees as required. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution’s human research committee.

RESULTS

Participants

Data were collected from 10 696 individuals between the ages of 40 and 60 years from the six research sites. Table 1 shows the number of men and women, and the age distributions for the entire cohort and according to site. The mean age of the total cohort was 49.9 years (SD 0.06) with no significant differences noted for age between the sites. Four of the sites (Soweto, Nairobi, Navrongo and Nanoro) each contributed 19% of the participants to the final sample, while Agincourt (14%) and Dikgale (10%) had fewer participants in this age range. Of the total participants, 63 did not have a full set of study data in the REDCap database (0.58% of the total), and were excluded from the data analysis.

Prevalence

The prevalence of hypertension in the entire cohort was 33.3% (see Table 2), being significantly higher in women (35%; 95% CI 33.7–36.2) than in men (31%; 95% CI 30.0–32.6). Given this difference, further analyses were stratified by sex. Table 2 shows a breakdown of the prevalence of hypertension according to the JNC7 definition stratified for sex and site. The highest prevalence was observed in the South African sites (ranging from 41.6% to 54.1%) and the lowest in Burkina Faso (15%). Sex stratified prevalence showed

that women had significantly higher rates of hypertension in Agincourt, Dikgale and Nairobi, whereas men had significantly higher rates in Nanoro. Navrongo and Soweto had no significant sex differences in hypertension. Hypertension significantly increased with age, both for the entire population, and for men and women separately (Table 2).

Awareness and Diagnosis

Of the 3543 individuals who were hypertensive, 51 did not answer the question on self-reported hypertension status, so awareness was reported on 3492 individuals. Among those hypertensive participants that answered the question about awareness of hypertension, only 1667 (47.7%) were aware of their status. By sex, a significantly higher percentage of women knew their status compared to men (53.8% vs 39.4% $p < 0.001$; Figure 1.A and 1.B).

Table 3 presents correctness of self-knowledge of being hypertensive with respect to sensitivity (how many of the hypertensive cases knew they were hypertensive) and specificity (what proportion of those who thought they were hypertensive actually were, i.e. controlled on medication or hypertensive at the time of measurement). We observe that across the entire cohort, specificity was very high for both men (97.9%) and women (91.2%). However, sensitivity was low in both men (39.4%) and women (53.8%). Comparison by site indicated higher levels of awareness in Agincourt (63.1%), Dikgale (60.8%) and Nairobi (60.0%), while levels of awareness were much lower in Nanoro (39.9%), Soweto (37.2%) and Navrongo (38.6%). There were also significant differences in awareness between men and women (Figure 3). In addition, Navrongo (21.4%) and Soweto (39.4%) had the lowest sensitivity among men, while Navrongo (34.6%), Soweto (35.1%) and Nanoro (37.0%) had the lowest sensitivity among women. Specificity was high ($>85.0\%$) in all sites for both sexes, with the exception being women in Soweto (50.8%).

Blood Pressure Control

To examine how well hypertension was being managed (Figure 1), we found that of the women that were aware of their hypertensive status, 87% were being treated. Of those, only 51% had controlled blood pressure. For men, 78% of those who were aware of their hypertensive condition were on treatment, but only 39% were controlled. When measuring how well hypertensive blood pressure was controlled by site, we observed that the rates for women (as a proportion of those on treatment) with well controlled blood pressure levels was 61% in Nanoro, 57% in Dikgale, 50% in Nairobi, 49% in Agincourt, 45% in Soweto and 40% in Navrongo (Figure 1.B). The rate of controlled blood pressure in men at these sites was 61% in Dikgale, 54% in Nanoro, 38% in Navrongo, 36% in Agincourt, 32% in Soweto and 30% in Nairobi (Figure 1.A).

Finally, hypertension prevalence (Figure 2.A), awareness and control data by sex and by sites (Figure 2.B) are shown, and demonstrate a high prevalence of hypertension in the South African sites compared to the general low levels of awareness and control in all sites.

DISCUSSION

In this study we found differences in the prevalence, awareness, and blood pressure control of hypertension at the study sites in west, east, and southern Africa. Our aim was to

document these differences and to more fully understand how hypertension affects SSA countries and could inform the need to customize awareness, treatment and prevention approaches according to the needs of each individual region and country. Such information is essential to the design of effective interventions that could impact the downstream consequences of the rising rates of hypertension, most particularly cardiovascular diseases (25).

Overall, only 39.4% of men and 53.8% of women who were hypertensive knew their status (Table 3); an observation that is common in many studies of hypertension in SSA (26). In all sites, except Nanoro and Soweto, women were more aware of their condition, compared to men. In terms of blood pressure control, we note that among all hypertensive women, 22% have controlled blood pressure compared to only 12% of men. This indicates that only a fifth (18%) of those found to be hypertensive in this study have controlled blood pressure; this is striking, but not surprising given the low levels of awareness. In contrast, three out of four participants who were aware of their status received treatment, which could indicate a high level of engagement with primary health care providers. However, that only half of these participants had controlled blood pressure is a concern. Similar results were found in other studies in Africa. Comparing our data to the SAGE study (6) where women were more aware of their status than men (54% vs 42%), women had significantly higher levels of controlled blood pressure than men (11% vs 9%) although the levels of control and the difference between sexes were smaller than in the AWI-Gen study. These data are congruent with a review on hypertension in sub-Saharan Africa which found levels of awareness under 40% and levels of controlled blood pressure under 20% (4). Interestingly, a study in Bangladesh and India in 2000 showed a hypertension awareness of 45% with controlled levels of 10% (27) suggesting a higher level of control in sub-Saharan Africa than in Bangladesh and India.

West Africa: Burkina Faso and Ghana

Across all study sites, we observed the lowest prevalence of hypertension among men (19.2%) and women (11.0%) in rural Nanoro, Burkina Faso. Interestingly, our data show that the other west African site in rural Navrongo (Ghana), approximately 250 km from Nanoro, has significantly higher rates among men (24.1%), and more than double the prevalence among women (24.8%).

In 2013, the STEPS study conducted in Burkina Faso reported an overall prevalence of hypertension of 17.6% (95% CI, 15.7–19.4), with a prevalence of 14.8% (95% CI, 13.1–16.5) in the rural areas and a prevalence of 14.2% (95% CI, 9.5–19.0) in the Centre-West region where Nanoro is based (28), which is similar to the prevalence of 15.1% (95% CI, 13.5–16.6) reported in the AWI-Gen study. The prevalence of hypertension in Nanoro is also similar to that reported in informal settings in Ouagadougou (29) but higher than the 9.4% observed in the semi-urban area of the Kaya Health and Demographic Surveillance (HDSS) site in the Center-North region (30). In this study, Nanoro is the only site to display a significant sex disparity in favor of men having hypertension, in line with the observation in the STEPS study with prevalence of 19.4% for men and 16.0% for women (28).

The Nanoro cohort had an overall awareness of hypertension of 41.4% for men and 30.7% for women, which although still low, is the highest awareness level reported recently in the country compared to the STEPS survey (31.8%), the Ouagadougou HDSS (27.4%) and the Kaya HDSS (26.8%). The levels of controlled blood pressure in those on treatment were 53.6% for men, and 61.1% for women, and were among the highest reported in the country, but when represented as the percentage of controlled blood pressure among all hypertensives, only 18.5 and 18.8% of men and women respectively are controlled. While awareness was low, treatment levels for those aware of hypertension (86.2% for men, 92.3% for women) was higher than had been recorded by previous studies (31–33).

In Ghana, over the last three to four decades, several population-based studies have examined the epidemiology of hypertension (33), but only one was conducted in the North where Navrongo is situated (34). This study, which examined 207 men and 367 women, reported the prevalence of high blood pressure at 19.3%. The AWI-Gen study is the first large-scale population-based study to examine prevalence and awareness of hypertension among older adults in rural northern Ghana. Overall, 24.1% of men and 24.8% of women participants from Navrongo were hypertensive. The rate is similar to previous studies in rural southern Ghana with a prevalence of 24.1% (33) but is an appreciable increase in hypertension prevalence from 19.3% observed in 2008 in the same study area (34). Our study demonstrates a significant change in measured hypertension with increasing age, which is consistent with other population-based surveys conducted in Ghana (32, 35).

Examination of the levels of awareness of hypertension in Ghana reveals that it has not changed significantly compared to previous studies in rural areas. Between 1979 and 2008, seven studies examined awareness of hypertension in Ghana, four of them were conducted in rural communities and three in urban communities. Awareness ranged from 22% to 54%, with minimal sex differences, and rural areas being lower than urban (33). Other studies reported awareness levels of between 25.4% and 34% for rural communities in southern Ghana (31, 32). That this study also showed low levels of awareness (21.4% for men and 34.6% for women) which indicates a continued need to intensify public education around regular blood pressure monitoring. An explanation for these low levels of awareness could be the high level of illiteracy in this Ghanaian rural area. The health seeking behavior of inhabitants in the study area is also a likely contributor since a considerable number of people still prefer to seek healthcare from traditional healers. It also reflects a limited health awareness and promotion program on non-communicable diseases (NCDs), and especially in a community that has disproportionate burden of communicable diseases (especially malaria) over NCDs.

In Navrongo, the majority of those who are aware of their hypertensive status are on treatment, and this is higher among women (90.0%) than men (80.0%). This high rate of treatment might be attributed to general improvements in access to healthcare in the study area. The introduction of the national health insurance scheme in Ghana in 2003 has also lead to significant improvements in access to medications for chronic diseases such as hypertension (36).

While the greater proportion of participants with hypertension are being treated, it is important to examine whether treatment is effective or not. This study indicated that only 4 out of 10 hypertensives on treatment were controlled (37.8% for men, 39.8% for women). Expressed as a percentage of total hypertensives in Navrongo, this means that only 1 out of 10 of the total hypertensive population in Navrongo is being adequately managed. This raises serious issues regarding the effectiveness of treatment as well as the extent to which people are following their treatment regimen. Similar studies in 2006 also found low rates of control among rural communities in Ghana (32), which documented 24% on treatment, yet just 6.2% controlled. Addo et al. (31) also documented a control rate of 16.7% in four rural areas in the Ga district of Ghana. These observations are consistent with a systematic review that showed a treatment level ranging from 7% to 31% with a controlled range of 4% to 13% (33).

West Africa therefore shows a similar trend, with significant heterogeneity in the prevalence, treatment and awareness of hypertension in two rural areas just about 250 km apart and may reflect general health awareness and approaches to healthcare.

East Africa: Nairobi

In a Nairobi informal settlement in east Africa we observed an overall prevalence of hypertension of 21.9% for men and 28.7% for women, which is similar to the national STEPS survey conducted in Kenya in 2015 (24%) (37), which included adults aged 18–69 years. However, looking only at the 45–59 year-old adults, the STEPS study reports hypertension at 38.7% in men and 48.5% in women. These rates of prevalence are much higher than those we found in the AWI-Gen study among individuals of a similar age group. This could be due to the fact that the AWI-Gen study was conducted among participants under a different socioeconomic and health profile to the rest of the population. In a larger population-based study conducted in 2008/09, of 5190 adults 18+ years in the same study area, the prevalence of hypertension was 12.3% (12.7% in women and 12.0% in men) (38). A baseline survey done in 2012 among all adults 35+ years in the same study area as part of an intervention (39) to reduce hypertension found a prevalence of 24% (40) which is similar to the findings of this study. In a study conducted in a different urban informal settlement - Kibera - in 2010, among 2200 adults, the crude prevalence of hypertension was 12.6% (41). In this study, it was also reported that prevalence increased with age and was significantly higher among females in the age strata of 35–44 years. These age differences by sex in the prevalence of hypertension could be associated with higher magnitudes of obesity and physical inactivity among the study population as reported by previous studies. Thus, a large population-based study conducted in 2008/09 in the same study area showed a prevalence of hypertension of 24.5% among women aged 40–49 years and 36.4% among women 50–59 years. For men the observed prevalence was 13.2% for those aged 40–49 and 27.9% among men aged 50–59 years (38).

At the Nairobi site, 41.1% of men and 71.8% of women with hypertension correctly reported their blood pressure status. The high level of awareness may be a result of several screening programs, health campaigns and intervention studies conducted in the study area since 2008 (42, 43).

Treatment levels were higher in women (87.7%) than men (74.1%) which is possibly related to higher levels of awareness among women. As age increased, both the percentage of hypertensive patients on treatment and those not on treatment increased. This is due to an increasing overall prevalence of hypertension. The Kenyan STEPS survey reported that only 2.7% of men and 6.9% of women with hypertension were receiving treatment. More specifically, 8.3% of hypertensive patients between the ages of 45–59 years were receiving treatment. The higher proportion found in the AWI-Gen study compared to the STEPS survey may be due to several programs to increase access to hypertension diagnostic and treatment services in the study area (42, 43).

A study describing the profile of people with hypertension in Nairobi's informal settlement has shown that this population has a high prevalence of overweight (30%) and obesity (21%) as well as behavioral risk factors for hypertension such as smoking (8%), high alcohol intake (8%) and a low vegetable and fruit intake (68%) (42). Besides, three quarters of hypertensive patients in Nairobi informal settlements do not take anti-hypertensive medication, and even the ones who do show little adherence (42).

South Africa

The prevalence of hypertension was highest in South Africa, but we did not observe a marked urban-rural difference, especially among women. In urban Soweto, 53.4% of women were hypertensive, and in rural Agincourt 52.1%. In Dikgale, another rural area about 300km northwest of Agincourt, the prevalence among women was 47.2%. However, men showed a significantly higher prevalence in Soweto (54.7%) compared to Agincourt (39.9%) and Dikgale (29.2%).

The National SAGE study on populations 50 years and over showed high levels of self-reported hypertension (75.3%) in South Africa (44), which is the highest prevalence of hypertension compared to other SAGE national studies in various low and middle income countries (LMIC) (6). In the AWI-Gen study, the three South African sites also had the highest proportion of hypertensives among 40 to 60 year olds across all sites, with 54% in Soweto, 47% in Agincourt and 42% in Dikgale. These percentages are considerably higher than in recent data from SANHANES (45) which reported a prevalence of 19.5% in the 45–54 years age group. A similar prevalence (19.8%) was found in the Modeling the Epidemiological Transition Study (METS) (46). Longitudinal data from the PURE study (47) found that 47.7% of black South Africans 30 years and older were classified as hypertensive, a prevalence similar to that reported here. In another study, the burden of hypertension was measured in an urban black population in Cape Town (CRIBSA) with the finding that age-standardized prevalence of hypertension of adults 25 to 64 years old was 38.9%, prevalence in 45–54 year old men was 58.2% and in women was 60.6% (48).

The CRIBSA study also found that among the participants with hypertension, significantly more women than men were detected, treated and controlled on treatment, with the proportion detected and treated increasing in the women but not the men between 1990 and 2008/9. In our study, the Agincourt site had the higher levels of awareness; however, Dikgale presented the highest levels of treatment and controlled blood pressure, with Soweto presenting low levels of awareness and controlled blood pressure. Given the urban context of

Soweto, with good accessibility to health services, this is a concern, indicating the need for greater communication around hypertension and screening in this population. It is important to further explore the socio-demographic determinants associated with these findings, and whether primary healthcare programs in neighboring provinces could help explain the variations observed between sites in South Africa.

CONCLUSION

Hypertension is a rising global health problem and it is affecting most African regions. Although related diseases such as stroke, heart failure, kidney failure and others are associated with higher mortality, there is still a low level of awareness of hypertension among African populations. Low awareness has a direct impact on the levels of the hypertensive population that receive the necessary treatment and achieve well controlled blood pressure levels. Controlled blood pressure in the majority of individuals has the potential to significantly lower the risk of several chronic NCDs in SSA.

Implications for public health

This study shows that hypertension is a critical health problem in different regions and settings in Africa. Our data show stark sex-specific and region-specific differences that will require further detailed understanding to inform effective intervention strategies. Moreover, given the low levels of awareness of hypertension shown in this study, and the related consequences of low levels of awareness in the control of hypertension, we believe that improvements in health promotion and system strengthening could help improve awareness, treatment and control of hypertension among African rural and urban populations. It is crucial to plan and implement more active campaigns aiming to increase awareness and diagnosis of hypertension. However, this alone would not be enough, as we observed that only half of those who are aware and who are on treatment have controlled blood pressure levels. This speaks to the need for programmatic systems to ensure effective adherence and follow up, in particular outreach and community based approaches. With increasing life expectancy due both to the roll out of antiretroviral treatment to all individuals with HIV, and general improvements in healthcare, it is expected that the prevalence of hypertension will continue to increase. Consequently, NCDs including hypertension, need to be prioritized and managed to reduce the public health burden.

Further Research Questions

There are several areas for future research. First, it is clear that a series of complex health transitions are occurring in Africa, and as such, this underlines the importance of a better understanding determinants of change such as social, biological and healthcare factors as this will influence policy responses. Second, in a changing society, it will become increasingly important to study the relationship between hypertension and associated risks, diseases, comorbidities, and mortality, in large SSA cohorts. Third with an improved understanding of the underlying changes, what would the effects of changes in lifestyle be? Will it be possible to reverse the incoming tide of hypertension? And finally, to what extent might the susceptibility of the population to hypertension be due to genetics: Are there specific gene-environment interactions that may lead to increases in hypertension related

diseases in Africa? These, are just some of the questions that the H3Africa AWI-Gen study aims to address.

Acknowledgments

The AWI-Gen Collaborative Centre is funded by the NIH (1U54HG006938) as part of the H3Africa Consortium. MR is a South African Research Chair in Genomics and Bioinformatics of African Populations hosted by the University of the Witwatersrand, funded by the Department of Science and Technology and administered by National Research Foundation of South Africa (NRF). OS is Executive Director of INDEPTH. The INDEPTH Network receives core support funding from the William and Flora Hewlett Foundation, Sida/Research Cooperation and Wellcome Trust. This paper describes the views of the authors and does not necessarily represent the official views of the National Institutes of Health or the NRF. The Agincourt HDSS received core support from the University of the Witwatersrand and the Medical Research Council, South Africa, and the Wellcome Trust, UK (Grants 058893/Z/99/A; 069683/Z/02/Z; 085477/Z/08/Z; 085477/B/08/Z). The Nairobi HDSS receives core support from Sida and the William and Flora Hewlett Foundation. The Soweto Cohort (DPHRU) is supported by the University of the Witwatersrand, the Medical Research Council, South Africa, DST-NRF Centre for Excellence in Human Development, and The Wellcome Trust. The Dikgale HDSS has received funding from the Norwegian Universities Committee for Development Research and Education (NUFU); the Free University of Amsterdam; the INDEPTH Network; the Flemish Inter-University Council (VLIR), Belgium; the National Research Foundation, South Africa; and the University of Limpopo.

References

1. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012; 380(9859):2224–60. [PubMed: 23245609]
2. Poulter NR, Prabhakaran D, Caulfield M. Hypertension. *Lancet*. 2015; 386(9995):801–12. [PubMed: 25832858]
3. Perkovic V, Huxley R, Wu Y, Prabhakaran D, MacMahon S. The burden of blood pressure-related disease: a neglected priority for global health. *Hypertension*. 2007; 50(6):991–7. [PubMed: 17954719]
4. Addo J, Smeeth L, Leon DA. Hypertension in sub-saharan Africa: a systematic review. *Hypertension*. 2007; 50(6):1012–8. [PubMed: 17954720]
5. Ibrahim MM, Damasceno A. Hypertension in developing countries. *Lancet*. 2012; 380(9841):611–9. [PubMed: 22883510]
6. Lloyd-Sherlock P, Beard J, Minicuci N, Ebrahim S, Chatterji S. Hypertension among older adults in low- and middle-income countries: prevalence, awareness and control. *International journal of epidemiology*. 2014; 43(1):116–28. [PubMed: 24505082]
7. Mendis S. Challenges for the management of hypertension in low-resource settings. *Ethn Dis*. 2003; 13(2 Suppl 2):S67–70. [PubMed: 13677416]
8. Juma PA, Mohamed SF, Wisdom J, Kyobutungi C, Oti S. Analysis of Non-communicable disease prevention policies in five Sub-Saharan African countries: Study protocol. *Arch Public Health*. 2016; 74:25. [PubMed: 27335640]
9. Pereira M, Lunet N, Azevedo A, Barros H. Differences in prevalence, awareness, treatment and control of hypertension between developing and developed countries. *Journal of hypertension*. 2009; 27(5):963–75. [PubMed: 19402221]
10. Lawes CM, Vander Hoorn S, Law MR, Elliott P, MacMahon S, Rodgers A. Blood pressure and the global burden of disease 2000. Part II: estimates of attributable burden. *Journal of hypertension*. 2006; 24(3):423–30. [PubMed: 16467640]
11. Lawes CM, Vander Hoorn S, Law MR, Elliott P, MacMahon S, Rodgers A. Blood pressure and the global burden of disease 2000. Part I: estimates of blood pressure levels. *Journal of hypertension*. 2006; 24(3):413–22. [PubMed: 16467639]
12. Sankoh O, Byass P. The INDEPTH Network: filling vital gaps in global epidemiology. *International journal of epidemiology*. 2012; 41(3):579–88. [PubMed: 22798690]

13. Rotimi C, Abayomi A, Abimiku A, Adabayeri VM, Adebamowo C, et al. H3Africa Consortium. Research capacity. Enabling the genomic revolution in Africa. *Science*. 2014; 344(6190):1346–8. [PubMed: 24948725]
14. Ramsay M, Sankoh O. as members of the AWI-Gen Studys, the H3Africa Consortium. African partnerships through the H3Africa Consortium bring a genomic dimension to longitudinal population studies on the continent. *International journal of epidemiology*. 2016; 45(2):305–8. [PubMed: 26659658]
15. Kahn K, Collinson MA, Gomez-Olive FX, Mokoena O, Twine R, Mee P, et al. Profile: Agincourt health and socio-demographic surveillance system. *International journal of epidemiology*. 2012; 41(4):988–1001. [PubMed: 22933647]
16. Alberts M, Dikotope SA, Choma SR, Masemola ML, Modjadji SE, Mashinya F, et al. Health & Demographic Surveillance System Profile: The Dikgale Health and Demographic Surveillance System. *International journal of epidemiology*. 2015; 44(5):1565–71. [PubMed: 26275454]
17. Richter L, Norris S, Pettifor J, Yach D, Cameron N. Cohort Profile: Mandela's children: the 1990 Birth to Twenty study in South Africa. *International journal of epidemiology*. 2007; 36(3):504–11. [PubMed: 17355979]
18. Beguy D, Elung'ata P, Mberu B, Oduor C, Wamukoya M, Nganyi B, et al. Health & Demographic Surveillance System Profile: The Nairobi Urban Health and Demographic Surveillance System (NUHDSS). *International journal of epidemiology*. 2015; 44(2):462–71. [PubMed: 25596586]
19. Oduro AR, Wak G, Azongo D, Debpuur C, Wontuo P, Kondayire F, et al. Profile of the Navrongo Health and Demographic Surveillance System. *International journal of epidemiology*. 2012; 41(4): 968–76. [PubMed: 22933645]
20. Derra K, Rouamba E, Kazienga A, Ouedraogo S, Tahita MC, Sorgho H, et al. Profile: Nanoro Health and Demographic Surveillance System. *International journal of epidemiology*. 2012; 41(5): 1293–301. [PubMed: 23045201]
21. Ramsay M, Crowther N, Tambo E, Agongo G, Baloyi V, Dikotope S, Gómez-Olivé X, Jaff N, Sorgho H, Wagner R, Khayeka-Wandabwa C, Choudhury A, Hazelhurst S, Kahn K, Lombard Z, Mukomana F, Soo C, Soodyall H, Wade A, Afolabi S, Agorinya I, Amenga-Etego L, Ali SA, Bognini JD, Boua RP, Debpuur C, Diallo S, Fato E, Kazienga A, Konkobo SZ, Kouraogo PM, Mashinya F, Micklesfield L, Nakanabo-Diallo S, Njamwea B, Nonterah E, Ouedraogo S, Pillay V, Somande AM, Tindana P, Twine R, Alberts M, Kyobutungi C, Norris SA, Oduro AR, Tinto H, Tollman S, Sankoh O. as members of AWI-Gen and the H3Africa Consortium. H3Africa AWI-Gen Collaborative Centre: a resource to study the interplay between genomic and environmental risk factors for cardiometabolic diseases in four sub-Saharan African countries. *Global Health and Epidemiology and Genomics*. 2016; 1e20:1–13.
22. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003; 289(19):2560–72. [PubMed: 12748199]
23. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009; 42(2):377–81. [PubMed: 18929686]
24. Klipin M, Mare I, Hazelhurst S, Kramer B. The process of installing REDCap, a web based database supporting biomedical research: the first year. *Appl Clin Inform*. 2014; 5(4):916–29. [PubMed: 25589907]
25. Nulu S, Aronow WS, Frishman WH. Hypertension in Sub-Saharan Africa: A Contextual View of Patterns of Disease, Best Management, and Systems Issues. *Cardiol Rev*. 2016; 24(1):30–40. [PubMed: 26284525]
26. Ataklte F, Erqou S, Kaptoge S, Taye B, Echouffo-Tcheugui JB, Kengne AP. Burden of undiagnosed hypertension in sub-saharan Africa: a systematic review and meta-analysis. *Hypertension*. 2015; 65(2):291–8. [PubMed: 25385758]
27. Hypertension Study G. Prevalence, awareness, treatment and control of hypertension among the elderly in Bangladesh and India: a multicentre study. *Bull World Health Organ*. 2001; 79(6):490–500. [PubMed: 11436469]
28. STEPS. Organisation mondiale de la Santé. 2013. Rapport de l'enquête nationale sur la prevalence des principaux facteurs de risques communs aux maladies non transmissible au Burkina Faso.

Enquête STEPS 2013. Ministère de la Santé. Communauté économique des États de l'Afrique de l'Ouest.

29. Doulougou B, Kouanda S, Rossier C, Soura A, Zunzunegui MV. Differences in hypertension between informal and formal areas of Ouagadougou, a sub-Saharan African city. *BMC Public Health*. 2014; 14:893. [PubMed: 25175061]
30. Doulougou B, Kouanda S, Ouedraogo GH, Meda BI, Bado A, Zunzunegui MV. Awareness, treatment, control of hypertension and utilization of health care services following screening in the North-central region of Burkina Faso. *Pan Afr Med J*. 2014; 19:259. [PubMed: 25852802]
31. Addo J, Amoah AG, Koram KA. The changing patterns of hypertension in Ghana: a study of four rural communities in the Ga District. *Ethn Dis*. 2006; 16(4):894–9. [PubMed: 17061743]
32. Agyemang C. Rural and urban differences in blood pressure and hypertension in Ghana, West Africa. *Public Health*. 2006; 120(6):525–33. [PubMed: 16684547]
33. Bosu WK. Epidemic of hypertension in Ghana: a systematic review. *BMC Public Health*. 2010; 10:418. [PubMed: 20626917]
34. Kunutsor S, Powles J. Descriptive epidemiology of blood pressure in a rural adult population in Northern Ghana. *Rural Remote Health*. 2009; 9(2):1095. [PubMed: 19508111]
35. Addo J, Agyemang C, Smeeth L, de-Graft Aikins A, Edusei AK, Ogedegbe O. A review of population-based studies on hypertension in Ghana. *Ghana Med J*. 2012; 46(2 Suppl):4–11. [PubMed: 23661811]
36. Blanchet NJ, Fink G, Osei-Akoto I. The effect of Ghana's National Health Insurance Scheme on health care utilisation. *Ghana Med J*. 2012; 46(2):76–84. [PubMed: 22942455]
37. WHO. Kenya STEPwise Survey for Non-Communicable Diseases Risk Factors 2015 Report. Ministry of Health (Division of Non-Communicable Diseases), Kenya National Bureau of Statistics, World Health Organization; 2015.
38. van de Vijver SJ, Oti SO, Agyemang C, Gomez GB, Kyobutungi C. Prevalence, awareness, treatment and control of hypertension among slum dwellers in Nairobi, Kenya. *Journal of hypertension*. 2013; 31(5):1018–24. [PubMed: 23425703]
39. Oti SO, van de Vijver S, Gomez GB, Agyemang C, Egondi T, Kyobutungi C, et al. Outcomes and costs of implementing a community-based intervention for hypertension in an urban slum in Kenya. *Bull World Health Organ*. 2016; 94(7):501–9. [PubMed: 27429489]
40. Oti SO, van de Vijver SJ, Kyobutungi C, Gomez GB, Agyemang C, Moll van Charante EP, et al. A community-based intervention for primary prevention of cardiovascular diseases in the slums of Nairobi: the SCALE UP study protocol for a prospective quasi-experimental community-based trial. *Trials*. 2013; 14:409. [PubMed: 24289751]
41. Joshi MD, Ayah R, Njau EK, Wanjiru R, Kayima JK, Njeru EK, et al. Prevalence of hypertension and associated cardiovascular risk factors in an urban slum in Nairobi, Kenya: a population-based survey. *BMC Public Health*. 2014; 14:1177. [PubMed: 25407513]
42. Hulzebosch A, van de Vijver S, Oti SO, Egondi T, Kyobutungi C. Profile of people with hypertension in Nairobi's slums: a descriptive study. *Global Health*. 2015; 11:26. [PubMed: 26116577]
43. Werner ME, van de Vijver S, Adhiambo M, Egondi T, Oti SO, Kyobutungi C. Results of a hypertension and diabetes treatment program in the slums of Nairobi: a retrospective cohort study. *BMC Health Serv Res*. 2015; 15:512. [PubMed: 26577953]
44. Phaswana-Mafuya N, Peltzer K, Chirinda W, Musekiwa A, Kose Z. Sociodemographic predictors of multiple non-communicable disease risk factors among older adults in South Africa. *Global health action*. 2013; 6:20680. [PubMed: 24044582]
45. Shisana, O., Labadarios, D., Rehle, T., Simbayi, L., Zuma, K., Dhansay, A., et al. South African National Health and Nutrition Examination Survey (SANHANES-1). Cape Town: Human Science Research Council Press; 2013.
46. Stringhini S, Forrester TE, Plange-Rhule J, Lambert EV, Viswanathan B, Riesen W, et al. The social patterning of risk factors for noncommunicable diseases in five countries: evidence from the modeling the epidemiologic transition study (METS). *BMC Public Health*. 2016; 16:956. [PubMed: 27612934]

47. Schutte AE, Schutte R, Huisman HW, van Rooyen JM, Fourie CM, Malan NT, et al. Are behavioural risk factors to be blamed for the conversion from optimal blood pressure to hypertensive status in Black South Africans? A 5-year prospective study. *International journal of epidemiology*. 2012; 41(4):1114–23. [PubMed: 22825590]
48. Peer N, Steyn K, Lombard C, Gwebushe N, Levitt N. A high burden of hypertension in the urban black population of Cape Town: the cardiovascular risk in Black South Africans (CRIBSA) study. *PLoS One*. 2013; 8(11):e78567. [PubMed: 24250798]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

HIGHLIGHTS

- Stark differences in hypertension prevalence, awareness and control across Africa.
- Hypertension prevalence ranges between 15.1% to 54.1% in sub-Saharan Africans.
- Only 47.7% of hypertensives knew their blood pressure status.
- Levels of blood pressure control ranged from 30.0% to 61.1%.
- Urgent need for localized monitoring and treatment of blood pressure.

Figure 1.

Diagnosis, awareness, treatment and control of hypertension in men (Figure 1.A) and women (Figure 1.B) by site in the AWI-Gen study. Please note that when the numbers do not add up to the previous category, this is due to missing data (see methods).

Figure 2.

Figure 2.A Comparison of the prevalence of hypertension at each of the six AWI-Gen study sites: Burkina Faso, Ghana, Kenya and South Africa (SA). There are significant differences between the proportions of men and women who have hypertension at the following sites: Agincourt, Dikgale and Nanoro (each with $p < 0.00001$) and Navrongo ($p < 0.0001$).

Figure 2.B Shows awareness and control of hypertension at each site. The pie charts represent all individuals with hypertension and show the proportion with controlled blood pressure (green), those who are aware that they are hypertensive, but who had high blood pressure at the time of measurement (light orange), and those who were not aware that they had high blood pressure (dark orange).

Demographic characteristics of the AWI-Gen participants (aged 40 to 60 years) per research site

Table 1

Sex	Agincourt N (%)	Dikgale N (%)	Nairobi N (%)	Nanoro N (%)	Navrongo N (%)	Soweto N (%)	Total N (%)
Men	573 (39)	343 (31)	924 (45)	1034 (50)	917 (46)	1025 (51)	4816 (45)
Women	891 (61)	774 (69)	1112 (55)	1029 (50)	1071 (54)	1002 (49)	5879 (55)
Total	1464	1117	2036	2063	1988	2027	10695 *
Age							
Mean age (SD)	50.81 (5.78)	50.41 (5.98)	48.56 (5.45)	49.80 (5.81)	51.02 (5.74)	49.27 (5.83)	49.9 (5.82)
Age group							
40–44	294 (20)	231 (21)	575 (28)	500 (24)	314 (16)	539 (27)	2453 (23)
45–49	281 (19)	253 (23)	625 (31)	499 (24)	510 (26)	539 (27)	2707 (25)
50–54	434 (30)	283 (25)	470 (23)	539 (26)	454 (23)	484 (24)	2664 (25)
55–60	455 (31)	350 (31)	366 (18)	526 (26)	710 (36)	465 (23)	2872 (27)
Total	1464	1117	2036	2064	1988	2027	10696

* There is a missing value for sex in one participant from Nanoro.

Table 2

Prevalence of hypertension based on the JNC7 definitions per site, age group and sex

Sex	Total sample ¹				Men			Women		
	Study population (%)	High Blood Pressure	Prevalence (95% CI)	Study population (%)	High Blood Pressure	Prevalence (95% CI)	Study population (%)	High Blood Pressure	Prevalence (95% CI)	
Men	4816 (45)	1503	31.3 (30.0–32.6)							
Women	5879 (55)	2040	35.0 (33.7–36.2)							
Total	10695 ¹	3543	33.3 (32.4–34.2)	4797 ²	1503	31.3 (30.0–32.6)	5835 ²	2040	35.0 (33.7–36.2)	
Site										
Agincourt	1432 (13.5)	678	47.3 (44.8–49.9)	556 (11.6)	222	39.9 (35.8–44.0)	876 (15.0%)	456	52.1 (48.7–55.4)	
Dikgale	1115 (10.5)	464	41.6 (38.7–44.5)	343 (7.1)	100	29.2 (24.3–34.0)	772 (13.2%)	364	47.2 (43.6–50.7)	
Nairobi	203 (19.1)	521	25.6 (23.7–27.5)	924 (19.3)	202	21.9 (19.2–24.5)	1111 (19.0%)	319	28.7 (26.0–31.4)	
Nanoro	2063 (19.4)	311	15.1 (13.5–16.6)	1034 (21.6)	198	19.1 (16.7–21.6)	1028 (17.6%)	113	11.0 (9.1–12.9)	
Navrongo	1988 (18.7)	487	24.5 (22.6–26.4)	917 (19.1)	221	24.1 (21.3–26.9)	1071 (18.4%)	266	24.8 (22.2–27.4)	
Soweto	2000 (18.8)	1082	54.1 (51.9–56.3)	1023 (21.3)	560	54.7 (51.7–57.8)	977 (16.7%)	522	53.4 (50.3–56.6)	
Age groups										
40–44	2434 (22.9)	553	22.7 (21.1–24.4)	1149 (24.0)	270	23.5 (21.0–26.0)	1284 (22.0)	283	22.0 (19.8–24.3)	
45–49	2692 (25.3)	789	29.3 (27.6–31.0)	1197 (25.0)	339	28.3 (25.8–30.9)	1495 (25.6)	450	30.1 (27.8–32.4)	
50–54	2648 (24.9)	945	35.7 (33.9–37.5)	1171 (24.3)	365	31.2 (28.5–33.9)	1478 (25.3)	580	39.2 (36.7–41.7)	
55–60	2859 (26.9)	1256	43.9 (42.1–45.8)	1281 (26.7)	529	41.3 (38.6–44.0)	1578 (27.0)	727	46.1 (43.6–48.5)	

¹Total sample was 10696 but one person had missing sex

²Sixty-three individuals (44 women and 19 men) were excluded from hypertension analysis because they did not have blood pressure measurements

Table 3

Awareness of hypertension in the AWI-Gen cohort by research site and sex.

	AWI-Gen Total		Agincourt		Dikgale		Nairobi		Nanoro		Navrongo		Soweto	
	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women	Men	Women
Sensitivity (%)	39.4	53.8	50.5	69.3	46.9	64.5	41.1	71.8	41.5	37.0	21.4	34.6	39.4	35.1
Specificity (%)	97.9	91.2	92.8	86.7	95.9	97.5	99.2	97.7	98.2	98.4	99.3	99.2	98.4	50.8
PPV	89.6	76.9	82.4	84.9	82.1	95.9	93.1	92.7	84.4	74.1	90.4	93.9	96.8	45.0
NPV	78.2	78.4	73.8	72.2	81.6	75.3	85.9	89.6	87.6	92.7	79.8	82.0	58	40.5

PPV indicates positive predictive value, NPV indicates negative predictive value