Published in final edited form as:

Nat Chem. 2016 October ; 8(10): 922–928. doi:10.1038/nchem.2594.

Dearomative dihydroxylation with arenophiles

Emma H. Southgate[†], Jola Pospech[†], Junkai Fu, Daniel R. Holycross, and David Sarlah^{*}

Roger Adams Laboratory, Department of Chemistry, University of Illinois, Urbana, Illinois 61801, USA

Abstract

Aromatic hydrocarbons are some of the most elementary feedstock chemicals, produced annually on a million metric ton scale, and are used in the production of polymers, paints, agrochemicals and pharmaceuticals. Dearomatization reactions convert simple, readily available arenes into more complex molecules with broader potential utility, however, despite substantial progress and achievements in this field, there are relatively few methods for the dearomatization of simple arenes that also selectively introduce functionality. Here we describe a new dearomatization process that involves visible-light activation of small heteroatom-containing organic molecules—arenophiles—that results in their *para*-cycloaddition with a variety of aromatic compounds. The approach uses N–N-arenophiles to enable dearomative dihydroxylation and diaminodihydroxylation of simple arenes. This strategy provides direct and selective access to highly functionalized cyclohexenes and cyclohexadienes and is orthogonal to existing chemical and biological dearomatization processes. Finally, we demonstrate the synthetic utility of this strategy with the concise synthesis of several biologically active compounds and natural products.

The dearomatization of aromatic compounds is a fundamental synthetic strategy that provides direct and efficient access to a wide range of valuable intermediates from simple and abundant sources of hydrocarbons^{1–5}. Numerous bioactive compounds, natural products, and drugs, such as the analgesic morphine^{6–8}, the broad-spectrum antibiotic doxycycline⁹ and the antiviral drug oseltamivir (Tamiflu)^{10–12}, have been synthesized utilizing dearomatization as a key step. Despite their strategic and widespread use, most dearomative strategies do not result in the introduction of additional functionality. Indeed, the venerable dissolving-metal reduction (Birch reduction)¹³, oxidative dearomatization of phenols¹⁴, and arene–alkene photocycloadditions¹⁵ are exceptionally powerful synthetic transformations; however, most of the dearomatized products have to be subjected to further manipulations to install the desired level of functionalization. To date, only certain stoichiometric reactions of

Reprints and permissions information is available online at www.nature.com/reprints.

^{*} sarlah@illinois.edu.

[†] These authors contributed equally to this work.

Correspondence and requests for materials should be addressed to D.S.

Author contributions

E.H.S., J.P., J.F. and D.R.H. conducted the experiments, analysed the data and prepared the Supplementary Information. E.H.S., J.P. and D.S. conceived and designed the project, analysed the data and wrote the manuscript.

Competing financial interests

The authors declare no competing financial interests.

Supplementary information and chemical compound information are available in the online version of the paper.

transition-metal complexes based on Os, Ru, Re, Cr and Mn can enable more elaborate functionalizations of the corresponding metal-bound arenes (Fig. 1a, left)^{2,16–19}. Both η^2 - and η^6 -coordination modes greatly reduce the aromatic character of arene ligands and, as a consequence, activate them towards reactions with electrophiles or nucleophiles. After additional functionalization, oxidative decomplexation liberates the dearomatized products. Although these methods provide rapid access to compounds that otherwise require long and tedious manipulations, the toxicity and cost of the above-mentioned metal complexes have been significant deterrents to their widespread synthetic use. In addition to stoichiometric methods, a catalytic dearomative polyhydroxylation of benzene is known and proceeds through photoinduced charge-transfer osmylation²⁰.

Aside from chemical processes, microbial arene oxidation converts arenes into the corresponding *cis*-1,2-dihydroxycyclohexa-3,5-dienes (2,3-dihydrodiols, Fig. 1a, right) with high levels of enantioselectivity²¹. Although exceptionally powerful and synthetically useful²², this biotransformation often involves the use of specific bacterial strains or recombinant organisms to effect substrate-specific transformations, and these can usually be obtained only from the laboratories in which they were first cultured.

The development of dearomative functionalization strategies for arenes is intrinsically challenging and remains a largely unsolved synthetic problem. The high resonance energy renders aromatic compounds particularly unreactive as starting materials, and reagents that can overcome this chemical inertness preferentially react with the more-reactive unsaturated dearomatized products. Ultimately, this problem leads to overreaction and decomposition of the starting material. We envisioned dearomative functionalization as a two-stage process, and thus we avoided restrictive reactivity differences between the starting arenes and partially unsaturated products. Key to the successful execution of this plan was the use of visible-light photoactivable 2π components, for which we introduce the term ‘arenophiles’, in analogy to thermal cycloaddition processes. Specifically, using heteroatom-containing arenophiles that could formally react in a [4+2] fashion would significantly expand the toolbox of dearomative chemistry, as they would simultaneously induce dearomatization, introduce functionality, create stereogenic centres and enable further functionalization. Subsequent retrocycloaddition or fragmentation of the arenophile moiety would then provide selective access to the corresponding dearomatized products. Herein we report the realization of this concept in the development of a method for the dearomative dihydroxylation and diaminodihydroxylation of simple arenes (Fig. 1a, bottom). By using an N–N-arenophile and osmium-catalysed dihydroxylation, a variety of aromatic hydrocarbon compounds were transformed selectively into the corresponding 3,4-dihydrodiols or diaminodihydrodiols. The synthetic value of this method was demonstrated through the synthesis of several highly functionalized small organic molecules from readily available starting arenes (Fig. 1b).

Results and discussion

Design of arenophiles and reaction development

Cycloaddition reactions that involve arenes encompass an important group of dearomatization strategies²³. Aromatic compounds, known for their stability in the ground

state, become exceptionally reactive upon photoexcitation and can undergo cycloaddition with a variety of alkenes¹⁵. For these processes, high-energy ultraviolet light is usually required to access the relatively high π, π^* -singlet state of the aromatic nucleus to enable reactivity. The resulting *meta*-photocycloaddition is well documented in the literature and has been used many times in organic synthesis^{24,25}, whereas the *ortho*- and, particularly, the *para*-photocycloadditions have not received much attention, because both types rarely occur with olefins, and typically are low-yielding side reactions²⁴.

A conceptually distinct approach towards dearomative cycloaddition involves photoexcitation of the other cycloaddend to engage the arene in the ground state during the reaction (Fig. 2a). Ideally, this alternative activation mode would provide complimentary periselectivity, as well as enable cycloaddition with partners other than alkenes. In this context, we were intrigued by a report by Hamrock and Sheridan²⁶ that indicated the existence of arenophile-type reactivity with benzene; however, because of its transient nature, the corresponding *para*-cycloadduct had not been isolated or chemically explored. Furthermore, arene–arenophile photo-cycloadditions are still mechanistically ambiguous and could occur via multiple reaction trajectories, including photoinduced electron- or charge-transfer complexes between the arene and the excited arenophile²³. Nevertheless, the energy provided by visible light is sufficient to excite only the arenophile, because of its much lower-lying and narrower HOMO–LUMO gap (HOMO, highest occupied molecular orbital; LUMO, lowest unoccupied molecular orbital) compared with that of an arene (Fig. 2a, inset).

A crucial electronic requirement for the photoreactivity of an arenophile is that both the HOMO and LUMO energies are within the range of the energy of the HOMO of the arene. To evaluate the viability of potential N–N-arenophiles for the photocycloaddition chemistry, we performed a computational frontier molecular orbital analysis²⁷ of several small organic molecules using benzene (**1a**) (HOMO = –9.9 eV) and naphthalene (**2a**) (HOMO = –8.4 eV) as benchmarks (Fig. 2b). Thus, a number of different 1,2,4-triazoline-3,5-diones, **A1** (HOMO = –11.2 eV, LUMO = –9.7 eV), **A2** (HOMO = –10.8 eV, LUMO = –9.7 eV), **A4** (HOMO = –10.8 eV, LUMO = –9.3 eV) and **A5** (HOMO = –10.4 eV, LUMO = –8.9 eV), and certain symmetric cyclic (*Z*)-diazo-containing compounds connected to electron-deficient groups, such as **A3** (HOMO = –10.9 eV, LUMO = –9.5 eV) and **A6** (HOMO = –9.9 eV, LUMO = –8.8 eV), were found to meet the electronic criteria to react with benzene (the complete list of compounds is given in Supplementary Information, page 67). Next, on the visible-light irradiation of dichloromethane solutions that contain the potential arenophiles (**A1**–**A6**) in the presence of benzene, we detected the formation of *para*-cycloadducts in three different cases. Although **A1**, **A2** and **A4** all showed the desired reactivity, we decided to continue our investigations with 4-methyl-1,2,4-triazoline-3,5-dione (**A2**) because of the ease of its preparation and its stability.

We commenced our studies on dearomative dihydroxylation by evaluating the optimal reaction parameters for the formation of the cycloadduct of benzene with **A2** and its *in situ* trapping with osmium tetroxide. Although cycloaddition occurred readily at –78 °C under the influence of visible light, as evidenced by the complete disappearance of the characteristic magenta colour of **A2** (Fig. 2c) and monitoring by ¹H-NMR spectroscopy, the

corresponding cycloadduct proved to be rather thermally unstable. Cycloreversion occurred slowly at temperatures above $-50\text{ }^{\circ}\text{C}$ and rapidly above $-10\text{ }^{\circ}\text{C}$. Furthermore, we found that a 10:1 molar ratio of arene to **A2** proved optimal, although ratios as low as 2:1 often gave similar yields, albeit at the expense of longer reaction times. In view of the low thermal stability of the intermediate product, the development of reaction conditions for cold-temperature dihydroxylation proved necessary. As a result, two different catalytic conditions were identified for the *in situ* dihydroxylation of **A2** cycloadducts with mononuclear arenes. Under the first set of conditions (Table 1, conditions A), the cycloaddition reaction was run in acetone, and the subsequent addition of osmium(VIII) oxide and a solution of 4-methylmorpholine *N*-oxide (NMO), water and *p*-toluenesulfonamide (*p*-TsNH₂) delivered the dihydroxylated benzene cycloadduct **3a** in 56% yield. The addition of *p*-TsNH₂ proved to be crucial for higher yields as it facilitated hydrolysis of the intermediate osmate ester²⁸. The second dihydroxylation process (conditions B), based on a modified Narasaka–Sharpless method²⁹, was performed using dichloromethane as the solvent. Thus, after the disappearance of the magenta colour, the addition of osmium(VIII) oxide, NMO and *n*-butylboronic acid afforded the cyclic boronate ester **4a** in 65% yield.

Reaction generality

With the optimal conditions in hand, we began an exploration of the scope of this dearomative tetrafunctionalization by examining simple, mononuclear arenes (Table 1). In addition to benzene (**1a**), a variety of monosubstituted derivatives proved to be suitable photocycloaddition partners. The tolerance of halogen (**1k**, **1l** and **1m**) and benzylic heteroatom (**1e**, **1f**, **1h**, **1i** and **1j**) substituents is interesting, as these type of substrates are not generally compatible with chemical-based dearomatizations. Traditionally, the use of ultraviolet irradiation in arene cycloaddition chemistry does not permit broader functional-group incorporation^{15,23}. In contrast, even benzyl chloride (**1l**) and bromobenzene (**1m**) underwent *para*-cycloaddition with **A2**. This dearomative protocol can be conducted on a larger scale, exemplified with a multigram conversion of benzene (**1a**) and bromobenzene (**1m**) without significant erosions in yields (Supplementary Information, pages 7 and 21). In addition to cycloaddition, we also observed that the abstraction of benzylic C–H bonds by photoexcited **A2** was a competitive process with certain substrates, to give formal C–H amidation products (limitations of the method are given in Supplementary Information, page 22)³⁰. For example, amidation proved to be a major reaction pathway with toluene (not shown) and a minor side process with cumene (**1b**). Importantly, all the substrates reacted in a highly stereo- and regioselective manner; the resulting products were consistently obtained as a single constitutional isomer and diastereoisomer (an X-ray structure of the acetonide-protected **3a** is given in Supplementary Information, page 162). At present, there is no conclusive explanation for the high observed regioselectivity of the cycloaddition process; however, computational studies are currently ongoing to elucidate the origins of this selectivity. Finally, at the current level of development, polysubstituted mononuclear arenes are not suitable substrates for the dearomatization reaction we describe (Supplementary Information, page 23).

With the dihydroxylated bicyclic adducts prepared, we turned our attention to the cycloreversion of the arenophile moiety to liberate the desired dihydrodiols. Although

similar unsaturated bicyclic urazoles are known to undergo thermal cycloreversion³¹, no reaction was observed on heating cycloadducts **3** or **4** to temperatures up to 250 °C. Therefore, we decided to examine a one-pot urazole hydrolysis/bicyclic hydrazine oxidation sequence that would generate dihydrodiols via the extrusion of molecular dinitrogen. Although the hydrolysis and oxidation sequence was expected to be reasonably straightforward, the lability of such dihydrodiols was potentially troublesome^{21,32}. Indeed, this turned out to be the case, as we observed significant amounts of phenol formation during the oxidation step. However, on examination of a range of protecting groups and conditions, we eventually found that using the corresponding acetonides **5** in combination with neat hydrazine or KOH in 2-propanol³³, followed by CuCl₂-mediated oxidation, successfully generated protected dihydrodiols **6** as stable compounds. This one-pot sequence proved to be highly efficient, as the corresponding diene diol products were prepared in high yields (Table 2, left side). Most of the functional groups tested proved resistant to urazole hydrolysis and hydrazine oxidation, with the exception that esters were converted into the corresponding carboxylic acids (**6j**) and alcohols (**6e**, **6f** and **6g**). Importantly, under the reaction conditions described, no potentially competitive re-aromatization process was observed and all the acetonide-protected dihydrodiols were stable to standard purification methods. Finally, all the substituted dihydrodiols were complementary constitutional isomers to those obtained by microbial arene oxidation²¹.

Next, we sought to investigate arenophile fragmentation to extend further the functional scope and utility of this method (Table 2, right side). This manipulation was achieved in two steps using hydrolysis of the urazole moiety under the above-mentioned conditions, followed by benzoylation and reductive cleavage of the N–N bond with samarium diiodide³⁴. A range of substituted cycloadducts was transformed into the respective unsaturated diaminodiols **7**. Moreover, vinyl bromide (**7m**) survived under these reductive conditions; only cycloadduct **5l**, derived from benzyl chloride, was converted into the corresponding amide **7l** during this reaction sequence. Importantly, this synthetic sequence provides direct and selective access to highly functionalized small molecules that are characterized by four contiguous heteroatom-containing stereogenic centres. The arenophile-mediated dihydroxylation with subsequent fragmentation was also applicable to site-selective dearomatization of polynuclear arenes (Table 3)^{35–38}. Thus, using a two-step protocol that involves **A2** cycloaddition/dihydroxylation and urazole hydrolysis/catalytic hydrogenation, a series of naphthalene derivatives was successfully converted into the corresponding diaminodiols. In addition to halogens **2b–2d**, amide (**2e**) and benzylic ketal (**2f**) substituents survived under the reaction conditions.

Moreover, this strategy proved suitable for trinuclear arenes, such as phenyl- and pyridine-containing naphthalenes (**2g** and **2h**) and phenanthrene (**2j**). For **2g** and **2h**, the observed site selectivity of dearomatization could be explained by the higher ionization potentials of phenyl and pyridine as compared with that of naphthalene³⁹. In all cases, a single constitutional isomer and diastereoisomer was obtained (X-ray structures of **8a** and tetra-acetylated **Ac-9a** are given in Supplementary Information, pages 163 and 164).

Applications of dearomative dihydroxylation

The presented dearomative dihydroxylation strategy can enable rapid access to small, highly functionalized organic molecules; three illustrative cases are demonstrated in Fig. 3. For example, conduramine A (ref. 40) (**11**) was prepared from the corresponding **A2**–benzene cycloadduct **5a** through a modified hydrolysis–oxidation sequence that installed an additional 1,4-*syn*-aminohydroxy functionality. Accordingly, one-pot successive urazole hydrolysis, hydrazine/oxamic acid oxidation⁴¹ and subsequent hetero Diels–Alder reaction delivered the bicyclic product **10** in 83% yield. Concurrent N–O cleavage and deprotection of the trichloroethoxycarbonyl (Troc) group under reductive conditions, followed by acid-mediated deprotection of the acetonide, furnished conduramine A (**11**). Next, the highly oxygenated cyclohexene herbicidal natural product MK7607 (**13**) (ref. 42) was expediently synthesized in three steps from benzyl-alcohol-derived dihydrodiol **6e** via protection, dihydroxylation (**6e** → **12**) and subsequent double deprotection. Similarly, bromodihydro-3,4-diol **6m** was concisely converted into the 3-*O*-desmethylated analogue of the potent herbicidal agent phomentrioloxin (**16**)⁴³ through a stereoselective dihydroxylation, followed by Sonogashira coupling between the resulting vinyl bromide **14** and acetylene **15**, followed by deprotection⁴⁴.

Conclusion

In summary, we disclose here a dearomative functionalization of simple arenes that provides access to dihydrodiols and diaminodihydrodiols. Significant features of this strategy include visible-light activation of the arenophile, *in situ* dihydroxylation of the corresponding arene–arenophile cycloadduct, and subsequent arenophile cycloreversion or fragmentation. This method permits the use of substrates that are normally not suitable for chemical-based dearomatizations, such as halogen- and benzylic-heteroatom-containing arenes, and provides products that are not obtained by microbial oxidation. In addition to engaging a broad range of monosubstituted benzene derivatives, this system was also found to be applicable to the site-selective dearomative diaminodihydroxylation of polynuclear arenes. Finally, we have showcased the utility of arenophile-based dihydroxylation as a starting point in the expedient preparation of highly functionalized small organic molecules. The study and expansion of the arenophile dearomative platform to other areas, including the application of different functionalization reactions and natural product synthesis, is currently underway and will be reported in due course.

Data availability

Crystallographic data for this paper have been deposited at the Cambridge Crystallographic Data Centre under deposition numbers CCDC 1455841–1455843. These data can be obtained free of charge from www.ccdc.cam.ac.uk/data_request/cif.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors acknowledge the University of Illinois for generous support of this work. E.H.S. is a Springborn Graduate Fellow and J.P. is a Fellow of the German Research Foundation (DFG). We thank S. E. Denmark and P. J. Hergenrother (University of Illinois) for critical proofreading of this manuscript. We dedicate this article to K. C. Nicolaou on the occasion of his 70th birthday.

References

1. Roche SP, Porco JA. Dearomatization strategies in the synthesis of complex natural products. *Angew Chem Int Ed.* 2011; 50:4068–4093.
2. Pape AR, Kaliappan KP, Kündig EP. Transition-metal-mediated dearomatization reactions. *Chem Rev.* 2000; 100:2917–2940. [PubMed: 11749310]
3. López Ortiz F, Iglesias MJ, Fernández I, Andújar Sánchez CM, Ruiz Gómez G. Nucleophilic dearomatizing (DNAr) reactions of aromatic C,H-systems. A mature paradigm in organic synthesis. *Chem Rev.* 2007; 107:1580–1691. [PubMed: 17488057]
4. Pouységu L, Deffieux D, Quideau S. Hypervalent iodine-mediated phenol dearomatization in natural product synthesis. *Tetrahedron.* 2010; 66:2235–2261.
5. Zhuo CX, Zhang W, You SL. Catalytic asymmetric dearomatization reactions. *Angew Chem Int Ed.* 2012; 51:12662–12686.
6. Parker KA, Fokas D. Convergent synthesis of ($\pm$)-dihydroisocodeine in 11 steps by the tandem radical cyclization strategy. A formal total synthesis of ($\pm$)-morphine. *J Am Chem Soc.* 1992; 114:9688–9689.
7. Varghese V, Hudlicky T. Short chemoenzymatic total synthesis of ent-hydromorphone: an oxidative dearomatization/intramolecular [4+2] cycloaddition/amination sequence. *Angew Chem Int Ed.* 2014; 53:4355–4358.
8. Tissot M, et al. Gram-scale enantioselective formal synthesis of morphine through an *ortho*–*para* oxidative phenolic coupling strategy. *Angew Chem Int Ed.* 2014; 53:13498–13501.
9. Charest MG, Lerner CD, Brubaker JD, Siegel DR, Myers AG. A convergent enantioselective route to structurally diverse 6-deoxytetracycline antibiotics. *Science.* 2005; 308:395–398. [PubMed: 15831754]
10. Zutter U, Iding H, Spurr P, Wirz B. New, efficient synthesis of oseltamivir phosphate (Tamiflu) via enzymatic desymmetrization of a meso-1,3-cyclohexanedicarboxylic acid diester. *J Org Chem.* 2008; 73:4895–4902. [PubMed: 18517254]
11. Shie JJ, Fang JM, Wong CH. A concise and flexible synthesis of the potent anti-influenza agents Tamiflu and Tamiphosphor. *Angew Chem Int Ed.* 2008; 47:5788–5791.
12. Sullivan B, Carrera I, Drouin M, Hudlicky T. Symmetry-based design for the chemoenzymatic synthesis of oseltamivir (Tamiflu) from ethyl benzoate. *Angew Chem Int Ed.* 2009; 48:4229–4231.
13. Rabideau PW, Marcinow Z. The Birch reduction of aromatic compounds. *Org React.* 1992; 42:1–334.
14. Moriarty RM, Prakash OM. Oxidation of phenolic compounds with organohypervalent iodine reagents. *Org React.* 2001; 57:327–415.
15. Cornelisse J. The meta photocycloaddition of arenes to alkenes. *Chem Rev.* 1993; 93:615–669.
16. Kündig EP, Pape A. Dearomatization via η^6 -arene complexes. *Top Organomet Chem.* 2004; 7:71–94.
17. Keane JM, Harman WD. A new generation of π -basic dearomatization agents. *Organometallics.* 2005; 24:1786–1798.
18. Smith PL, Chordia MD, Dean Harman W. Synthetic applications of the dearomatization agent pentaammineosmium(II). *Tetrahedron.* 2001; 57:8203–8225.
19. Todd MA, Sabat M, Myers WH, Smith TM, Harman WD. Stereoselective umpolung tandem addition of heteroatoms to phenol. *J Am Chem Soc.* 2008; 130:6906–6907. [PubMed: 18473458]
20. Jung PMJ, Motherwell WB, Williams AS. Stereochemical observations on the bromate induced monobromopentahydroxylation of benzene by catalytic photoinduced charge-transfer osmylation. A concise synthesis of ($\pm$)-pinitol. *Chem Commun.* 1997:1283–1284.

21. Johnson RA. Microbial arene oxidations. *Org React*. 2004; 63:117–264.
22. Boyd DR, Bugg TDH. Arene *cis*-dihydrodiol formation: from biology to application. *Org Biomol Chem*. 2006; 4:181–192. [PubMed: 16391757]
23. McCullough JJ. Photoadditions of aromatic compounds. *Chem Rev*. 1987; 87:811–860.
24. Streit U, Bochet CG. The arene–alkene photocycloaddition. *Beilstein J Org Chem*. 2011; 7:525–542. [PubMed: 21647263]
25. Wender PA, et al. Arene–alkene cycloadditions and organic synthesis. *Pure Appl Chem*. 2009; 62:1597–1602.
26. Hamrock SJ, Sheridan RS. *Para* photoaddition of *N*-methyltriazolinedione to benzene Synthesis of energy-rich azo compounds comprising benzene + nitrogen. *J Am Chem Soc*. 1989; 111:9247–9249.
27. Zhang G, Musgrave CB. Comparison of DFT methods for molecular orbital Eigenvalue calculations. *J Phys Chem A*. 2007; 111:1554–1561. [PubMed: 17279730]
28. Sharpless KB, et al. The osmium-catalyzed asymmetric dihydroxylation: a new ligand class and a process improvement. *J Org Chem*. 1992; 57:2768–2771.
29. Gypser A, Michel D, Nirschl DS, Sharpless KB. Dihydroxylation of polyenes using Narasaka's modification of the Upjohn procedure. *J Org Chem*. 1998; 63:7322–7327. [PubMed: 11672379]
30. Breton GW, Hoke KR. Application of radical cation spin density maps toward the prediction of photochemical reactivity between *N*-methyl-1,2,4-triazoline-3,5-dione and substituted benzenes. *J Org Chem*. 2013; 78:4697–4707. [PubMed: 23594077]
31. Rickborn B. The retro-Diels–Alder reaction. Part II Dienophiles with one or more heteroatoms. *Org React*. 1998; 53:223–629.
32. Lawlor DA, et al. Hyperaromatic stabilization of arenium ions: a remarkable *cis* stereoselectivity of nucleophilic trapping of β -hydroxyarenium ions by water. *J Am Chem Soc*. 2011; 133:19718–19728. [PubMed: 22023504]
33. Nelsen SF, Trieber DA, Wolff JJ, Powell DR, Rogers-Crowley S. Intramolecular electron transfer between doubly six σ -bond-linked trialkyldiazonium cation and diazenyl radical units. *J Am Chem Soc*. 1997; 119:6873–6882.
34. Burk MJ, Feaster JE. Enantioselective hydrogenation of the C=N group: a catalytic asymmetric reductive amination procedure. *J Am Chem Soc*. 1992; 114:6266–6267.
35. Kjell DP, Sheridan RS. Photochemical cycloaddition of *N*-methyltriazolinedione to naphthalene. *J Am Chem Soc*. 1984; 106:5368–5370.
36. Hamrock SJ, Sheridan RS. Photochemical Diels–Alder addition of *N*-methyltriazolinedione to phenanthrene. *Tetrahedron Lett*. 1988; 29:5509–5512.
37. Kjell DP, Sheridan RS. A photochemical Diels–Alder reaction of *N*-methyltriazolinedione. *J Photochem*. 1985; 28:205–214.
38. Breton GW, Newton KA. Further studies of the thermal and photochemical Diels–Alder reactions of *N*-methyl-1,2,4-triazoline-3,5-dione (MeTAD) with naphthalene and some substituted naphthalenes. *J Org Chem*. 2000; 65:2863–2869. [PubMed: 10814171]
39. Turner, DW. *Advances in Physical Organic Chemistry*. Gold, V., editor. Vol. 4. Academic Press; 1966. p. 31–71.
40. Balci M, Sütbeyaz Y, Secen H. Conduritols and related compounds. *Tetrahedron*. 1990; 46:3715–3742.
41. Frazier CP, Engelking JR, Read de Alaniz J. Copper-catalyzed aerobic oxidation of hydroxamic acids leads to a mild and versatile acylnitroso ene reaction. *J Am Chem Soc*. 2011; 133:10430–10433. [PubMed: 21678942]
42. Nobuji, Y., et al. Novel herbicidal MK7607 and its manufacture with *Curvularia*. Japan Kokai Tokkyo Koho, Japanese Patent. 06306000. 1994.
43. Cimmino A, et al. Phomentrioloxin, a fungal phytotoxin with potential herbicidal activity, and its derivatives: a structure–activity relationship study. *J Agric Food Chem*. 2013; 61:9645–9649. [PubMed: 24083323]

44. Ma X, Banwell MG, Willis AC. Chemoenzymatic total synthesis of the phytotoxic geranycyclohexentriol (–)-phomentrioloxin. *J Nat Prod.* 2013; 76:1514–1518. [PubMed: 23895674]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1. Current strategies in dearomative functionalization of arenes, which include this work
a, Left, transition-metal complexation to arenes results in significant distortions of the of the π -electron density and enables dearomative functionalization. Top right, arene dioxygenase (Protein Data Bank 3EN1) is a key enzyme in microbial arene oxidation that converts monosubstituted mononuclear arenes into the corresponding optically pure 2,3-dihydrodiols. Bottom right, visible-light activation of small organic molecules (arenophiles) with *in situ* dihydroxylation provides tetrafunctionalized bicyclic compounds (this work). Subsequent retrocycloaddition or fragmentation delivers racemic 3,4-dihydrodiols or diaminodihydrodiols, respectively. **b**, Small, highly functionalized molecules prepared using arenophile-based dihydroxylation. Conduramine synthesis was achieved via nitroso Diels–Alder cycloaddition and deprotection from the benzene-derived dihydrodiol. MK7607 synthesis was accomplished via dihydroxylation and deprotection of the corresponding dihydrodiol derivative of benzyl acetate. The synthesis of 3-*O*-desmethyl phomentrioloxin was completed via dihydroxylation and Sonogashira coupling of the bromobenzene-derived dihydrodiol. TM, transition metal; Nu, nucleophile; E, electrophile.

Figure 2. Mechanistic considerations and necessary criteria for arenophile reactivity

a, Mechanistic rationale for the visible-light activation of an arenophile in the presence of an arene. The excited state of the arenophile is capable of forming an electron- and/or charge-transfer-derived exciplex with the ground-state arene. The resulting exciplex subsequently collapses to form an arene-arenophile cycloadduct. **b**, Mechanism-guided computational (B3LYP/6-311+G(d,p)) discovery of potential arenophiles based on benzene (**1a**) and naphthalene (**2a**) as archetypical arenes. The requirements for arenophile reactivity are: (1) the HOMO-LUMO gap of the arenophile is narrow enough to permit visible-light excitation and (2) the HOMO of the arene is within the HOMO-LUMO gap of the arenophile. Three arenophiles (insets) showed the desired reactivity with benzene. **c**, Magenta colour of a solution of **A2** and benzene (**1a**, 10 equiv.) in dichloromethane before and after visible-light-mediated photocycloaddition. SOMO, singly occupied molecular orbital.

Figure 3. Synthetic applications of dearomative dihydroxylation

Further functionalization of the dihydroxylated **A2**–benzene cycloadduct **5a** and benzyl-alcohol- and bromobenzene-derived 3,4-dihydrodiols (**6e** and **6m**) provides rapid access to conduramine A (**11**) (**a**), MK7607 (**13**) (**b**) and 3-*O*-desmethyl phomentrioloxin (**16**) (**c**). imid., imidazole; Py, pyridine; r.t., room temperature; TBS, *t*-butyldimethylsilyl; TFA, trifluoroacetic acid; Troc, trichloroethoxycarbonyl.

Table 1

Substrate scope of mononuclear arenes.

Conditions A: arene (**1**, 10 equiv.), **A2** (1.0 equiv.), visible-light-emitting diodes (VLEDs), CH_2Cl_2 , -78°C , 4 h; then OsO_4 (5 mol%), NMO (1.2 equiv.), *p*-TsNH₂ (1.2 equiv.), H₂O (20 equiv.), acetone (0.1 M based on **A2**), -78°C to 0°C . Conditions B: arene (**1**, 10 equiv.), **A2** (1.0 equiv.), VLEDs, CH_2Cl_2 (0.1 M based on **A2**), -78°C , 4 h; then OsO_4 (5 mol%), NMO (1.2 equiv.), *n*BuB(OH)₂ (1.2 equiv.), CH_2Cl_2 , -78°C to 0°C . Isolated yields of the purified material based on **A2**. TMS, trimethylsilyl.

Table 2

Cycloreversion or fragmentation of the arenophile moiety.

Conditions for the preparation of 6: N_2H_4 (10 equiv.), 100 °C, 12 h; or KOH (10 equiv.), tPrOH (0.1 M), 100 °C, 12 h; then CuCl_2 , NH_4OH , 25 °C, 5 min. Conditions for the preparation of 7: (1) N_2H_4 (10 equiv.), 100 °C, 12 h; or KOH (10 equiv.), tPrOH (0.1 M), 100 °C, 12 h, then BzCl (5.0 equiv.); (2) SmI_2 (3.0 equiv.), MeOH , 25 °C, 1 h. Isolated yields of the purified material based on the A2-arene cycloadduct 5. Bz, benzoyl.

Table 3

Site-selective dearomative diaminodihydroxylation of polynuclear arenes.

Conditions for step 1: arene (**2**, 1.5 equiv.), **A2** (1.0 equiv.), VLEDs, acetone, -78°C , 3 h; then OsO_4 (5 mol%), NMO (2.0 equiv.), citric acid (0.5 equiv.), H_2O (20 equiv.), acetone (0.1 M based on **A2**), -78°C to 0°C . Conditions for step 2: N_2H_4 (10 equiv.), 100°C , 12 h; then Raney-Ni (0.5 equiv.), H_2 , EtOH, 50°C , 12 h. Isolated yields of purified material. Cat., catalyst.