CORRESPONDENCE

Polyneuropathies—Etiology, Diagnosis, and Treatment Options

by Prof. Dr. med. Claudia Sommer, PD Dr. med. Christian Geber, Prof. Dr. med. Peter Young, Prof. Dr. med. Raimund Forst, Prof. Dr. med. Frank Birklein, and Prof. Dr. med. Benedikt Schoser in issue 6/2018

Disturbed Microcirculation as Cause

This interesting overview of polyneuropathy (PNPs) does not give ischemia, as a pathophysiological agent, the weighting it deserves in the epidemiology of microvascular risk constellations (1). In light of the facts that the increasing prevalence of PNPs in the general population cannot be explained solely by the population pyramid and the prevalence of diabetes mellitus, and that 20%–46% of PNPs cannot be assigned to established etiologies (2), a disturbed microcirculation deserves to be considered as a cause of PNP, beyond the rare condition of vasculitis.

The functional significance of a supply deficiency to neurons due to capillary / arterial hypoperfusion comes from the microcirculatory changes on the acral part of lower extremities during prolonged static posture in an orthostatic body position. Numerous risk situations for this can be found in the industrial world, such as working in a concentrated manner seated at a computer desk over longer periods of time. The cause of the hypoxic reduced oxygen supply is the circulatory physiological event known as the venoarterial reflex (3). During prolonged sitting, postcapillary level of tissue oxygen saturation decreases from 40%–50% to values of 10%. This effect is part of the “savings principle” of the organism, setting on after only 2 to 4 minutes rather than hours, and persisting until onset of longer phases of movement. Chronic recurrent hypoxic malnutrition therefore appears to be worthy of discussion as a cause of PNP in individuals with relevant exposure, especially in times of epidemic proportions of prolonged fixed sitting at computer workplaces.

Exercise strategies to maintain foot and toe mobility are an option for prophylaxis without side effects, that is also free of cost (4). The efficiency of these measures, which appear trivial only to those who do not know about them, has not yet been proven for PNP in studies, but the treatment principle has already been used successfully elsewhere (for example, for therapy-refractory wounds) and has even been commercially implemented.

DOI: 10.3238/arztebl.2018.0295a

References

1. Sommer C, Geber C, Young P, Forst R, Birklein F, Schoser B: Polyneuropathies—etiology, diagnosis, and treatment options. *Dtsch Arztebl Int* 2018; 115: 83–90.
2. Hanewinkel R, Drenthen J, van Oijen M, Hofman A, van Doorn PA, Ikram MA: Prevalence of polyneuropathy in the general middle-aged and elderly population. *Neurology* 2016; 87: 1892–8.

3. Henriksen O, Sejrsen P: Local reflex in microcirculation in human skeletal muscle. *Acta Physiol Scand* 1977; 99: 19–26.
4. Brandl R, Stiegler H: Das diabetische Fußsyndrom – Pathophysiologie, Diagnostik und Therapie. *Deutsche Med Wochenschr* 2015; 140: 353–62.

Prof. Dr. med. Richard Brandl

Practice for Vascular Surgery and Venous Medicine am Marienplatz München
info@venenpraxis-muenchen-zentrum.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Occurrence in Stages of Prediabetes

I noticed that the article did not mention that (sometimes severe) polyneuropathies can occur even in stages of prediabetes (prediabetes, subclinical diabetes, and impaired glucose tolerance) (1). So what counts here is the diabetic diathesis, the predisposition, rather than the current blood sugar level. Every practicing neurologist knows this, and no recent research has shown a relationship in type 2 diabetes between good blood glucose control and neurological damage or symptoms. This differs from other sequelae of diabetes, such as vascular damage, for which blood sugar levels are very important. Why then is obtaining good blood glucose control still being repeatedly recommended (especially from internists) for the treatment of polyneuropathy? *Ut aliquid fiat?* DOI: 10.3238/arztebl.2018.0295b

Reference

1. Sommer C, Geber C, Young P, Forst R, Birklein F, Schoser B: Polyneuropathies—etiology, diagnosis, and treatment options. *Dtsch Arztebl Int* 2018; 115: 83–90.

Dr. med. Klaus Pillhatsch

Neurology Practice
Regensburg

Conflict of interest statement

The author declares that no conflict of interest exists.

Supplemental Information Necessary

The article provides a good overview of causes, pathophysiology, symptoms, and treatment of polyneuropathy (1). Nonetheless, I would like to make some additional comments. The review overlooks post-actinic, severe plexus polyneuropathies as well as delayed damage from radiation (X-rays, cobalt, etc.) to the small vessels of the spinal cord and the nerves (neurons, medullary sheaths, myelin-sheath gaps, etc.), which are associated with massive pain, sensory and motor failures in the thoracic and pelvic areas, as well

as tendency to edema (neuropathic, lymphogenic). With this in mind, mammary carcinomas, lymphomas, and solid tumors should be considered. As the frequently requested procedure of nerve biopsy is not risk-free, and as one would also expect specialists to show restraint in this, exclusion diagnostics should primarily be used.

With respect to therapy, the authors cite a meta-analysis (see reference [36] in the article) yet do not present a critical analysis of the often overwhelming massive side effects—which often lead to discontinuation of the preparations. The sometimes helpful high-frequency therapy or the implantation of an SCS (spinal cord stimulator) should be considered in patients whose peripheral spinal nerves are still intact.

DOI: 10.3238/arztebl.2018.0295c

References

1. Sommer C, Geber C, Young P, Forst R, Birklein F, Schoser B: Polyneuropathies—etiology, diagnosis, and treatment options. *Dtsch Arztebl Int* 2018; 115: 83–90.

Dr. med. Torsten Maurer
Dresden

Conflict of interest statement

The author declares that no conflict of interest exists.

Amiodarone-Associated Vestibulopathy

The successful review by C. Sommer and colleagues (1) should be supplemented with information about a recently discovered new facet of toxic polyneuropathy as a side effect of amiodarone (2). As has emerged in the past year, this common antiarrhythmic drug is responsible not only for sensorimotor symptoms in polyneuropathy, but also for sensory—and more specifically, vestibular—neuropathy, as it causes a bilateral loss of balance (bilateral vestibulopathy) (3). The clinical appearance of bilateral vestibulopathy is vertigo, gait instability, oscillopsia, and risk of falls. Thus, in one dizziness center, 43% of patients taking amiodarone had bilateral vestibulopathy (2). Further, a recent retrospective multi-center study showed that 12% of patients with “idiopathic” bilateral vestibulopathy were taking amiodarone (3). This phenomenon has not been previously reported in the literature. The probable explanation for this is that the affected patients usually also have sensory polyneuropathy (or other central nervous symptoms), which may also cause gait instability, resulting in the symptoms being attributed to these diseases and the balance function not being examined (4). Today, a wide range of functional tests for the balance system is available, which can quickly and reliably document vestibulopathy as the cause of dizziness and gait instability. Therefore, patients taking amiodarone who present with dizziness and/or gait instability should undergo vestibular function testing. Only then can this side effect be recognized early and be included in the benefit–risk assessment of amiodarone therapy.

DOI: 10.3238/arztebl.2018.0296a

References

1. Sommer C, Geber C, Young P, Forst R, Birklein F, Schoser B: Polyneuropathies—etiology, diagnosis, and treatment options. *Dtsch Arztebl Int* 2018; 115: 83–90.
2. Ruehl RM, Gürkov R: Amiodarone-induced gait unsteadiness is revealed to be bilateral vestibulopathy. *Eur J Neurol* 2017; 24: e7–e8.
3. Gürkov R, Manzari L, Blodow A, Wenzel A, Pavlovic D, Luis L: Amiodarone-associated bilateral vestibulopathy. *Eur Arch Otorhinolaryngol* 2017; 275: 823–25.
4. Gürkov R: Amiodarone: a newly discovered association with bilateral vestibulopathy. *Front Neurol* 2018; 9: 119.

Prof. Dr. med. Robert Gürkov

Klinik für Hals-Nasen-Ohrenheilkunde
Ludwig-Maximilians-Universität München
rguerkov@med.uni-muenchen.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Leprosy Neuritis

I would like to add a small supplement to the review article “Polyneuropathies”, which also takes into account rarer clinical pictures, and point out that cases of neuritis also occur in leprosy (albeit rarely) (1). In November 1992, a detailed article by Prof. Schliack was published on this topic (leprosy neuritis) in *Deutsches Ärzteblatt* (2). Although leprosy has declined since that publication, I would like to summarize a case study from practice: years ago, a patient requiring further neurological diagnostics presented with symmetrical sensations in the feet and hands and skin lesions on the face and trunk of body, which were indicative of an allergic reaction. Indeed, he had been dermatologically assessed a few days earlier and had been given the diagnosis of an unclear allergic reaction. Neurologically, the symptoms were summarized as “glove and stocking” hyperalgesia, weakened Achilles tendon reflexes, no pareses, and (in contrast to the typical symptoms of polyneuropathy) several 1-Euro (US quarter coin)-sized, well-reproducible analgesic areas in the trunk. Although symmetrical sensory disturbances are not typical of leprosy neuritis, and the prior dermatological assessment did not suggest leprosy, the symptoms of the analgesic areas on the trunk that were unusual for a “typical polyneuropathy” led to the suspicion of leprosy.

Presentation of the patient to the Department of Infectious Diseases of the responsible university hospital confirmed the suspicion of a borderline form of leprosy, and treatment was initiated. Based on this contact to the Department of Infectious Diseases, two further patients with leprosy (who had been in constant care of the infectious outpatient clinic) presented themselves for neurological-neurophysiological diagnostics in the following years, suggesting that the disease is probably not quite as rare in Germany as one might expect. With regard to migratory movements in Germany, one should perhaps keep this diagnosis in mind.

DOI: 10.3238/arztebl.2018.0296b

References

1. Sommer C, Geber C, Young P, Forst R, Birklein F, Schoser B: Polyneuropathies—etiology, diagnosis, and treatment options. *Dtsch Arztebl Int* 2018; 115: 83–90.
2. Schliack H: Die lepröse Neuritis. *Dtsch Arztebl* 1992; 89: A-3742.

Dr. med. Frank von Au
Bad Orb
dr.med.frank.von-au@gmx.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We are grateful for the five constructive contributions to the discussion. Prof. Brandl points out that a neurological microcirculation disorder, which among other things occurs as chronic ischemia in about 20% of the patients with peripheral arterial disease (PAD), should not be overlooked. Ischemia of the peripheral nerve often leads to neuropathic pain. In addition to anamnesis and examination, quantitative sensory testing (QST) and, if necessary, determination of intraepidermal nerve fiber density are helpful. In addition to drug therapy with non-opioid and opioid analgesics, or antidepressants and anticonvulsants, maintaining mobility of the foot and toes may be used to some degree as a prophylaxis, if necessary. However, controlled studies are lacking. Alternatively, there may be an indication for the use of spinal cord stimulation (SCS) with good evidence (1).

Dr. Pillhatsch suggests that pre-stages of diabetes may be associated with polyneuropathy, and that there is no direct correlation between optimization of management of type 2 diabetes and diabetic neuropathy. Indeed, this unexpected difference to type 1 diabetes-associated neuropathy underscores that these diseases must have distinct pathomechanisms, despite similar clinical syndromes (2). The differences in pathomechanisms concern the polyol pathway, the hexosamine pathway, protein kinase C isoforms, accumulation of “advanced glycation end products” (AGEs), and the proportion of glucose and fatty acids in diabetic nerves. Nonetheless, intensified diabetes therapy is important for vascular-associated and retinal events.

Dr. Maurer supplements the spectrum of polyneuropathy with post-actinic, severe plexus polyneuropathies and, in particular, with the important topic of delayed damage from radiation therapy (3). Performing a nerve biopsy is of course a surgical and diagnostic procedure, but in the hands of an experienced surgeon does not present a high risk. No matter the positive therapeutic effects shown for a drug therapy in studies,

the individual risk profile and possible side effects must of course be taken into account when giving information to a patient.

Dr. Gürkov comments on the importance of amiodarone-induced vestibulopathy in the differential diagnosis of a gait disorder. Therefore, in our opinion, the vestibulo-ocular reflex should be checked routinely during the initial clinical examination.

Dr. von Au recalls an important but rare cause of neuropathy, namely leprosy. This is mostly a multilocal distribution of hypoesthetic areas, which provides an important differential diagnosis to a classic distal symmetric polyneuropathy syndrome.

DOI: 10.3238/arztebl.2018.0297

References

1. Lang PM.: Ischämische schmerzhaftes Neuropathie. *Der Nervenarzt* 2015; 86: 151–5.
2. Callaghan BC, Hur J, Feldman EL.: Diabetic neuropathy: one disease or two? *Curr Opin Neurol* 2012; 25: 536–41.
3. Pradat PF, Delanian S.: Late radiation injury to peripheral nerves. *Handb Clin Neurol* 2013; 115: 743–58.
4. Sommer C, Geber C, Young P, Forst R, Birklein F, Schoser B.: Polyneuropathies—etiology, diagnosis, and treatment options. *Dtsch Arztebl Int* 2018; 115: 83–90.

Prof. Dr. med. Benedikt Schoser

Friedrich-Baur Institute, Department of Neurology
Ludwig-Maximilians-Universität München
benedikt.schoser@med.uni-muenchen.de

Prof. Dr. med. Claudia Sommer

Department of Neurology, University Hospital Würzburg

PD Dr. med. Christian Geber, DRK Pain Center Mainz

Prof. Dr. med. Raimund Forst, University Orthopedic Clinic Erlangen

Prof. Dr. med. Peter Young, Department of Sleep Medicine and Neuromuscular Disorders, Münster University

Prof. Dr. med. Frank Birklein, Klinik für Neurologie, Universitätsklinikum Mainz

Conflict of interest statement

Prof. Schoser has received conference fee and travel cost reimbursement as well as speaking honoraria from CSL Behring.

Prof. Sommer has received consultant honoraria from AirLiquide, Astellas, Baxter/Baxalta, CSL Behring, and Genzyme, LFB; speaking honoraria from Baxalta, Genzyme, Kedrion, Novartis, and Pfizer; and study support (third-party funds) from Kedrion and CSL Behring.

Dr. Geber has received consultant honoraria from Pfizer.

Prof. Young has received consultant honoraria, reimbursement for conference fees and travel costs, and study support (third-party funds) from Pharmext.

Prof. Birklein has received speaking honoraria from Pfizer and study support (third-party funds) from Pfizer and Lilly.

Prof. Forst declares that no conflict of interest exists.