Published in final edited form as:

ChemBiochem. 2018 January 18; 19(2): 121–125. doi:10.1002/cbic.201700424.

Synthesis and Immunological Evaluation of a Multicomponent Cancer Vaccine Candidate Containing a Long MUC1 Glycopeptide

Dr. Nitin T. Supekar^{[a],[b]}, Dr. Vani Lakshminarayanan^[c], Dr. Chantelle J. Capicciotti^[a], Dr. Anju Sirohiwal^[a], Cathy S. Madsen^[c], Dr. Margreet A. Wolfert^{[a],[d]}, Prof. Dr Peter A. Cohen^[c], Prof. Dr Sandra J. Gendler^[c], and Prof. Dr Geert-Jan Boons^{[a],[b],[d]}

^[a]Complex Carbohydrate Research Center, The University of Georgia, 315 Riverbend Road, Athens, Georgia 30602 (USA) ^[b]Department of Chemistry, The University of Georgia, 140 Cedar Street, Athens, Georgia 30602 (USA) ^[c]Departments of Biochemistry and Molecular Biology and Immunology, Mayo Clinic College of Medicine and Mayo Clinic Comprehensive Cancer Center, 13400 East Shea Boulevard, Scottsdale, AZ 85259 (USA) ^[d]Department of Chemical Biology and Drug Discovery, Utrecht Institute for Pharmaceutical Sciences and Bijvoet Center for Biomolecular Research, Utrecht University, Universiteitsweg 99, 3584 CG Utrecht (Netherlands)

Abstract

A fully synthetic MUC1-based cancer vaccine was designed and chemically synthesized containing an endogenous helper T-epitope (MHC class II epitope). The vaccine elicited robust IgG titers that could neutralize cancer cells by antibody-dependent cell-mediated cytotoxicity (ADCC). It also activated cytotoxic T-lymphocytes. Collectively, the immunological data demonstrate engagement of helper T-cells in immune activation. A synthetic methodology was developed for a penta-glycosylated MUC1 glycopeptide, and antisera of mice immunized by the new vaccine recognized such a structure. Previously reported fully synthetic MUC1-based cancer vaccines that elicited potent immune responses employed exogenous helper T-epitopes derived from microbes. It is the expectation that the use of the newly identified endogenous helper T-epitope will be more attractive, because it will activate cognate CD4⁺ T-cells that will provide critical tumor-specific help intratumorally during the effector stage of tumor rejection and will aid in the generation of sustained immunological memory.

Keywords

cancer; endogenous helper; glycopeptide; multicomponent vaccine; T-epitope

Correspondence to: Geert-Jan Boons.

Conflict of Interest

G.-J.B. is the founder of Viamune, a company that focuses on MUC1 cancer vaccine development.

Supporting information and the ORCID identification numbers for the authors of this article can be found under <https://doi.org/10.1002/cbic.201700424>.

During malignancy, biosynthesis of the mucin MUC1 is upregulated by as much as 50-fold, resulting in a loss of polarization.^[1] Furthermore, in tumor-associated MUC1, the variable number of tandem repeats (VNTR) of the extracellular domain, which harbor five potential sites of O-glycosylation, are underglycosylated, revealing peptide epitopes that are shielded by complex glycans in healthy tissue.^[2] The overexpression and aberrant glycosylation of MUC1 correlates with invasive cell growth and metastasis.^[3] It disrupts cell–cell adhesion, contributes to escape of tumor immune surveillance, and provides chemoresistance and anti-apoptotic properties. MUC1 is also an oncoprotein and can activate cell signaling events that contribute to cancer progression. Humoral, as well as cellular, immunity against tumor-associated MUC1 have been observed in cancer patients and correlate with favorable disease outcomes. These observations have created considerable interest in the development of immune-therapies that target tumor associated MUC1.^[4]

Previously, we reported a fully synthetic multicomponent vaccine composed of an aberrantly glycosylated MUC1 peptide, a peptide helper T-epitope derived from poliovirus, and the immunoadjuvant Pam₃CysSK₄ (**1**, Scheme 1), which can disrupt immune tolerance in human MUC1 transgenic mice.^[5] The candidate vaccine elicited robust IgG antibody responses that could neutralize cancer cells by antibody-dependent cell-mediated cytotoxicity (ADCC). It also activated cytotoxic T-lymphocytes (CTLs) and was efficacious in a mouse model for mammary cancer. Structure–activity relationship (SAR) studies showed that optimal immune responses were achieved when the three components were covalently linked and the B-epitope was presented at the C terminus of the glycolipopeptide. Furthermore, we found that an immunogen with the Pam₃CysSK₄ moiety replaced by CpG, which is a TLR9 agonist, elicited inferior immune responses.^[6] A fully synthetic multicomponent vaccine containing STn also elicited potent humoral and cellular immune responses.^[7] Several other multicomponent MUC1-based cancer vaccines that exhibit some of the properties of compound **1** have been described.^[8]

We envisaged that the long MUC1-derived glycopeptide, APGSTAPPAHGVTSAPDT(*O*-GalNAc)RPAP, having a number of unique properties, would offer a tantalizing antigen for cancer vaccine development. It was anticipated to contain multiple epitopes that can activate B-cells, cytotoxic T-lymphocytes, and helper T-cells. The glycopeptide SAPDT(*O*-GalNAc)RPAP is a well-established B-epitope that can elicit relevant humoral responses.^[9] It is also an MHC class I (Kb) epitope that can activate cytotoxic T-lymphocytes.^[10] Glycosylation of this peptide is important for eliciting antibodies and activating CD8⁺ cells that can neutralize MUC1-expressing cancer cells.^[5b] In addition, the peptides STAPPAHGV and PAHGVTSa are predicted MHC class I (Ab) epitopes.^[10] We analyzed the full-length tandem repeat of MUC1 by Rankpep,^[11] which indicated that the peptide STAPPAHGVTSa might function as a promiscuous helper T-epitope (class II). This was supported by the finding that MUC1-specific CD4⁺ T cells could be elicited by employing a MUC1-derived peptide of 100 amino acids or by tumor challenge.^[12] Although MHC class II epitopes derived from tumor antigens may not activate T-cell as effectively as exogenous epitopes such as the one derived from poliovirus used in vaccine **1**,^[5] they will induce cognate CD4⁺ T-cells that will provide important help during secondary immune responses.

A number of studies have confirmed the attractiveness of using self-epitopes for stimulation of CD4⁺ T-cell responses.^[13]

Vaccine candidate **2** was prepared to examine whether the MUC1-derived glycopeptide (APGSTAPPAHGVTSAPDT(*O*-Gal-NAc)RPAP) could activate CD4⁺ and CD8⁺ T-cells and induce IgG antibodies. The threonine moiety of the SAPDTRPAP epitope was modified by GalNAc, because previous studies have shown that glycosylation of this residue is important for eliciting tumor-relevant CTLs and antibodies.^[5b,14] The other serine and threonine moieties were not modified, because glycosylation can block proteolytic processing of the tandem repeat, which might compromise presentation of MHC1 and MHCII epitopes required for activation of CD4⁺ and CD8⁺ T-cells, respectively.^[15] In addition, the monoglycosylated glycopeptide **3** was prepared, which has an N-terminal cysteine moiety for conjugation to a carrier protein to evaluate the presence of serum antibodies by ELISA. A similar pentaglycosylated tandem repeat (**4**) was prepared to examine whether antibodies elicited by **2** could recognize highly glycosylated MUC1. In this respect, tumor-associated MUC1 is characterized by high-density glycosylation of the tandem repeat, and it is known that glycosylation of MUC1 epitopes can influence antibody recognition.^[16]

Synthesis of the monoglycosylated vaccine candidate **2** was straightforward and entailed coupling of the first four amino acids on a Rink amide AM LL resin (**5**) by using a CEM Liberty 12-channel automated microwave peptide synthesizer and 2-(1*H*-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU)/1-hydroxybenzotriazole (HOBt) as the activation protocol (Scheme 2 A). The glycosylated amino acid *N*-Fmoc-Thr(AcO₃-α-D-GalNAc (**7**) was introduced manually under microwave heating with 1-[dis(dimethyl amino)methylene]-1*H*-1,2,3-triazolo[4,5-*b*]pyridinium 3-oxid hexafluorophosphate (HATU)/1-hydroxy-7-azabenzotriazole (HOAt) as the activation reagent. The resulting resin, **8**, was elongated in the synthesizer under standard coupling conditions to give resin-bound glycopeptide **9**, which was treated with 60 % hydrazine in methanol to remove the acetyl esters^[5a] of GalNAc. Manually, Fmoc-Pam₂Cys (**10**) was coupled to the N terminus of **9** by using HATU/HOAt in the presence of *N,N*-diisopropylethylamine (DIPEA) in DMF under microwave heating, which was followed by removal of the Fmoc protecting group and further acylation of the resulting amine with palmitic acid to give **11**. The resin-bound glycolipopeptide was released from the resin by treatment with a mixture of TFA, phenol, H₂O, and TIPS (88:5:5:2, *v/v/v/v*), with simultaneous deprotection of the amino acid side chain protecting groups. Purification by HPLC on a reversed-phase C₄ column gave homogeneous **2**. A small amount of glycopeptide **9** was released from the resin prior to further coupling, and analysis by MS showed the absence of incomplete sequences. Glycopeptide **3** was prepared according to a similar coupling strategy (Scheme S1 in the Supporting Information).

Preparation of pentaglycosylated glycopeptide **4** was more challenging. Installation of the second glycosylated amino acid resulted in product **14** and by-product **15** (10–20 %), in which the N-terminal amino group of the starting peptide was acetylated (Scheme 2 B). Similar side reactions were observed during the coupling of the third, fourth, and fifth glycosylated amino acids, resulting in a very low yield of desired product. It is probable that the high reaction temperature employed in the microwave-assisted coupling step and the

longer reaction time caused acetyl migration from the glycosylated amino acids to the N-terminal amine of the starting peptide. This problem could be addressed by reducing the amount of glycosylated amino acid (1.1 equiv), lowering the reaction temperature from 75 to 60 °C, and shortening the reaction time (10 to 5 min) during the coupling steps of the glycosylated amino acids (Scheme S2). A Kaiser test was performed after coupling of each glycosylated amino acid, and unreacted amino groups were capped by using acetic anhydride. After assembly of the full tandem repeat, an N-terminal cysteine residue was introduced, and the resulting compound was deprotected and purified according to standard procedures to give compound **4** in good overall yield.

Glycopeptides **3** and **4**, containing an N-terminal cysteine residue, were conjugated to BSA-modified maleimides (BSA-MI) according to a standard protocol. The conjugates were purified by spin filtration, the coupling efficiency was determined by MALDI-TOF mass spectrometry (~ten copies of glycopeptide/ BSA), and protein concentration was determined by Bradford protein assay.

Humoral and cellular immune responses elicited by vaccine candidate **2** were explored in MUC1.Tg mice (C57BL/6; H-2^b) expressing human MUC1.^[17] A group of five mice were intradermally immunized with liposomal preparations of compound **2** four times at biweekly intervals. Pam₃CysSK₄ and empty liposomes were administered as negative controls. One week after the last immunization, the mice were sacrificed, and humoral and cellular immune responses were evaluated by examining titers of MUC1-specific antibodies, the ability of the antisera to lyse MUC1-bearing tumor cells, and ELISPOT assay with T-cells.

Mice immunized with compound **2** elicited robust IgG antibody titers compared to Pam₃CysSK₄ and empty liposomes. The elicited antibodies recognized both mono- and pentaglycosylated full-length MUC1 epitopes **3** and **4**, respectively (Table 1). IgG subtyping showed the presence of IgG1, IgG2a, and IgG2b antibodies, indicating a mixed Th1/Th2 response (IgG2a/b represents a Th1 response, and IgG1 represents a Th2 response). Very low titers of IgM antibodies were observed, demonstrating efficient class switching from low-affinity IgM to high-affinity IgG antibodies. These results indicated that the vaccine candidate embedded an MHCII epitope that could activate helper T-cells.

Only antisera obtained by immunization with **2** showed significant binding to C57mg mammary cancer cells transfected with MUC1 (Figure S2). Wild-type C57mg cells, known to lack MUC1 expression, did not show binding with antisera **2** (Figures S1 and S2). Antibody-dependent cell-mediated cytotoxicity (ADCC) was examined by labeling MUC1-expressing C57mg cells with ⁵¹Cr, followed by the addition of antisera and cytotoxic effector cells (NK cells) and measurement of released ⁵¹Cr. The antisera obtained by immunization with **2** significantly increased cancer cell lysis compared to the control groups immunized with empty liposomes or Pam₃CysSK₄ (Figure 1 A).

To assess the ability of **2** to activate CTLs, CD62L^{low} T-cells were isolated by magnetic cell sorting and incubated with dendritic cells (DCs) without in vitro stimulation and then analyzed for MUC1-specific IFN γ spot formation on ELISPOT plates. It was observed that vaccine candidate **2** had robustly activated T-cells compared to the control groups (Figure 1

B). The T-cell response was further evaluated for MUC1 epitope recognition of CD4⁺ and CD8⁺ T-cells by culturing T-cells derived from lymph nodes and expressing low levels of CD62L for seven days in the presence of DCs pulsed with the corresponding immunizing construct. The resulting cells were analyzed by intracellular cytokine (ICC) staining for the presence of CD4⁺ IFN γ ⁺ (Figure 1 C) and CD8⁺IFN γ ⁺ T-cells (Figure 1 D). As anticipated, both CD4⁺ and CD8⁺ T-cells derived from mice immunized with multicomponent vaccine **2** were responsive to the MUC1 epitope.

Collectively, our data demonstrated that vaccine candidate 2, which is devoid of any artificial linkers or exogenous helper T-epitopes, can activate T-cells and elicit robust cytotoxic IgG antibody responses. Previously, a number of synthetic MUC1 vaccines have been reported containing a full-length tandem repeating unit.^[8b,e,h,n,18] Although these vaccines were examined in wild-type mice that did not require breaking immune tolerance, they elicited relatively low titers of IgG antibodies. The modest immune activation was probably due to the fact that these vaccine candidates contain a MUC1 tandem repeat (e.g., AHGVT SAPDTRPAPGSTAPP) that does not embed the full putative helper T-epitope. In addition, some of these immunogens were glycosylated at all serine and threonine moieties of the MUC1 tandem repeat, which might compromise proteolytic processing to provide epitopes that can be presented by MHCI or MHCII.^[15,19] Linkers employed to connect the Pam₃CysSK₄ moiety with the MUC1 glycopeptide could also interfere with proteolytic processing. Thus, careful selection of the MUC1 tandem repeating unit and proper glycosylation are likely critical determinates of a multicomponent vaccine that can activate each arm of the immune system. It is the expectation that a MUC1-based vaccine that contains an endogenous helper T-epitope will promote induction of cognate CD4⁺ T-cells that could provide critical tumor-specific help intratumorally during the effector stage of tumor rejection and potentially aid in the generation of sustained immunological memory.^[13a,20] Future studies will focus on establishing the optimal glycosylation pattern of the MUC1 tandem repeating unit to elicit the most tumor-relevant immune responses.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by the National Cancer Institute of the US National Institutes of Health (R01CA88986 to G.-J.B.), the Mayo Breast Specialized Programs of Research Excellence (SPORE; P50CA116201 to S.J.G.) and the Mayo Pancreas SPORE (P50CA102701 to P.A.C. and S.J.G.). The Mayo Clinic Natalie Schafer Animal Care attendants are acknowledged for their animal care.

References

1. Apostolopoulos V, Stojanovska L, Gargosky SE. Cell Mol Life Sci. 2015; 72:4475–4500. [PubMed: 26294353]
2. a) Cascio S, Finn OJ. Biomolecules. 2016; 6:E39. [PubMed: 27754373] b) Hanson RL, Hollingsworth MA. Biomolecules. 2016; 6:E34. [PubMed: 27483328]
3. a) Deng J, Wang L, Chen H, Li L, Ma Y, Ni J, Li Y. Cancer Metastasis Rev. 2013; 32:535–551. [PubMed: 23609751] b) Nath S, Mukherjee P. Trends Mol Med. 2014; 20:332–342. [PubMed: 24667139]

4. a) Beatson RE, Taylor-Papadimitriou J, Burchell JM. Immunotherapy. 2010; 2:305–327. [PubMed: 20635898] b) Hossain MK, Wall KA. Vaccine. 2016; 4:E25.c) Roulois D, Gregoire M, Fonteneau JF. BioMed Res Int. 2013; 2013:871936. [PubMed: 23509794]
5. a) Ingale S, Wolfert MA, Gaekwad J, Buskas T, Boons GJ. Nat Chem Biol. 2007; 3:663–667. [PubMed: 17767155] b) Lakshminarayanan V, Thompson P, Wolfert MA, Buskas T, Bradley JM, Pathangey LB, Madsen CS, Cohen PA, Gendler SJ, Boons GJ. Proc Natl Acad Sci USA. 2012; 109:261–266. [PubMed: 22171012]
6. Abdel-Aal AB, Lakshminarayanan V, Thompson P, Supekar N, Bradley JM, Wolfert MA, Cohen PA, Gendler SJ, Boons GJ. ChemBioChem. 2014; 15:1508–1513. [PubMed: 24890740]
7. Thompson P, Lakshminarayanan V, Supekar NT, Bradley JM, Cohen PA, Wolfert MA, Gendler SJ, Boons GJ. Chem Commun. 2015; 51:10214–10217.
8. a) Renaudet O, BenMohamed L, Dasgupta G, Bettahi I, Dumy P. Chem-MedChem. 2008; 3:737–741.b) Kaiser A, Gaidzik N, Becker T, Menge C, Groh K, Cai H, Li YM, Gerlitzki B, Schmitt E, Kunz H. Angew Chem Int Ed. 2010; 49:3688–3692.Angew Chem. 2010; 122:3772–3776.c) Wilkinson BL, Malins LR, Chun CKY, Payne RJ. Chem Commun. 2010; 46:6249–6251.d) Cai H, Huang ZH, Shi L, Zhao YF, Kunz H, Li YM. Chem Eur J. 2011; 17:6396–6406. [PubMed: 21538615] e) Wilkinson BL, Day S, Malins LR, Apostolopoulos V, Payne RJ. Angew Chem Int Ed. 2011; 50:1635–1639.Angew Chem. 2011; 123:1673–1677.f) Abdel-Aal AB, El-Naggar D, Zaman M, Batzloff M, Toth I. J Med Chem. 2012; 55:6968–6974. [PubMed: 22800462] g) Cai H, Sun ZY, Huang ZH, Shi L, Zhao YF, Kunz H, Li YM. Chem Eur J. 2013; 19:1962–1970. [PubMed: 23280874] h) Sarkar S, Salyer AC, Wall KA, Sucheck SJ. Bioconjugate Chem. 2013; 24:363–375.i) Cai H, Sun ZY, Chen MS, Zhao YF, Kunz H, Li YM. Angew Chem Int Ed. 2014; 53:1699–1703.Angew Chem. 2014; 126:1725–1729.j) McDonald DM, Wilkinson BL, Corcilius L, Thaysen-Andersen M, Byrne SN, Payne RJ. Chem Commun. 2014; 50:10273–10276.k) Palitzsch B, Hartmann S, Stergiou N, Glaffig M, Schmitt E, Kunz H. Angew Chem Int Ed. 2014; 53:14245–14249.Angew Chem. 2014; 126:14469–14473.l) Zhou Z, Mondal M, Liao G, Guo Z. Org Biomol Chem. 2014; 12:3238–3245. [PubMed: 24728423] m) McDonald DM, Byrne SN, Payne RJ. Front Chem. 2015; 3:60. [PubMed: 26557640] n) Karmakar P, Lee K, Sarkar S, Wall KA, Sucheck SJ. Bioconjugate Chem. 2016; 27:110–120.o) Liu Y, Zhang W, He Q, Yu F, Song T, Liu T, Zhang Z, Zhou J, Wang PG, Zhao W. Chem Commun. 2016; 52:10886–10889.
9. a) von Mensdorff-Pouilly S, Petrakou E, Kenemans P, van Uffelen K, Verstraeten AA, Snijdwint FG, van Kamp GJ, Schol DJ, Reis CA, Price MR, Livingston PO, Hilgers J. Int J Cancer. 2000; 86:702–712. [PubMed: 10797294] b) Graves CRL, Robertson JFR, Murray A, Price MR, Chapman CJ. J Peptide Res. 2005; 66:357–363. [PubMed: 16316451]
10. a) Apostolopoulos V, Haurum JS, McKenzie IFC. Eur J Immunol. 1997; 27:2579–2587. [PubMed: 9368613] b) Apostolopoulos V, Karanikas V, Haurum JS, McKenzie IFC. J Immunol. 1997; 159:5211–5218. [PubMed: 9548459]
11. Reche PA, Glutting JP, Zhang H, Reinherz EL. Immunogenetics. 2004; 56:405–419. [PubMed: 15349703]
12. a) Hiltbold EM, Ciborowski P, Finn OJ. Cancer Res. 1998; 58:5066–5070. [PubMed: 9823312] b) Tempero RM, VanLith ML, Morikane K, Rowse GJ, Gendler SJ, Hollingsworth MA. J Immunol. 1998; 161:5500–5506. [PubMed: 9820526]
13. a) Knutson KL, Disis ML. Cancer Immunol Immunother. 2005; 54:721–728. [PubMed: 16010587] b) Kennedy R, Celis E. Immunol Rev. 2008; 222:129–144. [PubMed: 18363998]
14. a) Coltart DM, Royyuru AK, Williams LJ, Glunz PW, Sames D, Kuduk SD, Schwarz JB, Chen XT, Danishefsky SJ, Live DH. J Am Chem Soc. 2002; 124:9833–9844. [PubMed: 12175243] b) Dziadek S, Griesinger C, Kunz H, Reinscheid UM. Chem Eur J. 2006; 12:4981–4993. [PubMed: 16642533]
15. Ninkovic T, Hanisch FG. J Immunol. 2007; 179:2380–2388. [PubMed: 17675499]
16. Rangappa S, Artigas G, Miyoshi R, Yokoi Y, Hayakawa S, Garcia-Martin F, Hinou H, Nishimura SI. Med Chem Comm. 2016; 7:1102–1122.
17. Rowse GJ, Tempero RM, VanLith ML, Hollingsworth MA, Gendler SJ. Cancer Res. 1998; 58:315–321. [PubMed: 9443411]

18. a) Liu YF, Sun ZY, Chen PG, Huang ZH, Gao Y, Shi L, Zhao YF, Chen YX, Li YM. *Bioconjugate Chem.* 2015; 26:1439–1442. b) Sun ZY, Chen PG, Liu YF, Zhang BD, Wu JJ, Chen YX, Zhao YF, Li YM. *Chem Commun.* 2016; 52:7572–7575.
19. Madsen CB, Petersen C, Lavrsen K, Harndahl M, Buus S, Clausen H, Pedersen AE, Wandall HH. *PLoS One.* 2012; 7:e50139. [PubMed: 23189185]
20. Shedlock DJ, Shen H. *Science.* 2003; 300:337–339. [PubMed: 12690201]

Figure 1.

A) Induction of ADCC with C57mg.MUC1 tumor cells. B) Induction of cytotoxic CD62L^{low} T-cell response by analyzing MUC1-specific IFN γ spot formation without in vitro stimulation. Each data point represents an individual mouse, and horizontal lines indicate the mean for the group of mice. ICC staining for the presence of C) CD4⁺IFN γ ⁺ and D) CD8⁺IFN γ ⁺ T-cells. See Figure S3 for representative flow cytometry plots. Data are shown as mean $\pm$ SEM ($n=3$) for vaccine candidate 2 (Pam₃CysSK₄ APGSTAPPAHGVT-SAPDT(O-GalNAc)RPAP) and controls Pam₃CysSK₄ and empty liposomes (EL).

Scheme 1.

Glycopeptides and glycolipopeptides for immunization studies.

Scheme 2.

A) Chemical synthesis of glycolipopeptide **2** and B) chain termination during synthesis of glycopeptide **4**. a) 20 % 4-methylpiperidine, DMF, MW, 3 min; b) Fmoc-AA-OH, HOBt, HBTU, DIPEA, DMF, MW, 5 min; c) FmocNH-T*-COOH (**7**; 2 equiv), HATU, HOAt, DIPEA, DMF, MW, 10 min; d) 70 % hydrazine in MeOH; e) Fmoc-Pam₂Cys-OH (**10**), HATU, HOAt, DIPEA, DMF, MW, 10 min; f) palmitic acid, HATU, HOAt, DIPEA, DMF, MW, 10 min; g) TFA/phenol/H₂O/TIPS (88:5:5:2); h) FmocNH-S*-COOH (**13**; 2 equiv), HATU, HOAt, DIPEA, DMF, MW, 10 min.

Table 1

ELISA anti-LMUC1 antibody titers in endpoint serum samples.

	IgG total	IgG1	IgG2a	IgG2b	IgG3	IgM
BSA-MI-3 ^[a]						
2	23 200	29 200	1200	3600	5800	300
Pam ₃ CysSK ₄	300	0	0	0	0	0
EL	2000	600	300	800	500	0
BSA-MI-4 ^[b]						
2	21 700	29 900	2000	4800	3500	500
Pam ₃ CysSK ₄	0	0	0	0	0	100
EL	1200	500	600	400	300	0

Antibody titers are presented as median values for groups of mice. ELISA plates were coated with [a] BSA-MI-CAPGSTAPPAGHGVTSAPDT (αGalNAc)R-PAP (BSA-MI-3) conjugate for anti-LMUC1 (Tn18) antibody titers or [b] BSA-MI-CAPGSTAPPAGHGVTSAPDT (αGalNAc)S(αGalNAc)APDT-(αGalNAc)RPAP (BSA-MI-4) conjugate for anti-LMUC1 (5Tn) antibody titers. EL: empty liposomes.