HHS Public Access

Author manuscript

Conf Rec Asilomar Conf Signals Syst Comput. Author manuscript; available in PMC 2018 June 05.

Published in final edited form as:

Conf Rec Asilomar Conf Signals Syst Comput. 2016 November ; 2016: 1066–1073. doi:10.1109/ACSSC.

2016.7869533

Neuronal Network Models for Sensory Discrimination

Mohammad Samavat,

School of Electrical and Computer and Energy Engineering Arizona State University

Dori Luli, and

School of Human Evolution and Social Change Arizona State University

Sharon Crook

School of Mathematical and Statistical Sciences School of Life Sciences Arizona State University

Abstract

Previous modeling studies have demonstrated that lateral inhibition contributes to enhanced precision in sensory networks. That is, inhibitory connections reduce the spread of activity and repress neighboring cells, increasing the reliability of a sensory response. However, much less is understood about how connections that spread activity might contribute to the processing of sensory stimuli in the context of a sensory discrimination task. In this work, we examine the role of excitatory connections and gap junctions in network dynamics and some contributions to sensory discrimination.

I. Introduction

Sensory discrimination provides the ability to interpret sensory information. In this work, the fruit fly, *Drosophila melanogaster*, is used as a model organism. In the olfactory system of *Drosophila*, sensory transduction activates olfactory receptor neurons (ORNs) housed in the antennae and maxillary palps. These sensory neurons provide input to the antennal lobe (AL), where the structural unit is the glomerulus. Each glomerulus is associated with at least three classes of neurons: excitatory local neurons (eLNs), inhibitory local neurons (iLNs), and projection neurons (PNs). PNs provide the output from the antennal lobe, sending axon terminals to the mushroom bodies and lateral horn, while the local neuron arborizations are confined to the antennal lobe.

In this work, we construct a conductance-based neuronal network model of the *Drosophila* antennal lobe, based on experimental data for channel kinetics and constrained by specific odor stimulus/response characteristics described by Bhandawat et al. (1). We use computational studies to investigate the role of interglomerular and intraglomerular connectivity patterns in the antennal lobe, where ORN inputs to the antennal lobe are modeled with Poisson spike trains, and model synaptic connections are mediated by chemical synapses and gap junctions as described in the antennal lobe literature. We find that in the case of homogeneous synaptic connection strengths, the model suggests that the eLN network facilitates odor detection in the presence of weak stimuli. Excitatory local neurons can spread excitation from PNs, amplifying weak inputs. However, depending on connectivity strength, eLNs may decrease the ability of the network to discriminate between

odors. In addition, various electrical connectivity regimes are explored to elucidate the role of gap junctions in the dynamical behavior of the antennal lobe network.

II. Formalism for Conductance Based Models

Electrical activity in neurons is sustained and propagated via ionic currents through neuron membranes. The Hodgkin-Huxley model formalism for an excitable membrane (2) is one of the most important models in describing the kinetics of voltage dependent currents. As summarized by (3), an ionic current I_{ion} is modeled as

$$I_{ion} = g_{ion}p(V - E_{ion}), \quad (1)$$

where V is the cell membrane potential (in mV), g_{ion} is the maximal conductance for the channel type (in mS/cm^2), p is the average proportion of channels in the open state, and E_{ion} is the reversal (Nernst) potential for the corresponding active ion (in mV). The proportion of open channels for voltage gated channels can be represented by a product of activation and inactivation

$$p = m^y h^z, \quad (2)$$

where m and h are the activation and inactivation gating variables, respectively, and y and z determine the influence of the gating processes on the conductance. In some cases, only an activation variable is needed. The dynamics of the activation variable m is described by the equation

$$\frac{dm}{dt} = \frac{m_{\infty}(V) - m}{\tau_m(V)}, \quad (3)$$

where $m_{\infty}(V)$ is the steady-state activation function and has the form

$$m_{\infty}(V) = \frac{\alpha_m(V)}{\alpha_m(V) + \beta_m(V)}, \quad (4)$$

and $\alpha_m(V)$ and $\beta_m(V)$ describe the voltage dependent rates between open and closed states of the channels. The steady state activation function can be approximated by the Boltzmann function

$$m_{\infty}(V) = \frac{1}{1 + \exp[(V - \theta_m)/k_m]}, \quad (5)$$

where θ_m is the half activation voltage for the gating function (in mV), and k_m is the activation sensitivity (in mV). $\tau_m(V)$ determines the time scale of the exponential decay or saturation of the dynamic gating processes and has the form

$$\tau_m(V) = \frac{1}{\alpha_m + \beta_m}, \quad (6)$$

and can often be approximated by the Gaussian function

$$\tau_m(V) = C_{base} + C_{amp} \exp \frac{-(V_{max} - V)^2}{\sigma^2}, \quad (7)$$

where the function values lie between C_{base} and C_{amp} . V_{max} is the potential where the maximal τ_m value is achieved, and σ measures the characteristic width of the curve. Av-Ron and Rospars (4) express τ_w as

$$\tau_w(V) = \frac{1}{r_w \cosh\left(\frac{V - \theta_w}{k_{\tau_w}}\right)}, \quad (8)$$

where r_w is the basal rate of the gating reaction. In some cases when a gating variable, say m , approaches the steady state very quickly, we can replace the gating variable with its steady state function, $m_{\infty}(V)$. The dynamics of the inactivation variable h is described in a similar way to the activation variable m .

Using currents modeled as discussed above, the current balance equation for the excitable membrane is

$$C_m \frac{dV}{dt} = - \sum_{ion} I_{ion} - g_L(V - E_L) - I_{syn} + I_{app}, \quad (9)$$

where C_m denotes the membrane capacitance (in $\mu F/cm^2$), I_{syn} is the synaptic current if present, $g_L(V - E_L)$ describes the leak current, and I_{app} (in $\mu A/cm^2$) simulates a current injection that mimics the application of injected current in an experiment.

A. Calcium Gated Channels

While the original Hodgkin-Huxley model characterizes voltage-gated ion channels, there are many types of ion channels that are controlled by factors other than membrane potential, including ligand-gated channels, which vary their conductance in response to changes in the intracellular concentration of some other molecule. The most common signaling molecule is calcium, and a widely-used simple model for the changes in intracellular calcium concentration assumes that the concentration increases due to an inward flux of calcium ions through calcium ion channels when calcium currents are activated and that a linear pumping process through the plasma membrane removes calcium ions from the cell (5). The total calcium in a cell or cellular compartment consists of free calcium and calcium bound to a buffer. Thus, the equation that represents the dynamics of the free calcium concentration $[Ca]$ (in μM) is

$$\frac{d}{dt}[Ca] = \lambda(-\alpha_{Ca}I_{Ca} - r[Ca]), \quad (10)$$

where λ is the percent of free to bound calcium, α_{Ca} converts the ionic current, I_{Ca} , to a rate of change in calcium concentration, and r is the removal rate constant due to calcium pumps. In the model, the conductance due to calcium dependent channels is modeled as an instantaneous function of $[Ca]$ using a Hill equation in the standard manner.

B. Synapse Models

The synaptic current affects the electrical activity of the postsynaptic cell. Here faster synaptic currents, such as fast $GABA_A$ and acetylcholine, are governed by the equation

$$I_{fast} = g_{syn}[O](V - E_{syn}), \quad (11)$$

where $[O]$ is the fraction of open channels (6). Following the arrival of an action potential at the presynaptic terminal, neurotransmitter molecules are released into the synaptic cleft. These molecules are taken to bind to postsynaptic receptors. $[O]$ is calculated according to the equation

$$\frac{d[O]}{dt} = \alpha(1 - [O])[T] - \beta[O], \quad (12)$$

where α and β are the forward and backward rate constants for transmitter binding. $[T]$ is the concentration of transmitter and is given by

$$[T] = \frac{1}{1 + \exp[(V - \theta_T)/k_T]}, \quad (13)$$

as described by Bazhenov et al. (7).

Slower synaptic current receptors have a more complex scheme of activation. The activation of potassium channels by G proteins is involved in the case of $GABA_B$ receptors activation. The slower inhibitory synaptic current is given by the equation

$$I_{slow} = g_{syn} \frac{[G]^4}{[G]^4 + K} (V - E_K), \quad (14)$$

where $[G]$ is the concentration of G proteins determined by the equation

$$\frac{d[G]}{dt} = k_3[R] - k_4[G], \quad (15)$$

and K is the dissociation rate (6). Here $[R]$ is the fraction of activated receptors determined by the equation

$$\frac{d[R]}{dt} = k_1(1 - [R])[T] - k_2[R], \quad (16)$$

and k_1 , k_2 , k_3 , and k_4 are kinetic rate constants.

The gap junction synaptic current is described as

$$I_{gap} = g_c(V_{pre} - V_{post}), \quad (17)$$

where g_c is the coupling strength, and V_{pre} and V_{post} represent the membrane potentials of the presynaptic and postsynaptic model cells respectively.

A synaptic mechanism which consists of a postsynaptic conductance which changes as an alpha function over time is described as

$$g(t) = g_{syn} \frac{t - t_{sp}}{t_{peak}} \exp(1 - \frac{t - t_{sp}}{t_{peak}}), \quad (18)$$

where t_{sp} is time of previous spike, and t_{peak} controls both the decay and rise time.

In the case of depression of a synapse, the synaptic current is multiplied by a scaling factor d , where d is determined by the equation

$$d = 1 - sd \exp\left(-\frac{(t - t_{sp})}{\tau_d}\right). \quad (19)$$

Here, sd is the decrease in conductance strength, and τ_d is the time constant for recovery of d due to a presynaptic event at time t_{sp} .

III. MATHEMATICAL MODELS FOR *DROSOPHILA* ANTENNAL LOBE

Here we develop realistic minimal conductance based models for the iLN, eLN, and PN, where the models follow the formalism presented in Section II, and model currents are based on experimental data from *Drosophila* neurons (8). All of the computational studies that follow for both single cell models and network models rely on custom numerical routines written in MATLAB. Systems of differential equations are solved using MATLAB solver ode15s with relative error tolerance of 1e-6 and absolute error tolerance of 1e-6. ode15s is a variable order solver based on Gear's method.

A. Inhibitory Local Neuron Model

iLN behavior is represented by a two dimensional system of differential equations. The two variables are the membrane potential V_{iLN} and the recovery variable w_{iLN} , which represents the activation of the potassium current. There are four types of currents in this model, an inward sodium current I_{Na} , an outward potassium current I_K , a leak current I_L , and an applied current I_{app} . Based on studies described in Section I, the minimal model for the iLN is constructed so that it exhibits regular repetitive firing in the presence of I_{app} .

The iLN current balance and recovery differential equations are

$$C_m \frac{dV_{iLN}}{dt} = -I_{Na} - I_K - I_L + I_{app}, \quad (20)$$

$$\frac{dw_{iLN}}{dt} = \frac{w_{\infty}(V_{iLN}) - w_{iLN}}{\tau_w(V_{iLN})}. \quad (21)$$

The time varying currents are

$$I_{Na} = g_{Na} m_{\infty}^3(V_{iLN})(1 - w_{iLN})(V_{iLN} - E_{Na}), \quad (22)$$

$$I_K = g_{K_{iLN}} w_{iLN}^4 (V_{iLN} - E_K) \quad (23)$$

$$I_L = g_L (V_{iLN} - E_L). \quad (24)$$

Note that inactivation of sodium is expressed using $1 - w_{iLN}$. The steady state functions are

$$m_{\infty}(V_{iLN}) = \frac{1}{1 + \alpha_m^2(V_{iLN})}, \quad (25)$$

$$w_{\infty}(V_{iLN}) = \frac{1}{1 + \alpha_w^2(V_{iLN})}, \quad (26)$$

where

$$\alpha_m(V_{iLN}) = \exp\left(\frac{V_{iLN} - \theta_m}{k_m}\right), \quad (27)$$

$$\alpha_w(V_{iLN}) = \exp\left(\frac{V_{iLN} - \theta_w}{k_w}\right), \quad (28)$$

$$\tau_w(V_{iLN}) = \frac{1}{r_w \cosh\left(\frac{V_{iLN} - \theta_w}{k_{\tau}}\right)}. \quad (29)$$

The parameters of this model are given in Table I. Figure 1 Panel A depicts the firing behavior of the iLN when $I_{app}=0.2$ nA, where the iLN exhibits regular repetitive firing. Figure 1 Panel B shows the iLN frequency-current (F-I) curve, where the iLN firing frequency increases as the amount of applied current increases.

B. Excitatory Local Neuron Model

Based on experimental studies reviewed in (9), the minimal model for the eLN model should exhibit bursting for some values of I_{app} . Therefore, we add a calcium current and a calcium dependent potassium current to the model presented in Section III-A. The eLN current balance equation is

$$C_m \frac{dV_{eLN}}{dt} = -I_{Na} - I_K - I_{Ca} - I_{KCa} - I_L + I_{app}. \quad (30)$$

The sodium, potassium, and leak currents as well as the recovery equation are the same as in the iLN model. The added calcium and calcium dependent potassium currents are represented by

$$I_{Ca} = g_{Ca} n_{\infty}^2 (1 - w_{eLN}) (V_{eLN} - E_{Ca}), \quad (31)$$

$$I_{KCa} = g_{KCa} \frac{[Ca]}{0.04 + [Ca]} (V_{eLN} - E_K), \quad (32)$$

where

$$w_{Ca_{\infty}}(V_{eLN}) = \frac{1}{1 + \exp\left(\frac{V_{eLN} - \theta_{Ca}}{k_{Ca}}\right)}, \quad (33)$$

$$\frac{d[Ca]}{dt} = -\alpha_{Ca} I_{Ca} - r[Ca]. \quad (34)$$

The additional parameters associated with the eLN model are also shown in Table I. Figure 2 Panel A depicts the firing behavior of the eLN when $I_{app} = 0.2$ nA, Panel B depicts the calcium dynamics, and Panel C depicts the time course of the calcium dependent potassium current. The eLN exhibits bursting for low values of I_{app} . The calcium channel is activated for the duration of the burst. The intracellular calcium concentration increases during the burst, resulting in an outward calcium dependent potassium current, and upon sufficient activation, repetitive firing can't be sustained and the quiescence phase commences. During quiescence, calcium is removed, closing the calcium dependent potassium channels, and upon sufficient removal of calcium and closing of calcium dependent potassium channels, bursting occurs again. Figure 3 Panel A depicts the firing behavior of the eLN when $I_{app} = 0.3$ nA, Panel B depicts the calcium dynamics, and Panel C depicts the time course of the

calcium dependent potassium current. The eLN exhibits repetitive firing for higher values of I_{app} , where the calcium and calcium dependent potassium channels remain activated, but the hyperpolarizing current is not large enough to cause quiescence in the presence of this stronger depolarization.

C. Projection Neuron Model

The PN model is almost identical to the iLN model where the only change is in the value of the maximal conductance of potassium current, which is given in Table I. Based on studies described in Section I, the minimal model for the PN exhibits regular repetitive firing but of higher frequency than the iLN. Similar to the iLN model, PN firing frequency increases as the amount of applied current increases.

D. Network Model

In an effort to build a realistic minimal network model of the *Drosophila* antennal lobe, we describe a network model composed of iLNs, eLNs, and PNs, with the synaptic connections among cell types motivated by experimental studies that are summarized in Table 4.3 in (9). We model seven glomeruli, where each glomerulus is associated with one PN, one iLNs and one eLN. eLN, iLN, PN, and synaptic models used for the network model are described in Section II-B. When we include synaptic inputs, the current balance equations for the PN, iLN, and eLN in the *Drosophila* antennal lobe network model become

$$C_m \frac{dV_{PN_j}}{dt} = -I_{Na} - I_K - I_L - \sum_{k=1}^n I_{gap\_eLN_k} - \sum_{k=1}^n I_{GABA_{A-iLN_k}} - \sum_{k=1}^n I_{GABA_{B-iLN_k}} \quad (35)$$

$$- I_{ORN},$$

$$C_m \frac{dV_{iLN_j}}{dt} = -I_{Na} - I_K - I_L - \sum_{k=1}^n I_{gap\_eLN_k} - \sum_{k=1}^n I_{ACh\_eLN_k} - \sum_{k=1}^n I_{ACh\_PN_k} - I_{ORN},$$

(36)

$$\begin{aligned}
C_m \frac{dV_{eLN_j}}{dt} = & -I_{Na} - I_K - I_{Ca} - I_{KCa} - I_L - \sum_{s=1}^n I_{gap\_eLN_s} \quad (37) \\
& - \sum_{k=1}^n I_{ACh\_PN_k} - \sum_{k=1}^n I_{gap\_PN_k} - \sum_{k=1}^n I_{GABA\_A-iLN_k} \\
& - \sum_{k=1}^n I_{GABA\_B-iLN_k} - \sum_{k=1}^n I_{gap\_iLN_k} - I_{ORN},
\end{aligned}$$

where j is the cell number, n is the number of glomeruli in the network, and $s = j$. Here, I_{ORN} is determined by the Poisson spike train input as described in Section III-E. Parameter values for the network model are provided in Table I and Table II.

E. Input for the Network Model

Bhandawat et al. (1) perform electrophysiological recordings from *Drosophila* olfactory receptor neurons and projection neurons. An odor stimulus is presented in multiple trials while recording from the same cells. Spike counts are quantified for ORNs and PNs, and response profiles are obtained for different odors from seven ORN types and seven PN types corresponding to the same glomeruli. These response profiles include peri-stimulus time histograms for each odor and glomerulus, averaged across representations of a particular odor stimulus lasting 500 ms. In addition, responses are given where they are averaged over the entire 500 ms odor stimulus.

Many mathematical models express input from the ORNs as a constant current (10; 11; 12). Here, we model input from ORNs in a more realistic way using Poisson spike trains, where driving rates for spontaneous activity and during each odor for the input from ORNs to the AL are derived from the experimental data described by Bhandawat et al., where the rates vary for different odors and different glomeruli. The Poisson process is modeled such that the firing rate is time dependent and has an absolute refractory period of 3 ms (13). As an example, Figure 4 depicts model ORN responses during odor presentation and spontaneous activity as Poisson spike trains for the odor benzaldehyde. These model olfactory receptor neuron responses provide inputs to seven different glomeruli (DL1, DM1, DM2, DM3, DM4, VA2, and VM2). Simulations are shown for one trial, where the odor stimulus is presented from 500 ms to 1000 ms.

F. Data Analysis of Simulation Results

Spiking activity of ORN inputs and all model cells in the network can be visualized with raster plots, where spike times are indicated with a dot or short line for each cell type. Due to the large number of variables and therefore high dimensionality for model results, we also perform principal component analysis (PCA) on PN output. This allows for a lower dimensional visualization of the output of the AL network model. PCA is used for dimensionality reduction to project data on to a space with less dimensions while minimizing the mean-square error of the distance between the original vectors and their projections on to the principal components. In results shown here, we simulate the

application of a single odor across 10 trials, where each simulated trial corresponds to a different set of generated ORN Poisson spike train inputs. For each trial, spike times for all PNs are determined and used to obtain the firing rates in Hz of each PN for a particular odor every 10 ms over the course of a simulation.

IV. Results

In order to compare simulations with the experimental literature, we generate Poisson spike trains to represent input from olfactory receptor neurons, where ten trials are run for different odors (such as benzaldehyde, butyric acid, 1-butanol, and cyclohexanone). In this case we can examine average firing rates across trials for the model. Figure 5 depicts the averages of firing rates across ten trials for model neurons for the odor benzaldehyde. Odor stimulus is presented from 500 ms to 1000 ms. ORN activity and PN output activity compare favorably to experimental results from Bhandawat et al. (1). Note that the inclusion of eLNs allows for spread of activity to additional glomeruli.

Additional simulations with artificially generated ORN inputs allow us to examine network dynamics more carefully under various conditions. In the raster plot shown in Figure 6, the horizontal axis corresponds to time, and the vertical axis shows the spike times grouped for the four cell types. Within each cell type, each row corresponds to one of the seven model glomeruli. Ten ORN inputs are generated for each glomerulus so that each model cell in that glomerulus can integrate the ten inputs for a more realistic input profile. Results are for a simulated ramp-type input, where the odor increases in strength across time during the entire simulation. Simulated ORN inputs show increased activity over time for the ORNs that project to the glomerulus that favors that odor. Note increased PN firing rates in the glomerulus that responds to that odor. Increased ORN input and PN activity also recruits more iLN activity globally, suppressing activity in other glomeruli as activity increases.

We also examine network results for different values of gap junction conductance. Figure 7 summarizes these results in a PCA plot. For a very low value of the gap junction conductance ($g_{gap} = 0.001$), the gap junctions have no effect on network dynamics. The PCA trajectory depicts a rapid divergence from background spontaneous activity to form a small transient during odor evoked activity, and then following the odor offset, the activity returns to the background spontaneous activity. As the conductance is increased ($g_{gap} = 0.01$), the variance in the first PCA eigenvalue decreases, but the variance increases across the other eigenvalues, indicating more complexity in the spatiotemporal pattern of activity in the network. As the conductance is further increased ($g_{gap} = 0.05$), the PCA trajectory exhibits larger transient activity during the odor representation and increased variability in the first component. However, the variability is greatly decreased in the other components. This result corresponds to strong synchrony across the network due to the strengthened gap junction coupling.

Lastly, we perform network simulations for different odors under different network connectivity profiles. Figure 8 depicts the projections of the simulated responses of seven different glomeruli for an AL network with only iLNs (black curves) and for a model network with both iLNs and eLNs (red curves) for odors benzaldehyde, 1-butanol,

cyclohexanone, and butyric acid. Note the different temporal patterns across different glomeruli for different odors. Across all panels in Figure 8, we observe a rapid divergence from background spontaneous activity to form a transient during odor evoked activity, and following the odor offset the activity returns to the background spontaneous activity. Also, note the presence of longer transients in the network model with only iLNs compared to the network model when adding eLNs. These results suggest that eLNs decrease the ability of the network to discriminate among single odors.

V. Conclusion

In this paper, as a first step in building a realistic minimal network model of the *Drosophila* antennal lobe, we described a network model composed of iLNs, eLNs, and PNs, with the synaptic connections among cell types motivated by experimental studies. We modeled seven glomeruli, where each glomerulus is associated with one PN, one iLN, and one eLN. In this paper, the synapse models include mechanisms of chemical synapses and gap junctions.

Many local neurons innervate most or all glomeruli (14). Experimental studies show that there is a uniform global lateral inhibition in the *Drosophila* antennal lobe (15). We examined the role of this global network of inhibitory local neurons in shaping projection neuron output in Figures 6 and 8. Our simulation results show that inhibitory local neurons spread inhibition among all glomeruli and for all odors. Projection neuron responses are decreased relatively uniformly when lateral excitatory connections are spatially widespread and homogeneous in strength as reported in the literature (16). We investigated the role of excitatory local neurons in our network model in shaping glomeruli output. Our simulation results show that excitatory local neurons spread excitation among glomeruli for all odors, which is what one might expect with homogeneous connection strengths. Additionally, we show that gap junctions can spread excitation across glomeruli and regulate synchrony.

In further work, we will expand the number of cells modeled so that we can include inhomogeneity in connections. These more realistic heterogeneous connections are needed to understand the complex influences among glomeruli and will allow us to include both local and global inhibition in the network model.

Acknowledgments

The authors would like to thank Reuben Haynes for his help with data analysis and feedback on the project. This work was funded in part by NIH R01MH1006674 to SMC.

References

1. Bhandawat V, Olsen SR, Gouwens NW, Schlieff ML, Wilson RI. Sensory processing in the *Drosophila* antennal lobe increases reliability and separability of ensemble odor representations. *Nat Neurosci*. Nov; 2007 10(11):1474–82. [PubMed: 17922008]
2. Hodgkin AL, Huxley AF. A quantitative description of membrane current and its application to conduction and excitation in nerve. *J Physiol*. Aug; 1952 117(4):500–44. [PubMed: 12991237]
3. Izhikevich, EM. *Dynamical systems in Neuroscience: The geometry of excitability and bursting*. The MIT Press; 2007.

4. Av-Ron E, Parnas H, Segel LA. A basic biophysical model for bursting neurons. *Biol Cybern.* 1993; 69(1):87–95. [PubMed: 8334193]
5. Fall, CP., Marland, ES., Wagner, JM., Tyson, JJ. *Computational Cell Biology.* Springer; 2002.
6. Destexhe A, Bal T, McCormick DA, Sejnowski TJ. Ionic mechanisms underlying synchronized oscillations and propagating waves in a model of ferret thalamic slices. *J Neurophysiol.* Sep; 1996 76(3):2049–2070. [PubMed: 8890314]
7. Bazhenov M, Stopfer M, Rabinovich M, Abarbanel HD, Sejnowski TJ, Laurent G. Model of cellular and network mechanisms for odor-evoked temporal patterning in the locust antennal lobe. *Neuron.* May; 2001 30(2):569–81. [PubMed: 11395015]
8. Herrera-Valdez M, McKiernan E, Berger S, Ryglewski S, Duch C, Crook S. Relating ion channel expression, bifurcation structure, and diverse firing patterns in a model of an identified motor neuron. *Journal of Computational Neuroscience.* 2012
9. Luli, D. PhD dissertation. Arizona State University; 2013. A neuronal network model of drosophila antennal lobe.
10. Av-Ron E, Rospars JP. Modeling insect olfactory neuron signaling by a network utilizing disinhibition. *Biosystems.* 1995; 36(2):101–8. [PubMed: 8573691]
11. Bazhenov M, Stopfer M, Rabinovich M, Huerta R, Abarbanel HD, Sejnowski TJ, Laurent G. Model of transient oscillatory synchronization in the locust antennal lobe. *Neuron.* May; 2001 30(2):553–67. [PubMed: 11395014]
12. Martinez D, Montejo N. A model of stimulus-specific neural assemblies in the insect antennal lobe. *PLoS Comput Biol.* 2008; 4(8):e1000139. [PubMed: 18795147]
13. Dayan, P., Abbott, LF. *Theoretical Neuroscience.* The MIT Press; 2001.
14. Wilson R. Understanding the functional consequences of synaptic specialization: insight from the drosophila antennal lobe. *Curr Opin Neurobiol.* Mar.2011
15. Silbering AF, Galizia CG. Processing of odor mixtures in the drosophila antennal lobe reveals both global inhibition and glomerulus-specific interactions. *J Neurosci.* Oct; 2007 27(44):11 966–77.
16. Wilson RI. Eppendorf 2007 winner. neural circuits underlying chemical perception. *Science.* Oct; 2007 318(5850):584–585. [PubMed: 17962548]

Fig. 1. iLN firing behavior. Panel A: iLN membrane potential when a constant current, $I_{app} = 0.2$ nA, is applied. The iLN exhibits regular repetitive firing. Panel B: iLN F-I Curve. The iLN firing frequency increases as the amount of applied current increases.

Fig. 2.

eLN firing behavior and calcium concentration. Panel A: eLN membrane potential when $I_{app} = 0.2$ nA. The eLN exhibits bursting for low values of I_{app} . Panel B: The intracellular calcium concentration increases during the burst which eventually brings about the quiescence phase. During quiescence calcium is removed and upon sufficient removal, bursting occurs again. Panel C: Calcium dependent potassium current. During firing, the increase in intracellular calcium turns on I_{KCa} , upon sufficient activation, the active phase terminates. During quiescence, the decrease in intracellular calcium concentration closes potassium dependent calcium channels. Firing resumes when enough KCa channels are closed.

Fig. 3. eLN firing behavior and calcium concentration. Panel A: eLN membrane potential when $I_{app} = 0.3$ nA. The eLN exhibits repetitive firing for higher values of I_{app} . Panel B: Calcium dynamics. In the presence of enough applied current, calcium channels remain activated. Panel C: Calcium dependent potassium current. In the presence of enough applied current, calcium dependent potassium channels remain activated, but does not cause quiescence.

Fig. 4.

Spike times generated for model ORNs during odor and spontaneous activity. Figure depicts ORN Poisson spike trains for the odor benzaldehyde corresponding to projections to seven different glomeruli indicated by the vertical labels. The driving spike rates are described by (1). Simulations are shown for one trial, where the odor stimulus is presented from 500 ms to 1000 ms.

Fig. 5.

Averages of firing rates across ten trials for model ORNs and PNs. Odor stimulus is presented from 500 ms to 1000 ms. Poisson firing rates are based on data for the odor benzaldehyde for ORNs associated with seven different glomeruli. Results for only four glomeruli shown here. Black curves depict ORN responses, blue curves depict PN responses with only intraglomerular connections and no lateral connections, and the red curves depict PN responses with eLNs included. Compare to the experimental results (1).

Fig. 6.

The horizontal axis depicts time, and the vertical axis shows the number of spikes. Each row is one model cell where they are grouped into the four cell types. Within a cell type, the different rows correspond to the seven model glomeruli. ORN input to each glomerulus consists of ten different Poisson spike trains. As ORN input activity increases in one glomerulus, PN output activity will increase in that glomerulus. Moreover, global inhibition then serves to suppress activity in other glomeruli as activity increases.

Fig. 7. Temporal evolution of PN responses for different odors across PCA space and the equivalent eigenvalues.

Fig. 8.

Temporal evolution of PN responses for four different odors across PCA space. The black and red curves depict the projections of seven different glomeruli responses with only iLNs in the network model and with both iLNs and eLNs, respectively. The four panels show results for odors (a) benzaldehyde, (b) 1-butanol, (c) cyclohexanone, and (d) butyric acid for the first three principal components. During spontaneous activity before the odor is presented, the changes in the PCA values over time remain localized. Note that trajectories depict a rapid divergence from background spontaneous activity to form a transient during odor evoked activity, and then following the odor offset, the activity returns to the background spontaneous activity. Also, note the presence of longer transients in the network model with only iLNs compared to the network model when adding eLNs. These results suggest that eLNs may decrease the ability of the network to discriminate among single odors.

TABLE I

Parameter values for iLN, eLN, and PN cell models.

Symbol	Value	Unit	Description
C_m	0.13	μF	membrane capacitance
E_K	-72	mV	K^+ reversal potential
E_{Na}	55	mV	Na^+ reversal potential
E_L	-75	mV	L reversal potential
g_{Na}	1.96	mS	I_{Na} maximal conductance
g_L	0.01372	mS	L maximal conductance
$g_{K_{iLN}}$	11.76	mS	I_K maximal conductance iLN
θ_w	-44	mV	half-activation for b
θ_m	-32	mV	half-activation for m
k_m	-30.92	mV	activation sensitivity for m
k_w	-28.34	mV	activation sensitivity for b
k_τ	28.34	mV	activation sensitivity for τ
r_w	0.1	$1/ms$	basal rate for b
E_{Ca}	124	mV	Ca^{2+} reversal potential
θ_{Ca}	-10	mV	half-activation for n
g_{Ca}	20	mS	I_{Ca} maximal conductance
g_{KCa}	0.03	mS	I_{KCa} maximal conductance
α_{Ca}	0.00005		Ca conversion rate
r	0.01		Ca removal rate
k_{Ca}	-8.467	mV	activation sensitivity for n
$g_{K_{PN}}$	10.78	mS	I_K maximal conductance PN

TABLE II

Synaptic parameter values for the network model, Cond stands for the word conductance, and potential is shortened to pot.

Symbol	Value	Unit	Description
α_{GABA_A}	10	1/ms	forward rate constant
β_{GABA_A}	0.16	1/ms	backward rate constant
α_{ACh}	10	1/ms	forward rate constant
β_{ACh}	0.2	1/ms	backward rate constant
θ_T	-20	mV	half activation
k_T	-1.5	mV	activation sensitivity
E_{GABA_A}	-90	mV	$GABA_A$ reversal pot.
E_{GABA_B}	-95	mV	$GABA_B$ reversal pot.
E_{ACh}	0	mV	ACh reversal potential
K	100	μM^4	dissociation rate
k_1	0.5	1/ms	kinetic rate constant
k_2	0.0013	1/ms	kinetic rate constant
k_3	0.1	1/ms	kinetic rate constant
k_4	0.033	1/ms	kinetic rate constant
g_{AChPN_eLN}	0.0008	mS	PN to eLN ACh
g_{AChPN_iLN}	0.0001	mS	PN to iLN ACh
$g_{ACh eLN\_iLN}$	0.00001	mS	eLN to iLN ACh
$g_{GABA_A iLN\_eLN}$	0.00001	mS	iLN to eLN $GABA_A$
$g_{GABA_B iLN\_eLN}$	0.00001	mS	iLN to eLN $GABA_B$
$g_{GABA_A iLN\_PN}$	0.21	mS	iLN to PN $GABA_A$
$g_{GABA_B iLN\_PN}$	0.21	mS	iLN to PN $GABA_B$
g_{gapPN_eLN}	0.1	mS	eLN-PN coupling
$g_{gap eLN\_iLN}$	0.0001	mS	eLN-iLN coupling
$g_{gap eLN}$	0.0008	mS	eLN-eLN coupling
sd	0.08	mV	synaptic depression
τ_{dep}	100	mS	synaptic decay
g_{peak_ORN}	0.311	mS	ORN synaptic cond
t_{peak_ORN}	8	mS	ORN sensitivity
E_{ORN}	30	mV	synaptic reversal pot.