Membrane interactions of ionic liquids and imidazolium salts

Da Wang¹ · Hans-Joachim Galla¹ · Patrick Drücker²

Received: 5 October 2017 / Accepted: 13 December 2017 / Published online: 4 January 2018

© International Union for Pure and Applied Biophysics (IUPAB) and Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Room-temperature ionic liquids (RTILs) have attracted considerable attention in recent years due to their versatile properties such as negligible volatility, inflammability, high extractive selectivity and thermal stability. In general, RTILs are organic salts with a melting point below ~100 °C determined by the asymmetry of at least one of their ions. Due to their amphiphilic character, strong interactions with biological materials can be expected. However, rising attention has appeared towards their similarity and interaction with biomolecules. By employing structural modifications, the biochemical properties of RTILs can be designed to mimic lipid structures and to tune their hydrophobicity towards a lipophilic behavior. This is evident for the interaction with lipid-membranes where some of these compounds present membrane-disturbing effects or cellular toxicity. Moreover, they can form micelles or lipid-like bilayer structures by themselves. Both aspects, cellular effects and membrane-forming capacities, of a novel class of lipophilic imidazolium salts will be discussed.

Keywords Ionic liquid · Imidazolium salt · Micelle · Bilayer · Vesicle · Biomembrane

Introduction

Room-temperature ionic liquids (RTILs) are salts in a liquid state below ~100 °C, which have a hybrid, organic-ionic nature and exhibit remarkable properties, such as negligible volatility and thermal stability, as well as an impressive solvation ability. Some of these compounds show astonishing similarities to biomolecules, similar to lipids that promote attention to investigate their structure-to-function principles and biomolecule interactions. In general, many RTILs serve as designer solvents with specialized properties and applications like the liquid–liquid extraction of antibiotics (Cull et al. 2000) or function as solutes for transition metal catalysis (Wasserscheid and Keim 2000). Based on a large number of cation–anion combinations, RTILs can be tailored for a

variety of applications including organic synthesis (Welton 2004), separation (Visser et al. 2001) and electrochemical processes (Kamavaram et al. 2005). In the past few years, ionic liquids and N-heterocyclic carbenes (NHCs) have furthermore gained significant attention in scientific fields such as complex chemistry, material science, drugs and catalysis (Dröge and Glorius 2010; Hallett and Welton 2011; Hopkinson et al. 2014; Richter et al. 2014; Welton 1999). One of the most recent findings in this novel field is that some RTILs can adopt mono- or bilayer arrangements in carbide-derived carbon nanopores that show structural anomalies, such as partially broken columbic orderings, analyzed by X-ray scattering and Monte Carlo simulations (Futamura et al. 2017). The latter work, being at the frontier of recent science, proved the existence of a superionic state and might have further implications for electrochemical applications and the field of charge distributions in close vicinity of surfaces (Futamura et al. 2017; Kondrat and Kornyshev 2011).

Beside these current studies, we here scrutinize characteristics that are important for the interaction of RTILs with biological membranes and naturally occurring lipids. Among a wide chemical variety, an important class of RTILs are imidazolium cations, carrying aliphatic or aromatic residues at the nitrogen atoms. For instance, transmembrane anion transport properties of adamantyl-functionalized imidazolium salts have been reported in combination with the effect of the

This article is part of a Special Issue on 'Ionic Liquids and Biomolecules' edited by Antonio Benedetto and Hans-Joachim Galla.

✉ Patrick Drücker
patrick.druecker@ana.unibe.ch

¹ Institute of Biochemistry, Westfälische Wilhelms-University Münster, Wilhelm-Klemm-Strasse 2, D-48149 Münster, Germany

² Department of Cell Biology, Institute of Anatomy, University of Bern, Baltzerstrasse 2, 3000 Bern 9, Switzerland

flexibility of their linkers to the central imidazolium cation (Gravel and Schmitzer 2015). Interestingly, RTILs could serve as a substitute for aqueous solutions, which embed lipid bilayer membranes. Gayet et al. showed the formation of DPPC lipid vesicles that could be stabilized in the presence of several RTILs such as 1-butyl-3-methylimidazolium tetrafluoroborate (BmimBF_4) or RTIL–water mixtures (Gayet et al. 2011). They furthermore demonstrated that the bilayer structure and its stability depends on the RTIL content by applying small-angle neutron scattering and differential scanning calorimetry. Again, this RTIL was modified with alkyl residues at the nitrogen atoms. Instead of employing ionic liquids as a substitute for solvents, RTILs themselves have also been shown to form membranes in aqueous media. An early approach was the ability of RTILs to self-aggregate in relation to the electrostatic or hydrophobic contributions of the isolated cation in water (Blesic et al. 2007). They discovered mixed-micelles formed by $[\text{C}_n\text{mim}]\text{Cl} + \text{SDS}$ for $n \approx 6\text{--}8$ and reported on a rather “pure salt effect” for shorter chain lengths. Liu et al. showed by molecular dynamics simulations the capability of imidazolium-based RTILs $[\text{C}_n\text{mim}]\text{Br}$ ($n = 10, 12, 14$), that contain a single hydrophobic tail at one nitrogen atom, to form unilamellar vesicles in aqueous solutions without any supplement (Liu et al. 2015). In our approach to focus on similarities to natural lipids, the structure of $[\text{C}_{14}\text{mim}]^+$ already contained a hydrophilic head and a hydrophobic tail, mimicking to some extent the structure of lyso-phospholipids (Fig. 1a).

The structural flexibility of imidazolium salts was further improved as Richter et al. introduced 4,5-alkyl-*N*-heterocyclic imidazolium salts. In their work, the hydrophobic residues were connected to the hydrocarbon backbone of the imidazolium cation, whereas the 1,3-nitrogen positions remain accessible for further modifications (Richter et al. 2014). Although the initial design was tailor-made as a precursor for the stabilization of metal-nanoparticles in aqueous solutions, this structure allowed the independent design of the lipid head group-mimicking part and the hydrophobic tail-mimicking portion of the molecule. Employing ligand exchange reactions, the imidazolium salt, 1,3-dimethyl-4,5-dipentadecylimidazolium iodide ($\text{C}_{15}\text{-Ime}\cdot\text{HI}$), could be functionalized to a carbene moiety that associates to palladium-nanoparticles. The long alkyl chains of the molecules then generated a hydrophobic shell around the Pd-particles that prevents their aggregation (Richter et al. 2014). However, this molecule and derivatives thereof (Fig. 1b) exhibit a highly structural similarity to phospholipids, such as DPPC. Following this work, lipid mimetic properties and membrane interactions were further investigated, and the backbone-alkylated imidazolium salts revealed antibacterial, cytotoxic and activities against tumor cells (Wang et al. 2015a, b), to be discussed in detail here.

Additionally, further modifications of the 1,3-nitrogen residues on the imidazolium ring leads to a broad variety of

structure designs with manifold properties (Rühling et al. 2017). Interestingly, the introduction of bulky benzyl modifications leading to the 1,3-dibenzyl-4,5-dipentadecylimidazolium salt ($\text{C}_{15}\text{-IBn}\cdot\text{HBr}$) conserved some structural similarity to lipids, but showed unexpected properties in aqueous solutions. Depending on the bulkiness of the head group and the length of the hydrocarbon chains, lipid-droplet-like emulsions, probably stabilized by π -stacking effects of the aromatic benzyl residues, or even giant vesicles were formed (Drücker et al. 2017). Their membrane interactions were concomitantly studied via a series of artificial lipid bilayer models in vitro.

Cellular effect and structural features of 4,5-dialkylated imidazolium-based salts

To date, the best-known RTILs are imidazolium-based ionic liquids. In the past few years, a number of studies have focused on 1-alkyl-3-methylimidazolium species (Ranke et al. 2004; Cui et al. 2003; Kumar and Malhotra 2010; Samori et al. 2010; Stolte et al. 2007) which behave as amphipathic compounds and reveal surface activities analogous to quaternary ammonium compounds (QACs). The polar and hydrophilic imidazolium head group is beneficial for an electrostatic interaction with membrane lipids, showing a remarkable membrane activity (Samori et al. 2010; Anderson and Long 2010; Geng et al. 2010; Carson et al. 2009; Rawat and Bohidar 2012). In addition, the long alkyl chain attached to the N^1 atom has been shown to effectively enhance surface activity and self-assembly characteristics of 1-alkylimidazolium cations in aqueous solution (Tischer et al. 2012). This mono-alkylated imidazolium species has been reported to exhibit potent cellular toxicity against a panel of tumor cell lines, including human origin (MCF-7, HeLa, HT-29 and Caco-2 cells) (Kumar et al. 2009; Stepnowski et al. 2004; Frade et al. 2007, 2009; García-Lorenzo et al. 2008) as well as animal tumor cell lines (IPC-81 and C6 Glioma) (Ranke et al. 2004, 2007; Stolte et al. 2006). Generally, their cytotoxicity could be directly correlated with their lipophilic properties (Łuczak et al. 2010; Ranke et al. 2004). Hitherto uncharacterized 4,5-dialkylated imidazolium cations, which represent a novel class of imidazolium-based RTILs, have been designed and synthesized (Wang et al. 2015a, b), as mentioned above. However, further structural diversity was introduced by head group modifications to monitor lipid mimetic properties in relation to head group size and hydrophobicity (Wang et al. 2016; Rühling et al. 2017; Drücker et al. 2017). These backbone-alkylated imidazolium RTILs consist, for instance, of a hydrophilic $\text{N,N}'$ -disubstituted imidazolium head group and two hydrophobic alkyl chains located at the 4- and 5-positions of the imidazole ring (Fig. 2). Either an *N*-methyl- or *N*-benzyl-substituent was

Fig. 1 Lipid mimetic ionic liquids. **a** $[\text{C}_n\text{mim}]^+$ ($n = 10, 12, 14$) salts imitating the structure of lyso-phosphatidylcholine. **b** $(\text{C}_n\text{-Ime-HI}, n = 7, 11, 15)$ imidazolium salts resembling the structure of DPPC

employed as a tunable wingtip group of the imidazole core, which allowed modification of steric properties and the hydrophobicity of imidazolium RTILs. In particular, the N-methylated imidazolium head group was additionally equipped with C^2 -methylation. Varying alkyl chain lengths of $n = 7, 11$ and 15 were taken into account to cover the different cation lipophilicities. This work combines the structural features of phospholipids and imidazolium cations to mimic the structure and biophysical behavior of membrane lipids (Fig. 2) (Rühling et al. 2017).

The cellular toxicity of these compounds was evaluated in vitro using a panel of cell lines, including the rat C6 Glioma tumor cell line and cells such as Madin–Darby canine kidney (MDCK II), as well as the mouse embryonic fibroblast cell line (NIH3T3). The half-maximal effective concentrations (EC_{50}) are summarized in Table 1. Remarkably, compared to general mono-alkylated imidazolium species with an $\text{EC}_{50} > 1000 \mu\text{M}$ (Ranke et al. 2004), all studied 4,5-dialkylated imidazolium RTILs reveal an approximate three orders of magnitude higher anti-tumor activity against the C6 glioma

cell line. The most potent one is $\text{C}_7\text{-IBn-HBr}$, displaying the lowest EC_{50} of $\sim 1.0 \mu\text{M}$ (Wang et al. 2015a, b). This indicates that the introduction of two hydrophobic alkyl chains into the backbone contributes greatly to improve the activity against tumor-derived cell lines of imidazolium-based RTILs.

Further, for a given type of head group, the cellular toxicity is decreased with increasing alkyl chain length, which is true for all tested cell lines. In other words, the cytotoxic effect of the 4,5-dialkylimidazolium RTILs carrying the same head group is negatively correlated with their lipophilicity, which is apparently inverse to the earlier reported findings for 1-alkylimidazolium species (Łuczak et al. 2010; Ranke et al. 2004).

In order to analyze a possible structure-activity relationship for $\text{C}_n\text{-Ime-HI}$ and $\text{C}_n\text{-IBn-HBr}$ species to natural occurring lipids, their critical micelle concentration (CMC) values, which also characterize surface activity, were co-plotted with the EC_{50} values of their toxicity on cells (Fig. 2a, b) (Wang et al. 2015a, b). It appeared that the CMC values are negatively correlated with the EC_{50} values for both the $\text{C}_n\text{-Ime-HI}$ and

Fig. 2 Chemical structures of investigated 4,5-dialkylated imidazolium-based RTILs. The imidazolium RTILs are abbreviated as $\text{C}_n\text{-IR-R'A}$, where n represents the number of carbon atoms in each alkyl chain, I the imidazolium, R the substituent at the N-position, R' the substituent at the 2-position and A the anion. Modified from Rühling et al. (2017)

Table 1 CMC [μM] and EC_{50} [μM] and confidence intervals of all tested 4,5-dialkylimidazolium RTILs

Abbreviations	CMC/ μM	$\text{EC}_{50} \pm \text{SD}/\mu\text{M}$		
		C6 glioma	MDCK II	NIH3T3
$\text{C}_7\text{-Ime-HI}$	65.9 (50–80)	2.3 ± 0.5	6.7 ± 0.8	4.7 ± 0.8
$\text{C}_7\text{-IBn-HBr}$	17.9 (10–20)	1.0 ± 0.1	12.9 ± 2.1	2.5 ± 0.4
$\text{C}_{11}\text{-Ime-HI}$	8.5 (5–10)	10.5 ± 0.3	8.9 ± 0.8	5.0 ± 0.5
$\text{C}_{11}\text{-Ime-MeI}$	7.9 (5–10)	5.3 ± 0.5	8.4 ± 0.5	1.0 ± 0.2
$\text{C}_{11}\text{-IBn-HBr}$	2.3 (2.0–5.0)	2.0 ± 0.1	16.0 ± 5.7	4.2 ± 0.5
$\text{C}_{15}\text{-IH-NH}_2\text{Cl}$	4.1 (2.5–5)	67.2 ± 3.9	32.2 ± 3.3	12.2 ± 1.7
$\text{C}_{15}\text{-Ime-HI}$	1.4 (1.0–5.0)	28.1 ± 0.7	15.9 ± 0.5	7.7 ± 0.2
$\text{C}_{15}\text{-Ime-MeI}$	0.9 (0.7–1.0)	16.3 ± 0.6	23.6 ± 1.1	3.5 ± 0.9
$\text{C}_{15}\text{-IBn-HBr}$	0.5 (0.4–1.3)	11.3 ± 2.0	20.6 ± 4.2	8.7 ± 0.7

the $\text{C}_n\text{-IBn-HBr}$ compound series, showing that the elongation of alkyl chains in the backbone result in a declined tendency of their activity against tumor-derived cell lines (Fig. 3a; Wang et al. 2015a, b).

Interestingly, the EC_{50} value ($\sim 10.5 \mu\text{M}$) of $\text{C}_{11}\text{-Ime-HI}$ (Fig. 3a) is very close to its CMC value ($\sim 8.5 \mu\text{M}$) and almost overlapping in the case of the $\text{C}_{11}\text{-IBn-HBr}$ compound with an EC_{50} ($\sim 2.0 \mu\text{M}$) and CMC ($\sim 2.3 \mu\text{M}$) (Fig. 2b). This suggests that the micellization plays a critical role in their cellular

toxicity (Wang et al. 2015a, b). The micelle formation and the CMC values might indicate an effective adsorption for C_{11} derivatives at the cell-media interface, thus facilitating the fusion into the cell membrane. A strong membrane destabilization or disruption is potentially given by the hydrophobic mismatch between the medium length C_{11} alkyl chains and long alkyl chains of membrane lipids. This effect, presumably rising at a specific chain length range of 10–12 carbon atoms, is well supported by earlier results with respect to the biological activities of long-chain imidazolium cationic surfactants (Pernak et al. 2003). In order to visually present the cellular toxicity of C_{11} imidazolium salt species by wide-field microscopy using phase-contrast imaging, C6 glioma cells were treated with $100 \mu\text{M}$ $\text{C}_{11}\text{-Ime-HI}$ for 24 h (Fig. 3c–e). The intact cell membranes of untreated C6 glioma cells are presented in Fig. 3c. However, after lysis of cell membranes by Triton X-100, only the nuclei of the tumor cells can be identified (Fig. 3d). After treatment with $100 \mu\text{M}$ $\text{C}_{11}\text{-Ime-HI}$, the plasma membranes of C6 glioma cells have nearly disappeared (Fig. 3e), which display a similar cell morphology compared to the control shown in Fig. 3d (Wang et al. 2015a, b). This demonstrates that $\text{C}_{11}\text{-Ime-HI}$ act as a detergent and exhibits strong lytic activity on the cell membranes.

As reported by Ranke et al. (2004), the single, long alkyl chain at the imidazolyl-1-position may intercalate into the cell membranes, which could then promote a strong membrane

Fig. 3 Cellular interactions of $\text{C}_n\text{-Ime-HI}$ and $\text{C}_n\text{-IBn-HBr}$ ($n = 7, 11, 15$). Both CMC and EC_{50} values against C6 glioma cell line plotted as a function of the number of carbon atoms (n) in each alkyl chain for **a** $\text{C}_n\text{-Ime-HI}$ and **b** $\text{C}_n\text{-IBn-HBr}$ ($n = 7, 11, 15$). Microscopy-phase-

contrast images of C6 glioma cells: **c** untreated, **d** treated by 1% Triton X-100 for 10 min and **e** treated by $100 \mu\text{M}$ $\text{C}_{11}\text{-Ime-HI}$ for 24 h. Modified from Wang et al. (2015a, b)

disruption. However, backbone-alkylated imidazolium RTILs are structurally even closer to the natural phospholipid framework. Thus, it could be speculated that they might exert their potent cellular effects (mentioned above) along distinct pathways differing from those of 1-alkylimidazolium salts. To verify this hypothesis, a series of investigations of the interaction between C_n -Ime·HI ($n = 7, 11, 15$) and monolayers using the Langmuir technique *in vitro* were carried out (Fig. 4) (Wang et al. 2016). DPPC, a zwitterionic phospholipid, was used as a representative model membrane lipid to establish binary monolayers at the air-water interface. It appears that C_{15} -Ime·HI components generate significant concentration-dependent modifications of the monolayer structure below 15 mN/m (Fig. 4a.). With increasing addition of C_{15} -Ime·HI (molar ratio from 0.05 to 0.3), a drastic rigidifying effect on the DPPC monolayer is shown by the increasing appearance of lipid condensed domains (dark area). At the 0.3 M fraction of the C_{15} compound, the most remarkable rigidification is reached, accompanied by the formation of a Langmuir film with the best thermodynamic stability (minimal $\Delta G^{\text{ex}} = -1849$ J/mol, Fig. 4d) (Wang et al. 2016). This implies that C_{15} -Ime·HI has good affinity to mix with DPPC, leading to a readily miscible lipid monolayer at the air-water interface. Furthermore, its capability to mix well with membrane lipids (such as DPPC) might correlate with a lower cellular toxicity (i.e. by higher EC_{50} values) (Table 1; Wang et al. 2015a, b, 2016). Instead, the integrity of the lipid film is severely

disrupted by the incorporation of C_{11} -Ime·HI, showing lowered line tensions between domains (irregular appearance of domains) (Fig. 4b). These findings thus suggest, that the capability of these imidazolium salts to interact, i.e. intercalate into the membrane, may play along with the cellular toxicity and membrane disturbance. In fact, the medium length C_{11} compound shows a stronger cellular effect than the long chain C_{15} , which is in line with the domain disturbance mentioned above (Fig. 4). This is also in agreement with the findings from the cytotoxicity assays by Wang et al. (2015a, b) (Fig. 3c–e) showing membrane lysis. However, the short chain C_7 -Ime·HI exhibits negligible membrane activity (Fig. 4c), although it is the most toxic compound in the C_n -Ime·HI series. Therefore, it can be concluded that, for a given type of head group (particularly -Me), a negative correlation between the cellular toxicity of 4,5-dialkylated imidazolium RTILs and their lipophilicity, i.e. the capability to intercalate in dependence of alkyl chain length, is evident.

Further, their cellular toxic effect is moderated by another important factor, the hydrophobicity and size of the imidazolium head group. Here, a guanidinium RTIL (C_{15} -IH-NH₂Cl) bearing a highly hydrophilic imidazolium head group (Fig. 2, fourth structure), where the nitrogen atoms are reduced and the carbon C² is modified by an amine group, showed the lowest toxicity ($EC_{50} \sim 67.2$ μ M), accompanied by a relatively high water solubility (CMC ~ 4.1 μ M) (Rühling et al. 2017). However, C_n -Ime·MeI ($n = 11, 15$), with a

Fig. 4 Interactions of C_n -Ime·HI ($n = 7, 11, 15$) on a DPPC containing monolayer. Epi-fluorescence images of the monolayers of DPPC mixed with different molar fractions of **a** C_{15} -Ime·HI, **b** C_{11} -Ime·HI, and **c** C_7 -Ime·HI at the air-water interface at 293 K. All samples were mixed with 0.5 mol% of the fluorescent dye BODIPY-PC. The frame size of all images is 120×120 μ m. **d** The excess Gibbs free energy (ΔG^{ex}) of mixing versus composition plots of the binary mixed monolayers of DPPC/ C_{15} -Ime·HI at several, constant surface pressures at 293 K. Modified from Wang et al. (2016)

trimethylated and hence more hydrophobic imidazolium head group, have a relatively high activity against C6 glioma cells in comparison to the structurally related C_n -IMe·HI compounds. Here, the cytotoxic effect is enhanced 1.72-fold and 1.98-fold in the cases of $n = 11$ and $n = 15$, respectively (Rühling et al. 2017, oral communication). By contrast, the N-benylation of the imidazole structure did more effectively improve the anti-tumor activity towards C6 glioma cells (Table 2) (Rühling et al. 2017).

The most significant effect of N-benylation on cellular toxicity appears for a alkyl chain length of $n = 11$, where it is 5.3-fold higher in comparison to N-methylation of the imidazolium ring (Table 2). This might be interpreted by the different stability of micelles in solution (Table 1). These results suggest that the head group hydrophobicity and size of imidazolium-based RTILs could influence the probability to interact and intercalate into cellular bilayers. A head group with enhanced hydrophobicity but lower size favors a higher partitioning probability into the hydrophobic interior of the cell membrane, followed by membrane disturbance or disruption.

Taken together, the cellular toxicity effect of the 4,5-dialkylated imidazolium RTILs discussed here strongly depend on both the alkyl chain length as well as the head group type. Specifically, their activity against C6 glioma tumor-derived cells in vitro can be summarized in the following order:

Membrane interactions of 4,5-dialkylated imidazolium-based salts

By employing different techniques, it could be demonstrated that the 1,3-dimethyl-4,5-dialkylimidazolium salts (C_n -IMe·HI, $n = 7, 11, 15$) strongly interact with artificial and cellular bilayer membranes (Drücker et al. 2017; Wang et al. 2015a, b). The combination of the quartz crystal microbalance technique and fluorescent membrane probes allowed further determination of fundamental membrane interactions and bilayer properties of the pure compound. Lipid monolayer insertion properties and monolayer-domain rigidifications were probed by surfactant mimicking Langmuir film measurements as discussed before (Wang et al. 2016; Rühling et al. 2017). Additionally, the application of laser scanning confocal microscopy demonstrated vesicle formation and bilayer membrane domain fluidization of the long alkyl chain C_{15} -IMe·HI.

The quartz crystal microbalance technique with dissipation monitoring (QCM-D) is a versatile method to analyze surface

interactions at the fluid/solid interface. It has been frequently used to analyze the contact of biomolecules, such as peptide nucleic acids or hemoglobin with membrane surfaces (Höök et al. 1998; Janshoff et al. 2000). The simultaneous measurement of the sensor's frequency shift ($\Delta F = F(t) - F_0$), representing changes in associated mass, and dissipation ($\Delta D = D(t) - D_0$), which provides information about the viscoelastic properties delivers two important parameters to characterize the interaction of RTILs with membrane models (Sauerbrey 1959; Janshoff et al. 2000; Fattison et al. 2011).

Since C_{15} -IMe·HI has a strong structural resemblance to naturally occurring phospholipids (Fig. 1), a charged, polar head group and two hydrophobic alkyl chains, it is interesting to gain further insights into functional similarities. Therefore, negatively charged POPC/POPS (4:1) vesicle membranes were incubated with suspensions of C_n -IMe·HI ($n = 7, 11, 15$) imidazolium salts in a tris-buffered milieu, pH 7.4 (TBS) that roughly mimics cellular conditions (Fig. 5).

Following the equilibration of the sensor's surface in TBS buffer, 200-nm POPC/POPS (4:1) vesicles were tethered on the surface via streptavidin/biotin linkers. The adsorption showed a frequency shift of $\Delta\Delta F = 311 \pm 16$ Hz (Drücker et al. 2017; Wang et al. 2015a, b), accompanied by a moderate change in dissipation (Fig. 5, arrows 1). The latter results from the fact that surface-associated vesicles show a much larger loss of resonance energy due to their viscoelastic properties than, for instance, solid supported bilayers (Drücker et al. 2014). With respect to that comparison, the subsequent incubation of C_n -IMe·HI ($n = 7, 11, 15$) imidazolium salts revealed a much more enhanced effect. First, the long chain C_{15} and medium length C_{11} imidazolium compounds showed a huge additional increase in the frequency shift ($\Delta\Delta F_{C_{15}} = 445 \pm 8$ Hz), which demonstrates the accumulation of a large mass in relation to the initial mass of liposomes (Fig. 5, arrows 2). Second, the interaction of these imidazolium salts with the vesicle membrane showed an even larger gain in the dissipation shift than the adsorption of vesicles in the first place. This effect was mostly pronounced for the C_{15} compound and appeared to be proportional to chain length (Drücker et al. 2017). It is likely that the simple intercalation of the imidazolium salts into the vesicle bilayer, which leads to the formation of a mixed membrane, is not sufficient to explain such large

Table 2 Comparison of cellular toxic efficiency between C_n -IMe·HI and C_n -IBn·HBr species

$\frac{EC_{50}(C_n\text{-IMe}\cdot\text{HI})}{EC_{50}(C_n\text{-IBn}\cdot\text{HBr})}$	C6 glioma	MDCK II	NIH3T3
$n = 7$	2.3	0.52	1.9
$n = 11$	5.3	0.56	1.2
$n = 15$	2.5	1.2	0.89

Fig. 5 Bilayer interactions of C_n -IMe·HI on POPC/POPS (4:1) vesicles probed by QCM-D. Large unilamellar vesicles of POPC/POPS (4:1, 200 nm) were surface-tethered by biotin-streptavidin linkers on a self-assembled monolayer coated sensor surface (arrows 1). Following equilibration with tris-buffered-saline, pH 7.4, suspensions of 0.1 mM (C_n -IMe·HI, $n = 7, 11, 15$) imidazolium salts were incubated, which induced diverse membrane interactions (arrows 2). Remarkably, the C_{15} (red) and C_{11} (blue) compounds lead to an increase in surface associated mass, indicated by $\Delta\Delta F$ and an increased viscoelasticity of the associated layers ($\Delta\Delta D$), whereas the short chain C_7 (green)

imidazolium salt did not much alter vesicle membranes. After another equilibration step with buffer for 30–60 min, rinsing with pure water applied hypo-osmotic conditions (arrows 3). The osmotic gradient across the vesicle membrane resulted in a severe loss of vesicle mass after interaction with the C_{11} compound. Vesicles incubated with the C_{15} imidazolium salt partially lost some mass and vesicles treated by the C_7 compound did not considerably change their mass. As a control, ~0.5 mg/ml POPC:POPS (4:1), 200-nm vesicles were also applied (black). The cartoon represents the effects of imidazolium salt on the vesicles (e.g. C_{11} -IMe·HI). Modified from Drücker et al. (2017) and Wang et al. (2015a, b)

shifts. Rather than detecting only the addition of membrane mass, it seems to be reasonable that the intercalation of these imidazolium salts moreover altered the viscoelastic properties by generating packing defects in the bilayer. This is in good agreement with the fact that the C_{11} and C_7 imidazolium compounds cannot fully penetrate the membrane and hence led to packing defects as shown for a DPPC bilayer simulated by molecular dynamics (Wang et al. 2016). Whereas in contrast to that, the long tail C_{15} -IMe·HI compound induced to the formation of a gel-phase in the DPPC mixed bilayer. Following the interactions of C_n -IMe·HI ($n = 7, 11, 15$) imidazolium salts with the POPC/POPS (4:1) vesicle membranes, the application of hypo-osmotic conditions (Fig. 5, arrows 3), showed the almost complete loss of vesicle membranes in case of prior C_{11} compound interactions. Vesicle membranes, which have been incubated with the C_{15} compound, were partially disrupted, whereas the C_7 imidazolium

salt did not much affect the membranes at all. This is in good agreement with the difference in head/tail size proportions of these compounds in relation to phosphatidylcholine lipids. The long tail C_{15} -IMe·HI provided the least hydrophobic mismatch between alkyl-chains and the palmitoyl- or oleoyl-groups of the membrane lipids, whereas C_{11} -IMe·HI exhibits a much larger hydrophobic mismatch, resulting in a stronger disintegration of vesicle bilayers.

In contrast to POPC/POPS (4:1) membranes, a neutrally charged model of pure POPC liposomes showed less prominent interactions with these imidazolium salts, therefore suggesting that the positive charge of C_n -IMe·HI salts facilitates surface contact to the bilayer and promotes membrane intercalation (Drücker et al. 2017).

The C_{15} -IMe·HI imidazolium salt interacted starkly with vesicle membranes, thus proposing that the structure-to-function relationship of this compound matches best to natural

phospholipids. Surprisingly, it could be further demonstrated that it incorporates fluorescent membrane probes, such as 1,6-diphenyl-1,3,5-hexatriene (DPH) or Laurdan. These probes change their fluorescence intensity and anisotropy in dependence of water contact, and report on membrane dynamics and fluidity when they are embedded within the bilayer (Peng et al. 2012; Parasassi et al. 1998).

The fluorescence spectra of these two probes incorporated in pure C₁₅-IMe-HI, exhibit strong similarity to spectra of the same probes embedded in a DOPC reference bilayer (Fig. 6a, b). This demonstrates two interesting findings. First, the fluorescence intensity suggests that it can stabilize bilayer membranes and, furthermore, that these membranes show more similarity to fluid DOPC rather than ordered DPPC bilayers. Moreover, a change in the fluorescence anisotropy of DPH showed a phase transition at 28–32 °C (Drücker et al. 2017).

Visual proof that the C_n-IMe-HI imidazolium salts ($n = 11, 15$) can indeed form bilayer membranes was provided by the attempt to generate giant vesicles in sucrose or HBS, pH 7.4, employing confocal laser scanning microscopy (Fig. 7a, b). The vesicle membranes appeared to be uniform and consisted of one homogeneous phase at room temperature, indicated by phase selective dyes like Bodipy-PC or DiI_{C18} (Drücker et al. 2017). If domains in a membrane exist, it is expected that those dyes exhibit a liquid ordered (l_o) or liquid disordered (l_d) domain preference (Baumgart et al. 2007; Klymchenko et al. 2014). Although vesicles formed by C₁₅-IMe-HI appeared to be stable, the C₁₁-IMe-HI compound forms kinetically driven, unstable giant vesicles. The latter could be observed, when vesicles of C₁₁-IMe-HI were formed in sucrose, diluted 1:4 by HBS, and instantly captured

by fluorescence microscopy (Fig. 7c). Note the equidistant yellow lines, highlighting the shrinkage in diameter during 110 s (Fig. 7c). The total amount of the C₁₁-IMe-HI membrane surface that could be observed by microscopy was decreasing, hence suggesting that micelle nucleation leads to the formation of non-observable smaller structures. This phenomenon could be explained by the relatively high CMC of ~8.5 μM for this compound (Wang et al. 2015a, b). Consequently, the C₁₁ compound vesicles become more unstable after dilution than micelles. The hydrophobic tail is shorter than that of the C₁₅ compound, which could explain the preferential transformation. The shrinking process possibly starts with C₁₁ imidazolium salt protrusions at the surface of giant vesicles, which eventually leads to micelle nucleation, elongation and the subsequent detachment of micelles (Fig. 7d).

A somewhat similar process was determined by Yoo et al. using coarse-grained molecular dynamics simulations of 1-butyl-3-methylimidazolium chloride, [C₄mim]Cl, in interaction with POPC bilayers (Yoo et al. 2016). They concluded that the short tail cationic [C₄mim]Cl could insert into one leaflet of a planar bilayer altering the leaflet strain and the membrane's bending modulus (RTIL to lipid ratio 2000:2304). The asymmetric insertion of [C₄mim]Cl promotes a monotonic increase in leaflet strain, which then causes bilayer buckling until micelle nucleation occurs at the saturation limit (Fig. 7e). Furthermore, they also simulated POPC vesicles, which could also insert [C₄mim]Cl, leading to a discontinuity in the vesicle sphericity. This then promoted nucleation and elongation of mixed POPC-[C₄mim]Cl-containing micelles (Yoo et al. 2016).

Fig. 6 Characterization of membranes formed by pure C₁₅-IMe-HI using fluorophore probes. **a** Fluorescence spectra of DPH incorporated into C₁₅-IMe-HI membranes. Vesicles of DOPC and DPPC, which incorporated the probe under similar conditions, were used as a reference. Membranes formed by C₁₅-IMe-HI are likely to have similar properties as DOPC, because their spectra show similar intensity and

appearance. The fluorescence anisotropy of DPH showed a transition at 28–32 °C. **b** Fluorescence spectra of Laurdan in C₁₅-IMe-HI and DOPC and DPPC reference membranes. The inset shows normalized intensities to present the difference in the Laurdan maximum intensity shift. Modified from Drücker et al. (2017)

Fig. 7 Vesicles formed by C_n -IMe·HI imidazolium salts and models of RTIL-membrane interactions. **a** Giant vesicle formed by pure C_{15} -IMe·HI in hepes buffered saline, pH 7.4, monitored by confocal laser scanning microscopy. **b** Giant vesicle formed by C_{11} -IMe·HI in 320 mM sucrose. **c** Shrinkage of C_{11} -IMe·HI vesicles after dilution (1:4) with hepes buffer. Total duration: 110 s. Note the equidistant *yellow lines*, marking the vesicle size at the beginning of that measurement and the relative shrinkage over time. **d**: Model of the probable mechanism of C_{11} -IMe·HI vesicle instability after dilution. **e** Incubation of the ionic liquid $[C_4mim]Cl$ (green) on POPC membranes (cartoon). Modified from Drücker et al. (2017) and Yoo et al. (2016)

Another intriguing property of RTILs and their salt-precursors is their impact on membranes containing spatial domains. It has been shown that C_n -IMe·HI ($n = 7, 11, 15$) imidazolium salts exhibit strong activity against tumor-derived cells, and that 100 μM C_{11} -IMe·HI led to the disruption of C6 glioma cell plasma membranes (Fig. 3) (Wang et al. 2015a, b). Beside membrane transduction of C_n -IMe·HI ($n = 7, 11, 15$) imidazolium salts and other as yet unknown intracellular factors that promote activities against tumor cell lines, the disintegration of functional cell membrane platforms or domains (lipid rafts) may be another determinant of cell death. It is commonly discussed that cellular plasma membranes contain spatial and timely fluctuating lipid platforms that were enriched in certain lipids, proteins and cholesterol (Sezgin

et al. 2017; Simons and Gerl 2010). Moreover, Wang et al. could also demonstrate, that C_n -IMe·HI ($n = 7, 11, 15$) and other structurally related imidazolium salts change the interfacial line tensions of DPPC domains on a Langmuir monolayer model as mentioned above (Fig. 4) (Wang et al. 2016; Rühling et al. 2017).

Employing giant vesicles of DOPC/N-stearoyl-D-erythro-sphingophosphorylcholine (SSM)/cholesterol (1:1:1) and DOPC/SSM/Chol/ C_{15} -IMe·HI (33:23:33:10) at $T = 38.0^\circ C$ showed an intriguing fluidization of lipid domains (Fig. 8a,b).

Doped with Bodipy-PC and DiI $_{C18}$, vesicles of DOPC/SSM/cholesterol (1:1:1), showed small domains which protrude out of the surface and exhibit co-localization of both markers (Fig. 8a). After incorporation of 10% C_{15} -

Fig. 8 Membrane fluidization induced by C_{15} -IMe-HI on giant vesicles. **a** Giant vesicles of DOPC/SSM/Chol (1:1:1) doped with 0.4 mol% Bodipy-PC and 0.4 mol% $\text{DiI}_{C_{18}}$ at $T = 38^\circ\text{C}$ using confocal laser scanning microscopy. Note the small protrusions marked by both dyes. **b** The incorporation of 10% C_{15} -IMe-HI results in giant vesicles of DOPC/SSM/Chol/ C_{15} -IMe-HI (33:23:33:10), which present fluidized, larger domains rich in $\text{DiI}_{C_{18}}$ at $T = 38^\circ\text{C}$. Note that the specificity of the dyes changed. $\text{DiI}_{C_{18}}$ shows high preference for one domain wherein Bodipy-PC is slightly excluded. Modified from Drücker et al. (2017)

IMe-HI, domains appeared to be larger, flat within the bilayer plane, and the dyes showed opposing domain preference. Therefore, a hydrophobic mismatch of the imidazolium salt at domain borders reduced the interfacial energy and provoked bilayer fluidization (Fig. 8b).

Based on that wealth of evidence, another pathway of how the interaction of C_n -IMe-HI ($n = 7, 11, 15$) imidazolium salts can injure tumor cells (Fig. 3) (C6 glioma; Wang et al. 2015a, b) became apparent. The intercalation of C_{15} -IMe-HI into laterally separated bilayer membranes changed the interfacial

energy at domain borders and hence may result in a loss of function on plasma membrane raft domains. Since raft domain integrity is vital for cellular signaling and endo- or exocytosis-related trafficking (Sezgin et al. 2017; Simons and Gerl 2010), the change of cell domain morphology and hence the loss functionality may stimulate cell death.

The major interactions of C_n -IMe-HI ($n = 7, 11, 15$) imidazolium salts with naturally occurring lipid bilayer membranes is briefly summarized by the following cartoon (Fig. 9). The long chain C_{15} -IMe-HI likely had the strongest

Fig. 9 Model representing conceivable interactions of C_n -IMe-HI imidazolium salts with bilayer membranes. The long chain C_{15} -IMe-HI imidazolium salt forms vesicles under buffered conditions and can fuse with membranes. Charge interactions of positive imidazolium salt head groups and zwitterionic or negatively charged lipids led to the intercalation into the membrane. This is likely to be accompanied by

the formation of mixed micelles. The medium chain length C_{11} -IMe-HI imidazolium salt can form vesicles and micelles which disintegrate and lyse bilayer membranes by fusion and incorporation. It also may form mixed micelles. The short chain C_7 -IMe-HI imidazolium salt forms micelles and cannot stabilize giant vesicles in solution. It is likely to pass membranes without disruption. Modified from Drücker et al. (2017)

intermolecular and intramolecular Van der Waals interactions with itself and lipid alkyl chains, promoting the formation of giant vesicles and the fusion to bilayer membranes. It could intercalate, probably form domains, and induce the creation of mixed micelles. The medium chain length C₁₁-IMe-HI compound forms thermodynamically unstable vesicles, which can transiently transform into micelles after dilution. According to the imperfect hydrophobic interactions and its cationic charged head group, it stimulates disintegration of bilayer and plasma membranes. Based on QCM-D results with tethered vesicles and the cytotoxicity *in vitro*, discussed above, the short tail C₇-IMe-HI imidazolium salt evidently passes bilayer membranes without disruption. It cannot form giant vesicles.

Conclusion

This overview summarizes some of the new interesting findings of imidazolium salts in the rising field of ionic liquids and their biomembrane interactions as well as biomolecule, lipid- and membrane-mimetic properties. Motivated by the biological activities of imidazolium-based ionic liquids and the structural features of natural membrane lipids, a novel class of 4,5-dialkylated imidazolium salts are discussed. The investigation of their biophysical properties greatly assists in capturing some details that determine their potential biological effects. Recent work has opened a new path to apply a series of biomimetic, 4,5-dialkylimidazolium salts as a new class of anti-tumor candidates in medical chemistry. Although some of the imidazolium salts discussed here showed strong activity against a few tumor-derived cell lines, a further investigation employing non-tumor cells as well as additional cancer types may answer whether a more sophisticated clinical approach, and drug discovery studies of RTILs could become interesting in future approaches. Additionally, attention deserves to be paid towards the detailed investigation of potential mechanisms that govern their biological effects and selectivity.

Additionally, new emerging approaches, such as the catalysis in drug production and the novel “active pharmaceutical ingredient-ionic liquid” concept that suggest the employment of traditional drugs in the form of ionic liquids (Egorova et al. 2017), provide powerful potentials for upcoming sophisticated applications.

Compliance with ethical standards

Conflict of interest Da Wang declares that he has no conflict of interest. Hans-Joachim Galla declares that he has no conflict of interest. Patrick Drücker declares that he has no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Anderson EB, Long TE (2010) Imidazole- and imidazolium-containing polymers for biology and material science applications. *Polymer* 51: 2447–2454
- Baumgart T, Hunt G, Farkas ER, Webb WW, Feigenson GW (2007) Fluorescence probe partitioning between Lo/Ld phases in lipid membranes. *Biochim Biophys Acta* 1768(9):2182–2194
- Blesic M, Marques MH, Plechkova NV, Seddon KR, Rebelo LPN, Lopes A (2007) Self-aggregation of ionic liquids: micelle formation in aqueous solution. *Green Chem* 9(5):481–490
- Carson L, Chau PKW, Earle MJ, Gilea MA, Gilmore BF, Gorman SP, McCann MT, Seddon KR (2009) Antibiofilm activities of 1-alkyl-3-methylimidazolium chloride ionic liquids. *Green Chem* 11:492–497
- Cui B, Zheng BL, He K, Zheng QY (2003) Imidazole alkaloids from *Lepidium meyenii*. *J Nat Prod* 66:1101–1103
- Cull SG, Holbrey JD, Vargas-Mora V, Seddon KR, Lye GJ (2000) Room-temperature ionic liquids as replacements for organic solvents in multiphase bioprocess operations. *Biotechnol Bioeng* 69(2):227–233
- Dröge T, Glorius F (2010) The measure of all rings - N-heterocyclic Carbenes. *Angew Chem Int Ed* 49(39):6940–6952
- Drücker P, Rühling A, Grill D, Wang D, Draeger A, Gerke V, Glorius F, Galla H-J (2017) Imidazolium salts mimicking the structure of natural lipids exploit remarkable properties forming lamellar phases and giant vesicles. *Langmuir* 33(6):1333–1342
- Drücker P, Grill D, Gerke V, Galla H-J (2014) Formation and characterization of supported lipid bilayers containing phosphatidylinositol-4,5-bisphosphate and cholesterol as functional surfaces. *Langmuir* 30(49):14877–14886
- Egorova KS, Gordeev EG, Ananikov VP (2017) Biological activity of ionic liquids and their application in pharmaceuticals and medicine. *Chem Rev* 117(10):7132–7189
- Fatissou J, Azari F, Tufenkji N (2011) Real-time QCM-D monitoring of cellular responses to different cytomorphic agents. *Biosens Bioelectron* 26(7):3207–3212
- Frade RFM, Matias A, Branco LC, Afonso CAM, Duarte CMM (2007) Effect of ionic liquids on human colon carcinoma HT29 and CaCo-2 cell lines. *Green Chem* 9:873–877
- Frade RFM, Rosatella AA, Marques CS, Branco LC, Kulkarni PS, Mateus NMM, Afonso CAM, Duarte CMM (2009) Toxicological evaluation on human colon carcinoma cell line (CaCo-2) of ionic liquids based on imidazolium, guanidinium, ammonium, phosphonium, pyridinium and pyrrolidinium cations. *Green Chem* 11:1660–1665
- Futamura R, Iiyama T, Takasaki Y, Gogotsi Y, Biggs MJ, Salanne M, Segalini J, Simon P, Kaneko K (2017) Partial breaking of the Coulombic ordering of ionic liquids confined in carbon nanopores. *Nat Mater*. <https://doi.org/10.1038/nmat4974>. Advance online publication
- García-Lorenzo A, Tojo E, Tojo J, Teixeira M, Rodríguez-Berrocá FJ, González MP, Martínez-Zorzano VS (2008) Cytotoxicity of selected imidazolium-derived ionic liquids in the human Caco-2 cell line. Sub-structural toxicological interpretation through a QSAR study. *Green Chem* 10:508–516
- Gayet F, Marty J-D, Brület A, Viguerie N L-d (2011) Vesicles in ionic liquids. *Langmuir* 27(16):9706–9710
- Geng F, Zheng L, Yu L, Li G, Tung C (2010) Interaction of bovine serum albumin and long-chain imidazolium ionic liquid measured by fluorescence spectra and surface tension. *Process Biochem* 45:306–311
- Gravel J, Schmitzer AR (2015) Transmembrane anion transport mediated by adamantyl-functionalized imidazolium salts. *Supramol Chem* 27(5–6):364–371
- Hallett JP, Welton T (2011) Room-temperature ionic liquids: solvents for synthesis and catalysis. 2. *Chem Rev* 111:3508–3576

- Hopkinson MN, Richter C, Schedler M, Glorius F (2014) An overview of N-heterocyclic carbenes. *Nature* 510(7506):485–496
- Höök F, Rodahl M, Kasemo B, Brzezinski P (1998) Structural changes in hemoglobin during adsorption to solid surfaces: effects of pH, ionic strength, and ligand binding. *Proc Natl Acad Sci U S A* 95(21):12271–12276
- Janshoff A, Galla H-J, Steinem C (2000) Piezoelectric mass-sensing devices as biosensors—an alternative to optical biosensors? *Angew Chem Int Ed* 39(22):4004–4032
- Kamavaram V, Mantha D, Reddy RG (2005) Recycling of aluminum metal matrix composite using ionic liquids: effect of process variables on current efficiency and deposit characteristics. *Electrochim Acta* 50:3286–3295
- Klymchenko, Andrey S, Kreder R (2014) Fluorescent probes for lipid rafts: from model membranes to living cells. *Chem Biol* 21(1):97–113
- Kondrat S, Kornyshev AA (2011) Superionic state in double-layer capacitors with nanoporous electrodes. *J Phys Condens Matter* 23:022201
- Kumar RA, Papaiconomou N, Lee JM, Salminen J, Clark DS, Prausnitz JM (2009) In vitro cytotoxicities of ionic liquids effect of cation rings functional groups and anions. *Environ Toxicol* 24:388–395
- Kumar V, Malhotra SV (2010) Ionic liquids as pharmaceutical salts: a historical perspective. *ACS Symp Ser* 1038:1–12
- Liu X, Zhou G, Huo F, Wang J, Zhang S (2015) Unilamellar vesicle formation and microscopic structure of ionic liquids in aqueous solutions. *J Phys Chem C* 120:659–667
- Łuczak J, Jungnickel C, Łacka I, Stolte S, Hupka J (2010) Antimicrobial and surface activity of 1-alkyl-3-methylimidazolium derivatives. *Green Chem* 12:593–601
- Parasassi T, Krasnowska EK, Bagatolli L, Gratton E (1998) Laurdan and Prodan as polarity-sensitive fluorescent membrane probes. *J Fluoresc* 8(4):365–373
- Peng A, Pisal DS, Doty A, Balu-Iyer SV (2012) Phosphatidylinositol induces fluid phase formation and packing defects in phosphatidylcholine model membranes. *Chem Phys Lipids* 165(1):15–22
- Pernak J, Sobaszekiewicz K, Mirska I (2003) Anti-microbial activities of ionic liquids. *Green Chem* 5:52–56
- Ranke J, Moelter K, Stock F, Bottin-Weber U, Poczubutt J, Hoffmann J, Ondruschka B, Filser J, Jastorff B (2004) Biological effects of imidazolium ionic liquids with varying chain lengths in acute *Vibrio fischeri* and WST-1 cell viability assays. *Ecotoxicol Environ Saf* 58:396–404
- Ranke J, Müller A, Bottin-Weber U, Stock F, Stolte S, Arning J, Stoermann R, Jastorff B (2007) Lipophilicity parameters for ionic liquid cations and their correlation to in vitro cytotoxicity. *Ecotoxicol Environ Saf* 67:430–438
- Rawat K, Bohidar HB (2012) Universal charge quenching and stability of proteins in 1-methyl-3-alkyl(Hexyl/Octyl) imidazolium chloride ionic liquid solutions. *J Phys Chem B* 116:11065–11074
- Richter C, Schaepe K, Glorius F, Ravoo BJ (2014) Tailor-made N-heterocyclic carbenes for nanoparticle stabilization. *Chem Commun* 50(24):3204–3207
- Rühling A, Wang D, Ernst JB, Wulff S, Honeker R, Richter C, Ferry A, Galla H-J, Glorius F (2017) Influence of the headgroup of azolium-based lipids on their biophysical properties and cytotoxicity. *Chem Eur J* 23(25):5920–5924
- Samori C, Malferrari D, Valbonesi P, Montecavalli A, Moretti F, Galletti P, Sartor G, Tagliavini E, Fabbri E, Pasteris A (2010) Introduction of oxygenated side chain into imidazolium ionic liquids: evaluation of the effects at different biological organization levels. *Ecotoxicol Environ Saf* 73:1456–1464
- Sauerbrey G (1959) Verwendung von Schwingquarzen zur Wägung dünner Schichten und zur Mikrowägung. *Z Phys A* 155(2):206–222
- Sezgin E, Levental I, Mayor S, Eggeling C (2017) The mystery of membrane organization: composition, regulation and roles of lipid rafts. *Nat Rev Mol Cell Biol* 18(6):361–374
- Simons K, Gerl MJ (2010) Revitalizing membrane rafts: new tools and insights. *Nat Rev Mol Cell Biol* 11(10):688–699
- Stepnowski P, Skladanowski AC, Ludwiczak A, Laczynska E (2004) Evaluating the cytotoxicity of ionic liquids using human cell line HeLa. *Hum Exp Toxicol* 23:513–517
- Stolte S, Arning J, Bottin-Weber U, Matzke M, Stock F, Thiele K, Uerdingen M, Welz-Biermann U, Jastorff B, Ranke J (2006) Anion effects on the cytotoxicity of ionic liquids. *Green Chem* 8:621–629
- Stolte S, Arning J, Bottin-Weber U, Müller A, Pitner WR, Welz-Biermann U, Jastorff B, Ranke J (2007) Effects of different head groups and functionalised side chains on the aquatic toxicity of ionic liquids. *Green Chem* 9:760–767
- Tischer M, Pradel G, Ohlsen K, Holzgrabe U (2012) Quaternary ammonium salts and their antimicrobial potential: targets or nonspecific interactions? *ChemMedChem* 7:22–31
- Visser A, Swatloski RP, Reichert WM, Mayton R, Sheff S, Wierzbicki A, Davis JH, Rogers RD (2001) Task-specific ionic liquids for the extraction of metal ions from aqueous solutions. *Chem Commun* 1:135–136
- Wang D, Richter C, Rühling A, Drücker P, Siegmund D, Metzler-Nolte N, Glorius F, Galla H-J (2015a) A remarkably simple class of Imidazolium-based lipids and their biological properties. *Chem Eur J* 21(43):15123–15126
- Wang D, de Jong DH, Rühling A, Lesch V, Shimizu K, Wulff S, Heuer A, Glorius F, Galla H-J (2016) Imidazolium-based lipid analogues and their interaction with Phosphatidylcholine membranes. *Langmuir* 32(48):12579–12592
- Wang D, Richter C, Rühling A, Hüwel S, Glorius F, Galla H-J (2015b) Anti-tumor activity and cytotoxicity in vitro of novel 4,5-dialkylimidazolium surfactants. *Biochem Biophys Res Commun* 467:1033–1038
- Wasserscheid P, Keim W (2000) Ionic liquids—new “solutions” for transition metal catalysis. *Angew Chem Int Ed* 39(21):3772–3789
- Welton T (1999) Room-temperature ionic liquids. Solvents for synthesis and catalysis. *Chem Rev* 99(8):2071–2084
- Welton T (2004) Ionic liquids in catalysis. *Coord Chem Rev* 248:2459–2477
- Yoo B, Zhu Y, Maginn EJ (2016) Molecular mechanism of ionic-liquid-induced membrane disruption: morphological changes to bilayers, multilayers, and vesicles. *Langmuir* 32:5403–5411