Published in final edited form as:

Semin Cell Dev Biol. 2018 October ; 82: 127–136. doi:10.1016/j.semcdb.2017.11.037.

Physiological functions of non-apoptotic caspase activity in the nervous system

Emilie Hollville¹ and Mohanish Deshmukh^{1,2,#}

¹Neuroscience Center, UNC Chapel Hill, Chapel Hill NC, USA

²Department of Cell Biology and Physiology, UNC Chapel Hill, Chapel Hill NC, USA

Abstract

Caspases are cysteine proteases that play important and well-defined roles in apoptosis and inflammation. Increasing evidence point to alternative functions of caspases where restricted and localized caspase activation within neurons allows for a variety of non-apoptotic and non-inflammatory processes required for brain development and function. In this review, we highlight sublethal caspase functions in axon and dendrite pruning, neurite outgrowth and dendrite branches formation, as well as in long-term depression and synaptic plasticity. Importantly, as non-apoptotic activity of caspases is often confined in space and time in neurons, we also discuss the mechanisms that restrict caspase activity in order to maintain the neuronal networks in a healthy and functional state.

Keywords

caspase; XIAP; pruning; neurite branching; synaptic plasticity; neuron

Introduction

Caspases belong to a family of cysteine proteases which have important functions in apoptosis, a form of programmed cell death, and inflammation. Caspases are expressed in a wide range of organisms. Initially identified in worms, 18 mammalian homologs of the *C. elegans* Cell death protein 3 (CED-3) have been described to date. However, the set of caspases expressed within mammals is heterogeneous [1]. Pro-apoptotic caspases are responsible for the proteolytic cleavage of hundreds of caspase substrates in response to pro-apoptotic stimuli, ultimately leading to the controlled fragmentation of cellular components, a process essential for the removal of unwanted or damaged cells by specialized phagocytes [2]. In the context of inflammation, a subset of caspases are responsible for the proteolytic maturation of well-defined pro-inflammatory cytokines, as well as the initiation of an inflammation specific form of cell death called pyroptosis [3]. In healthy cells, caspases are

[#]Corresponding author: mohanish@med.unc.edu; (919) 843-6005.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

expressed as inactive zymogen and their activation, which is usually initiated by proteolytic cleavage, is tightly regulated.

While the mechanisms controlling caspases activation and their targets are well established in the context of apoptosis and inflammation, accumulating evidence also support a non-apoptotic and non-inflammatory function of caspases. These functions comprise differentiation and cell fate determination (including differentiation of stem cells and terminal differentiation of keratinocytes, erythroblasts and myoblasts) as well as cell proliferation and tissue regeneration as a result of non-cell autonomous effect on survival and proliferation [4, 5]. In addition, caspases exerts non-apoptotic function in the nervous system [5–9]. Over the recent years, studies have demonstrated that restricted and localized caspase activation within neurons allows for a variety of processes that are relevant to neuronal development and function. In this review, we briefly summarize our understanding of caspase activation before exploring the physiological sublethal roles of caspases in the nervous system. We illustrate the roles of caspases in shaping neuronal networks during development and reshaping neuronal connectivity during maturation of the nervous system. We finally discuss the mechanisms that potentially confine and restrict caspase activation in the nervous system.

1. Pathways for caspase activation

Apoptotic caspases can be subdivided into initiator and effector caspases where initiator caspases (caspase-2, -8, -9 and -10) are activated within molecular platforms and are responsible for the direct proteolytic activation of effector caspases (caspase-3, -6 and -7). These effector caspases are responsible for the cleavage of hundreds of cellular substrates and are the real effectors of the apoptotic program. Two molecular platforms, activated by two different pro-apoptotic pathways, are known to activate the apoptotic initiator caspases [4].

The extrinsic, or death receptor, pathway is initiated by ligand-dependent stimulation of cell surface receptors of the tumor necrosis factor (TNF) superfamily, including TNF receptor-1 (TNFR1), Fas/CD95, TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (TRAILR1, TRAILR2), death receptor 3 (DR3), and DR6. Upon ligand-induced trimerization, these receptors, identified by the presence of a death domain and commonly called death receptors, engage into the formation of a death-inducing signaling complex called DISC. This platform consists in the trimerized receptor, the adaptors Fas associated via death domain (FADD) and/or TNFR1-associated death domain protein (TRADD) and the pro-caspase-8 or -10. Within the DISC, pro-caspase-8 homodimerizes and undergoes auto-proteolytic activation. Fully activated caspase-8 can in turn, cleave and activate the effector caspase-3, -6 and -7 [10].

Alternatively, the intrinsic, or mitochondrial, pathway originates from the detection of intracellular stress and involves a signaling cascade leading to mitochondrial outer membrane permeabilization (MOMP) and release of proteins, such as cytochrome *c* and second mitochondria-derived activator of caspases (SMAC), from the mitochondrial intermembrane space. Members of the B cell lymphoma-2 (Bcl-2) family, which are

characterized by the presence of Bcl-2 homology (BH) domains, play a key role in the control of MOMP. The pro-apoptotic members Bax and Bak are the effectors of this family, forming pores in the mitochondrial outer membrane upon oligomerization, allowing the release of mitochondrial proteins. Their activation is inhibited by interactions with anti-apoptotic members of the family (e.g. Bcl-2, Bcl-xl, Mcl-1, Bcl-b, Bcl-w, A1), while members containing a single BH domain, the so-called BH3-only proteins (e.g. Bid, Bim, Puma, Noxa, Bad, Bmf, Hrk, Bik), promote the activation of Bax and Bak by either direct interaction and/or inhibition of the anti-apoptotic Bcl-2 proteins [11, 12]. In the cytosol, cytochrome *c* promotes the assembly of another caspase activating platform: the apoptosome. The adaptor apoptotic protease activating factor 1 (APAF-1) oligomerizes upon binding to cytosolic cytochrome *c* and ATP, promoting the recruitment and activation of pro-caspase-9. Formation of the apoptosome subsequently promotes the proteolytic activation of pro-caspase-3, -6 and -7 [13].

2. Pruning of Axons and Dendrites

During development, neurons extend their axons to innervate their target regions often resulting in superfluous connections. This outgrowth phase is followed by a regressive phase where excessive or inappropriate axons, dendrites and synapses are eliminated while suitable connections are maintained. The selective elimination of unwanted axons, dendrites, and synapses is known as pruning and occurs without the death of the parent neuron [14, 15]. Pruning is essential for the refinement of neuronal connectivity and establishment of a mature and functional network. In vertebrates, pruning takes place largely during early postnatal development. Classical examples of developmental pruning occur in response to limited neurotrophic factors for sensory and sympathetic neurons, at the neuromuscular junction, in the midbrain where retinal ganglion cells project their axons as well as in the cortex and hippocampus [14, 15]. During insect metamorphosis, large-scale pruning allows larval processes to remodel and form adult-specific connections [16]. Evidence that caspase activity is required for developmental pruning has emerged from both genetic and biochemical studies in multiple models.

Caspase function in pruning was first shown in the *Drosophila* model. During metamorphosis, remodelling of class IV sensory neurons innervating the epidermis involves the elimination of larval dendrites and the subsequent regrowth of adult-specific dendrites. Mutants of the initiator caspase Death regulator Nedd2-like caspase (DRONC) fail to prune the larval dendrites of these sensory neurons [17, 18]. In this context, pruning requires Death-associated APAF1-related killer (DARK), the fly homolog of APAF-1 that is necessary for activating caspases *via* the apoptosome complex. Moreover, overexpression of p35, the baculovirus inhibitor of effector caspases, also inhibits dendritic pruning, suggesting that effector caspases are also important for dendrite pruning during metamorphosis [18]. Importantly, unlike in the context of cell death, caspase activity is spatially restricted to the dendrites of neurons undergoing pruning [17, 18]. Effector caspases have also been implicated in the large-scale pruning of axons of the retinal ganglion cells (RGC) that occurs in the midbrain of mammals. During embryonic development, RGC extend their axons in the superior colliculus but largely overshoot their targets. Neuronal activity promotes the elimination of these inappropriate extensions during

the first postnatal week in order to refine the eye-specific projection map [19, 20]. In mice deficient in caspase-3, -6, or the caspase-3 target calpastatin, RGC axon projections remain outside of their targeted area in the superior colliculus beyond this refinement period [21, 22].

The role of caspases in axon pruning has also been studied in the context of axon degeneration induced by neurotrophic factor deprivation in sensory and sympathetic neurons (Figure 1). During development, neurotrophin-responsive neurons extend their axons to innervate target regions producing neurotrophins such as Neuronal Growth Factor (NGF). *In vitro*, depletion of NGF induces the apoptotic death of Dorsal Root Ganglia (DRG) and Superior Cervical Ganglia (SCG) neurons [23, 24]. However, deprivation of NGF from only the distal axons of these neurons cultivated in compartmentalized chambers (referred to as “local NGF deprivation”), promotes axon pruning without causing neuronal death [25]. The involvement of caspases in axon pruning, initially reported using the caspases inhibitors zVEID and zDEVD [26], was confirmed with the observation that axons of DRG and SCG derived from caspase-3, caspase-6 or caspase-9-deficient mice are protected against axonal NGF deprivation [21, 27]. Although caspases are present in both axons and cell bodies [28–31], caspase activation is spatially restricted to the axonal compartment where cleaved caspase-3 and caspase-6 can be detected during pruning [21, 27].

The modalities and sequence of caspase activation during axon pruning is not yet well understood. Since addition of the transcription inhibitor Actinomycin D in the soma compartment preserves axons from degeneration, caspase activation during pruning is thought to be regulated by gene transcription [32]. The transcription factor c-Jun was found to be phosphorylated in a JNK- and DLK-dependent manner in the cell bodies of neurons locally deprived of NGF [28, 33]. Moreover, chemical inhibition of the MAP kinases p38, JNK or DLK in the axonal compartment prevents axonal pruning, suggesting that these kinases are activated locally at the site of NGF deprivation [32–34]. This signalling pathway most likely triggers the intrinsic mitochondrial pathway of apoptosis as JNK/c-Jun has previously been involved in transcriptional upregulation of the BH3-only proteins Hrk, Bim and Puma that activate the pro-apoptotic members of the Bcl-2 family [23]. Consistent with this model, anti-apoptotic Bcl-2 family members Bcl-xl and Bcl-w protect against axonal degeneration while the BH3-only protein Puma and pro-apoptotic Bcl-2 family member Bax are required for axon pruning [26, 27, 34–36]. Moreover, cytochrome *c* is released from mitochondria in response to local NGF withdrawal and genetic analysis suggests that caspase-6 activation occurs downstream of caspase-3, caspase-9 and Bax engagement [27]. However, the adaptor APAF-1, is dispensable for axon pruning [27], raising the possibility that caspase activation during axon pruning occurs independently of the apoptosome (Figure 1).

In mammals, a role for the death receptor DR6 has been implicated in pruning. Genetic studies suggest that DR6 regulates dendritic spine density in cortical neurons in young mice [37, 38] and also RGC axon pruning in the superior colliculus [26]. However, the modalities of DR6 activation and its involvement in pruning are still unclear. The extracellular domain of DR6 was initially reported to bind the cleaved amino terminal fragment of amyloid precursor protein (APP), the precursor of Amyloid β in Alzheimer's disease, but the authors

subsequently reported that DR6 was rather binding to the extracellular domain 2 of APP [26, 39, 40]. Interestingly, RGC from APP-deficient mice present pruning defects similar to the DR6-deficient mice and epistasis analysis revealed that RGC pruning defect in DR6-deficient mice was not enhanced by loss of APP, suggesting that APP and DR6 function in the same pathway [40]. Although, in other systems, DR6 was shown to recruit the adaptor protein TRADD and activate Bax in a caspase-8-independent manner [41, 42], there have not been any reports of the mechanism by which DR6 may activate caspases in the context of axon pruning. Moreover, the modalities regulating APP-DR6 interaction and the consequences of this interaction during pruning remain to be resolved.

3. Neurite outgrowth and arborization (branching, axonal guidance)

Axon outgrowth, pathfinding and arborization are essential during development in order to establish functional neuronal connectivity. Guidance proteins play a crucial role in this process and mutations in guidance molecules or their receptors have been associated with several neurological disorders including congenital axon guidance disorders, autism and neurodegenerative diseases [43]. Guidance proteins are either secreted or membrane bound factors which are recognized by specific receptors on the surface of neuronal growth cones. They act as attractant or repellent to guide growing axons to their specific target regions during development [43]. Their action is localized, as growth cones of severed axons are still repelled or attracted by these factors [44]. Caspases were first reported to participate in neurite outgrowth and guidance in the context of Netrin-1 chemoattractant's effect on *Xenopus* retinal ganglion cells (Figure 2) [45]. Netrin-1 is a secreted guidance protein which, upon binding to its receptors DCC and UNC5, promotes growth cone attraction or repulsion, respectively [46]. *In vitro*, caspase-3 is rapidly activated in growth cones of retinal neurons stimulated with Netrin-1 and participates in the Netrin-1-mediated attractive chemotrophic effect. Mouse hippocampal neurons also have been reported to express active caspase-3 in their growth cone *in vitro*, and NCAM clustering triggers neurite outgrowth in a caspase-3/8-dependent manner [47]. In addition, developing olfactory bulb of rodents activate caspase-3 in a caspase-9-dependent manner. Interestingly, deficiency in APAF-1 and caspase-9 does not affect the number of olfactory neurons, it rather results in aberrant projection and axon trajectories in olfactory sensory neurons [48].

Slit constitute another family of guidance molecules that are able to activate caspases upon binding to their Robo receptors. Similar to many guidance proteins, the outcome of Slit-Robo signalling is highly dependent on the context. It has been reported to have both attractive and repulsive effect on growing axons, as well as both permissive and repressive effect on arborization during development [49–51]. *In vivo*, Slit1a and its receptor Robo2 stimulate local caspase-3 activity in zebrafish embryonic retinal ganglion cells, in absence of neuronal death (Figure 2). In this context, caspase-3 activity is restricted spatially - at branching points but not in dendrites or cell bodies - and temporally - over a short period of time when arborization is most dynamic. Time-lapse imaging revealed that caspase-3 is activated upon branch formation and promotes both branch formation and retraction. Further, Slit1a, Robo2, caspase-9 and caspase-3 cooperate in a pathway required for maintaining newly formed branch tips and presynaptic terminal in a dynamic, unstable phase [52]. A Bax/Bak/caspase pathway seems to be also involved in controlling axonal branching

at postnatal stages. Skilled movement acquisition during early postnatal development requires motor circuit reorganization through refinement of corticospinal axons connectivity. A recent study found caspase-3 to be activated in a Bax/Bak-dependent manner in axons of corticospinal neurons in the spinal cord of neonatal mice. Early postnatal inactivation of Bax/Bak in motor cortex results in increased axonal branches in the spine of these animals once they reach adult age. As a result, refinement of connectivity is lost and animals fail to acquire fine voluntary movements [53]. Although neuronal activity is known to be a prerequisite for branch formation [54], the upstream signalling pathways responsible for caspase activation in this context is currently unknown.

To the best of our knowledge, there is currently no report of activation of non-apoptotic caspase function in response to other type of guidance proteins, including Semaphorins, Ephrins and Repulsive Guidance Molecules (RGMs) in neurons. However, several members of these proteins or their receptors have been reported to induce caspase activation and apoptosis. For instance, Semaphorin 3A induces caspase-dependent cell death in sympathetic neurons through binding to its receptors PlexinA3 and Neuropilin-1 [55, 56]. Additionally, overexpression of Neogenin, the receptor for RGMs, has been shown to induce apoptosis in neural tube of chick embryos and immortalized neuroblasts [57]. Overexpression of Ephrin-A5 in neurons expressing its receptor EphA7 induces caspase-3 activation and apoptosis in cortical progenitors [58]. Therefore, it is possible that in a context of local activation, these guidance molecules trigger caspase activation to promote their repulsive or attractive effect. Differential outcome of local *versus* global receptor stimulation has been observed in the case of the TNF receptor superfamily member lymphotoxin beta receptor (LT- β R). LIGHT stimulate cell death upon binding to its receptor LT- β R in non-neuronal and motoneuron cultures [59, 60]. Conversely, when locally applied to the axons of motoneurons, LIGHT promotes axonal outgrowth and branching [60]. Moreover, the outgrowth and branching promoting effect of LIGHT can be inhibited by the use of the caspase-9 inhibitor LEHD. Rather surprisingly however, the DEVD inhibitor has no effect in the same experimental conditions [60].

Axonal injury often permits neurons to survive and is sometimes associated with sprouting at the axon tip, allowing formation of a growth cone and axon extension. Axon segments which are separated from their cell bodies are able to regrow, suggesting that axonal regeneration most likely is a local process [61]. Although inhibition of caspases has been reported to promote regeneration by inhibiting apoptosis [62–64], there is evidence that axon stabilization and axon regeneration are antagonistic phenomenon [65]. Rather counterintuitively, caspases have been reported to promote regenerative neurite outgrowth in response to injury. For instance, caspase-3 inhibition reduces growth cone formation and axonal regeneration following *in vitro* axotomy in dorsal root ganglion cells from embryonic and postnatal rats [66]. In *C. elegans*, axonal outgrowth in response to *in vivo* axotomy of motor and mechanosensory neurons is reduced in the APAF-1 homolog CED-4 and in the caspase homolog CED-3 inactive mutants [67]. Also, upon sciatic nerve crush in mice, LIGHT expression is stimulated in B lymphocytes and LIGHT is involved in neuromuscular junction regeneration. However, it remains unknown whether caspases are required for this regenerative function of LIGHT as observed during LIGHT-induced outgrowth and branching [60].

Exactly how caspases are activated in the context of axonal outgrowth and arborization and how they promote outgrowth and arborization is largely unknown. A few guidance receptors have been reported to recruit caspases, including the EphrinA receptors EphA7 and EphA4 which are able to recruit caspase-8 [68] and the Netrin-1 receptor DCC which is able to interact with caspase-3 and caspase-9 [69]. However, NCAM is the only receptor that has been shown to recruit caspase-8 in the context of neurite outgrowth [47]. Although members of the UNC5 receptors family possess a death domain in their cytoplasmic portion [70], it is currently unknown if this domain play any role in caspase activation.

4. Long-term depression and synapse plasticity

Synaptic plasticity plays a major role in developmental maturation of neuronal circuits as well as in experience-dependent remodelling of neuronal circuits. Durable synapse modification in response to neuronal stimulation can result in enhanced or reduced synaptic strength and is known as Long-Term Potentiation (LTP) or Long-Term Depression (LTD), respectively [71]. Activity-dependent change in synaptic strength is largely recognized as a mechanism associated with spatial learning and memory in the hippocampus, fear memory in the amygdala, task memory in the cortex and learning in the cerebellum [72]. The best characterized form of LTP and LTD is experienced by excitatory synapses in response to L-glutamate. Glutamate binds notably to N-methyl-D-aspartate receptor (NMDA) receptors (NMDARs), α -amino-3-hydroxy-5-methylisoxazole-4-propionic acid (AMPA) receptors (AMPA receptors) and metabotropic glutamate receptors (mGluRs) and triggers an increase in calcium in the stimulated spine. The outcome of calcium entry is a gain in AMPAR expression at the synapses experiencing LTP [73] while AMPA receptors are internalized at synapses undergoing LTD [74]. Modulation of synaptic strength during LTP and LTD is also associated with remodelling of dendritic spine size and density which are increased during LTP and reduced during LTD. Similar to synaptic strength, spine shrinkage during activity-dependent LTD requires NMDAR and calcium-calmodulin signalling [75, 76].

The use of caspase inhibitors *in vivo* has uncovered a functional role for caspases in long-term spatial memory formation [77], in avoidance learning [78] and in auditory memory formation [79]. Consistently with a role for caspases in memory formation, active caspase can be detected in the hippocampus [77] and cortex [78] of mice. As would be anticipated, caspase activation seems to be transient and restricted to postsynaptic structures in these contexts [79, 80]. Mechanistically, NMDAR-LTD is inhibited in hippocampal neurons from caspase-3-deficient mice, suggesting that this effector caspase plays an active role in NMDAR-dependent LTD [81] and caspase-3 is sufficient to reduce synaptic strength [80]. The role of caspases in synapse plasticity might not be restricted to excitatory glutamatergic synapses as there is also genetic evidence that CED-4/APAF-1 participates in inhibitory GABAergic synapse elimination in worms [82]. Although caspase-3 deficiency has been shown to have no impact on high frequency stimulation-induced LTP in mice hippocampal neurons [81], it is possible that caspases regulate LTP in different contexts. Long exposure to caspase inhibitors have been reported to inhibit LTP in rat hippocampal neurons [83] but also in snail parietal neurons [84], reports that have not been confirmed genetically. Contradictory studies have however suggested that caspases could negatively regulate LTP in hippocampal neurons, either directly using inhibitors in the context of high-frequency

stimulation [85] or in response to LTP antagonists such as amyloid β [86] or aluminium [87].

Caspase-3-dependent control of synaptic plasticity relies on Bax-mediated mitochondrial membrane permeabilization in order to promote NMDAR-LTD (Figure 3) [80, 81] as well as the amyloid β -mediated inhibition of LTP, which is also NMDAR-dependent [88]. Similar to apoptotic triggers [89], NMDA induces Bad dephosphorylation in a calcineurin-dependent manner to promote its activation and relocation to mitochondria in hippocampal slices [80]. Calcium influx through NMDAR, which is essential for LTP and LTD, therefore also triggers a signalling pathway leading to caspase-3 activation to promote NMDAR-LTD or inhibition of LTP. Consistent with a transient activation of caspase-3 in response to synaptic activity, activated Bad and Bax, and also released cytosolic cytochrome *c*, are similarly short lived in this context [80]. The transitory nature of this pathway may be required to maintain caspase-3 activity at a low level in order to prevent neurons from undergoing apoptosis [80] and it is possible that neurological synapses activate a negative feedback loop to restrain caspase-3 activation in these situations. In other cellular context such as immunological synapse between activated dendritic cells and T lymphocytes, a similar glutamate-NMDAR-calcium pathway triggers sustained caspase-3 activation and apoptosis [90].

Several possible roles for effector caspases have been suggested in the context of synaptic plasticity. While the outcome of LTP/LTD converges on controlling the expression of AMPAR at the surface of synapses, there is evidence to support a role of caspase-3 in the regulation of AMPAR distribution. Deficiency in the pro-apoptotic proteins that have been implicated in LTD, including, Bad, Bax and caspase-3, result in impairment of NMDAR-induced AMPAR internalisation in hippocampal neurons [80, 81]. How caspase-3 could mediate this effect is not exactly known. Han and colleagues have identified the synaptic protein Gap43 as a caspase-3 substrate for which its caspase-dependent cleavage seems to be required for LTD and AMPAR internalisation in hippocampal neurons [91]. Caspase-3 could also promote AMPAR internalisation *via* increased activation of its substrate calcineurin [92], which is required for NMDAR-induced endocytosis of AMPAR during LTD [74]. As an alternative mechanism to caspase-mediated AMPAR internalisation, it has been suggested that caspases directly cleave AMPAR subunits as this has been observed in apoptotic hippocampal neurons in response to amyloid β , staurosporine or starvation [93, 94]. Another substrate of caspase-3 recognized to influence LTD and LTP is Akt1. During apoptosis, caspase-mediated cleavage of Akt1 results in loss of Akt1 kinase activity [95]. In the context of synaptic plasticity, cleavage of Akt1 is required for NMDAR-LTD as well as for amyloid β -mediated repression of LTP [81, 86]. The PI3K-Akt1 pathway is known to antagonize LTD and to repress the activity of GSK3 β [96]. Because GSK3 β is activated and required for NMDAR-LTD [96] as well for amyloid β -mediated inhibition of LTP [86], it is possible that caspase-3 promotes LTD and suppress LTP *via* repression of Akt1 and promotion of GSK3 β activity (Figure 3).

Dendritic spine density remodelling that occurs during LTP and LTD [75, 76] is another aspect of synaptic plasticity involving caspase-3 activity [97]. In response to neuronal activity, the neurotrophin BDNF is secreted as a precursor (proBDNF), which binds the neurotrophin receptor p75NTR. ProBDNF is processed by extracellular proteases to its

mature form (mBDNF), which binds the TrkB receptor [98]. BDNF has been reported to play a role in hippocampal memory both in humans and in rodents [99]. ProBDNF and mBDNF have opposite effects on synaptic strength and synaptic density remodelling. Mature BDNF increases spine density and promotes LTP in a TrkB-dependent manner [100, 101]. Conversely, proBDNF has been shown to promote NMDAR-LTD through its receptor p75NTR, which localizes at synapses [102]; proBDNF can also decrease dendrite arbor complexity and spine density [97, 103]. ProBDNF signalling triggers the mitochondrial pathway and caspase-3 activation that is required for its ability to promote the reduction in spine density [97]. Although the role of caspase-3 in proBDNF-induced LTD has not been directly assessed, these combined studies strongly support a role for caspases in synaptic plasticity and memory formation.

5. How is non-apoptotic caspase activity confined and restricted in neurons?

As illustrated in the case of pruning, dendrite branches formation and synaptic plasticity, non-apoptotic caspase function is often restricted in time and space whether to dendrites during pruning, branching points during arborization or postsynaptic buttons during LTD. Localized activation of caspase in living neurons contrasts with global caspase activity observed during apoptosis and can only be achieved if regulatory mechanisms limit caspase activity in order to prevent the loss of neurons that would occur in response to widespread caspase activation [5].

Several mechanisms have been described to restrict caspase function in neurons (Figure 4). The best characterized regulators of caspase function in neurons belongs to the family of Inhibitor of Apoptosis Proteins (IAP). IAPs are identified by the presence of Baculovirus IAP Repeat (BIR) domains which mediate protein-protein interactions. Several IAPs, including those involved in caspase regulation, also possess a Really Interesting New Gene (RING) domain that confers IAPs an E3 ligase activity [104]. IAPs interfere with caspase function either by direct binding or by promoting their ubiquitination and occasionally their degradation. For example, *Drosophila* DIAP1 restricts both the initiator caspase DRONC, either in its monomeric inactive form or as part of the apoptosome [105–107], and the activated effector caspase death related ICE-like caspase (DRICE) [108–110], through ubiquitination. DIAP2, on the other hand, only ubiquitinates activated DRICE [111, 112]. Protein ubiquitination is often associated with proteasomal degradation. However, degradation of caspases is rarely observed as a consequence of ubiquitination by DIAPs. DRONC targeting to the proteasome pathway only seems to occur in the context of the apoptosome following ubiquitination by DIAP1 [113]. Otherwise, DIAP1 or DIAP2 rather promote non-degradative ubiquitination and inactivation of DRICE or monomeric DRONC [105, 109, 111]. Mammals express several IAPs, of which only X-linked inhibitor of apoptosis (XIAP) can efficiently interfere with a specific set of caspases, namely, caspase-9, -3 and -7 [114]. Unlike DIAP1 and DIAP2, XIAP does not seem to target caspases for ubiquitination but rather inhibits caspases activity by direct binding, obstructing the catalytic site of caspase-3 and -7 or the homo-dimerization domain of caspase-9 [115–118].

The efficiency of IAPs at preventing caspase activation and apoptosis in different mammalian cells is variable. In neurons however, IAPs play a major role in keeping caspases inactive and protecting neurons from death [30, 119, 120]. Interestingly, the increased effectiveness of XIAP in restricting caspases in neurons is not because of high XIAP expression but rather due to low levels of the adaptor APAF-1 in neurons, which limits the extent of caspase activation allowing for strict control by XIAP [121]. IAPs also protect the cell bodies when local caspase activity is triggered in the dendrites, thus keeping caspase activity localized during dendrite elimination [120]. XIAP was also shown to restrict caspase-3 activation and axon degeneration in the NGF deprivation model of axon pruning *in vitro* [27]. Consistent with this function of XIAP, neurons innervating the skin of mice deficient in XIAP have reduced axonal length, while overexpression of XIAP prevents pruning [30]. Finally, XIAP was shown to restrict AMPA receptor internalization and caspase-3 activation during long-term depression [81, 122, 123].

Interestingly, one mechanism that allows for the restricted caspase activation is by degradation of IAPs in localized region. For example, caspase-3 activation during pruning of dendrites from sensory neurons during fly metamorphosis requires degradation of DIAP1 by the proteasome [17]. XIAP stability was recently shown to be enhanced by a brain-specific isoform of Fas Apoptotic Inhibitory Molecule (FAIM-L). Endogenous FAIM-L notably interferes with AMPA receptor internalization in the context of LTD, while overexpression of FAIM-L is able to stabilize XIAP, inhibit caspase-3 activation and prevent LTD as well as pruning [122].

These observations suggest that IAPs function is tightly regulated in neurons. IAPs are stabilized when and where caspase function needs to be repressed, and conversely, IAPs are inactivated when and where limited caspase activity is required. Although proteasomal degradation of XIAP and DIAP1 has been shown to be required for effector caspases activation during pruning [17], very little is known about the mechanisms controlling XIAP or DIAP1 to allow for local caspase activity under physiological conditions in neurons. In the context of apoptosis, IAPs activity and stability is regulated by their inhibitory proteins, which in mammals include the protease Omi/HtrA2 and the IAP antagonist Smac/DIABLO, and in flies, Grim, Hid, Reaper and Sickie. These mitochondrial proteins are characterized by the presence of a conserved IAP-binding motif (IBM), necessary for binding and antagonizing IAPs [124]. In addition, the septin ARTS (Apoptosis-Related protein in the TGF- β Signaling pathway) was found to antagonize XIAP through a different mechanism [125]. ARTS, which also localizes at mitochondria, binds XIAP via a different motif and recruits the E3 ligase Siah1 to XIAP. Ubiquitination of XIAP by Siah1 results in XIAP degradation and promotion of apoptosis [126]. The role of these IAP antagonists in the activation of non-apoptotic function of caspases remains obscure. Interestingly, Hid has been implicated in non-apoptotic activation of caspases in neurons as tight control of Hid levels during larval remodelling of class IV sensory neurons is required to attain an balanced degree of dendritic growth and retraction [127].

Other modifications of XIAP or DIAP1 have also been shown to alleviate their inhibition on caspase activity. For instance, XIAP and DIAP1 can be phosphorylated by the IKK ϵ kinase. In *Drosophila*, inactivation of I κ B kinase ϵ (IKK ϵ) suppress Reaper-induced cell death,

suggesting that IKK ϵ is activated downstream of the IAP antagonist [128]. Both mammalian and fly IKK ϵ interact with, phosphorylate and mediate proteasomal degradation of XIAP or DIAP1 respectively, therefore promoting caspase-3/DRONC activation [128, 129]. Interestingly, fly IKK ϵ was suggested to affect the non-apoptotic function of caspases during development rather than developmental cell death [128]. Further studies will be required to determine if IKK ϵ is also activated and recruited to XIAP in response to release of IAP antagonists in the cytosol and if this kinase participate in caspase activation in neuronal physiology. Another posttranslational modification that can promote the inactivation of XIAP is S-nitrosylation. S-nitrosylation of XIAP impairs its ability to inhibit caspase-3. It is however unclear how S-nitrosylation affects XIAP function as it was shown to either not affect or reduce XIAP E3 ligase function [130, 131]. Functionally, XIAP S-nitrosylation reduces XIAP capacity to inactivate caspase-3 in response to NMDA-induced excitotoxicity [131], suggesting that local S-nitrosylation of XIAP might be involved in restricted caspase activation during long-term depression. Consistently, Alzheimer's disease which is associated with increase LTD, synaptic loss and synaptic caspase-3 activity [86, 132], is also associated with increased S-nitrosylated XIAP [131].

6. Perspectives for the future

Nearly 25 years after the discovery of caspase function during apoptosis [133], the most unexpected aspects of caspase function that have emerged is the finding that many of the same caspases that can induce cell death are also activated in physiological contexts of neuronal pruning, axon guidance, and synaptic plasticity. While the function of caspases in these non-apoptotic situations in the brain is now firmly established, several important question remain unanswered. Exactly how are caspases activated in these physiological contexts? Are the mechanisms the same as seen during apoptosis? Are these differences in how caspases are activated during dendritic or axonal pruning, or axon guidance, or synaptic plasticity? How is caspase activity restricted, both spatially and temporally, in these situations so as to avoid apoptosis? While the importance of the XIAP in restricting caspases in several of these situations has become clear, are there other mechanisms? Most importantly, do these caspase restraining mechanisms in non-apoptotic physiological contexts become less effective and ultimately contribute to the neuronal loss in the pathological contexts of neurodegeneration?

Acknowledgments

Due to space limitations, we apologize for not including all the papers relevant to this literature. This work was supported by a grant from National Institutes of Health (GM118331) to MD.

References

1. Eckhart L, Ballaun C, Hermann M, VandeBerg JL, Sipos W, Uthman A, Fischer H, Tschachler E. Identification of Novel Mammalian Caspases Reveals an Important Role of Gene Loss in Shaping the Human Caspase Repertoire. *Mol Bio Evol.* 2008; 25(5):831–841. [PubMed: 18281271]
2. Taylor RC, Cullen SP, Martin SJ. Apoptosis: controlled demolition at the cellular level. *Nat Rev Mol Cell Biol.* 2008; 9(3):231–241. [PubMed: 18073771]
3. Broz P, Dixit VM. Inflammasomes: mechanism of assembly, regulation and signalling. *Nat Rev Immunol.* 2016; 16(7):407–420. [PubMed: 27291964]

4. Shalini S, Dorstyn L, Dawar S, Kumar S. Old, new and emerging functions of caspases. *Cell Death Differ.* 2015; 22:526–539. [PubMed: 25526085]
5. Unsain N, Barker Philip A. New views on the misconstrued: executioner caspases and their diverse non-apoptotic roles. *Neuron.* 2015; 88(3):461–474. [PubMed: 26539888]
6. Hyman BT, Yuan J. Apoptotic and non-apoptotic roles of caspases in neuronal physiology and pathophysiology. *Nat Rev Neurosci.* 2012; 13(6):395–406. [PubMed: 22595785]
7. Mukherjee A, Williams DW. More alive than dead: non-apoptotic roles for caspases in neuronal development, plasticity and disease. *Cell Death Differ.* 2017; 24(8):1411–1421. [PubMed: 28644437]
8. D'Amelio M, Sheng M, Cecconi F. Caspase-3 in the central nervous system: beyond apoptosis. *Trends Neurosci.* 2012; 35(11):700–709. [PubMed: 22796265]
9. Li Z, Sheng M. Caspases in synaptic plasticity. *Mol Brain.* 2012; 5(1):15. [PubMed: 22583788]
10. Ashkenazi A, Salvesen G. Regulated Cell Death: Signaling and Mechanisms. *Annu Rev Cell Dev Biol.* 2014; 30(1):337–356. [PubMed: 25150011]
11. Luna-Vargas MPA, Chipuk JE. Physiological and Pharmacological Control of BAK, BAX, and Beyond. *Trends Cell Biol.* 2016; 283:2676–2689.
12. Czabotar PE, Lessene G, Strasser A, Adams JM. Control of apoptosis by the BCL-2 protein family: implications for physiology and therapy. *Nat Rev Mol Cell Biol.* 2014; 15(1):49–63. [PubMed: 24355989]
13. Green DR, Kroemer G. The Pathophysiology of Mitochondrial Cell Death. *Science.* 2004; 305(5684):626–629. [PubMed: 15286356]
14. Riccomagno MM, Kolodkin AL. Sculpting neural circuits by axon and dendrite pruning. *Annu Rev Cell Dev Biol.* 2015; 31:779–805. [PubMed: 26436703]
15. Schuldiner O, Yaron A. Mechanisms of developmental neurite pruning. *Cell Mol Life Sci.* 2014;1–19.
16. Yu F, Schuldiner O. Axon and dendrite pruning in *Drosophila*. *Curr Opin Neurobiol.* 2014; 27(0): 192–198. [PubMed: 24793180]
17. Kuo CT, Zhu S, Younger S, Jan LY, Jan YN. Identification of E2/E3 ubiquitinating enzymes and caspase activity regulating *Drosophila* sensory neuron dendrite pruning. *Neuron.* 2006; 51(3):283–290. [PubMed: 16880123]
18. Williams DW, Kondo S, Krzyzanowska A, Hiromi Y, Truman JW. Local caspase activity directs engulfment of dendrites during pruning. *Nat Neurosci.* 2006; 9(10):1234–1236. [PubMed: 16980964]
19. McLaughlin T, Torborg CL, Feller MB, O'Leary DD. Retinotopic map refinement requires spontaneous retinal waves during a brief critical period of development. *Neuron.* 2003; 40(6): 1147–1160. [PubMed: 14687549]
20. Luo L, O'Leary DD. Axon retraction and degeneration in development and disease. *Annu Rev Neurosci.* 2005; 28:127–156. [PubMed: 16022592]
21. Simon DJ, Weimer RM, McLaughlin T, Kallop D, Stanger K, Yang J, O'Leary DD, Hannoush RN, Tessier-Lavigne M. A caspase cascade regulating developmental axon degeneration. *J Neurosci.* 2012; 32(49):17540–17553. [PubMed: 23223278]
22. Yang J, Weimer Robby M, Kallop D, Olsen O, Wu Z, Renier N, Uryu K, Tessier-Lavigne M. Regulation of axon degeneration after injury and in development by the endogenous calpain inhibitor calpastatin. *Neuron.* 2013; 80(5):1175–1189. [PubMed: 24210906]
23. Kristiansen M, Ham J. Programmed cell death during neuronal development: the sympathetic neuron model. *Cell Death Differ.* 2014; 21:1025–1035. [PubMed: 24769728]
24. Buss RR, Sun W, Oppenheim RW. Adaptive roles of programmed cell death during nervous system development. *Annu Rev Neurosci.* 2006; 29:1–35. [PubMed: 16776578]
25. Geden MJ, Deshmukh M. Axon degeneration: context defines distinct pathways. *Curr Opin Neurobiol.* 2016; 39:108–115. [PubMed: 27197022]
26. Nikolaev A, McLaughlin T, O'Leary DD, Tessier-Lavigne M. APP binds DR6 to trigger axon pruning and neuron death via distinct caspases. *Nature.* 2009; 457(7232):981–989. [PubMed: 19225519]

27. Cusack CL, Swahari V, Hampton Henley W, Michael Ramsey J, Deshmukh M. Distinct pathways mediate axon degeneration during apoptosis and axon-specific pruning. *Nat Commun.* 2013; 4:1876. [PubMed: 23695670]
28. Mok SA, Lund K, Campenot RB. A retrograde apoptotic signal originating in NGF-deprived distal axons of rat sympathetic neurons in compartmented cultures. *Cell Res.* 2009; 19(5):546–560. [PubMed: 19188931]
29. Schoenmann Z, Assa-Kunik E, Tiomny S, Minis A, Haklai-Topper L, Arama E, Yaron A. Axonal degeneration is regulated by the apoptotic machinery or a NAD⁺-sensitive pathway in insects and mammals. *J Neurosci.* 2010; 30(18):6375–6386. [PubMed: 20445064]
30. Unsain N, Higgins Julia M, Parker Kristen N, Johnstone Aaron D, Barker Philip A. XIAP regulates caspase activity in degenerating axons. *Cell Rep.* 2013; 4(0):751–763. [PubMed: 23954782]
31. Akpan N, Serrano-Saiz E, Zacharia BE, Otten ML, Ducruet AF, Snipas SJ, Liu W, Velloza J, Cohen G, Sosunov SA, Frey WH, Salvesen GS, Connolly ES, Troy CM. Intranasal Delivery of Caspase-9 Inhibitor Reduces Caspase-6-Dependent Axon/Neuron Loss and Improves Neurological Function after Stroke. *J Neurosci.* 2011; 31(24):8894–8904. [PubMed: 21677173]
32. Chen M, Maloney JA, Kallop DY, Atwal JK, Tam SJ, Baer K, Kissel H, Kaminker JS, Lewcock JW, Weimer RM, Watts RJ. Spatially coordinated kinase signaling regulates local axon degeneration. *J Neurosci.* 2012; 32(39):13439–13453. [PubMed: 23015435]
33. Ghosh AS, Wang B, Pozniak CD, Chen M, Watts RJ, Lewcock JW. DLK induces developmental neuronal degeneration via selective regulation of proapoptotic JNK activity. *J Cell Biol.* 2011; 194(5):751–764. [PubMed: 21893599]
34. Simon David J, Pitts J, Hertz Nicholas T, Yang J, Yamagishi Y, Olsen O, Teši Mark M, Molina H, Tessier-Lavigne M. Axon degeneration gated by retrograde activation of somatic pro-apoptotic signaling. *Cell.* 2016; 164(5):1031–1045. [PubMed: 26898330]
35. Cosker KE, Pazyra-Murphy MF, Fenstermacher SJ, Segal RA. Target-Derived Neurotrophins Coordinate Transcription and Transport of Bclw to Prevent Axonal Degeneration. *J Neurosci.* 2013; 33(12):5195–5207. [PubMed: 23516285]
36. Maor-Nof M, Romi E, Shalom Hadas S, Ulisse V, Raanan C, Nof A, Leshkowitz D, Lang R, Yaron A. Axonal Degeneration Is Regulated by a Transcriptional Program that Coordinates Expression of Pro- and Anti-degenerative Factors. *Neuron.* 2016; 92(5):991–1006. [PubMed: 27889097]
37. Kallop DY, Meilandt WJ, Gogineni A, Easley-Neal C, Wu T, Jubb AM, Yaylaoglu M, Shamloo M, Tessier-Lavigne M, Searce-Levie K, Weimer RM. A death receptor 6-amyloid precursor protein pathway regulates synapse density in the mature CNS but does not contribute to Alzheimer's disease-related pathophysiology in murine models. *J Neurosci.* 2014; 34(19):6425–6437. [PubMed: 24806669]
38. Marik SA, Olsen O, Tessier-Lavigne M, Gilbert CD. Death Receptor 6 Regulates Adult Experience-Dependent Cortical Plasticity. *J Neurosci.* 2013; 33(38):14998–15003. [PubMed: 24048829]
39. Xu K, Olsen O, Tzvetkova-Robev D, Tessier-Lavigne M, Nikolov DB. The crystal structure of DR6 in complex with the amyloid precursor protein provides insight into death receptor activation. *Genes Dev.* 2015; 29(8):785–790. [PubMed: 25838500]
40. Olsen O, Kallop DY, McLaughlin T, Huntwork-Rodriguez S, Wu Z, Duggan CD, Simon DJ, Lu Y, Easley-Neal C, Takeda K, Hass PE, Jaworski A, O'Leary DDM, Weimer RM, Tessier-Lavigne M. Genetic analysis reveals that amyloid precursor protein and death receptor 6 function in the same pathway to control axonal pruning independent of β -secretase. *J Neurosci.* 2014; 34(19):6438–6447. [PubMed: 24806670]
41. Hu R, Du Q, Yin X, Li J, Wang T, Zhang L. Agonist antibody activates death receptor 6 downstream signaling involving TRADD recruitment. *FEBS Lett.* 2014; 588(3):401–407. [PubMed: 24374337]
42. Zeng L, Li T, Xu DC, Liu J, Cui MZ, Fu X, Xu X. Death Receptor 6 Induces Apoptosis Not Through Type I or Type II Pathways, but Via a Unique Mitochondria-dependent Pathway by Interacting with Bax. *J Biol Chem.* 2012
43. Van Battum EY, Brignani S, Pasterkamp RJ. Axon guidance proteins in neurological disorders. *Lancet Neurol.* 2015; 14(5):532–546. [PubMed: 25769423]

44. Campbell DS, Holt CE. Chemotropic responses of retinal growth cones mediated by rapid local protein synthesis and degradation. *Neuron*. 2001; 32(6):1013–1026. [PubMed: 11754834]
45. Campbell DS, Holt CE. Apoptotic pathway and MAPKs differentially regulate chemotropic responses of retinal growth cones. *Neuron*. 2003; 37(6):939–952. [PubMed: 12670423]
46. Hong K, Hinck L, Nishiyama M, Poo MM, Tessier-Lavigne M, Stein E. A ligand-gated association between cytoplasmic domains of UNC5 and DCC family receptors converts netrin-induced growth cone attraction to repulsion. *Cell*. 1999; 97(7):927–941. [PubMed: 10399920]
47. Westphal D, Sytnyk V, Schachner M, Leshchynska I. Clustering of the neural cell adhesion molecule (NCAM) at the neuronal cell surface induces caspase-8- and -3-dependent changes of the spectrin meshwork required for NCAM-mediated neurite outgrowth. *J Biol Chem*. 2010; 285(53):42046–42057. [PubMed: 20961848]
48. Ohsawa S, Hamada S, Kuida K, Yoshida H, Igaki T, Miura M. Maturation of the olfactory sensory neurons by Apaf-1/caspase-9-mediated caspase activity. *Proc Natl Acad Sci U S A*. 2010; 107(30):13366–71. [PubMed: 20624980]
49. Wang KH, Brose K, Arnott D, Kidd T, Goodman CS, Henzel W, Tessier-Lavigne M. Biochemical purification of a mammalian slit protein as a positive regulator of sensory axon elongation and branching. *Cell*. 1999; 96(6):771–784. [PubMed: 10102266]
50. Campbell DS, Stringham SA, Timm A, Xiao T, Law MY, Baier H, Nonet ML, Chien CB. Slit1a inhibits retinal ganglion cell arborization and synaptogenesis via Robo2-dependent and -independent pathways. *Neuron*. 2007; 55(2):231–245. [PubMed: 17640525]
51. Kidd T, Bland KS, Goodman CS. Slit is the midline repellent for the robo receptor in *Drosophila*. *Cell*. 1999; 96(6):785–794. [PubMed: 10102267]
52. Campbell DS, Okamoto H. Local caspase activation interacts with Slit-Robo signaling to restrict axonal arborization. *J Cell Biol*. 2013; 203(4):657–672. [PubMed: 24385488]
53. Gu Z, Serradj N, Ueno M, Liang M, Li J, Bacceti ML, Martin JH, Yoshida Y. Skilled Movements Require Non-apoptotic Bax/Bak Pathway-Mediated Corticospinal Circuit Reorganization. *Neuron*. 2017; 94(3):626–641. [PubMed: 28472660]
54. Sretavan DW, Shatz CJ, Stryker MP. Modification of retinal ganglion cell axon morphology by prenatal infusion of tetrodotoxin. *Nature*. 1988; 336(6198):468–471. [PubMed: 2461517]
55. Shirvan A, Shina R, Ziv I, Melamed E, Barzilai A. Induction of neuronal apoptosis by Semaphorin3A-derived peptide. *Mol Brain Res*. 2000; 83(1):81–93. [PubMed: 11072098]
56. Wehner AB, Abdesslem H, Dickendeshier TL, Imai F, Yoshida Y, Giger RJ, Pierchala BA. Semaphorin 3A is a retrograde cell death signal in developing sympathetic neurons. *Development*. 2016; 143(9):1560–1570. [PubMed: 27143756]
57. Matsunaga E, Tauszig-Delamasure S, Monnier PP, Mueller BK, Strittmatter SM, Mehlen P, Chedotal A. RGM and its receptor neogenin regulate neuronal survival. *Nat Cell Biol*. 2004; 6(8):749–755. [PubMed: 15258591]
58. Depaape V, Suarez-Gonzalez N, Dufour A, Passante L, Gorski JA, Jones KR, Ledent C, Vanderhaeghen P. Ephrin signalling controls brain size by regulating apoptosis of neural progenitors. *Nature*. 2005; 435(7046):1244–1250. [PubMed: 15902206]
59. Rooney IA, Butrovich KD, Glass AA, Borboroglu S, Benedict CA, Whitbeck JC, Cohen GH, Eisenberg RJ, Ware CF. The lymphotoxin-beta receptor is necessary and sufficient for LIGHT-mediated apoptosis of tumor cells. *J Biol Chem*. 2000; 275(19):14307–14315. [PubMed: 10799510]
60. Otsmane B, Moumen A, Aebischer J, Coque E, Sar C, Sunyach C, Salsac C, Valmier J, Salinas S, Bowerman M, Raoul C. Somatic and axonal LIGHT signaling elicit degenerative and regenerative responses in motoneurons, respectively. *EMBO Rep*. 2014; 15(5):540–547. [PubMed: 24668263]
61. Baas PW, White LA, Heidemann SR. Microtubule polarity reversal accompanies regrowth of amputated neurites. *Proc Natl Acad Sci*. 1987; 84(15):5272–5276. [PubMed: 3299383]
62. Monnier PP, D'Onofrio PM, Magharious M, Hollander AC, Tassew N, Szydlowska K, Tymianski M, Koeberle PD. Involvement of caspase-6 and caspase-8 in neuronal apoptosis and the regenerative failure of injured retinal ganglion cells. *J Neurosci*. 2011; 31(29):10494–10505. [PubMed: 21775595]

63. Koch JC, Tonges L, Barski E, Michel U, Bahr M, Lingor P. ROCK2 is a major regulator of axonal degeneration, neuronal death and axonal regeneration in the CNS. *Cell Death Dis.* 2014; 5:e1225. [PubMed: 24832597]
64. Hu J, Zhang G, Rodemer W, Jin LQ, Shifman M, Selzer ME. The role of RhoA in retrograde neuronal death and axon regeneration after spinal cord injury. *Neurobiol Dis.* 2017; 98:25–35. [PubMed: 27888137]
65. Chen L, Nye DM, Stone MC, Weiner AT, Gheres KW, Xiong X, Collins CA, Rolls MM. Mitochondria and Caspases Tune Nmnat-Mediated Stabilization to Promote Axon Regeneration. *PLoS Genet.* 2016; 12(12):e1006503. [PubMed: 27923046]
66. Verma P, Chierzi S, Codd AM, Campbell DS, Meyer RL, Holt CE, Fawcett JW. Axonal protein synthesis and degradation are necessary for efficient growth cone regeneration. *J Neurosci.* 2005; 25(2):331–342. [PubMed: 15647476]
67. Pinan-Lucarre B, Gabel CV, Reina CP, Hulme SE, Shevkoplyas SS, Slone RD, Xue J, Qiao Y, Weisberg S, Roodhouse K, Sun L, Whitesides GM, Samuel A, Driscoll M. The core apoptotic executioner proteins CED-3 and CED-4 promote initiation of neuronal regeneration in *Caenorhabditis elegans*. *PLoS Biol.* 2012; 10(5):e1001331. [PubMed: 22629231]
68. Lee H, Park S, Kang YS, Park S. EphA receptors form a complex with caspase-8 to induce apoptotic cell death. *Molecules and cells.* 2015; 38(4):349–355. [PubMed: 25855521]
69. Forcet C, Ye X, Granger L, Corset V, Shin H, Bredesen DE, Mehlen P. The dependence receptor DCC (deleted in colorectal cancer) defines an alternative mechanism for caspase activation. *Proc Natl Acad Sci.* 2001; 98(6):3416–3421. [PubMed: 11248093]
70. Hofmann K, Tschopp J. The death domain motif found in Fas (Apo-1) and TNF receptor is present in proteins involved in apoptosis and axonal guidance. *FEBS Lett.* 1995; 371(3):321–323. [PubMed: 7556620]
71. Artola A, Brocher S, Singer W. Different voltage-dependent thresholds for inducing long-term depression and long-term potentiation in slices of rat visual cortex. *Nature.* 1990; 347(6288):69–72. [PubMed: 1975639]
72. Collingridge GL, Peineau S, Howland JG, Wang YT. Long-term depression in the CNS. *Nat Rev Neurosci.* 2010; 11(7):459–473. [PubMed: 20559335]
73. Esteban JA, Shi SH, Wilson C, Nuriya M, Hugarir RL, Malinow R. PKA phosphorylation of AMPA receptor subunits controls synaptic trafficking underlying plasticity. *Nat Neurosci.* 2003; 6(2):136–143. [PubMed: 12536214]
74. Beattie EC, Carroll RC, Yu X, Morishita W, Yasuda H, von Zastrow M, Malenka RC. Regulation of AMPA receptor endocytosis by a signaling mechanism shared with LTD. *Nat Neurosci.* 2000; 3(12):1291–1300. [PubMed: 11100150]
75. Zhou Q, Homma KJ, Poo MM. Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron.* 2004; 44(5):749–757. [PubMed: 15572107]
76. Nagerl UV, Eberhorn N, Cambridge SB, Bonhoeffer T. Bidirectional activity-dependent morphological plasticity in hippocampal neurons. *Neuron.* 2004; 44(5):759–767. [PubMed: 15572108]
77. Dash PK, Blum S, Moore AN. Caspase activity plays an essential role in long-term memory. *Neuroreport.* 2000; 11(12):2811–2816. [PubMed: 10976968]
78. Stepanichev MY, Kudryashova IV, Yakovlev AA, Onufriev MV, Khaspekoy LG, Lyzhin AA, Lazareva NA, Gulyaeva NV. Central administration of a caspase inhibitor impairs shuttle-box performance in rats. *Neuroscience.* 2005; 136(2):579–591. [PubMed: 16198488]
79. Huesmann GR, Clayton DF. Dynamic role of postsynaptic caspase-3 and BIRC4 in zebra finch song-response habituation. *Neuron.* 2006; 52(6):1061–1072. [PubMed: 17178408]
80. Jiao S, Li Z. Nonapoptotic Function of BAD and BAX in Long-Term Depression of Synaptic Transmission. *Neuron.* 2011; 70(4):758–772. [PubMed: 21609830]
81. Li Z, Jo J, Jia JM, Lo SC, Whitcomb DJ, Jiao S, Cho K, Sheng M. Caspase-3 activation via mitochondria is required for long-term depression and AMPA receptor internalization. *Cell.* 2010; 141(5):859–871. [PubMed: 20510932]
82. Miller-Fleming TW, Petersen SC, Manning L, Matthewman C, Gornet M, Beers A, Hori S, Mitani S, Bianchi L, Richmond J, Miller DM. The DEG/ENaC cation channel protein UNC-8 drives

- activity-dependent synapse removal in remodeling GABAergic neurons. *Elife*. 2016; 5:e14599. [PubMed: 27403890]
83. Gulyaeva NV, Kudryashov IE, Kudryashova IV. Caspase activity is essential for long-term potentiation. *J Neurosci Res*. 2003; 73(6):853–864. [PubMed: 12949912]
 84. Bravarenko NI, Onufriev MV, Stepanichev MY, Ierusalimsky VN, Balaban PM, Gulyaeva NV. Caspase-like activity is essential for long-term synaptic plasticity in the terrestrial snail *Helix*. *Eur J Neurosci*. 2006; 23(1):129–140. [PubMed: 16420423]
 85. Lu C, Wang Y, Furukawa K, Fu W, Ouyang X, Mattson MP. Evidence that caspase-1 is a negative regulator of AMPA receptor-mediated long-term potentiation at hippocampal synapses. *J Neurochem*. 2006; 97(4):1104–1110. [PubMed: 16573645]
 86. Jo J, Whitcomb DJ, Olsen KM, Kerrigan TL, Lo SC, Bru-Mercier G, Dickinson B, Scullion S, Sheng M, Collingridge G, Cho K. Abeta(1–42) inhibition of LTP is mediated by a signaling pathway involving caspase-3, Akt1 and GSK-3beta. *Nat Neurosci*. 2011; 14(5):545–547. [PubMed: 21441921]
 87. Zhang H, Yang X, Qin X, Niu Q. Caspase-3 is Involved in Aluminum-Induced Impairment of Long-Term Potentiation in Rats Through the Akt/GSK-3beta Pathway. *Neurotoxicity research*. 2016; 29(4):484–494. [PubMed: 26787483]
 88. Olsen KM, Sheng M. NMDA receptors and BAX are essential for Abeta impairment of LTP. *Sci Rep*. 2012; 2:225. [PubMed: 22355739]
 89. Wang HG, Pathan N, Ethell IM, Krajewski S, Yamaguchi Y, Shibasaki F, McKeon F, Bobo T, Franke TF, Reed JC. Ca²⁺-induced apoptosis through calcineurin dephosphorylation of BAD. *Science*. 1999; 284(5412):339–343. [PubMed: 10195903]
 90. Affaticati P, Mignen O, Jambou F, Potier MC, Klingel-Schmitt I, Degrouard J, Peineau S, Gouadon E, Collingridge GL, Liblau R, Capiod T, Cohen-Kaminsky S. Sustained calcium signalling and caspase-3 activation involve NMDA receptors in thymocytes in contact with dendritic cells. *Cell Death Differ*. 2011; 18(1):99–108. [PubMed: 20577261]
 91. Han MH, Jiao S, Jia JM, Chen Y, Chen CY, Gucek M, Markey SP, Li Z. The novel caspase-3 substrate Gap43 is involved in AMPA receptor endocytosis and long-term depression. *Molecular & cellular proteomics: MCP*. 2013; 12(12):3719–3731. [PubMed: 24023391]
 92. Mukerjee N, McGinnis KM, Park YH, Gnegy ME, Wang KK. Caspase-mediated proteolytic activation of calcineurin in thapsigargin-mediated apoptosis in SH-SY5Y neuroblastoma cells. *Arch Biochem Biophys*. 2000; 379(2):337–343. [PubMed: 10898953]
 93. Chan SL, Griffin WS, Mattson MP. Evidence for caspase-mediated cleavage of AMPA receptor subunits in neuronal apoptosis and Alzheimer's disease. *J Neurosci Res*. 1999; 57(3):315–323. [PubMed: 10412022]
 94. Lu C, Fu W, Salvesen GS, Mattson MP. Direct cleavage of AMPA receptor subunit GluR1 and suppression of AMPA currents by caspase-3: implications for synaptic plasticity and excitotoxic neuronal death. *Neuromol Med*. 2002; 1(1):69–79.
 95. Widmann C, Gibson S, Johnson GL. Caspase-dependent cleavage of signaling proteins during apoptosis. A turn-off mechanism for anti-apoptotic signals. *J Biol Chem*. 1998; 273(12):7141–7147. [PubMed: 9507028]
 96. Peineau S, Taghibiglou C, Bradley C, Wong TP, Liu L, Lu J, Lo E, Wu D, Saule E, Bouchet T, Matthews P, Isaac JT, Bortolotto ZA, Wang YT, Collingridge GL. LTP inhibits LTD in the hippocampus via regulation of GSK3beta. *Neuron*. 2007; 53(5):703–717. [PubMed: 17329210]
 97. Guo J, Ji Y, Ding Y, Jiang W, Sun Y, Lu B, Nagappan G. BDNF pro-peptide regulates dendritic spines via caspase-3. *Cell Death Dis*. 2016; 7:e2264. [PubMed: 27310873]
 98. Park H, Poo MM. Neurotrophin regulation of neural circuit development and function. *Nat Rev Neurosci*. 2013; 14(1):7–23. [PubMed: 23254191]
 99. Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A, Zaitsev E, Gold B, Goldman D, Dean M, Lu B, Weinberger DR. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell*. 2003; 112(2):257–269. [PubMed: 12553913]

100. Pang PT, Teng HK, Zaitsev E, Woo NT, Sakata K, Zhen S, Teng KK, Yung WH, Hempstead BL, Lu B. Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science*. 2004; 306(5695):487–491. [PubMed: 15486301]
101. Sakuragi S, Tominaga-Yoshino K, Ogura A. Involvement of TrkB- and p75(NTR)-signaling pathways in two contrasting forms of long-lasting synaptic plasticity. *Sci Rep*. 2013; 3:3185. [PubMed: 24212565]
102. Woo NH, Teng HK, Siao CJ, Chiaruttini C, Pang PT, Milner TA, Hempstead BL, Lu B. Activation of p75NTR by proBDNF facilitates hippocampal long-term depression. *Nat Neurosci*. 2005; 8(8):1069–1077. [PubMed: 16025106]
103. Yang J, Harte-Hargrove LC, Siao CJ, Marinic T, Clarke R, Ma Q, Jing D, Lafrancois JJ, Bath KG, Mark W, Ballon D, Lee FS, Scharfman HE, Hempstead BL. proBDNF negatively regulates neuronal remodeling, synaptic transmission, and synaptic plasticity in hippocampus. *Cell Rep*. 2014; 7(3):796–806. [PubMed: 24746813]
104. Silke J, Meier P. Inhibitor of apoptosis (IAP) proteins-modulators of cell death and inflammation. *Cold Spring Harb Perspect Biol*. 2013; 5(2):a008730. [PubMed: 23378585]
105. Wilson R, Goyal L, Ditzel M, Zachariou A, Baker DA, Agapite J, Steller H, Meier P. The DIAP1 RING finger mediates ubiquitination of Dronc and is indispensable for regulating apoptosis. *Nat Cell Biol*. 2002; 4(6):445–450. [PubMed: 12021771]
106. Chai J, Yan N, Huh JR, Wu JW, Li W, Hay BA, Shi Y. Molecular mechanism of Reaper-Grim-Hid-mediated suppression of DIAP1-dependent Dronc ubiquitination. *Nat Struct Biol*. 2003; 10(11):892–898. [PubMed: 14517550]
107. Shapiro PJ, Hsu HH, Jung H, Robbins ES, Ryoo HD. Regulation of the Drosophila apoptosome through feedback inhibition. *Nat Cell Biol*. 2008; 10(12):1440–1446. [PubMed: 19011620]
108. Yan N, Wu JW, Chai J, Li W, Shi Y. Molecular mechanisms of DrICE inhibition by DIAP1, removal of inhibition by Reaper, Hid and Grim. *Nat Struct Mol Biol*. 2004; 11(5):420–428. [PubMed: 15107838]
109. Ditzel M, Broemer M, Tenev T, Bolduc C, Lee TV, Rigbolt KT, Elliott R, Zvelebil M, Blagoev B, Bergmann A, Meier P. Inactivation of effector caspases through nondegradative polyubiquitylation. *Mol Cell*. 2008; 32(4):540–553. [PubMed: 19026784]
110. Li X, Wang J, Shi Y. Structural mechanisms of DIAP1 auto-inhibition and DIAP1-mediated inhibition of drICE. *Nat Commun*. 2011; 2:408. [PubMed: 21811237]
111. Ribeiro PS, Kuranaga E, Tenev T, Leulier F, Miura M, Meier P. DIAP2 functions as a mechanism-based regulator of drICE that contributes to the caspase activity threshold in living cells. *J Cell Biol*. 2007; 179(7):1467–1480. [PubMed: 18166655]
112. Leulier F, Ribeiro PS, Palmer E, Tenev T, Takahashi K, Robertson D, Zachariou A, Pichaud F, Ueda R, Meier P. Systematic in vivo RNAi analysis of putative components of the Drosophila cell death machinery. *Cell Death Differ*. 2006; 13(10):1663–1674. [PubMed: 16485033]
113. Muro I, Hay BA, Clem RJ. The Drosophila DIAP1 Protein Is Required to Prevent Accumulation of a Continuously Generated, Processed Form of the Apical Caspase DRONC. *Journal of Biological Chemistry*. 2002; 277(51):49644–49650. [PubMed: 12397080]
114. Eckelman BP, Salvesen GS, Scott FL. Human inhibitor of apoptosis proteins: why XIAP is the black sheep of the family. *EMBO Rep*. 2006; 7(10):988–994. [PubMed: 17016456]
115. Deveraux QL, Takahashi R, Salvesen GS, Reed JC. X-linked IAP is a direct inhibitor of cell-death proteases. *Nature*. 1997; 388(6639):300–304. [PubMed: 9230442]
116. Deveraux QL, Leo E, Stennicke HR, Welsh K, Salvesen GS, Reed JC. Cleavage of human inhibitor of apoptosis protein XIAP results in fragments with distinct specificities for caspases. *EMBO J*. 1999; 18(19):5242–5251. [PubMed: 10508158]
117. Shiozaki EN, Chai J, Rigotti DJ, Riedl SJ, Li P, Srinivasula SM, Alnemri ES, Fairman R, Shi Y. Mechanism of XIAP-mediated inhibition of caspase-9. *Mol Cell*. 2003; 11(2):519–527. [PubMed: 12620238]
118. Scott FL, Denault JB, Riedl SJ, Shin H, Renatus M, Salvesen GS. XIAP inhibits caspase-3 and -7 using two binding sites: evolutionarily conserved mechanism of IAPs. *EMBO J*. 2005; 24(3):645–655. [PubMed: 15650747]

119. Potts PR, Singh S, Knezek M, Thompson CB, Deshmukh M. Critical function of endogenous XIAP in regulating caspase activation during sympathetic neuronal apoptosis. *J Cell Biol.* 2003; 163:789–799. [PubMed: 14623868]
120. Ertürk A, Wang Y, Sheng M. Local Pruning of Dendrites and Spines by Caspase-3-Dependent and Proteasome-Limited Mechanisms. *J Neurosci.* 2014; 34(5):1672–1688. [PubMed: 24478350]
121. Wright KM, Linhoff MW, Potts PR, Deshmukh M. Decreased apoptosome activity with neuronal differentiation sets the threshold for strict IAP regulation of apoptosis. *J Cell Biol.* 2004; 167(2): 303–313. [PubMed: 15504912]
122. Martinez-Marmol R, Barneda-Zahonero B, Soto D, Andres RM, Coccia E, Gasull X, Planells-Ferrer L, Moubarak RS, Soriano E, Comella JX. FAIM-L regulation of XIAP degradation modulates Synaptic Long-Term Depression and Axon Degeneration. *Sci Rep.* 2016; 6:35775. [PubMed: 27767058]
123. Gibon J, Unsain N, Gamache K, Thomas RA, De Leon A, Johnstone A, Nader K, Seguela P, Barker PA. The X-linked inhibitor of apoptosis regulates long-term depression and learning rate. *FASEB J.* 2016; 30(9):3083–3090. [PubMed: 27189977]
124. Saelens X, Festjens N, Vande Walle L, van Gurp M, van Loo G, Vandenabeele P. Toxic proteins released from mitochondria in cell death. *Oncogene.* 2004; 23(16):2861–74. [PubMed: 15077149]
125. Gottfried Y, Rotem A, Lotan R, Steller H, Larisch S. The mitochondrial ARTS protein promotes apoptosis through targeting XIAP. *EMBO J.* 2004; 23(7):1627–1635. [PubMed: 15029247]
126. Garrison JB, Correa RG, Gerlic M, Yip KW, Krieg A, Tumble CM, Shi R, Welsh K, Duggineni S, Huang Z, Ren K, Du C, Reed JC. ARTS and Siah Collaborate in a Pathway for XIAP Degradation. *Mol Cell.* 2011; 41(1):107–116. [PubMed: 21185211]
127. Bhogal B, Plaza-Jennings A, Gavis ER. Nanos-mediated repression of hid protects larval sensory neurons after a global switch in sensitivity to apoptotic signals. *Development.* 2016; 143(12): 2147–2159. [PubMed: 27256879]
128. Kuranaga A, Kanuka H, Tonoki A, Takemoto K, Tomioka T, Kobayashi M, Hayashi S, Miura M. Drosophila IKK-related kinase regulates nonapoptotic function of caspases via degradation of IAPs. *Cell.* 2006; 126(3):583–596. [PubMed: 16887178]
129. Nakhaei P, Sun Q, Solis M, Mesplede T, Bonneil E, Paz S, Lin R, Hiscott J. IkappaB kinase epsilon-dependent phosphorylation and degradation of X-linked inhibitor of apoptosis sensitizes cells to virus-induced apoptosis. *J Virol.* 2012; 86(2):726–737. [PubMed: 22072751]
130. Tsang AH, Lee YI, Ko HS, Savitt JM, Pletnikova O, Troncoso JC, Dawson VL, Dawson TM, Chung KK. S-nitrosylation of XIAP compromises neuronal survival in Parkinson's disease. *Proc Natl Acad Sci.* 2009; 106(12):4900–4905. [PubMed: 19273858]
131. Nakamura T, Wang L, Wong CC, Scott FL, Eckelman BP, Han X, Tzitzilonis C, Meng F, Gu Z, Holland EA, Clemente AT, Okamoto S, Salvesen GS, Riek R, Yates JR 3rd, Lipton SA. Transnitrosylation of XIAP regulates caspase-dependent neuronal cell death. *Mol Cell.* 2010; 39(2):184–195. [PubMed: 20670888]
132. D'Amelio M, Cavallucci V, Middei S, Marchetti C, Pacioni S, Ferri A, Diamantini A, De Zio D, Carrara P, Battistini L, Moreno S, Bacci A, Ammassari-Teule M, Marie H, Cecconi F. Caspase-3 triggers early synaptic dysfunction in a mouse model of Alzheimer's disease. *Nat Neurosci.* 2011; 14(1):69–76. [PubMed: 21151119]
133. Yuan J, Shaham S, Ledoux S, Ellis HM, Horvitz HR. The *C. elegans* cell death gene *ced-3* encodes a protein similar to mammalian interleukin-1 β -converting enzyme. *Cell.* 1993; 75(4): 641–652. [PubMed: 8242740]

Figure 1. Non apoptotic role of caspases in axon and dendrite pruning

In absence of neurotrophic factors (i.e. NGF), activation of the JNK pathway by DLK promotes the transcriptional upregulation of Puma, a pro-apoptotic BH3-only members of the Bcl-2 family. This ultimately results in the activation of Bax, mitochondrial depolarization and caspases, leading to local degeneration of the axon. Additionally, the death receptor DR6 might play a role in local activation of the caspase cascade.

Figure 2. Non apoptotic role of caspases in neurite outgrowth, axon guidance and branching
Caspases are activated in response to various receptors implicated in neurite outgrowth (NCAM), guidance (DCC in response to its ligand Netrin1- with its chemoattractive ability depicted as +) and branching (lymphotoxin beta receptor (LT-βR) when bound to its ligand LIGHT and Robo2 stimulated by its ligand Slit1a).

Figure 3. Non apoptotic role of caspases in long-term depression (LTD)

Caspases participate in synaptic plasticity of glutamatergic synapses by promoting the downregulation of AMPA receptor expression from the synaptic surface. In response to NMDA receptor-induced calcium entry, calcineurin dephosphorylates Bad, thereby stimulating Bax-induced mitochondrial depolarization and caspase-3 activation. Possible roles for caspase-3 in LTD include direct role in AMPAR endocytosis either by activation of GSK3 β via proteolysis of Akt1, or by GAP43 cleavage. Alternatively, caspase-3 may directly process and inactivate AMPAR.

Figure 4. XIAP restricts caspase activity during non apoptotic contexts

Caspases-3, -7 and -9 are kept inactivated by XIAP in neurons therefore restricting caspase activation where and when it is needed. Proposed mechanisms for releasing caspases from XIAP control include proteasomal degradation of XIAP stimulated by IKK ϵ and XIAP inactivation by S-nitrosylation.