Published in final edited form as:

AIDS. 2017 August 24; 31(13): 1839–1845. doi:10.1097/QAD.0000000000001540.

Impact of alemtuzumab on HIV persistence in an HIV-infected individual on antiretroviral therapy with Sezary syndrome

Thomas A. Rasmussen^{1,2}, James McMahon³, J. Judy Chang¹, Jori Symons¹, Michael Roche¹, Ashanti Dantanarayana¹, Afam Okoye⁴, Bonnie Hiener⁵, Sarah Palmer⁵, Wen Shi Lee¹, Stephen Kent¹, Carrie Van Der Weyden⁶, H. Miles Prince⁷, Paul U. Cameron^{1,3}, and Sharon R. Lewin^{1,3}

¹The Peter Doherty Institute for Infection and Immunity, The University of Melbourne and Royal Melbourne Hospital, Melbourne, Australia

²Department of Infectious Diseases, Aarhus University Hospital, Aarhus, Denmark

³Department of Infectious Diseases, Alfred Hospital and Monash University, Melbourne, Australia

⁴Vaccine and Gene Therapy Institute, Oregon Health & Science University, Beaverton, Oregon National Primate Research Center, Oregon, USA

⁵The Westmead Institute, University of Sydney, Westmead, NSW, 2145, Australia

⁶Division of Cancer Medicine, Peter MacCallum Cancer Centre, Melbourne, Australia

⁷Sir Peter MacCallum Department of Oncology, University of Melbourne and Epworth Healthcare, East Melbourne, Australia

Introduction

Despite the great advances in antiretroviral therapy (ART), treatment is life-long and there is no cure. HIV can persist on ART as a latent infection in multiple long-lived T-cell subsets including central memory and naïve T-cells^[1]. In addition, latently infected CD4+ T-cells can undergo homeostatic proliferation^[2] and clonal expansion^[3-6].

Sezary syndrome is a leukemic form of cutaneous T-cell lymphoma (CTCL) and is considered a malignancy of CD4+ central memory T-cells (Tcm)^[7]. Treatment options for Sezary syndrome have recently included low-dose alemtuzumab^[8], a humanized anti-CD52 monoclonal antibody licensed for B-cell chronic lymphocytic leukaemia and relapsing-remitting multiple sclerosis (RRMS)^[9, 10]. CD52 is expressed on all T-cell subsets including long-lived Tcm and naïve T-cells but also on B-cells, macrophages, natural killer (NK) cells and dendritic cells^[11, 12]. Alemtuzumab depletes CD52 positive circulating cells but has minimal effect on non-circulating skin resident effector memory T-cells (Tem), required to maintain mucosal immunity and protection from opportunistic infections. Although

Correspondence: Thomas Aagaard Rasmussen, Department of Infectious Diseases, Aarhus University Hospital, Palle Juul-Jensens Boulevard 99, 8200 Aarhus N, Denmark, thomasm@rm.dk.

Conflicts of interest

For the remaining authors none were declared.

associated with some risk, complete depletion of Tcm and naïve T-cells followed by immune reconstitution on ART, may potentially perturb or even eliminate HIV persistence on ART.

Here, we describe the effects of alemtuzumab treatment in an HIV infected individual on ART with Sezary syndrome. We aimed to characterise 1) whether malignant CD4+ T-cells in Sezary syndrome were a consequence of clonal expansion of HIV-infected cells; 2) the effect of alemtuzumab on malignant and non-malignant T-cells including memory subsets; and 3) how alemtuzumab impacted the frequency of latently infected CD4+ T-cells.

Methods

Sample collection

We collected blood and isolated peripheral blood mononuclear cells (PBMCs) and plasma at regular intervals prior to and following the diagnosis of Sezary syndrome and prior to and during alemtuzumab treatment (Fig. 1). The protocol was approved by the Ethics Committee of the Alfred Hospital and the patient provided written informed consent for participation and for use of stored PBMC collected from a previous study.

Flow cytometry for malignant and non-malignant CD4+ T cells

We used flow cytometry and fluorescence activated cell sorting (FACS) to quantify and isolate malignant CD4+ T cells, defined by lack of expression of CD7 and CD26^[13] and expression of T-cell receptor (TCR)-VBeta2. In some experiments, we used a less stringent definition of non-malignant cells as CD3+CD4+TCR-Vbeta2- (Fig.S1). Memory T cell subsets were defined as naïve (CD45RA+CCR7+), terminally differentiated (TD; CD45RA+CCR7-), Tcm (CD45RA-CCR7+), Tem (CD45RA-CCR7-CD27-) and transitional memory (Ttm; CD45RA-CCR7-CD27+) T-cells.

HIV-DNA quantification

Cryopreserved PBMCs were sorted into total CD4+ T-cells, malignant (CD3+CD4+TCR-Vbeta2+CD7-CD26-) or non-malignant (CD3+CD4+TCR-Vbeta2-) CD4+ T-cells using FACS. Cells were then lysed and cell-associated HIV-DNA was measured by quantitative PCR (qPCR) using primers and probes as previously described^[14].

HIV-DNA full-length sequencing

To address whether there was clonal expansion of HIV-infected cells, we employed a full-length HIV sequencing method based on next-generation sequencing techniques (Hiener et al. CROI 2017). Proviral sequences were diluted to a single genome and a near full-length 9 Kb region of HIV-DNA amplified using a nested PCR protocol. Each full-length HIV genome was fragmented and a sequence index or tag was added to the representative fragments of each genome for unique identification in the final data analysis. After sequencing, the individual proviruses were *de novo* assembled using a specifically designed workflow in CLC Genomics. Proviruses were characterized as defective (containing INDELs, stop codons or APOBEC3G hypermutation) or intact (full-length; lacking defects).

Gp120-specific antibodies and antibody-dependent natural killer (NK) cell activation assay

As previously described, we used an ELISA to determine the concentrations of gp120-specific antibodies in plasma^[15] and performed a plate-bound antibody-dependent NK cell activation assay to measure intracellular NK cell IFN γ and CD107a expression^[16].

Results

Clinical course

A 62-year old male with chronic hepatitis B virus (HBV) and HIV co-infection on ART presented with generalised rash and was diagnosed with Sezary syndrome. He also had co-existent primary cutaneous anaplastic large cell lymphoma (pcALCL), another separate cutaneous T cell lymphoma well recognized to occur in patients with co-existent mycosis fungoides/Sezary syndrome^[17]. Both HBV and HIV infections were diagnosed 4 years prior to presentation with Sezary syndrome at age 58. At HIV diagnosis, HIV RNA was 228,100 copies/mL and nadir CD4+ T-cell count was 110 cells/uL. Combination ART with tenofovir disoproxil fumarate, emtricitabine and dolutegravir was commenced immediately after HIV diagnosis. Suppression of plasma HIV RNA to <50 copies/mL was achieved after one month on ART, but over the following 4 years frequent low-level plasma HIV RNA was detected, in the range of 50-387 copies/mL, while CD4+ T-cells reached >500 cells/uL after approximately 3 years on ART (Fig. 1a).

Three months prior to diagnosis with Sezary syndrome, the patient initially observed an eczematous-like rash over his trunk, which rapidly progressed to generalised erythroderma with pruritus, scaling and acral keratoderma (Fig. 1a). Skin biopsies and peripheral blood flow cytometry were consistent with a diagnosis of Sezary syndrome. He also had multi-focal tumours over his neck, axilla and trunk and biopsies of these demonstrated co-existent CD30+ pcALCL. Five months after diagnosis of Sezary syndrome, his CD4 count had risen to 2599 cells/uL from 515 cells/uL three months prior to diagnosis. The rise was assumed to be an increase in the malignant CD4+ clone (Fig. 1a).

Treatment of the Sezary Syndrome was initiated 10 months after diagnosis with low-dose alemtuzumab combined with extra-corporeal photophoresis and tumour-directed radiotherapy. The patient received 10 mg alemtuzumab subcutaneously three times per week for four weeks. After an initial excellent symptomatic treatment response with marked reduction of pruritus and improvement in skin integrity and erythroderma, he subsequently experienced worsening liver function followed by multi-organ failure with progressive lymphadenopathy and died 6 weeks after commencing alemtuzumab.

HIV-DNA was detected in non-malignant but not malignant CD4+ T cells

Given recent reports of clonal expansion of latently infected CD4+ T-cells in individuals on ART^[3, 6, 18], we first asked whether the presentation of Sezary syndrome was associated with malignant clonal expansion of a latently infected CD4+ T-cell. We quantified HIV-DNA in malignant (CD7-CD26-TCR-Vbeta2+) and non-malignant (CD7+CD26+TCR-Vbeta2-) CD4+ T-cells collected 10 months after the diagnosis of Sezary syndrome but prior to alemtuzumab. HIV-DNA was detected in non-malignant CD4+ T-cells (1998 copies/

million cells) but not in the malignant CD4+ T-cells (<4 copies/million cells) (Fig. 1b). These findings were consistent with expansion of a CD4+ T-cell clone not infected with HIV. Malignant cells expressed both CXCR4 and CCR5 so we hypothesise that resistance to infection was secondary to an event downstream of co-receptor binding.

Because HIV-DNA in purified non-malignant CD4+ T-cells collected 10 months after the diagnosis of Sezary syndrome (pre-alemtuzumab) was 2.7 to 5.0-fold higher than in total CD4+ T-cells collected 20 and 8 months before diagnosis, respectively, we next asked whether there was tumour-antigen driven expansion of HIV-infected CD4+ T cells, similar to a recent case report of metastatic squamous cell carcinoma in an HIV-infected individual^[4]. We performed full-length sequencing of HIV-DNA in CD4+ T-cells collected before and after presentation with Sezary syndrome. We obtained and sequenced 25-26 single HIV genomic sequences and found 8-10 full-length HIV genomes from each time point; however, the majority of these genomes contained deletions prior to the HIV gag gene region and did not represent replication-competent virus. We identified only one clonal expansion and this sequence contained hypermutant genomes from all time points (Fig. 1c). Taken together, we demonstrated that the malignant CD4+ T-cells were not infected and there was no evidence of clonal expansion of HIV-DNA in the non-malignant cells.

Effect of alemtuzumab on T-cells, T-cell subsets and HIV persistence

Alemtuzumab led to a dramatic depletion of total T-cells within 18 days including a >1 log reduction in total CD3+ T-cells and >3 log reduction in CD4+ T cells (Fig. 2a). There was a greater decline in malignant compared to non-malignant CD4+ T-cells (Fig. 2b). We quantified memory CD4+ and CD8+ T-cells in cells enriched for malignant (CD7-) and non-malignant (CD7+) markers prior to and 4 and 11 days during alemtuzumab. We observed depletion of central memory but not effector memory CD4+ T-cells not expressing CCR7. This pattern was not seen for CD8+ memory T-cells or for CCR7+ cells (Fig. 2c and Fig. 2d). We did not quantify T-cell subsets 18 and 25 days post alemtuzumab due to very low T-cell numbers at these time points.

We next quantified HIV-DNA in total CD4+ T-cells prior to and following alemtuzumab. We observed a 57% reduction in HIV-DNA per million cells from pre-alemtuzumab (day 0) to day 25 on-alemtuzumab (Fig. 2e), but even when the total lymphocyte count was as low as 80 cells/uL on-alemtuzumab, we were still able to detect HIV-DNA. To exclude the possibility that the HIV-DNA was a contaminant, we sequenced the virus and found no alignment with any laboratory strains of HIV (data not shown). The rapid decay of malignant CD4+ T-cells after starting alemtuzumab allowed for quantification of HIV DNA in sorted non-malignant CD4+ T-cells at only one time point four days after starting alemtuzumab. We did not detect a decrease in HIV DNA in non-malignant CD4+ T cells at this early phase of alemtuzumab treatment (1998 and 2098 copies per million non-malignant CD4+ T cells at day 0 and day 4, respectively).

Finally, as CD52 is also expressed on NK cells, we investigated whether alemtuzumab affected HIV-specific antibody-mediated NK cell activation. We found no changes in anti-gp120 IgG concentrations during the four weeks of alemtuzumab treatment and also did not

see any change in the ability of the patient's plasma to induce NK cell expression of interferon gamma (IFN γ) or CD107a (Fig. S2).

Discussion

In an HIV infected individual on ART with Sezary syndrome, the malignant CD4 $^{+}$ T-cells were not infected with HIV and full-length sequencing of HIV-DNA demonstrated no clonal expansion in the non-malignant HIV infected CD4 $^{+}$ T-cells. Alemtuzumab caused extensive depletion of T-cells and decreased the frequency of latently infected CD4 $^{+}$ T-cells, but HIV-DNA remained detectable even after extensive CD4 $^{+}$ T-cell depletion.

Alemtuzumab is rarely used in HIV infected individuals and no previous study has analysed its effect on HIV persistence. In a case series of HIV infected patients receiving alemtuzumab as part of renal transplant conditioning, there were no opportunistic infections despite a slow recovery of CD4 $^{+}$ T-cells but effects on latent HIV were not investigated^[19]. Our data confirm the extensive depletion of multiple T-cell subsets also seen in other studies^[12, 20] and we also detected a decrease in the frequency of HIV-DNA in total CD4 $^{+}$ T-cells during alemtuzumab. The detection of HIV-DNA, even when the total CD4 $^{+}$ T-cell count was <1 cell/uL, was surprising and could potentially be explained by persistence of virus in other long-lived cells such as T memory stem cells. These have been shown to be preferentially infected with HIV in individuals on ART^[21], have an extremely long half-life and it is unknown if these cells are depleted by alemtuzumab.

Lymph node T follicular helper (Tfh) cells were recently shown to be an important source of persistent HIV in patients on suppressive ART^[22] but the direct effects of alemtuzumab in lymph node tissue including effects on Tfh cells could not be evaluated in our study. However, as CD52 is ubiquitously expressed on all mature T-cells^[11, 12], we assume CD52 is also expressed on Tfh cells as well as other long-lived CD4 $^{+}$ T-cells that serve as important reservoirs for HIV. Notably, in malignant transformation of Tfh cells, as seen in angioimmunoblastic T-cell lymphoma (AITL), high levels of CD52 expression in both lymph node aspirates and peripheral blood was demonstrated^[23-25]. Given that alemtuzumab-mediated depletion of CD52 positive cells occurs via antibody-dependent cell-mediated cytotoxicity (ADCC) and complement-dependent cytotoxicity^[11], which is dependent on the presence of immune effector cells such as NK cells and neutrophils^[26], tissue-resident cells like Tfh, may potentially avoid depletion in the absence of those effector functions. For example, in a study of mycosis fungoides, alemtuzumab was shown to potently deplete circulating long-lived central memory and naïve T-cells but had minimal effect on non-circulating skin resident effector memory T-cells (Tem)^[27]. Similarly, in cynomolgus monkeys who received alemtuzumab following cardiac transplant, there was a much more profound reduction of CD4 $^{+}$ T-cells in peripheral blood (99% reduction) than in lymph nodes (70% reduction)^[28].

We hypothesised that following immune reconstitution after alemtuzumab, the frequency of latently infected CD4 $^{+}$ T-cells could be substantially reduced if there was preferential recovery and expansion of uninfected cells. Unfortunately, the patient died 6 weeks post-alemtuzumab, which prevented analyses following T-cell reconstitution. We are currently

testing this hypothesis in a simian immunodeficiency virus (SIV)-infected rhesus macaque model.

In conclusion, alemtuzumab extensively depleted multiple T-cell subsets and also decreased the frequency of but did not eliminate CD4⁺ T-cells harbouring HIV-DNA in blood. Studying the effects of alemtuzumab in blood and lymphoid tissue as well as following immune recovery in HIV-infected individuals who require alemtuzumab for the treatment of malignancy or RRMS or in animal studies may provide further insights into novel HIV cure strategies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are very grateful to the patient described in this case report who agreed to increased visits and blood collections for this research protocol. We thank the clinical research team at Alfred Hospital, Melbourne, Australia for their help with coordinating study visits and collecting clinical samples. We also wish to thank Ajantha Rhodes (Doherty Institute, University of Melbourne, Australia) and Anni Winckelmann (Department of Infectious Diseases, Aarhus University Hospital, Aarhus Denmark). Ajantha Rhodes performed PCR analyses to quantify HIV DNA in total CD4⁺ T-cells and sorted subsets. Anni Winckelmann helped preparing Fig. 1d.

SRL, TAR, PUC, HMP, JM, JS, MR and AO conceived of and designed the study and its analyses; HMP, CW, JM, SRL and TAR were responsible for study visits and all clinical interaction; JJC and AD performed cell sorting and flow cytometry analyses; AR performed PCR analyses; BH and SP performed HIV DNA full-length sequences and related analyses; SK and WSL designed and performed analyses of antibody-mediated cellular toxicity; MR performed HIV DNA sequencing to rule out contamination; SRL, PUC, TAR, JM, MR, JJC and JS performed data analyses and interpretation; SRL, PUC, TAR, JJC, SP, BH and AO contributed to the graphical presentation of study results; SRL, TAR and CW drafted the manuscript; all authors reviewed, edited and approved the final manuscript.

The study was funded by the National Health and Medical Research Council (NHMRC) of Australia program grant (PI Lewin). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

SRL's institution has received funding from the National Health and Medical Research Council (NHMRC) of Australia, National Institutes for Health, American Foundation for AIDS Research; Merck, Viiv, Gilead and Tetralogic for investigator initiated research; Merck, Viiv and Gilead for educational activities.

References

1. Deeks SG, Lewin SR, Ross AL, Ananworanich J, Benkirane M, Cannon P, et al. International AIDS Society global scientific strategy: towards an HIV cure 2016. *Nat Med*. 2016
2. Chomont N, El-Far M, Ancuta P, Trautmann L, Procopio FA, Yassine-Diab B, et al. HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation. *NatMed*. 2009; 15(8):893–900.
3. Wagner TA, McLaughlin S, Garg K, Cheung CY, Larsen BB, Styrchak S, et al. HIV latency. Proliferation of cells with HIV integrated into cancer genes contributes to persistent infection. *Science*. 2014; 345(6196):570–573. [PubMed: 25011556]
4. Simonetti FR, Sobolewski MD, Fyne E, Shao W, Spindler J, Hattori J, et al. Clonally expanded CD4⁺ T cells can produce infectious HIV-1 in vivo. *Proc Natl Acad Sci U S A*. 2016; 113(7):1883–1888. [PubMed: 26858442]
5. Cohn LB, Silva IT, Oliveira TY, Rosales RA, Parrish EH, Learn GH, et al. HIV-1 integration landscape during latent and active infection. *Cell*. 2015; 160(3):420–432. [PubMed: 25635456]

6. Maldarelli F, Wu X, Su L, Simonetti FR, Shao W, Hill S, et al. HIV latency. Specific HIV integration sites are linked to clonal expansion and persistence of infected cells. *Science*. 2014; 345(6193):179–183. [PubMed: 24968937]
7. Campbell JJ, Clark RA, Watanabe R, Kupper TS. Sezary syndrome and mycosis fungoides arise from distinct T-cell subsets: a biologic rationale for their distinct clinical behaviors. *Blood*. 2010; 116(5):767–771. [PubMed: 20484084]
8. de Masson A, Guitera P, Brice P, Moulouguet I, Mouly F, Bouaziz JD, et al. Long-term efficacy and safety of alemtuzumab in advanced primary cutaneous T-cell lymphomas. *Br J Dermatol*. 2014; 170(3):720–724. [PubMed: 24438061]
9. Zinzani PL, Corradini P, Gallamini A, Grossi A, Lazzarino M, Marchetti M, et al. Overview of alemtuzumab therapy for the treatment of T-cell lymphomas. *Leukemia & lymphoma*. 2012; 53(5): 789–795. [PubMed: 21980956]
10. Riera R, Porfirio GJ, Torloni MR. Alemtuzumab for multiple sclerosis. *Cochrane Database Syst Rev*. 2016; 4:CD011203. [PubMed: 27082500]
11. Gribben JG, Hallek M. Rediscovering alemtuzumab: current and emerging therapeutic roles. *Br J Haematol*. 2009; 144(6):818–831. [PubMed: 19183194]
12. Bouvy AP, Klepper M, Betjes MG, Weimar W, Hesselink DA, Baan CC. Alemtuzumab as Antirejection Therapy: T Cell Repopulation and Cytokine Responsiveness. *Transplant Direct*. 2016; 2(6):e83. [PubMed: 27500273]
13. Boonk SE, Zoutman WH, Marie-Cardine A, van der Fits L, Out-Luiting JJ, Mitchell TJ, et al. Evaluation of Immunophenotypic and Molecular Biomarkers for Sezary Syndrome Using Standard Operating Procedures: A Multicenter Study of 59 Patients. *J Invest Dermatol*. 2016; 136(7):1364–1372. [PubMed: 26930587]
14. Elliott JH, Wightman F, Solomon A, Ghneim K, Ahlers J, Cameron MJ, et al. Activation of HIV Transcription with Short-Course Vorinostat in HIV-Infected Patients on Suppressive Antiretroviral Therapy. *PLoS pathogens*. 2014; 10(10):e1004473. [PubMed: 25393648]
15. Parsons MS, Madhavi V, Ana-Sosa-Batiz F, Center RJ, Wilson KM, Bunupuradah T, et al. Brief Report: Seminal Plasma Anti-HIV Antibodies Trigger Antibody-dependent Cellular Cytotoxicity: Implications for HIV Transmission. *J Acquir Immune Defic Syndr*. 2016; 71(1):17–23. [PubMed: 26761269]
16. Jegaskanda S, Job ER, Kramski M, Laurie K, Isitman G, de Rose R, et al. Cross-reactive influenza-specific antibody-dependent cellular cytotoxicity antibodies in the absence of neutralizing antibodies. *J Immunol*. 2013; 190(4):1837–1848. [PubMed: 23319732]
17. Kempf W, Pfaltz K, Vermeer MH, Cozzio A, Ortiz-Romero PL, Bagot M, et al. EORTC, ISCL, and USCLC consensus recommendations for the treatment of primary cutaneous CD30-positive lymphoproliferative disorders: lymphomatoid papulosis and primary cutaneous anaplastic large-cell lymphoma. *Blood*. 2011; 118(15):4024–4035. [PubMed: 21841159]
18. Wagner, T., McLaughlin, S., Garg, K., Huang, H., Styrchak, S., Mullins, J., et al. Proliferation of Cells With HIV Integrated Into Regulatory Genes Is a Mechanism of Persistence. *Conference of Retroviruses and Opportunistic Infections (CROI)*; Boston, MA. 2014.
19. Tan HP, Kaczorowski DJ, Basu A, Khan A, McCauley J, Marcos A, et al. Living-related donor renal transplantation in HIV+ recipients using alemtuzumab preconditioning and steroid-free tacrolimus monotherapy: a single center preliminary experience. *Transplantation*. 2004; 78(11): 1683–1688. [PubMed: 15591960]
20. Thomas K, Eisele J, Rodriguez-Leal FA, Hainke U, Ziemssen T. Acute effects of alemtuzumab infusion in patients with active relapsing-remitting MS. *Neurol Neuroimmunol Neuroinflamm*. 2016; 3(3):e228. [PubMed: 27213173]
21. Buzon MJ, Sun H, Li C, Shaw A, Seiss K, Ouyang Z, et al. HIV-1 persistence in CD4(+) T cells with stem cell-like properties. *Nat Med*. 2014; 20(2):139–142. [PubMed: 24412925]
22. Banga R, Procopio FA, Noto A, Pollakis G, Cavassini M, Ohmiti K, et al. PD-1 and follicular helper T cells are responsible for persistent HIV-1 transcription in treated aviremic individuals. *Nat Med*. 2016

23. Jiang L, Yuan CM, Hubacheck J, Janik JE, Wilson W, Morris JC, et al. Variable CD52 expression in mature T cell and NK cell malignancies: implications for alemtuzumab therapy. *Br J Haematol*. 2009; 145(2):173–179. [PubMed: 19236377]
24. Geissinger E, Bonzheim I, Roth S, Rosenwald A, Muller-Hermelink HK, Rudiger T. CD52 expression in peripheral T-cell lymphomas determined by combined immunophenotyping using tumor cell specific T-cell receptor antibodies. *Leukemia & lymphoma*. 2009; 50(6):1010–1016. [PubMed: 19479612]
25. Cortes JR, Palomero T. The curious origins of angioimmunoblastic T-cell lymphoma. *Curr Opin Hematol*. 2016; 23(4):434–443. [PubMed: 27177312]
26. Siders WM, Shields J, Garron C, Hu Y, Boutin P, Shankara S, et al. Involvement of neutrophils and natural killer cells in the anti-tumor activity of alemtuzumab in xenograft tumor models. *Leukemia & lymphoma*. 2010; 51(7):1293–1304. [PubMed: 20377308]
27. Clark RA, Watanabe R, Teague JE, Schlapbach C, Tawa MC, Adams N, et al. Skin effector memory T cells do not recirculate and provide immune protection in alemtuzumab-treated CTCL patients. *Sci Transl Med*. 2012; 4(117):117ra117.
28. Marco MR, Dons EM, van der Windt DJ, Bhama JK, Lu LT, Zahorchak AF, et al. Post-transplant repopulation of naive and memory T cells in blood and lymphoid tissue after alemtuzumab-mediated depletion in heart-transplanted cynomolgus monkeys. *Transpl Immunol*. 2013; 29(1-4): 88–98. [PubMed: 24120957]

Fig. 1. Clinical course and HIV-DNA in malignant and non-malignant cells

A. Chronological overview of key clinical events, plasma HIV RNA and CD4+ T cell counts since start of ART. B. HIV-DNA in sorted malignant (CD3+CD4+CD7-CD26-TCR-Vbeta2+, shown as closed circles) and non-malignant CD4+ T cells (CD3+CD4+TCR-Vbeta2-, shown as open circles) collected 2 weeks before starting alemtuzumab. The DNA PCR was done in three replicates, lower limit of detection was 1 copy per well, and a total of 220,000 malignant CD4+ T cells were analysed. C. Phylogenetic tree of full-length HIV-DNA sequences obtained from CD4+ T-cells isolated from blood collected prior to (blue and

green squares) and following (purple squares) the diagnosis of Sezary syndrome. The majority of sequences were defective and contained major deletions in HIV genes as illustrated by the coloured bars in the right hand side of the figure.

ART, antiretroviral therapy; TDF, tenofovir disoproxil fumarate; FTC, emtricitabine; DTG, dolutegravir; PHA, phytohaemagglutinin; GFP, green fluorescent protein.

Fig. 2. Depletion of T cells and HIV-DNA during alemtuzumab

A. Absolute counts of total lymphocytes and T-cell subsets in blood prior to and following alemtuzumab treatment. **B.** The change in frequency of CD7- (malignant) or CD7+ (non-malignant) CD4+ T cells following alemtuzumab. **C.** The distribution of memory T cell subsets, stratified by CD7 expression, shown for both CD4+ T cells and **D.** CD8+ T cells.; T-cell subset analysis was only performed during the first 2 weeks of alemtuzumab because of extensive T cell depletion at subsequent time points. **E.** Levels of HIV-DNA in total CD4+ T cells for samples obtained before diagnosis with Sezary syndrome, before starting

alemtuzumab and at several time points during alemtuzumab. Horizontal bars show mean values of 3 replicate measurements at each time point. Tcm, central memory T-cells; Ttm, transitional memory T-cells; Tem, effector memory T-cells, TD, terminally differentiated.